

Original Article

Comparison of Anti-inflammatory Activities of Six *Curcuma* Rhizomes: A Possible Curcuminoid-independent Pathway Mediated by *Curcuma phaeocaulis* Extract

Chihiro Tohda¹, Natsuki Nakayama¹, Fumiyuki Hatanaka¹ and Katsuko Komatsu^{2,3}

¹Division of Biofunctional Evaluation, Research Center for Ethnomedicine, Institute of Natural Medicine, University of Toyama, Toyama 930-0194, Japan, ²Division of Pharmacognosy, Institute of Natural Medicine, University of Toyama, Toyama 930-0194, Japan and ³21st Century COE Program, Institute of Natural Medicine, University of Toyama, Toyama 930-0194, Japan

We aimed to compare the anti-inflammatory activities of six species of *Curcuma* drugs using adjuvant arthritis model mice. When orally administered 1 day before the injection of adjuvant, the methanol extract of *Curcuma phaeocaulis* significantly inhibited paw swelling and the serum haptoglobin concentration in adjuvant arthritis mice. Also when orally administered 1 day after the injection of adjuvant, the methanol extract of *Curcuma phaeocaulis* significantly inhibited paw swelling. Other *Curcuma* species (*Curcuma longa*, *Curcuma wenyujin*, *Curcuma kwangsiensis*, *Curcuma zedoaria* and *Curcuma aromatica*) had no significant inhibitory effects on adjuvant-induced paw swelling. Cyclooxygenase (COX)-2 activity was significantly inhibited by the methanol extract of *C. phaeocaulis*. Curcuminoids' (curcumin, bis-demethoxycurcumin and demethoxycurcumin) were rich in *C. longa*, but less in *C. phaeocaulis* and *C. aromatica*, not in *C. wenyujin*, *C. kwangsiensis* and *C. zedoaria*, suggesting that curcuminoids' contents do not relate to inhibition of arthritis swelling. Therefore, *C. phaeocaulis* may be a useful drug among *Curcuma* species for acute inflammation, and the active constituents of *C. phaeocaulis* are not curcuminoids.

Keywords: adjuvant – arthritis – COX-2 – *Curcuma* – haptoglobin

Introduction

Many reports have suggested useful pharmacological properties of *Curcuma* drugs such as anti-inflammatory (1), anti-tumor (2) and immunological effects (3). Traditionally, *Curcuma* drugs called 'Ukon' and 'Gajutsu' in Japanese have been used in Oketsu syndromes (caused by the obstruction of blood circulation) in Chinese medicine (4). Since the pharmacological effects of curcuminoids, especially curcumin, have been investigated, such as radical scavenging (5), the inhibition of nitric oxide (NO) (6,7), anti-inflammation (8),

anti-tumor (9), anti-allergy (10) and anti-dementia (11), the usefulness of 'Ukon' derived from *Curcuma longa* has been intensively studied. Pharmacological studies of other *Curcuma* species were very few, because botanical origins of *Curcuma* drugs could not be easily identified due to similarity of morphology, and variety of naming derived from used parts and producing areas. At present, four *Curcuma* drugs are prescribed in Chinese Pharmacopoeia; Yujin (the tubers of *Curcuma wenyujin*, *C. longa*, *Curcuma kwangsiensis* or *Curcuma phaeocaulis*); Jianghuang (the rhizome of *C. longa*); Pian-Jianghuang (the rhizome of *C. wenyujin*) and Ezhu (the rhizomes of *C. phaeocaulis*, *C. kwangsiensis* or *C. wenyujin*). In addition, the rhizomes of *Curcuma zedoaria* and *Curcuma aromatica* have been used in Japan as a medicine or supplement. We previously compared five types of *Curcuma* drugs (rhizomes of *C. longa*, *C. kwangsiensis*, *C. phaeocaulis*,

For reprints and all correspondence: Dr Chihiro Tohda, Institute of Natural Medicine, Toyama Medical and Pharmaceutical University, 2630 Sugitani, Toyama 930-0194, Japan. Tel: +81-76-434-7646; Fax: +81-76-434-5068; e-mail: chihiro@inm.u-toyama.ac.jp

C. wenyujin and *C. zedoaria*), which were correctly identified by molecular biological analysis (12), on vasomotion in isolated rat aortas, and we found that the methanol extracts of all species had NO-independent relaxation effects (13). However, potencies of these vasomotion effects were not significantly different between species. In this study, we focused on evaluating differences among the anti-inflammatory activities of six *Curcuma* drugs (*C. longa*, *C. wenyujin*, *C. phaeocaulis*, *C. kwangsiensis*, *C. zedaria* and *C. aromatica*).

Materials and Methods

The six *Curcuma* drugs were used as shown in Table 1 and Fig. 1, which were correctly identified by the molecular biological method previously reported (12). All drugs were stored in the Museum of Materia Medica, Institute of Natural Medicine, University of Toyama (TMPW), Japan.

Preparation of Extracts

Methanol extracts were prepared as follows: 400 g of powdered drug was placed in methanol (1 l × 2) for 12 h at room temperature. The combined supernatants were evaporated on a water bath to obtain the methanol extracts. The extracts were dissolved in dimethyl sulfoxide (DMSO) (stock solution) and then suspended in olive oil in animal experiments. The extract was administered orally [500 mg kg⁻¹, 400 μl per mouse (olive oil : stock solution in DMSO = 320 μl : 80 μl)].

Induction of Arthritis

The mice were handled in accordance with the Guide for Animal Experiments, University of Toyama. Arthritis was induced in male ddY mice (6 week old; SLC, Shizuoka, Japan)

by injecting 50 μg per 50 μl⁻¹ of Complete Freund's adjuvant (CFA) (Sigma, St Louis, USA) into the right-hind footpad. The vehicle injection was 50 μl paraffin oil. In Fig. 2, immediately before and 1 day after injection of CFA, right footpad swelling




Figure 1. Morphologies of used *Curcuma* drugs.

Table 1. *Curcuma* drugs used in this study and the yields of extracts

Herbal drug name		Scientific name	Abbreviation	Part used	Cultivated area	Yield (%)	TMPW No. ^a
Japanese	Chinese						
Ukon	Jianghuang	<i>Curcuma longa</i> L.	CL	Rhizome	Guangdong, China	14.0	22 295 ^b
Henkyouou	Pian-Jianghuang	<i>C. wenyujin</i> Y.H. Chen and C. Ling	CW	Rhizome	Zhejiang, China	14.8	19 910 ^c
						12.1	22 292 ^b
						12.3	19 911 ^c
Gajutsu	Ezhu	<i>C. phaeocaulis</i> Val.	CP	Rhizome	Sichuan, China	6.5	22 297 ^b
						9.2	20 237 ^c
Gajutsu	Ezhu	<i>C. kwangsiensis</i> S.G. Lee and C.F. Liang	CK	Rhizome	Guangxi, China	2.0	22 471 ^b
						1.9	19 912 ^c
Gajutsu	—	<i>C. zedoaria</i> Rosc	CZ	Rhizome	Kagoshima, Okinawa, Japan	8.9	22 473 ^b
						7.0	20 285 ^c
Haruukon	—	<i>C. aromatica</i>	CA	Rhizoma	Okinawa, Japan	7.5	22 312 ^b
						8.9	20 284 ^c

^aThe number shows the registration number of the Museum of Materia Medica, Institute of Natural Medicine, University of Toyama (TMPW). These drugs were identified by the molecular biological method previously reported (13).

^bUsed in Figs 2–4.

^cUsed in Table 2.




Figure 2. Effects of pretreatment with methanol extracts of *Curcuma* drugs on paw swelling and increase in serum haptoglobin induced by adjuvant injection. Extracts of *Curcuma* drugs (500 mg kg^{-1} , hatched columns), dexamethasone (5 mg kg^{-1}) or were administered once 1 day before the adjuvant injection. Arthritis was induced by the injection of Complete Freund's adjuvant (CFA) into the right-hind footpad. Paraffin oil was injected in control mice. (A) Immediately before, and 1 day after injection of the adjuvant, swelling of the right footpad was measured with slide calipers. Change rate of the swelling of Day 2 compared with Day 1 was calculated. (B) After footpad measurement, serum was collected. The concentration of haptoglobin in serum was measured. The time schedule of treatments is shown in the bottom. The values represent the means and SEM of 5 mice. * $P < 0.05$ when compared with Veh.

(length \times wide) was measured with slide calipers. In Fig. 3, immediately before, 1 day after and 2 day after the injection of CFA, right footpad swelling (length \times wide) was measured with slide calipers. Extracts were administered once 1 day after the adjuvant injection. The change rate of paw swelling was




Figure 3. Post-treatment with methanol extracts of *Curcuma* drugs and its effect on paw swelling and increase in serum haptoglobin induced by adjuvant injection. Extracts of *Curcuma* drugs (500 mg kg^{-1}), dexamethasone (5 mg kg^{-1}) or 20% DMSO in olive oil (Veh) were administered once 1 day after the adjuvant injection. Arthritis was induced by the injection of Complete Freund's adjuvant (CFA) into the right-hind footpad. Paraffin oil was injected in control mice. (A) Immediately before, 1 day after and 2 days after injection of the adjuvant, swelling of the right footpad was measured with slide calipers. Change rates of the swelling of Day 1 (open columns) and Day 2 (hatched columns) compared with Day 0 were calculated. (B) After footpad measurement, serum was collected at Day 2. The concentration of haptoglobin in serum was measured. The time schedule of treatments is shown in the bottom. The values represent the means and SEM of five mice. * $P < 0.05$ when compared between before and after extract treatment in (A); * $P < 0.05$ when compared with Veh in (B).

calculated. The paw sizes did not change before and after extract administration.

Measurement of Haptoglobin

After measuring the footpads, blood (1 ml) was collected from the vena cava and added to a serum collection tube. After centrifugation (3500 r.p.m., 20 min), the supernatant was transferred into a new tube as a serum sample (400 μl). The concentration of haptoglobin in serum was measured using PHASE RANGE (Tridelta Development Limited, Maynooth, Ireland).

Measurement of COX-1 and COX-2

Cyclooxygenase (COX) activity was measured using a Colorimetric COX inhibitor assay kit (Cayman, Ann Arbor, USA).

The methanol extracts or indomethacin were dissolved in DMSO and added to the enzyme reaction mixture.

Quantifying Curcuminoid Content of *Curcuma* Drugs

Standard curcumin, bis-demethoxycurcumin and demethoxycurcumin were isolated and identified as pure compounds by spectra data in our laboratory (13). Each standard curcuminoid (1 mg) was accurately weighed and dissolved in methanol at a concentration of 0.2 mg ml⁻¹. To draw calibration curves, a series of standard solutions were prepared from the stock solution and filtered through a 0.2 µm Millipore filter (Advantec, Tokyo, Japan). Dried *Curcuma* drugs were pulverized and powdered. Methanol extracts of 20 mg were accurately weighed and dissolved in methanol at a concentration of 0.4 mg ml⁻¹. After ultrasonication for 30 min, supernatants were obtained by centrifugation at 2500 r.p.m. for 10 min. The supernatants were transferred into volumetric flasks, and methanol was added to obtain a final volume of 100 ml. After filtration through a 0.2 µm Millipore filter (Advantec), 5 µl was injected into the HPLC system for analysis. The JASCO HPLC system (Jasco, Tokyo, Japan) is composed of a PU-1580 intelligent HPLC pump, a DG-1580-53 3-line degasser, a LG-1580-02 ternary gradient unit, a CO-1565 intelligent column oven, an AS-2057 plus intelligent sampler and an MD-1510 diode array detector. Comparative analysis was carried out using a Mightysil RP-18GP (15 mm, 250 mm × 4.6 mm i.d.) with a column temperature of 40°C. The mobile phase was acetonitrile:water:acetic acid = 45 : 55 : 1. The flow rate was 1.0 ml min⁻¹ and the detection wavelength was 410 nm. The chromatographic data were collected and processed using BORWIN-PDA APPLICATION and BORWIN CHROMATOGRAPHY Software (version 1.5, Jasco). The values of retention time of curcumin, bis-demethoxycurcumin and demethoxycurcumin were 15.44, 11.75 and 13.48 min, respectively. The standard curves of curcuminoids were made using a dose of 2–100 µg ml⁻¹.

Statistical Analysis

Statistical comparisons were carried out using one-way analysis of variance followed by Dunnett's *post hoc* test or paired *t*-test. Values of $P < 0.05$ were considered significant. The means of the data are presented together with the SEM.

Results

We previously investigated the time-course of the paw edema and confirmed that swelling peaked at 24 h post-injection under our experimental conditions. In Fig. 2, methanol extracts (500 mg kg⁻¹) were orally administered 1 day before the adjuvant injection. Treatment with the *C. phaeocaulis* (CP) extract significantly reduced the paw edema (Fig. 2A). Although treatments with extracts of *C. longa* (CL), *C. kwangsiensis* (CK) and *C. zedoaria* (CZ) tended to reduce slightly the paw swelling, treatment with *C. wenyujin* (CW) and *C. aromatica* (CA) did not. A steroidal anti-

inflammatory drug, dexamethasone, was used as a reference. At a dose of 5 mg kg⁻¹, dexamethasone significantly inhibited paw swelling. Although indomethacin (5 and 10 mg kg⁻¹), a non-steroidal anti-inflammatory drug, was also used in this experiment, several mice died after severe body weight loss. Extract-treated groups showed no adverse effects. After measuring the paw swelling, serum haptoglobin was detected as an acute inflammation marker (14). The concentration of serum haptoglobin intensively increased (10-fold the control) by adjuvant injection (Fig. 2B). Serum haptoglobin was significantly reduced in a CP extract-treated group (inhibition rate: 72.5%) and CK extract-treated one (inhibition rate: 59.9%); however, no significant reduction was seen in other *Curcuma* drug extract-treated groups. Dexamethasone (5 mg kg⁻¹) did not reduce serum haptoglobin.

Paw Edema Significantly Reduced by *C. phaeocaulis*

In Fig. 3, methanol extracts (500 mg kg⁻¹) were orally administered 1 day after the adjuvant injection. Treatment with the *C. phaeocaulis* (CP) extract significantly reduced paw edema (Fig. 3A). Although treatments with extracts of CL, CW, CK, CZ and CA tended to reduce the paw swelling, these were not significant. Dexamethasone (5 mg kg⁻¹) significantly inhibited paw swelling. Serum haptoglobin was slightly reduced in CP and CK extracts-treated groups, but not significantly. Dexamethasone (5 mg kg⁻¹) also did not reduce serum haptoglobin significantly either.

Curcuma Extracts Showed Inhibitory Activity on Enzymatic Activities *In Vitro*

The inhibitory effects of *Curcuma* extracts on *in vitro* enzymatic activities were measured against COX-2 and COX-1 (Fig. 4). The methanol extract of CP (500 µg/ml) significantly inhibited COX-2 activity (inhibition rate: 24.4%). However, CL, CW, CK, CZ and CA extracts demonstrated no significant inhibition. For COX-1, CP, CL and CZ extracts showed inhibitory activity, and the inhibition rate with the CP extract was the most remarkable (inhibition rate: 38.4%). Indomethacin, a COX inhibitor, inhibited COX-2 (inhibition rate: 45.5% inhibition at 100 µM) and COX-1 (inhibition rate: 37.2% at 100 µM) activities dose dependently. The inhibitory efficacy of COX-2 with the methanol extract of CP was weaker than that of 100 µM indomethacin.

Testing of *Curcuma* Samples from China and Japan Confirmed That Curcumin was Rich in CL

The content of three curcuminoids in the methanol extracts of *Curcuma* drugs was quantified as shown in (Table 2). In CP and CA, a very small amount of curcumin was contained in the methanol extract, in addition to bis-demethoxycurcumin and demethoxycurcumin were hardly detected. None of the three curcuminoids was present in CW, CK and CZ, whereas large amounts of curcuminoids were detected in CL. In particular, curcumin concentrations were high in the CL




Figure 4. Inhibitory activities of methanol extracts of *Curcuma* drugs on COX-2 and COX-1 activities. The methanol extracts of *Curcuma* drugs ($500 \mu\text{g ml}^{-1}$, hatched columns), indomethacin (10 and 100 μM , closed columns), or the vehicle (0.1% DMSO, open columns) were added to the enzyme reaction mixture. COX-2 (A) and COX-1 (B) activities were measured using the Colorimetric COX inhibitor assay kit. The values represented the means and SEM. $n = 3$. * $P < 0.05$ when compared with Veh.

Table 2. Curcuminoid contents in the methanol extracts

	Contents (% in the methanol extract)		
	Curcumin	Bisdemethoxy-curcumin	Demethoxy-curcumin
<i>C. longa</i>	12.30	4.86	3.62
<i>C. wenyujin</i>	ND	ND	ND
<i>C. phaeocaulis</i>	0.89	ND	t
<i>C. kwangsiensis</i>	ND	ND	ND
<i>C. zedoaria</i>	ND	ND	ND
<i>C. aromatica</i>	0.11	ND	ND

ND, Not detected; t, trace.

methanol extract. We tested many *Curcuma* samples from China and Japan, and confirmed that curcumin was rich in CL (3.9–12.3% in the methanol extract), but not in CP, CW, CK, CZ and CA.

Discussion

The methanol extract of CP significantly reduced paw swelling (Fig. 2A) and the expression of an inflammation marker, hap-

toglobin in serum (Fig. 2B), of mice when it was administered orally 1 day before the adjuvant injection. Also in case of treatment with the methanol extracts 1 day after the adjuvant injection when the paw was maximally swollen, CP significantly reduced paw swelling (Fig. 3A). However, treatment with the methanol extracts of CL, CW, CZ and CA had no clear effects on inflammation. Therefore, CP is to be expected the most effective in reducing arthritis swelling among *Curcuma* drugs.

The methanol extract of CP had inhibitory activity on inflammation-related enzymes, COX-2 (Fig. 4A). A selective COX-2 inhibitor, SC-58125, rapidly reverses paw edema, the level of PGE₂, the expression of COX-2, and serum IL-6 and paw IL-6 levels in arthritis rats induced by CFA (15). Upregulated COX-2 also enhances PG production, and inflammation progresses further (15). Therefore, the inhibitory effect of CP on COX-2 activity may be useful to prevent inflammation. Both CP and indomethacin were not selective inhibitors of COX-2 (Fig. 4). Some indomethacin-treated mice died (data not shown). It is known that indomethacin treatment has a high risk of several adverse effects (gastrointestinal, hepatic and kidney disorders), which may be caused mainly by the inhibitory effect of COX-1. Considering that CP had anti-inflammatory effects on arthritic mice (Figs 1 and 2) with no adverse effects in spite of its COX-1 inhibitory action; some constituents which relieve the adverse effects may be contained in the methanol extract of CP. Dexamethasone did not inhibit the serum haptoglobin concentration (Figs 1 and 2). Each class of anti-inflammatory drugs has a specific effect on the regulation of acute phase proteins (16), and dexamethasone is known not to reduce the haptoglobin expression (17).

In an arthritis animal model, IL-1 β (18) and TNF- α (15) are increased by inflammatory stimulation, and IL-1 β induces COX-2 transcription in inflammatory cells (19). At the same time, increased IL-1 β and TNF- α in inflammation upregulate the IL-6 expression (20), and IL-6 stimulation in the liver induces the translocation of activated STAT3 molecules to nuclei (21,22). Transcription of the haptoglobin gene is enhanced through multiple IL-6 response elements, and haptoglobin increases in the liver and serum (23). Since the methanol extract of CP reduced serum haptoglobin in arthritis, it may inhibit anywhere in the cytokine pathway. We will investigate whether the expression levels of these cytokines in serum, and the COX-2 level in paw tissue, are changed by treatment with CP extract.

The pharmacological activities of curcumin have been studied intensively from many viewpoints, suggesting that curcumin inhibits LPS-induced NO production (5) and iNOS expression (6), COX-2 expression and activation (24), and expression of several inflammatory markers. In addition, curcumin is cytotoxic in several cell types (25,26), not only in cancer cells (27,28). Hepatotoxicity was reported in animal experiments using curcumin-containing *Curcuma* drugs (29,30), and although curcumin has many useful pharmacological and possible therapeutic activities, its safety should be considered carefully. However, CP contains very low amounts of curcuminoids (Table 2), but showed anti-inflammatory

activity (Figs 1 and 2), suggesting that some active constituents other than curcuminoids may exist in CP. Since a recent study showed that significant amounts of furanodienone and curcumenol were contained in CP (31), these compounds may be candidates for active principles. As very little basic pharmacological research or chemical analyses of CP have been performed, the usefulness of CP should be investigated in future studies.

Acknowledgments

We particularly thank Mrs K. Hayashi for her technical support. This work was partially supported by Tamura Foundation, and Grant-in-Aid for Scientific Research (B), No. 14406030 in 2002–04 and No. 17406004 in 2005–07, from Japan Society for the Promotion of Science, and by the 21st Century COE Program from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- Ahmed S, Anuntiyo J, Charles J, Malemud CJ, Haqqi TM. Biological basis for the use of botanicals in osteoarthritis and rheumatoid arthritis. *Evid Based Complement Altern Med* 2005;2:301–8.
- Ozaki Y. Antiinflammatory effect of *Curcuma xanthorrhiza* Roxb., and its active principles. *Chem Pharm Bull* 1990;38:1045–8.
- Gonda R, Tomoda M, Ohara N, Takada K. Arabinogalactan core structure and immunological activities of ukonan C, an acidic polysaccharide from the rhizome of *Curcuma longa*. *Biol Pharm Bull* 1993;16:235–8.
- Li SZ. 'Ben-cao gang-mu'. In: Jiangxi (ed.). Vol. 2, The People's Health Pub., House, Beijing, 1977, 880–5.
- Sreejayan N, Rao MN. Free radical scavenging activity of curcuminoids. *Arzneimittelforschung* 1996;46:169–71.
- Pan MH, Lin-Shiau SY, Lin JK. Comparative studies on the suppression of nitric oxide synthase by curcumin and its hydrogenated metabolites through down-regulation of I κ B kinase and NF κ B activation in macrophages. *Biochem Pharmacol* 2000;60:1665–76.
- Onoda M, Inano H. Effect of curcumin on the production of nitric oxide by cultured rat mammary gland. *Nitric Oxide* 2000;4:505–15.
- Banerjee M, Tripathi LM, Srivastava VM, Puri A, Shukla R. Modulation of inflammatory mediators by ibuprofen and curcumin treatment during chronic inflammation in rat. *Immunopharmacol Immunotoxicol* 2003;25:213–24.
- Khar A, Ali AM, Pardhasaradhi BV, Begum Z, Anjum R. Antitumor activity of curcumin is mediated through the induction of apoptosis in AK-5 tumor cells. *FEBS Lett* 1999;19:165–8.
- Ram A, Das M, Ghosh B. Curcumin attenuates allergen-induced airway hyperresponsiveness in sensitized guinea pigs. *Biol Pharm Bull* 2003;26:1021–4.
- Lim GP, Chu T, Yang F, Beech W, Frautschy SA, Cole GM. The curry spice curcumin reduces oxidative damage and amyloid pathology in an Alzheimer transgenic mouse. *J Neurosci* 2001;21:8370–7.
- Sasaki Y, Fushimi H, Cao H, Cai SQ, Komatsu K. Sequence analysis of Chinese and Japanese *Curcuma* drugs on the 18S rRNA gene and *tmK* gene and the application of amplification-refractory mutation system analysis for their authentication. *Biol Pharm Bull* 2002;25:1593–9.
- Sasaki Y, Goto H, Tohda C, Hatanaka F, Shibahara N, Shimada Y, et al. Effects of *Curcuma* drugs on vasomotion in isolated rat aorta. *Biol Pharm Bull* 2003;26:1135–43.
- Giffen PS, Turton J, Andrews CM, Barrett P, Clarke CJ, Fung KW, et al. Markers of experimental acute inflammation in the Wistar Han rat with particular reference to haptoglobin and C-reactive protein. *Arch Toxicol* 2003;77:392–402.
- Anderson GD, Hauser SD, McGarity KL, Bremer ME, Isakson PC, Gregory SA. Selective inhibition of cyclooxygenase (COX)-2 reverses inflammation and expression of COX-2 and interleukin 6 in rat adjuvant arthritis. *J Clin Invest* 1996;97:2672–9.
- do Nascimento CO, Hunter L, Trayhurn P. Regulation of haptoglobin gene expression in 3T3-L1 adipocytes by cytokines, catecholamines, and PPAR γ . *Biochem Biophys Res Commun* 2004;313:702–8.
- Lewis EJ, Bishop J, Cashin CH. Automated quantification of rat plasma acute phase reactants in experimental inflammation. *J Pharmacol Methods* 1989;21:183–94.
- Silva JC, Rocha MF, Lima AA, Brito GA, de Menezes DB, Rao VS. Effects of pentoxifylline and nabumetone on the serum levels of IL-1 β and TNF α in rats with adjuvant arthritis. *Inflamm Res* 2000;49:14–19.
- Crofford LJ, Tan B, McCarthy CJ, Hla T. Involvement of nuclear factor kappa B in the regulation of cyclooxygenase-2 expression by interleukin-1 in rheumatoid synoviocytes. *Arthritis Rheum* 1997;40:226–36.
- Guerne PA, Zuraw BL, Vaughan JH, Carson DA, Lotz M. Synovium as a source of interleukin 6 *in vitro*. Contribution to local and systemic manifestations of arthritis. *J Clin Invest* 1989;83:585–92.
- Naka T, Nishimoto N, Kishimoto T. The paradigm of IL-6: from basic science to medicine. *Arthritis Res* 2002;4:S233–42.
- Heinrich PC, Behrmann I, Haan S, Hermanns HM, Muller-Newen G, Schaper F. Principles of interleukin (IL)-6-type cytokine signalling and its regulation. *Biochem J* 2003;347:1–20.
- Wang Y, Kinzie E, Berger FG, Lim SK, Baumann H. Haptoglobin, an inflammation-inducible plasma protein. *Redox Rep* 2001;6:379–85.
- Hong J, Bose M, Ju J, Ryu JH, Chen X, Sang S, et al. Modulation of arachidonic acid metabolism by curcumin and related β -diketone derivatives: effects on cytosolic phospholipase A(2), cyclooxygenases and 5-lipoxygenase. *Carcinogenesis* 2004;25:1671–9.
- Donatus IA, Sardjoko, Vermeulen NP. Cytotoxic and cytoprotective activities of curcumin. Effects on paracetamol-induced cytotoxicity, lipid peroxidation and glutathione depletion in rat hepatocytes. *Biochem Pharmacol* 1990;39:1869–75.
- Matsuda H, Ninomiya K, Morikawa T, Yoshikawa M. Inhibitory effect and action mechanism of sesquiterpenes from *Zedoariae Rhizoma* on D-galactosamine/lipopolysaccharide-induced liver injury. *Bioorg Med Chem Lett* 1998;8:339–44.
- Woo JH, Kim YH, Choi YJ, Kim DG, Lee KS, Bae JH, et al. Molecular mechanisms of curcumin-induced cytotoxicity: induction of apoptosis through generation of reactive oxygen species, down-regulation of Bcl-XL and IAP, the release of cytochrome c and inhibition of Akt. *Carcinogenesis* 2003;24:1199–208.
- Dorai T, Cao YC, Dorai B, Buttyan R, Katz AE. Therapeutic potential of curcumin in human prostate cancer. III. Curcumin inhibits proliferation, induces apoptosis, and inhibits angiogenesis of LNCaP prostate cancer cells *in vivo*. *Prostate* 2001;47:293–303.
- Deshpande SS, Lalitha VS, Ingle AD, Raste AS, Gadre SG, Maru GB. Subchronic oral toxicity of turmeric and ethanolic turmeric extract in female mice and rats. *Toxicol Lett* 1998;95:183–93.
- Kandarkar SV, Sawant SS, Ingle AD, Deshpande SS, Maru GB. Subchronic oral hepatotoxicity of turmeric in mice-histopathological and ultrastructural studies. *Indian J Exp Biol* 1998;36:675–9.
- Yang EQ, Li SP, Chen Y, Lao SC, Wang YT, Dong TTX, Tsim KWK. Identification and quantitation of eleven sesquiterpenes in three species of *Curcuma* rhizomes by pressurized liquid extraction and gas chromatography-mass spectrometry. *J Pharm Biomed Anal* 2005;39:552–8.

Received December 28, 2005; accepted February 10, 2006



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

