



University of  
Zurich<sup>UZH</sup>

Zurich Open Repository and  
Archive

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2022

---

## Complete Genome Sequence of *Hafnia paralvei* Isolate AVS0177, Harboring *mcr* – 9 on a Plasmid

Biggel, Michael ; Zurfluh, Katrin ; Hoehn, Sarah ; Schmitt, Kira ; Frei, Andrea ; Jans, Christoph ; Stephan, Roger

Abstract: Here, we report the complete genome sequence of a *Hafnia paralvei* strain isolated from a lake in Switzerland in 2020. The genome consists of a 4.7-Mbp chromosome, a large plasmid (213 kb) harboring *mcr*-9, and a small plasmid (6 kb).

DOI: <https://doi.org/10.1128/mra.00966-21>

Other titles: Complete Genome Sequence of *Hafnia paralvei* Isolate AVS0177, Harboring *mcr*-9 on a Plasmid

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-231172>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Biggel, Michael; Zurfluh, Katrin; Hoehn, Sarah; Schmitt, Kira; Frei, Andrea; Jans, Christoph; Stephan, Roger (2022). Complete Genome Sequence of *Hafnia paralvei* Isolate AVS0177, Harboring *mcr* – 9 on a Plasmid. *Microbiology Resource Announcements*, 11(1):e0096621.

DOI: <https://doi.org/10.1128/mra.00966-21>



# Complete Genome Sequence of *Hafnia paralvei* Isolate AVS0177, Harboring *mcr-9* on a Plasmid

Michael Biggel,<sup>a</sup> Katrin Zurfluh,<sup>a</sup> Sarah Hoehn,<sup>b</sup> Kira Schmitt,<sup>a</sup> Andrea Frei,<sup>b</sup> Christoph Jans,<sup>b</sup> Roger Stephan<sup>a</sup>

<sup>a</sup>Institute for Food Safety and Hygiene, Vetsuisse Faculty University of Zürich, Zürich, Switzerland

<sup>b</sup>Amt für Verbraucherschutz, Kanton Zug, Steinhausen, Switzerland

**ABSTRACT** Here, we report the complete genome sequence of a *Hafnia paralvei* strain isolated from a lake in Switzerland in 2020. The genome consists of a 4.7-Mbp chromosome, a large plasmid (213 kb) harboring *mcr-9*, and a small plasmid (6 kb).

Since the initial recognition of *Enterobacteriaceae* strains harboring the plasmid-mediated colistin resistance gene *mcr-1* in 2015 (1), nine additional *mcr* variants (*mcr-2* to *mcr-10*) have been described in various species (2–4). Here, we describe the occurrence of *mcr-9* in a *Hafnia* isolate, an enterobacterial genus naturally resistant to colistin (5).

AVS0177, showing phenotypical resistance to colistin (MIC = 8 mg/mL) and a positive PCR result for *mcr-9*, was isolated in November 2020 from a water sample collected at a depth of 120 m in Lake Zug, Switzerland. The water sample (100 mL) was filtered through a 0.45- $\mu$ m membrane filter (Millipore). The filter was incubated in 10 mL *Enterobacteriaceae* enrichment (EE) broth (BD) at 37°C for 24 h. One loopful of the enrichment broth was spread onto cystine lactose electrolyte-deficient (CLED) agar (Oxoid) supplemented with 4 mg/mL colistin, 10 mg/mL vancomycin, and 5 mg/mL amphotericin and incubated at 37°C for 24 h. Matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI-TOF MS) (Bruker Daltonics) was used for preliminary genus identification. Different PCR primers were used to detect the genes *mcr-1* to *mcr-10* (6–8). Colistin susceptibility testing was performed by broth dilution and interpreted according to EUCAST v11.0 protocols ([https://www.eucast.org/clinical\\_breakpoints/](https://www.eucast.org/clinical_breakpoints/)). DNA was isolated from a subculture obtained from a single colony grown for 24 h at 37°C on sheep blood agar. For short-read sequencing, DNA was extracted using the DNeasy blood and tissue kit (Qiagen). Libraries were prepared using the Nextera DNA Flex library preparation kit (Illumina) and sequenced on the Illumina MiniSeq platform (2 × 150 bp). For long-read sequencing, DNA was obtained using the MasterPure Complete DNA and RNA purification kit (Lucigen) (no size selection/shearing). Multiplex libraries were prepared using the SQK-LSK109 kit with the EXP-NBD114 barcodes and sequenced using a MinION device on a FLO-MIN106 flow cell (Oxford Nanopore Technologies). Base calling was performed using Guppy CPU v4.2.2+effbaf8. Illumina and ONT adapters were trimmed using Trim Galore v0.6.6 (9) and Porechop v0.2.4 (10) and quality assessed using FastQC v0.11.9 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) and LongQC v1.2.0 (11), respectively. A hybrid assembly was generated from 290 Mbp long-read (16,912 reads; read  $N_{50}$ , 45 kb; coverage, 59×) and 222 Mbp short-read (754,150 paired-end reads; coverage, 45×) data using the Unicycler v0.4.8 pipeline (12), which includes assembly polishing, circularization, and rotation. The assembly was annotated using Prokaryotic Genome Annotation Pipeline (PGAP) (13). Resistance genes and plasmid replicons were identified using ABRicate v1.0.1 (14) (coverage/identity, >70%/>90%) with the ResFinder (15) and PlasmidFinder (16) databases, respectively. Default parameters were used for all software unless otherwise specified.

The complete genome of AVS0177 consisted of a circular 4,727,161-bp chromosome (rotated to start at *dnaA*) and the circular plasmids pAVS0177-a (212,640 bp) and

**Editor** Catherine Putonti, Loyola University Chicago

**Copyright** © 2022 Biggel et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Roger Stephan, [roger.stephan@uzh.ch](mailto:roger.stephan@uzh.ch).

The authors declare no conflict of interest.

**Received** 28 September 2021

**Accepted** 5 December 2021

**Published** 6 January 2022

pAVS0177-b (5,733 bp), with a GC content of 48.07%. rMLST analysis (17) assigned AVS0177 to the species *Hafnia paralvei*. Two antimicrobial resistance genes were detected: the chromosomally encoded *bla*<sub>ACC-1a</sub> (AmpC beta-lactamase) and *mcr-9*, located on plasmid pAVS0177-a. pAVS0177-a also harbored the *fecABCDE* gene cluster, encoding a ferric citrate transport system (18), and the *sil* gene cluster, encoding copper and silver resistance (19, 20). The incompatibility (Inc) type of pAVS0177-a could not be identified, and no similar plasmids (i.e., query coverage of >50%) were found in a BLASTn (MegaBLAST) search of the NCBI nucleotide collection. The 12.5-kb genetic region encompassing *mcr-9* (spanning from the transcriptional repressor *mcrR* to the insertion sequence *IS15DII* and including *IS903B* directly upstream of *mcr-9*) was structurally identical to the genetic context of *mcr-9* described in other species (21, 22). In conclusion, we present the complete genome sequence of an *mcr*-harboring *Hafnia* isolate and demonstrate that intrinsically resistant species may serve as reservoirs for transferable *mcr* genes.

**Data availability.** The complete genome sequence of AVS0177 has been deposited in GenBank under the accession numbers CP083737.1 (chromosome), CP083738.1 (pAVS0177-a), and CP083739.1 (pAVS0177-b) and the BioProject accession number PRJNA761908. The raw data were deposited in the NCBI Sequence Read Archive (SRA) under the accession numbers SRR15833986 (ONT reads) and SRR15833987 (Illumina reads).

## ACKNOWLEDGMENT

We thank Nicole Cernela for her technical support with Illumina sequencing.

## REFERENCES

- Liu Y-Y, Wang Y, Walsh TR, Yi L-X, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu L-F, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu J-H, Shen J. 2016. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* 16:161–168. [https://doi.org/10.1016/S1473-3099\(15\)00424-7](https://doi.org/10.1016/S1473-3099(15)00424-7).
- Gogry FA, Siddiqui MT, Sultan I, Haq QMR. 2021. Current update on intrinsic and acquired colistin resistance mechanisms in bacteria. *Front Med (Lausanne)* 8:677720. <https://doi.org/10.3389/fmed.2021.677720>.
- Ling Z, Yin W, Shen Z, Wang Y, Shen J, Walsh TR. 2020. Epidemiology of mobile colistin resistance genes *mcr-1* to *mcr-9*. *J Antimicrob Chemother* 75:3087–3095. <https://doi.org/10.1093/jac/dkaa205>.
- Wang C, Feng Y, Liu L, Wei L, Kang M, Zong Z. 2020. Identification of novel mobile colistin resistance gene *mcr-10*. *Emerg Microbes Infect* 9:508–516. <https://doi.org/10.1080/22221751.2020.1732231>.
- Jayol A, Saly M, Nordmann P, Ménard A, Poirel L, Dubois V. 2017. *Hafnia*, an enterobacterial genus naturally resistant to colistin revealed by three susceptibility testing methods. *J Antimicrob Chemother* 72:2507–2511. <https://doi.org/10.1093/jac/dkx154>.
- Borowiak M, Baumann B, Fischer J, Thomas K, Deneke C, Hammerl JA, Szabo I, Malorny B. 2020. Development of a novel *mcr-6* to *mcr-9* multiplex PCR and assessment of *mcr-1* to *mcr-9* occurrence in colistin-resistant *Salmonella enterica* isolates from environment, feed, animals and food (2011–2018) in Germany. *Front Microbiol* 11:80. <https://doi.org/10.3389/fmicb.2020.00080>.
- Rebelo AR, Bortolaia V, Kjeldgaard JS, Pedersen SK, Leekitcharoenphon P, Hansen IM, Guerra B, Malorny B, Borowiak M, Hammerl JA, Battisti A, Franco A, Alba P, Perrin-Guyomard A, Granier SA, De Frutos Escobar C, Malhotra-Kumar S, Villa L, Carattoli A, Hendriksen RS. 2018. Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Euro Surveill* 23:17-00672. <https://doi.org/10.2807/1560-7917.ES.2018.23.6.17-00672>.
- Xu T, Zhang C, Ji Y, Song J, Liu Y, Guo Y, Zhou K. 2021. Identification of *mcr-10* carried by self-transmissible plasmids and chromosome in *Enterobacter roggenkampii* strains isolated from hospital sewage water. *Environ Pollut* 268:115706. <https://doi.org/10.1016/j.envpol.2020.115706>.
- Krueger F, Trim Galore. <https://github.com/FelixKrueger/TrimGalore>.
- Wick RR. Porechop. <https://github.com/rrwick/Porechop>.
- Fukasawa Y, Ermini L, Wang H, Carty K, Cheung M-S. 2020. LongQC: a quality control tool for third generation sequencing long read data. *G3 (Bethesda)* 10:1193–1196. <https://doi.org/10.1534/g3.119.400864>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Tatusova T, DiCuccio M, Badretdin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Seemann T. Abricate. <https://github.com/tseemann/abricate>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
- Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
- Jolley KA, Bray JE, Maiden MCJ. 2018. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res* 3:124. <https://doi.org/10.12688/wellcomeopenres.14826.1>.
- Staudenmaier H, Van Hove B, Yaraghi Z, Braun V. 1989. Nucleotide sequences of the *fecBCDE* genes and locations of the proteins suggest a periplasmic-binding-protein-dependent transport mechanism for iron(III) dicitrate in *Escherichia coli*. *J Bacteriol* 171:2626–2633. <https://doi.org/10.1128/jb.171.5.2626-2633.1989>.
- Staehlin BM, Gibbons JG, Rokas A, O'Halloran TV, Slot JC. 2016. Evolution of a heavy metal homeostasis/resistance island reflects increasing copper stress in Enterobacteria. *Genome Biol Evol* 8:evw031. <https://doi.org/10.1093/gbe/evw031>.
- Gupta A, Matsui K, Lo J-F, Silver S. 1999. Molecular basis for resistance to silver cations in *Salmonella*. *Nat Med* 5:183–188. <https://doi.org/10.1038/5545>.
- Kieffer N, Royer G, Decousser J-W, Bourrel A-S, Palmieri M, Ortiz De La Rosa J-M, Jacquier H, Denamur E, Nordmann P, Poirel L. 2019. *mcr-9*, an inducible gene encoding an acquired phosphoethanolamine transferase in *Escherichia coli*, and its origin. *Antimicrob Agents Chemother* 63:e00965-19. <https://doi.org/10.1128/AAC.00965-19>.
- Ribeiro TG, Izdebski R, Urbanowicz P, Carmeli Y, Gniadkowski M, Peixe L. 2021. *Citrobacter telavivum* sp. nov. with chromosomal *mcr-9* from hospitalized patients. *Eur J Clin Microbiol Infect Dis* 40:123–131. <https://doi.org/10.1007/s10096-020-04003-6>.