

# Complete renormalization of QCD at five loops

---

Thomas Luthe,<sup>a</sup> Andreas Maier,<sup>b</sup> Peter Marquard<sup>c</sup> and York Schröder<sup>d</sup>

<sup>a</sup>*Faculty of Physics, University of Bielefeld,  
33501 Bielefeld, Germany*

<sup>b</sup>*Institute for Particle Physics Phenomenology, Durham University,  
Durham, United Kingdom*

<sup>c</sup>*Deutsches Elektronen Synchrotron (DESY),  
Platanenallee 6, Zeuthen, Germany*

<sup>d</sup>*Grupo de Física de Altas Energías, Universidad del Bío-Bío,  
Casilla 447, Chillán, Chile*

*E-mail:* [tluthe@physik.uni-bielefeld.de](mailto:tluthe@physik.uni-bielefeld.de), [andreas.maier@durham.ac.uk](mailto:andreas.maier@durham.ac.uk),  
[peter.marquard@desy.de](mailto:peter.marquard@desy.de), [yschroder@ubiobio.cl](mailto:yschroder@ubiobio.cl)

**ABSTRACT:** We present new analytical five-loop Feynman-gauge results for the anomalous dimensions of ghost field and -vertex, generalizing the known values for  $SU(3)$  to a general gauge group. Together with previously published results on the quark mass and -field anomalous dimensions and the Beta function, this completes the 5-loop renormalization program of gauge theories in that gauge.

**KEYWORDS:** Perturbative QCD, Renormalization Group

**ARXIV EPRINT:** [1701.07068](https://arxiv.org/abs/1701.07068)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Setup</b>	<b>2</b>
2.1	Renormalization constants	2
2.2	Computational strategy	3
2.3	Notation for group invariants	4
<b>3</b>	<b>Renormalization of ghost field and <math>\gamma</math>-vertex</b>	<b>5</b>
3.1	Ghost field anomalous dimension	5
3.2	Ghost-gluon vertex anomalous dimension	7
<b>4</b>	<b>Complete set of renormalization constants</b>	<b>8</b>
<b>5</b>	<b>Conclusions</b>	<b>9</b>
<b>A</b>	<b>Beta function</b>	<b>10</b>

---

## 1 Introduction

Non-Abelian quantum field theories have become a cornerstone of our description of nature, ever since it has been understood that they allow for the interaction strength to become weaker as the energy rises, a property called ‘asymptotic freedom’ [1–3]. Providing for an explanation of quark confinement and hence a successful theory of the strong interactions that bind together quarks into protons, Quantum Chromodynamics (QCD), much theoretical work has been focused on this central part of the Standard Model. The change (or ‘running’) of its couplings and masses with energy is governed by renormalization group equations, which in turn need the theory’s anomalous dimensions as an input. In the light of ever-increasing precision of collider experiments conducted at various energies, a first-principles determination of these parameters with the highest achievable theoretical accuracy has become necessary.

One of the hallmarks of perturbative renormalization is the so-called Beta function, governing the scale dependence of the renormalized strong coupling constant. For scalar  $\phi^4$  theory, for example, renormalization has recently been pushed to the 6-loop level [4, 5]. For the case of QCD, the current precision frontier is at five loops [6], and most recently, results for a generalization from the gauge group SU(3) to general simple Lie groups have appeared [7, 8]. The Beta function represents one of the above-mentioned anomalous dimensions that are contained in the renormalization group equations. In total there are five linearly independent ones, such as the quark mass anomalous dimension (for 5-loop results, see [9, 10]) that is of key phenomenological importance when evolving the masses typically measured at a few GeV to the electroweak scale, as well as three further non-physical coefficients that we will discuss in the following section.

In the present paper, our aim is to push the knowledge of the remaining renormalization constants to the same level. We follow up on our earlier works [7, 10] and complete the renormalization program by evaluating the anomalous dimension of the ghost field, and that of the ghost-gluon vertex. Using the same general techniques as in our previous works, we have been able to compute these coefficients in Feynman gauge (although a result for general covariant gauges is conceptionally not more involved, and could be obtained in a straightforward way given more computational resources).

The structure of the paper is as follows. After briefly summarizing our setup and introducing some notation in sections 2, we give our results for the anomalous dimensions of the ghost sector in 3. In section 4 we display the relations needed to obtain the remaining anomalous dimensions, and to derive all corresponding renormalization constants. All these results are lengthy and given in ancillary files for convenience. We conclude in section 5. In the appendix, the known Beta function coefficients are summarized in our notation.

## 2 Setup

In this section, we start by fixing some necessary conventions and notation, and comment on the computational strategy that we employ for our five-loop computation.

### 2.1 Renormalization constants

The fields and parameters of the gauge theory are renormalized via

$$\psi_b = \sqrt{Z_2}\psi_r, \quad A_b = \sqrt{Z_3}A_r, \quad c_b = \sqrt{Z_3^c}c_r, \quad (2.1)$$

$$m_b = Z_m m_r, \quad g_b = \mu^\epsilon Z_g g_r, \quad \xi_{L,b} = Z_\xi \xi_{L,r}, \quad (2.2)$$

where the subscript  $b$  ( $r$ ) stands for bare (renormalized) quantities, and all  $Z_i = 1 + \mathcal{O}(g_r^2)$ . There is no need to renormalize the gauge-fixing term  $\sim (\partial A)^2/\xi_L$ , such that setting  $Z_\xi = Z_3$  leaves five independent renormalization constants (RCs)  $Z_i$ . It is sometimes convenient to consider products of the  $Z_i$  as ‘vertex RCs’, such as those multiplying the 3-gluon, 4-gluon, ghost-gluon and quark-gluon vertex. These are usually denoted as  $Z_1^j$ , where  $j \in \{3g, 4g, ccg, \psi\psi g\}$ , and we will find it convenient to evaluate the combination  $Z_1^{ccg} = \sqrt{Z_3} Z_3^c Z_g$  instead of  $Z_g$ . For a complete set of relations among these constants, see section 4 below.

Instead of explicitly listing the renormalization constants  $Z_i$ , we will for simplicity only give the corresponding anomalous dimensions, defined by

$$\gamma_i = -\partial_{\ln \mu^2} \ln Z_i. \quad (2.3)$$

Two of them,  $\gamma_m$  and  $\gamma_2$ , corresponding to the renormalization of the quark mass and wave function, have already been given in [10]. Three more are required, in order to complete the set of independent RCs. A popular choice is to evaluate  $\gamma_3$ ,  $\gamma_3^c$  and  $\gamma_1^{ccg}$ , the latter two of which we provide in the present paper, while  $\gamma_3$  can then be reconstructed from the Beta function given in [8], see eq. (4.1) below.

Note that  $\gamma_3$  is conveniently traded for the Beta function, since the latter is a physical gauge invariant object and hence much more compact. Following usual conventions, instead of considering  $Z_g$ , one renormalizes the gauge coupling squared (which in our notation is

$a \sim g^2$ , see eq. (3.1) below) with the factor  $Z_a \equiv Z_g^2$  and calls the corresponding anomalous dimension the Beta function,  $\beta \equiv \gamma_a = 2\gamma_g$ . Note that, due to the renormalization scale independence of the bare gauge coupling, using eqs. (2.2) and (2.3) this immediately implies

$$\beta = \varepsilon + \partial_{\ln \mu^2} \ln a \quad \Leftrightarrow \quad \partial_{\ln \mu^2} a = -a[\varepsilon - \beta], \quad (2.4)$$

where  $a$  is the renormalized coupling of eq. (3.1).

## 2.2 Computational strategy

In order to evaluate the coefficients of the perturbative expansions for the renormalization constants and anomalous dimensions, we rely on the setup that we have developed and successfully tested in our previous works. While we refer to the literature for more detailed descriptions of our chain of computer-algebra programs [7, 10] and for technical details concerning our procedure for integral reduction and evaluation [11–14], let us briefly summarize the main components of our strategy here.

As is standard procedure in perturbative multiloop calculations, we start with expressions produced by the diagram generator `qgraf` [15, 16] linked with some own `FORM` [17–19] codes. Taking the required set of 2- and 3-point functions, we perform the group algebra with the help of `color` [20]. Next, we introduce a common mass term into all our massless propagators [21–23], in order to regulate the infrared behavior of the dimensionally regularized [24, 25]  $d = 4 - 2\varepsilon$  dimensional momentum-space integrals. While this does not change the ultraviolet (UV) behavior that we are interested in when extracting the (mass-independent) UV counterterms in the  $\overline{\text{MS}}$  scheme [26], it allows us to perform a systematic expansion in external momenta, which can eventually be nullified. This leaves us with expansions coefficients that belong to the well-studied class of fully massive vacuum integrals.

At five loops, this class of integrals can be labelled by 15 indices (corresponding to maximally 12 propagators plus 3 scalar products) [12]. To tame the enormous number of integrals that enter our calculation, we choose to perform a reduction to a small set of master integrals. To this end, we make use of our own codes `crusher` [11] and `TIDE` [12], which are based on integration-by-parts (IBP) identities [27] and use Laporta-type algorithms [28]. Both `C++` codes are largely independent, utilize `GiNaC` [29] and `Fermat` [30] for simple and complicated algebraic manipulations, respectively, and in conjunction provide us with a welcome verification of the time- and resource-consuming integral reduction process.

After reduction, we end up with a set of 110 five-loop master integrals. Their high-precision numerical  $\varepsilon$ -expansion has been studied previously [12, 13], much along the lines of previous work on the four-loop case [21, 31–34], relying again on IBP reductions, generating large coupled systems of linear difference equations that can be solved formally with factorial series [28]. A truncated version of these series then delivers high-precision numerical results for the coefficients of the  $\varepsilon$ -expansion of each individual master integral. This allows to employ the integer-relation finding algorithm `PSLQ` [35], testing for relations between some of these numbers, and discovering the analytic content of others.

Our high-precision evaluation of all 5-loop master integrals has not yet produced results for the 12-line families [7]. Fortunately, it turns out that in all our results, only three

independent linear combinations of 12-line master integrals contribute. While standard integration over the Feynman parametric representation (see e.g. [36]) with subsequent sector decomposition, using the strategy explained in [37, 38] and as implemented in FIESTA [39] as well as own code [14] gives 3-6 digits only, we were able to fix the three unknown linear combinations to 260 digits, as explained in [10]. Owing to this last step, we are able to provide analytic expressions for the specific combinations of master integrals that appear in all of our results below.

### 2.3 Notation for group invariants

In order to render the present paper self-contained, we wish to recall some group-theoretic notation that we had already utilized in our previous works [7, 10], and which we employ to present all of our results below. We study a Yang-Mills theory coupled to fermions, working over a semi-simple Lie algebra with hermitian generators  $T^a$ . The commutation relations  $T^a T^b - T^b T^a = i f^{abc} T^c$  define the real and antisymmetric structure constants  $f^{abc}$ . Following standard conventions, the quadratic Casimir operators of the fundamental and adjoint representations (of dimensions  $N_F$  and  $N_A$ , respectively) are defined via  $T^a T^a = C_F \mathbb{1}$  and  $f^{acd} f^{bcd} = C_A \delta^{ab}$ . The trace normalization reads  $\text{Tr}(T^a T^b) = T_F \delta^{ab}$ , the number of quark flavors is denoted by  $N_f$ , and we will make use of the following normalized combinations of group invariants:

$$n_f = \frac{N_f T_F}{C_A} \quad , \quad c_f = \frac{C_F}{C_A} . \tag{2.5}$$

In loop diagrams, higher-order group invariants arise when one encounters traces of more than two group generators. It is useful to classify these higher-order traces in terms of combinations of symmetric tensors [20], of which we presently need the following three (rewriting the generators of the adjoint representation as  $[F^a]_{bc} = -i f^{abc}$ , and again normalizing conveniently):

$$d_1 = \frac{[\text{sTr}(T^a T^b T^c T^d)]^2}{N_A T_F^2 C_A^2} , \quad d_2 = \frac{\text{sTr}(T^a T^b T^c T^d) \text{sTr}(F^a F^b F^c F^d)}{N_A T_F C_A^3} , \quad d_3 = \frac{[\text{sTr}(F^a F^b F^c F^d)]^2}{N_A C_A^4} . \tag{2.6}$$

In the above,  $\text{sTr}$  denotes a fully symmetrized trace (such that  $\text{sTr}(ABC) = \frac{1}{2} \text{Tr}(ABC + ACB)$  etc.).

As a concrete example, picking  $\text{SU}(N)$  as gauge group (and setting  $T_F = \frac{1}{2}$  and  $C_A = N$ ), our set of normalized invariants reads [20]

$$\text{SU}(N) : \quad n_f = \frac{N_f}{2N} , \quad c_f = \frac{N^2 - 1}{2N^2} , \quad d_1 = \frac{N^4 - 6N^2 + 18}{24N^4} , \quad d_2 = \frac{N^2 + 6}{24N^2} , \quad d_3 = \frac{N^2 + 36}{24N^2} . \tag{2.7}$$

In the case of  $\text{SU}(3)$  (corresponding to physical QCD), we therefore have

$$\text{SU}(3) : \quad n_f = \frac{N_f}{6} , \quad c_f = \frac{4}{9} , \quad d_1 = \frac{5}{216} , \quad d_2 = \frac{5}{72} , \quad d_3 = \frac{5}{24} . \tag{2.8}$$

Results for the group  $\text{U}(1)$  (corresponding to QED) can be obtained by setting

$$\text{U}(1) : \quad C_A = 0 , \quad C_F = 1 , \quad T_F = 1 , \quad N_A = 1 , \quad \text{sTr}(T^a T^b T^c T^d) = 1 , \quad \text{sTr}(F^a F^b F^c F^d) = 0 , \tag{2.9}$$

which, due to our normalization, is sometimes only possible after multiplying with the corresponding power of the gauge coupling (that we normalize with a positive power of  $C_A$ , see eq. (3.1) below), eliminating all inverse powers of  $C_A$ .

### 3 Renormalization of ghost field and -vertex

In this section, we present our new results for the ghost field and ghost-gluon vertex anomalous dimensions at five loops. Since these are gauge-dependent quantities, let us stress once more that we have worked in Feynman gauge only. For lower loops, we display the full gauge parameter dependence, where  $\xi = 0/1$  corresponds to Feynman/Landau gauge.

#### 3.1 Ghost field anomalous dimension

In contrast to the physical and gauge-independent Beta function and quark mass anomalous dimension, the anomalous dimension of the ghost field depends on the gauge parameter  $\xi$ . Its structure is

$$\gamma_3^c = -a \left[ -\frac{1}{4}(2 + \xi) + \gamma_{31}^c a + \gamma_{32}^c a^2 + \gamma_{33}^c a^3 + \gamma_{34}^c a^4 + \dots \right], \quad a \equiv \frac{C_A g^2(\mu)}{16\pi^2}, \quad (3.1)$$

where  $g(\mu)$  is the gauge coupling constant that depends on the renormalization scale  $\mu$ , and we work in the  $\overline{\text{MS}}$  scheme, in  $d = 4 - 2\epsilon$  space-time dimensions. The 2- and 3-loop coefficients are known to be (see, e.g. [40])

$$2^5 3^1 \gamma_{31}^c = 5[16n_f] - 2(98 - 3\xi), \quad (3.2)$$

$$2^8 3^3 \gamma_{32}^c = 35[16n_f]^2 + (324(15 - 16\zeta_3)c_f + 2(5 + 189\xi + 1944\zeta_3))[16n_f] - 4(14656 + 1485\xi - 405\xi^2 + 81\xi^3) - 648(4 - \xi)(2 - \xi)\zeta_3. \quad (3.3)$$

The 4-loop coefficient  $\gamma_{33}^c$  is known for the gauge group  $\text{SU}(N)$  [40], while for a general Lie group the result is only available up to the linear term in an expansion around Feynman gauge  $\xi = 0$  [33]. Unfortunately, due to the  $\text{SU}(N)$  degeneracies  $2d_2 = 7/12 - c_f$  and  $d_3 = 37/24 - 3c_f$ , it is not possible to uniquely reconstruct its remaining gauge dependence (up to three loops the reconstruction works, since only quadratic Casimir operators contribute). We have therefore computed  $\gamma_{33}^c$  in general covariant gauge from scratch, obtaining (to clearly expose the group structure of the coefficients, we employ a notation resembling scalar products with vectors in curly brackets, such as e.g.  $\{c_f, 1\} \cdot \{a, b\} = c_f a + b$ )

$$\begin{aligned} 2^{11} 3^4 \gamma_{33}^c &= (83 - 144\zeta_3)[16n_f]^3 \\ &+ \{c_f, 1\} \cdot \{24(1080\zeta_3 - 648\zeta_4 - 115), 2(779\xi - 8315)/3 - 432(43 + 2\xi)\zeta_3 + 11664\zeta_4\} [16n_f]^2 \\ &+ \{c_f^2, d_2, c_f, 1\} \cdot \{-864(271 + 888\zeta_3 - 1440\zeta_5), 124416(4\zeta_3 - 5\zeta_5), \\ &24(22517 + 3825\xi - 864(43 + \xi)\zeta_3 + 1296(23 - \xi)\zeta_4 - 25920\zeta_5), 432(2983 + 42\xi - 6\xi^2)\zeta_3 \\ &- 648(846 - 46\xi + \xi^2)\zeta_4 - 570240\zeta_5 + 14(128354 - 722\xi - 837\xi^2)/3\} [16n_f] \\ &+ \{d_3, 1\} \cdot \{1296(12(28 - 6\xi + \xi^2) - 4(2392 + 108\xi - 63\xi^2 - 17\xi^3 + 16\xi^4))\zeta_3 \\ &+ 5(1696 + 544\xi - 252\xi^2 + 42\xi^3 + 7\xi^4)\zeta_5, \\ &- 4(8202784 + 512546\xi - 111402\xi^2 + 28107\xi^3 - 3888\xi^4)/3 \\ &- 36(159040 - 19104\xi - 162\xi^2 + 1092\xi^3 - 123\xi^4)\zeta_3 + 1296(492 - 376\xi + 91\xi^2 - 9\xi^3)\zeta_4 \\ &+ 270(28832 + 320\xi - 732\xi^2 + 186\xi^3 - 7\xi^4)\zeta_5\}. \end{aligned} \quad (3.4)$$

We observe that eq. (3.4) agrees, in the  $SU(N)$  limit and for all powers of  $\xi$ , with the 4-loop results of [40]. Furthermore, its terms of order  $\xi^0$  and  $\xi^1$  coincide exactly with the corresponding terms given in [33], leading us to the conclusion that it represents the correct generalization of the known results to general covariant gauge. As a side note, we notice that the structure  $c_f n_f^0$  is absent, and that the coefficient of  $d_2 n_f$  does not depend on the gauge parameter. In retrospect, since the same could have been observed — at least to NLO in  $\xi$  — already in [33], assuming this pattern to hold for all powers of the gauge parameter would have allowed for a correct lift of the  $SU(N)$  results of [40] to a general gauge group.

We have evaluated the five-loop contribution in Feynman gauge ( $\xi = 0$ ) as

$$2^{14} 3^5 \gamma_{34}^c = \gamma_{344}^c [16n_f]^4 + \gamma_{343}^c [16n_f]^3 + \gamma_{342}^c [16n_f]^2 + \gamma_{341}^c [16n_f] + \gamma_{340}^c + \mathcal{O}(\xi), \quad (3.5)$$

$$\gamma_{344}^c = 3(65 + 80\zeta_3 - 144\zeta_4), \quad (3.6)$$

$$\gamma_{343}^c = \left\{ c_f, 1 \right\} \cdot \left\{ -2(14765 + 12528\zeta_3 - 38880\zeta_4 + 20736\zeta_5), \right. \\ \left. -3(8325 + 15664\zeta_3 + 12240\zeta_4 - 33408\zeta_5) \right\}, \quad (3.7)$$

$$\gamma_{342}^c = \left\{ c_f^2, c_f, d_1, d_2, 1 \right\} \cdot \left\{ -72(53927 - 182112\zeta_3 + 48384\zeta_3^2 + 42768\zeta_4 + 144000\zeta_5 - 86400\zeta_6), \right. \\ -4(364361 + 484488\zeta_3 - 1804032\zeta_3^2 + 1868184\zeta_4 - 2239488\zeta_5 + 777600\zeta_6), \\ 20736(107 - 109\zeta_3 - 96\zeta_3^2 - 36\zeta_4 + 180\zeta_5), \\ -41472(52\zeta_3 + 18\zeta_3^2 - 36\zeta_4 - 125\zeta_5 + 75\zeta_6), \\ \left. 2(239495 - 3082212\zeta_3 - 1721088\zeta_3^2 + 3863376\zeta_4 - 156384\zeta_5 - 1425600\zeta_6) \right\}, \quad (3.8)$$

$$\gamma_{341}^c = \left\{ c_f^3, c_f^2, c_f d_2, c_f, d_2, d_3, 1 \right\} \cdot \left\{ 746496(7 + 26\zeta_3 + 490\zeta_5 - 560\zeta_7), \right. \\ 576(24617 - 301866\zeta_3 - 196560\zeta_3^2 + 177066\zeta_4 + 274680\zeta_5 - 491400\zeta_6 + 725760\zeta_7), \\ 165888(4 + 66\zeta_3 + 216\zeta_3^2 - 705\zeta_5 + 357\zeta_7), 16(4796303 - 9571932\zeta_3 + 6399648\zeta_3^2 \\ + 11100240\zeta_4 - 16127424\zeta_5 + 8845200\zeta_6 - 10809288\zeta_7), \\ -5184(4192 - 87152\zeta_3 + 21432\zeta_3^2 + 5616\zeta_4 + 89300\zeta_5 - 27300\zeta_6 - 20139\zeta_7), \\ -864(2805 - 86018\zeta_3 - 15960\zeta_3^2 + 43542\zeta_4 - 70360\zeta_5 - 68700\zeta_6 + 192906\zeta_7), \\ 2(52725013 + 136974540\zeta_3 + 1505088\zeta_3^2 - 118046052\zeta_4 - 226012536\zeta_5 \\ \left. + 84380400\zeta_6 + 143718624\zeta_7) \right\}, \quad (3.9)$$

$$\gamma_{340}^c = \left\{ d_3, 1 \right\} \cdot \left\{ -6912(5326 + 771746\zeta_3 - 17934\zeta_3^2 - 209916\zeta_4 - 1172870\zeta_5 + 377625\zeta_6 \right. \\ \left. + 396669\zeta_7), -8(192342607 + 174080040\zeta_3 + 36201384\zeta_3^2 - 103216464\zeta_4 - 855002232\zeta_5 \right. \\ \left. + 222650100\zeta_6 + 492202872\zeta_7) \right\}, \quad (3.10)$$

where eq. (3.6) (as well as the leading- $N_f$  terms at lower loops given in eqs. (3.2)–(3.4)) agrees with the respective term of the known all-loop large- $N_f$  Landau-gauge expression<sup>1</sup> of [41], which in full form can be written as

$$\gamma_3^c|_{\xi=1} = -\eta(a_f)/n_f + \mathcal{O}(1/n_f^2), \quad a_f = 4an_f/3, \quad \eta(\varepsilon) = \frac{(2\varepsilon-3)\Gamma(4-2\varepsilon)}{16\Gamma^2(2-\varepsilon)\Gamma(3-\varepsilon)\Gamma(\varepsilon)}. \quad (3.11)$$

<sup>1</sup>Note that the 1-loop fermion bubble is transverse, such that the leading- $N_f$  does not pick up the gauge-parameter dependence of the bare gluon propagator. Hence, Feynman- and Landau-gauge expressions coincide.

As a second check, specializing to SU(3) allows us to compare with the Feynman-gauge 5-loop result given in eq. (3.2) of [42]. Again, we find full agreement.

### 3.2 Ghost-gluon vertex anomalous dimension

The anomalous dimension of the ghost-gluon vertex has the structure

$$\gamma_1^{cgg} = -a(1 - \xi) \left[ \frac{1}{2} + \frac{6 - \xi}{8} a + \gamma_{12}^{cgg} a^2 + \gamma_{13}^{cgg} a^3 + \gamma_{14}^{cgg} a^4 + \dots \right], \quad (3.12)$$

where the prefactor is consistent with the known fact that the ghost vertex in Landau gauge is finite [43, 44] and hence does not need to be renormalized,  $\gamma_1^{cgg}|_{\xi=1} = 0$ . The 3-loop coefficient can be found in [40]

$$2^7 \gamma_{12}^{cgg} = -15[16n_f] + 2(250 - 59\xi + 10\xi^2). \quad (3.13)$$

The 4-loop coefficient suffers from the same degeneracy of color factors as mentioned above, obstructing a direct generalization from SU( $N$ ) to a general gauge group. We have therefore computed it from scratch, in general covariant gauge, obtaining

$$\begin{aligned} 2^7 3^5 \gamma_{13}^{cgg} &= (-251 + 324\zeta_3)[16n_f]^2 \\ &+ (324(96\zeta_3 + 36\zeta_4 - 161)c_f - 6166 + 4077\xi/2 - 162(164 - 5\xi)\zeta_3 - 8748\zeta_4)[16n_f] \\ &+ 1944((272 - 60\xi + 3\xi^2 + 7\xi^3)\zeta_3 - 5(56 - 12\xi + 3\xi^2 + \xi^3)\zeta_5)d_3 \\ &+ 751120 - 27\xi(5434 - 1332\xi + 171\xi^2) + 81(2528 - 548\xi + 99\xi^2 - \xi^3)\zeta_3 \\ &+ 1458(4 - \xi)(2 - \xi)\zeta_4 - 405(496 - 72\xi + 9\xi^2 + 2\xi^3)\zeta_5. \end{aligned} \quad (3.14)$$

Once again, the SU( $N$ ) limit reproduces all orders of  $\xi$  as known from [40], while the terms of order  $\xi^0$  and  $\xi^1$  coincide with those in the the linear combination  $\gamma_1^{cgg} = \gamma_1^{\psi\psi g} - \gamma_2 + \gamma_3^c$  assembled from [33]. As above, in retrospect, one could have observed the absence of a  $c_f n_f^0$  term to NLO in  $\xi$  from the results of [33], and by conjecturing this to hold for the full gauge-dependent result as well one could have correctly reconstructed the full 4-loop coefficient eq. (3.14) from the SU( $N$ ) results of [40].

At five loops, we have obtained the new Feynman gauge ( $\xi = 0$ ) result

$$2^{14} 3^5 \gamma_{14}^{cgg} = \gamma_{143}^{cgg} [16n_f]^3 + \gamma_{142}^{cgg} [16n_f]^2 + \gamma_{141}^{cgg} [16n_f] + \gamma_{140}^{cgg} + \mathcal{O}(\xi), \quad (3.15)$$

$$\gamma_{143}^{cgg} = -2989 - 1440\zeta_3 + 5184\zeta_4, \quad (3.16)$$

$$\begin{aligned} \gamma_{142}^{cgg} &= \left\{ c_f, 1 \right\} \cdot \left\{ 1296(557 - 736\zeta_3 + 108\zeta_4 + 192\zeta_5), \right. \\ &\quad \left. 251891 + 1591056\zeta_3 - 335016\zeta_4 - 717984\zeta_5 \right\}, \end{aligned} \quad (3.17)$$

$$\begin{aligned} \gamma_{141}^{cgg} &= \left\{ c_f^2, c_f, d_2, d_3, 1 \right\} \cdot \left\{ 5184(3731 + 9588\zeta_3 - 1440\zeta_3^2 + 1332\zeta_4 - 10800\zeta_5 - 3600\zeta_6), \right. \\ &\quad - 1296(45129 - 14192\zeta_3 - 4032\zeta_3^2 + 5616\zeta_4 - 19296\zeta_5 - 7200\zeta_6), \\ &\quad - 31104(1360\zeta_3 + 168\zeta_3^2 + 144\zeta_4 - 1260\zeta_5 - 300\zeta_6 - 441\zeta_7), \\ &\quad - 10368(1126\zeta_3 + 150\zeta_3^2 - 567\zeta_4 - 1200\zeta_5 + 975\zeta_6 - 441\zeta_7), -42165410 \\ &\quad \left. - 432(145015\zeta_3 + 3564\zeta_3^2 - 9168\zeta_4 - 114001\zeta_5 - 10950\zeta_6 + 17640\zeta_7) \right\}, \end{aligned} \quad (3.18)$$

$$\begin{aligned} \gamma_{140}^{cgg} &= \left\{ d_3, 1 \right\} \cdot \left\{ 20736(70330\zeta_3 + 11076\zeta_3^2 - 8856\zeta_4 - 81380\zeta_5 + 16500\zeta_6 - 12607\zeta_7 - 2451), \right. \\ &\quad 8(114251711 + 54643392\zeta_3 + 7060608\zeta_3^2 - 7531704\zeta_4 - 143288568\zeta_5 + 9023400\zeta_6 \\ &\quad \left. + 52599078\zeta_7) \right\}. \end{aligned} \quad (3.19)$$



As an important check, we find full agreement with eq. (40) of [45], where the 5-loop term had been given for SU(3) and in Feynman gauge. Compared to the other anomalous dimensions, note that in  $\gamma_1^{cg}$  there are no terms proportional to  $a^\ell n_f^{\ell-1}$  at  $\ell$  loops; these would correspond to renormalon contributions, which are absent in this case due to consistency with the vanishing of  $\gamma_1^{cg}$  in Landau gauge, as has been mentioned already above.

#### 4 Complete set of renormalization constants

Now that the minimal set of renormalization constants is known, all remaining anomalous dimensions can be reconstructed easily, since they are related via gauge invariance of the QCD action (see e.g. [40]). From the results listed here, the anomalous dimensions of the gluon field, the gluon vertices as well as the quark-gluon vertex can be obtained from the linear relations

$$\gamma_3 = 2(\gamma_1^{cg} - \gamma_3^c) - \beta, \quad \gamma_1^{3g} = 3(\gamma_1^{cg} - \gamma_3^c) - \beta, \quad (4.1)$$

$$\gamma_1^{4g} = 4(\gamma_1^{cg} - \gamma_3^c) - \beta, \quad \gamma_1^{\psi\psi g} = \gamma_1^{cg} - \gamma_3^c + \gamma_2, \quad (4.2)$$

with  $\gamma_2$  from [10]. For the convenience of the reader, we attach an electronic version of the complete set of anomalous dimensions to the present paper.<sup>2</sup>

To reconstruct the renormalization constants  $Z_i$  from the set of anomalous dimensions  $\gamma_i$ , one starts from eq. (2.3), recalling that in general, renormalization scale dependence enters  $Z_i(a, \xi_L)$  through both of its variables. Therefore,

$$\begin{aligned} \gamma_i &= -(\partial_{\ln \mu^2} a)(\partial_a \ln Z_i) - (\partial_{\ln \mu^2} \ln \xi_L)(\partial_{\ln \xi_L} \ln Z_i) \\ &= -a(\beta - \varepsilon)(\partial_a \ln Z_i) - \gamma_3(\xi - 1)(\partial_\xi \ln Z_i) \\ &= -a(\beta - \varepsilon)(\partial_a \ln Z_i) - (2\gamma_1^{cg} - 2\gamma_3^c - \beta)(\xi - 1)(\partial_\xi \ln Z_i), \end{aligned} \quad (4.3)$$

where in the second line we have been careful to use the  $d$ -dimensional version of the Beta function of eq. (2.4), exploited that the bare gauge parameter renormalizes as the gluon field  $\xi_{L,b} = Z_3 \xi_{L,r}$  and changed the gauge parameter to our preferred notation  $\xi$  whose powers correspond to an expansion around Feynman gauge, the relation to  $\xi_L$  being  $\xi_L + \xi = 1$ . Finally, we have for convenience traded the gluon field anomalous dimension for the ones that we have given explicitly above. Writing the renormalization constants as  $Z_i = 1 + \sum_{n>0} z_i^{(n)}/\varepsilon^n$ , the coefficients  $z_i^{(n)}$  then follow by solving eq. (4.3), requiring  $\gamma_3$  (viz  $\gamma_1^{cg}$ ,  $\gamma_3^c$  and  $\beta$ ) at one loop lower only. Since the expressions for the complete 5-loop renormalization constants are somewhat large, we refrain from listing them here, but provide electronic versions thereof.<sup>2</sup>

Once the RCs  $Z_i$  are known, the corresponding anomalous dimensions can simply be extracted from the single poles, as  $\gamma_i = a\partial_a z_i^{(1)}$ .

---

<sup>2</sup>Electronic version of the complete set of 5-loop results for  $\gamma_i$  and  $Z_i$ , to be obtained as online resource of this article.

## 5 Conclusions

In recent years, technical progress has made possible the first complete five-loop computations in non-Abelian gauge theories. Adding to the already available set of renormalization constants for general Lie groups in  $\overline{\text{MS}}$ -like schemes [7, 8, 10], we have presented new analytic 5-loop Feynman-gauge results for the two missing anomalous dimensions, which we have chosen to be those of the ghost field eqs. (3.5)–(3.10) and ghost-gluon vertex eqs. (3.15)–(3.19). This completes the five-loop renormalization program for general groups, providing a generalization — and independent confirmation — of the previously known SU(3) coefficients [6, 9, 42, 45], relevant for physical QCD.

Along the way, we have closed a gap in the literature and provided full gauge-dependent expressions for the renormalization constants of the ghost sector at four loops, see eqs. (3.4) and (3.14) above. Together with our new results, we have prepared computer-readable versions of the complete set of anomalous dimensions and renormalization constants, available online.<sup>2</sup>

The methods we have employed here are well suited to be applied to the gluon propagator as well, and the corresponding computation of the gluon field anomalous dimension  $\gamma_3$  is under way [46]. The anticipated result would give an important independent check on the Feynman-gauge expression that we have provided in the ancillary files (which was derived using eq. (4.1)), and hence on the correctness of the Beta function from the independent calculation of [8].

For completeness, it might be interesting to evaluate the gauge parameter dependence of ghost field and -vertex (as well as the quark field) in the future. While this would, for example, provide a further independent check of the correctness of the Beta function (as well as the quark mass anomalous dimension  $\gamma_m$ ) due to gauge-parameter cancellation, given the strong constraints already discussed above we do not think this a pressing issue. However, from the viewpoint of truly completing the 5-loop renormalization program, knowledge of the full gauge dependence of all renormalization constants is certainly desirable.

In passing, we note that the analytic structure of the 5-loop Beta function (as well as that of the corresponding renormalization constant  $Z_a$ ), containing the Zeta values  $\{\zeta_3, \zeta_4, \zeta_5\}$  only, is considerably simpler than that of the other anomalous dimensions and RCs, which in addition need the weight-6 and weight-7 constants  $\{\zeta_3^2, \zeta_6, \zeta_7\}$ . Regarding the group-theoretic structure, the only outlier is  $\gamma_1^{ccg}$  (and  $Z_1^{ccg}$ ), where the factor  $d_1$  is absent.

## Acknowledgments

The work of T.L. has been supported in part by DFG grants GRK 881 and SCHR 993/2; he is grateful to the theory group of the University of Bielefeld for hospitality, and for continued access to parts of their computer clusters. A.M. is supported by a European Union COFUND/Durham Junior Research Fellowship under EU grant agreement number 267209. P.M. was supported in part by the EU Network HIGGSTOOLS PITN-GA-2012-316704. Y.S. acknowledges support from FONDECYT project 1151281 and UBB project GI-152609/VC.

## A Beta function

Being an important fundamental parameter of gauge theories, considerable effort has gone into evaluating the Beta function over the past four decades. After groundbreaking work at one loop [2, 3], establishing the asymptotically free nature of the strong coupling constant, the 2-loop [47, 48], 3-loop [49, 50] and 4-loop [21, 33] perturbative corrections have been evaluated, made possible by major technological developments that were pushed ahead in parallel. Five-loop results have appeared over the past 8 years or so, first for the case of QED [51–53], later for SU(3) [6, 54], and finally for general Lie groups [7, 8].

In order to translate the recent results of [8] to our notation (as introduced in section 2.3 and eq. (3.1)), we define the  $L$ -loop coefficients  $b_{L-1}$  of the Beta function as

$$\partial_{\ln \mu^2} a = -a \left[ \varepsilon - \beta \right] = -a \left[ \varepsilon + b_0 a + b_1 a^2 + b_2 a^3 + b_3 a^4 + b_4 a^5 + \dots \right]. \quad (\text{A.1})$$

The coefficients are polynomials in  $n_f$ , and up to four loops read

$$3^1 b_0 = [-4] n_f + 11, \quad (\text{A.2})$$

$$3^2 b_1 = [-36c_f - 60] n_f + 102, \quad (\text{A.3})$$

$$3^3 b_2 = [132c_f + 158] n_f^2 + [54c_f^2 - 615c_f - 1415] n_f + 2857/2, \quad (\text{A.4})$$

$$3^5 b_3 = [1232c_f + 424] n_f^3 + 432(132\zeta_3 - 5)d_3 + (150653/2 - 1188\zeta_3) + [72(169 - 264\zeta_3)c_f^2 + 64(268 + 189\zeta_3)c_f + 1728(24\zeta_3 - 11)d_1 + 6(3965 + 1008\zeta_3)] n_f^2 + [11178c_f^3 + 36(264\zeta_3 - 1051)c_f^2 + (7073 - 17712\zeta_3)c_f + 3456(4 - 39\zeta_3)d_2 + 3(3672\zeta_3 - 39143)] n_f, \quad (\text{A.5})$$

which the five-loop coefficient can be represented as

$$3^5 b_4 = b_{44} n_f^4 + b_{43} n_f^3 + b_{42} n_f^2 + b_{41} n_f + b_{40}, \quad (\text{A.6})$$

$$b_{44} = \left\{ c_f, 1 \right\} \cdot \left\{ -8(107 + 144\zeta_3), 4(229 - 480\zeta_3) \right\}, \quad (\text{A.7})$$

$$b_{43} = \left\{ c_f^2, c_f, d_1, 1 \right\} \cdot \left\{ -6(4961 - 11424\zeta_3 + 4752\zeta_4), -48(46 + 1065\zeta_3 - 378\zeta_4), 1728(55 - 123\zeta_3 + 36\zeta_4 + 60\zeta_5), -3(6231 + 9736\zeta_3 - 3024\zeta_4 - 2880\zeta_5) \right\}, \quad (\text{A.8})$$

$$b_{42} = \left\{ c_f^3, c_f^2, c_f d_1, c_f, d_2, d_1, 1 \right\} \cdot \left\{ -54(2509 + 3216\zeta_3 - 6960\zeta_5), 9(94749/2 - 28628\zeta_3 + 10296\zeta_4 - 39600\zeta_5), 25920(13 + 16\zeta_3 - 40\zeta_5), 3(5701/2 + 79356\zeta_3 - 25488\zeta_4 + 43200\zeta_5), -864(115 - 1255\zeta_3 + 234\zeta_4 + 40\zeta_5), -432(1347 - 2521\zeta_3 + 396\zeta_4 - 140\zeta_5), 843067/2 + 166014\zeta_3 - 8424\zeta_4 - 178200\zeta_5 \right\}, \quad (\text{A.9})$$

$$b_{41} = \left\{ c_f^4, c_f^3, c_f^2, c_f d_2, c_f, d_3, d_2, 1 \right\} \cdot \left\{ -81(4157/2 + 384\zeta_3), 81(11151 + 5696\zeta_3 - 7480\zeta_5), -3(548732 + 151743\zeta_3 + 13068\zeta_4 - 346140\zeta_5), -25920(3 - 4\zeta_3 - 20\zeta_5), 8141995/8 + 35478\zeta_3 + 73062\zeta_4 - 706320\zeta_5, 216(113 - 2594\zeta_3 + 396\zeta_4 + 500\zeta_5), 216(1414 - 15967\zeta_3 + 2574\zeta_4 + 8440\zeta_5), -5048959/4 + 31515\zeta_3 - 47223\zeta_4 + 298890\zeta_5 \right\}, \quad (\text{A.10})$$

$$b_{40} = \left\{ d_3, 1 \right\} \cdot \left\{ -162(257 - 9358\zeta_3 + 1452\zeta_4 + 7700\zeta_5), 8296235/16 - 4890\zeta_3 + 9801\zeta_4/2 - 28215\zeta_5 \right\}. \quad (\text{A.11})$$

Of these 5-loops coefficients,  $b_{44}$  has been known already for a long time from a large- $N_f$  analysis [55, 56], while  $b_{43}$  was given in [7], as a proof-of-concept of our setup that we have used above to determine the anomalous dimensions of the ghost sector. The three coefficients  $b_{42}$ ,  $b_{41}$  and  $b_{40}$  have recently been derived as well [8], using the background field method and relying on infrared rearrangement [57] and the  $R^*$  operation [58] to map the ultraviolet divergences onto massless four-loop two-point functions which were evaluated via their new code FORCER [59]. As an important check on the 5-loop expressions given above, setting the group invariants to their SU(3) values eq. (2.8), all coefficients coincide with the results given in [6].

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] G. 't Hooft, *Example of a gauge field theory*, talk at the *Marseille conference on renormalization of Yang-Mills fields and applications to particle physics*, Marseille, France, June 1972, unpublished.
- [2] D.J. Gross and F. Wilczek, *Ultraviolet Behavior of Nonabelian Gauge Theories*, *Phys. Rev. Lett.* **30** (1973) 1343 [[INSPIRE](#)].
- [3] H.D. Politzer, *Reliable Perturbative Results for Strong Interactions?*, *Phys. Rev. Lett.* **30** (1973) 1346 [[INSPIRE](#)].
- [4] D.V. Batkovich, K.G. Chetyrkin and M.V. Kompaniets, *Six loop analytical calculation of the field anomalous dimension and the critical exponent  $\eta$  in  $O(n)$ -symmetric  $\phi^4$  model*, *Nucl. Phys. B* **906** (2016) 147 [[arXiv:1601.01960](#)] [[INSPIRE](#)].
- [5] M. Kompaniets and E. Panzer, *Renormalization group functions of  $\phi^4$  theory in the MS-scheme to six loops*, *PoS(LL2016)038* [[arXiv:1606.09210](#)] [[INSPIRE](#)].
- [6] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Five-Loop Running of the QCD coupling constant*, [arXiv:1606.08659](#) [[INSPIRE](#)].
- [7] T. Luthe, A. Maier, P. Marquard and Y. Schröder, *Towards the five-loop  $\beta$ -function for a general gauge group*, *JHEP* **07** (2016) 127 [[arXiv:1606.08662](#)] [[INSPIRE](#)].
- [8] F. Herzog, B. Ruijl, T. Ueda, J.A.M. Vermaseren and A. Vogt, *The five-loop  $\beta$ -function of Yang-Mills theory with fermions*, [arXiv:1701.01404](#) [[INSPIRE](#)].
- [9] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Quark Mass and Field Anomalous Dimensions to  $\mathcal{O}(\alpha_s^5)$* , *JHEP* **10** (2014) 076 [[arXiv:1402.6611](#)] [[INSPIRE](#)].
- [10] T. Luthe, A. Maier, P. Marquard and Y. Schröder, *Five-loop quark mass and field anomalous dimensions for a general gauge group*, *JHEP* **01** (2017) 081 [[arXiv:1612.05512](#)] [[INSPIRE](#)].
- [11] P. Marquard and D. Seidel, *crusher*, unpublished.
- [12] T. Luthe, *Fully massive vacuum integrals at 5 loops*, Ph.D. Thesis, Bielefeld University (2015) [[INSPIRE](#)].
- [13] T. Luthe and Y. Schröder, *Fun with higher-loop Feynman diagrams*, *J. Phys. Conf. Ser.* **762** (2016) 012066 [[arXiv:1604.01262](#)] [[INSPIRE](#)].

- [14] T. Luthe and Y. Schröder, *Five-loop massive tadpoles*, [PoS\(LL2016\)074](#) [[arXiv:1609.06786](#)] [[INSPIRE](#)].
- [15] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279 [[INSPIRE](#)].
- [16] P. Nogueira, *Abusing qgraf*, *Nucl. Instrum. Meth. A* **559** (2006) 220 [[INSPIRE](#)].
- [17] J.A.M. Vermaseren, *New features of FORM*, [math-ph/0010025](#) [[INSPIRE](#)].
- [18] M. Tentyukov and J.A.M. Vermaseren, *The Multithreaded version of FORM*, *Comput. Phys. Commun.* **181** (2010) 1419 [[hep-ph/0702279](#)] [[INSPIRE](#)].
- [19] J. Kuipers, T. Ueda, J.A.M. Vermaseren and J. Vollinga, *FORM version 4.0*, *Comput. Phys. Commun.* **184** (2013) 1453 [[arXiv:1203.6543](#)] [[INSPIRE](#)].
- [20] T. van Ritbergen, A.N. Schellekens and J.A.M. Vermaseren, *Group theory factors for Feynman diagrams*, *Int. J. Mod. Phys. A* **14** (1999) 41 [[hep-ph/9802376](#)] [[INSPIRE](#)].
- [21] T. van Ritbergen, J.A.M. Vermaseren and S.A. Larin, *The Four loop  $\beta$ -function in quantum chromodynamics*, *Phys. Lett. B* **400** (1997) 379 [[hep-ph/9701390](#)] [[INSPIRE](#)].
- [22] M. Misiak and M. Münz, *Two loop mixing of dimension five flavor changing operators*, *Phys. Lett. B* **344** (1995) 308 [[hep-ph/9409454](#)] [[INSPIRE](#)].
- [23] K.G. Chetyrkin, M. Misiak and M. Münz,  *$\beta$ -functions and anomalous dimensions up to three loops*, *Nucl. Phys. B* **518** (1998) 473 [[hep-ph/9711266](#)] [[INSPIRE](#)].
- [24] C.G. Bollini and J.J. Giambiagi, *Dimensional Renormalization: The Number of Dimensions as a Regularizing Parameter*, *Nuovo Cim. B* **12** (1972) 20 [[INSPIRE](#)].
- [25] G. 't Hooft and M.J.G. Veltman, *Regularization and Renormalization of Gauge Fields*, *Nucl. Phys. B* **44** (1972) 189 [[INSPIRE](#)].
- [26] G. 't Hooft, *Dimensional regularization and the renormalization group*, *Nucl. Phys. B* **61** (1973) 455 [[INSPIRE](#)].
- [27] K.G. Chetyrkin and F.V. Tkachov, *Integration by Parts: The Algorithm to Calculate  $\beta$ -functions in 4 Loops*, *Nucl. Phys. B* **192** (1981) 159 [[INSPIRE](#)].
- [28] S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, *Int. J. Mod. Phys. A* **15** (2000) 5087 [[hep-ph/0102033](#)] [[INSPIRE](#)].
- [29] C.W. Bauer, A. Frink and R. Kreckel, *Introduction to the GiNaC framework for symbolic computation within the C++ programming language*, *J. Symb. Comput.* **33** (2000) 1 [[cs/0004015](#)] [[INSPIRE](#)].
- [30] R.H. Lewis, *fermat*, <http://home.bway.net/lewis/>.
- [31] S. Laporta, *High precision  $\epsilon$ -expansions of massive four loop vacuum bubbles*, *Phys. Lett. B* **549** (2002) 115 [[hep-ph/0210336](#)] [[INSPIRE](#)].
- [32] Y. Schröder, *Automatic reduction of four loop bubbles*, *Nucl. Phys. Proc. Suppl.* **116** (2003) 402 [[hep-ph/0211288](#)] [[INSPIRE](#)].
- [33] M. Czakon, *The Four-loop QCD  $\beta$ -function and anomalous dimensions*, *Nucl. Phys. B* **710** (2005) 485 [[hep-ph/0411261](#)] [[INSPIRE](#)].
- [34] Y. Schröder and A. Vuorinen, *High-precision  $\epsilon$ -expansions of single-mass-scale four-loop vacuum bubbles*, *JHEP* **06** (2005) 051 [[hep-ph/0503209](#)] [[INSPIRE](#)].

- [35] H.R.P. Ferguson, D.H. Bailey and S. Arno, *Analysis of PSLQ, an integer relation finding algorithm*, *J. Math. Comput.* **68** (1999) 351.
- [36] V.A. Smirnov, *Analytic tools for Feynman integrals*, *Springer Tracts Mod. Phys.* **250** (2012) 1 [INSPIRE].
- [37] M. Roth and A. Denner, *High-energy approximation of one loop Feynman integrals*, *Nucl. Phys. B* **479** (1996) 495 [hep-ph/9605420] [INSPIRE].
- [38] T. Binoth and G. Heinrich, *Numerical evaluation of multiloop integrals by sector decomposition*, *Nucl. Phys. B* **680** (2004) 375 [hep-ph/0305234] [INSPIRE].
- [39] A.V. Smirnov, *FIESTA4: Optimized Feynman integral calculations with GPU support*, *Comput. Phys. Commun.* **204** (2016) 189 [arXiv:1511.03614] [INSPIRE].
- [40] K.G. Chetyrkin, *Four-loop renormalization of QCD: Full set of renormalization constants and anomalous dimensions*, *Nucl. Phys. B* **710** (2005) 499 [hep-ph/0405193] [INSPIRE].
- [41] J.A. Gracey, *Quark, gluon and ghost anomalous dimensions at  $O(1/N(f))$  in quantum chromodynamics*, *Phys. Lett. B* **318** (1993) 177 [hep-th/9310063] [INSPIRE].
- [42] K.G. Chetyrkin, P.A. Baikov and J.H. Kühn, *Towards QCD running in 5 loops: quark mass anomalous dimension*, *PoS(RADCOR 2013)056* [arXiv:1402.6606] [INSPIRE].
- [43] J.C. Taylor, *Ward Identities and Charge Renormalization of the Yang-Mills Field*, *Nucl. Phys. B* **33** (1971) 436 [INSPIRE].
- [44] A. Blasi, O. Piguet and S.P. Sorella, *Landau gauge and finiteness*, *Nucl. Phys. B* **356** (1991) 154 [INSPIRE].
- [45] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Massless Propagators,  $R(s)$  and Multiloop QCD*, *Nucl. Part. Phys. Proc.* **261-262** (2015) 3 [arXiv:1501.06739] [INSPIRE].
- [46] T. Luthe, A. Maier, P. Marquard and Y. Schröder, *The five-loop Beta function for a general gauge group*, in preparation.
- [47] W.E. Caswell, *Asymptotic Behavior of Nonabelian Gauge Theories to Two Loop Order*, *Phys. Rev. Lett.* **33** (1974) 244 [INSPIRE].
- [48] D.R.T. Jones, *Two Loop Diagrams in Yang-Mills Theory*, *Nucl. Phys. B* **75** (1974) 531 [INSPIRE].
- [49] O.V. Tarasov, A.A. Vladimirov and A.Yu. Zharkov, *The Gell-Mann-Low Function of QCD in the Three Loop Approximation*, *Phys. Lett. B* **93** (1980) 429 [INSPIRE].
- [50] S.A. Larin and J.A.M. Vermaseren, *The three loop QCD  $\beta$ -function and anomalous dimensions*, *Phys. Lett. B* **303** (1993) 334 [hep-ph/9302208] [INSPIRE].
- [51] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Massless propagators: Applications in QCD and QED*, *PoS(RAD COR 2007)023* [arXiv:0810.4048] [INSPIRE].
- [52] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Adler Function, Bjorken Sum Rule and the Crewther Relation to Order  $\alpha_s^4$  in a General Gauge Theory*, *Phys. Rev. Lett.* **104** (2010) 132004 [arXiv:1001.3606] [INSPIRE].
- [53] P.A. Baikov, K.G. Chetyrkin, J.H. Kühn and J. Rittinger, *Vector Correlator in Massless QCD at Order  $O(\alpha_s^4)$  and the QED  $\beta$ -function at Five Loop*, *JHEP* **07** (2012) 017 [arXiv:1206.1284] [INSPIRE].

- [54] K. Chetyrkin, P. Baikov and J. Kühn, *The  $\beta$ -function of Quantum Chromodynamics and the effective Higgs-gluon-gluon coupling in five-loop order*, [PoS\(LL2016\)010](#) [[INSPIRE](#)].
- [55] A. Palanques-Mestre and P. Pascual, *The  $1/N_f$  Expansion of the  $\gamma$  and  $\beta$ -functions in QED*, *Commun. Math. Phys.* **95** (1984) 277 [[INSPIRE](#)].
- [56] J.A. Gracey, *The QCD  $\beta$ -function at  $O(1/N_f)$* , *Phys. Lett. B* **373** (1996) 178 [[hep-ph/9602214](#)] [[INSPIRE](#)].
- [57] A.A. Vladimirov, *Methods of Multiloop Calculations and the Renormalization Group Analysis of  $\phi^4$  Theory*, *Theor. Math. Phys.* **36** (1979) 732 [[INSPIRE](#)].
- [58] K.G. Chetyrkin and V.A. Smirnov,  *$R^*$  operation corrected*, *Phys. Lett. B* **144** (1984) 419 [[INSPIRE](#)].
- [59] T. Ueda, B. Ruijl and J.A.M. Vermaseren, *Forcer: a FORM program for 4-loop massless propagators*, [PoS\(LL2016\)070](#) [[arXiv:1607.07318](#)] [[INSPIRE](#)].