

B03

CERN LIBRARIES, GENEVA



SCAN-9508198

E4-95-80

SC9534

N.V.Antonenko, E.A.Cherepanov, A.K.Nasirov,
V.P.Permjakov, V.V.Volkov

COMPOUND NUCLEUS FORMATION
IN REACTIONS BETWEEN MASSIVE NUCLEI.
FUSION BARRIER

Submitted to «Physical Review C»

I. INTRODUCTION

The mechanism of the compound nucleus formation (CNF) in heavy ion-induced reactions is of considerable interest. However, its experimental investigation is very difficult since during the complete fusion the system does not give signals which allow one to judge about this process unambiguously. In experiments, the decay products of an excited compound nucleus are registered, however, they contain no information about the mechanism of the CNF. A consistent theoretical analysis of the complete fusion process of two multinucleon interacting systems is a very complicated problem. Therefore, a number of models have been developed for the description of the experimental data. These models are based on simplifying assumptions about the fusion process.

The critical distance model [1], the optical model [2] and the surface friction model [3,4] widely used for the calculation of fusion cross sections do not consider the mechanism of CNF itself. It is postulated usually that after the capture of a projectile by a target nucleus the complete fusion occurs inevitably. One can say sometimes that the process of compound nucleus formation looks like the nuclear collapse. The macroscopic dynamic model [5-7] allows one to trace the evolution of the fusion system in time. However, such important properties of nuclei as their nucleon composition and shell structure are not taken into account. In replacing the real atomic nuclei by homogeneous and structureless drops of a hypothetical nuclear liquid, the real process of the compound nucleus formation is inevitably distorted.

In [8] a new approach to the analysis of the complete fusion process has been suggested. This approach is based on the information about the interaction of two complex nuclei being in close contact, which has been obtained in the study of deep inelastic transfer reactions. In the framework of this approach the complete fusion of the nuclei is interpreted in the following way. At the capture stage after the full dissipation of the collision kinetic energy a dinuclear system (DNS) is formed. The DNS evolves to a compound nucleus by the nucleon transfer from a light nucleus to a heavy one. An important peculiarity of

the DNS evolution is the retaining of the individuality of nuclei through their way to the compound nucleus formation. During the DNS evolution all nucleons of the donor-nucleus occur to be transferred shell by shell to the acceptor-nucleus. This approach can be called the “DNS-approach”.

How to reveal the real mechanism of compound nucleus formation? Our calculations of compound nucleus formation cross sections in the framework of different models have demonstrated, that the complete fusion of massive nuclei ($A \geq 100$) can be used as a good test of validity of various complete fusion models.

As an illustration, for the reactions $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$ the experimental data [9] on evaporation residue cross sections are compared with results calculated in the framework of the standard models: the optical model [2], the surface friction model [1], the macroscopic dynamic model [7] (Sec. II). The calculations include the determination of the cross section of compound nucleus formation and the analysis of the competition between various de-excitation channels. The calculated results are in dramatic discrepancy with the experimental data. For the reaction $^{110}\text{Pd}+^{110}\text{Pd}$ the calculated data are several orders of magnitude larger than the experimental one.

In our opinion, this discrepancy is the result of quasi fission of a massive DNS formed in these reactions after full dissipation of the kinetic energy of collision (Sec. III) [10]. In the existing models of complete fusion the competition between complete fusion and quasi fission in the initial DNS is not taken into account. On the basis of the DNS-approach a model is proposed to calculate the competition between complete fusion and quasi fission in massive symmetric DNS formed at the collision energies above the Coulomb barrier. This model seems to be applicable also for almost symmetric system $^{124}\text{Sn}+^{96}\text{Zr}$ (Sec. IV). The evaporation residue cross section ($\sigma_{ER}(E)$) values calculated for the reactions $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$ by the proposed model are close to the experimental data (Sec. V). This fact can be considered as an indication of the validity of the interpretation of the compound nucleus formation mechanism suggested in the DNS-approach.

II. CALCULATION OF $\sigma_{ER}(E)$ IN THE FRAMEWORK OF STANDARD MODELS OF COMPLETE FUSION

A. Calculation of the compound nucleus formation cross section in the framework of the optical model

The CNF cross section $\sigma_{CN}(E)$ was estimated by one of the variants of the optical model that was used for the description of the experimental data on the synthesis of transuranium elements [2]. The model parameters have been systematized in a wide region of the product of atomic numbers of colliding nuclei, $Z_1 \cdot Z_2$ by comparing results of the calculation with the experimental data.

The cross section σ_{CN} is a part of the total reaction cross section σ_R

$$\sigma_R = \pi \lambda_0^2 \sum_{l=0}^{\infty} (2l+1) T(l, E_{c.m.}), \quad (1)$$

Here λ_0 is the de Broglie wave length of the relative motion of interacting nuclei, $E_{c.m.}$ is the bombarding energy in the center-of-mass system, T is the penetration coefficient of the l -th partial wave through the potential barrier. $T(l, E_{c.m.})$ is approximated by the penetration factor of a parabolic barrier. The potential describing the nucleus-nucleus interaction includes nuclear, Coulomb and centrifugal potentials

$$V(R) = V_N + V_C + V_r, \quad (2)$$

$$V_N = V_0 \left\{ 1 + \exp \left[\frac{R - r_{0v}(A_1^{1/3} + A_2^{1/3})}{d} \right] \right\}^{-1}, \quad (3)$$

$$V_C = \begin{cases} Z_1 Z_2 e^2 / R, & \text{if } R > R_C, \\ Z_1 Z_2 e^2 / 2R_C (3 - R^2 / R_C^2), & \text{if } R \leq R_C, \end{cases} \quad (4)$$

$$V_r = \hbar^2 l(l+1) / 2\mu R^2, \quad (5)$$

where R is the distance between nuclear centres; $R_C = 1.3(A_1^{1/3} + A_2^{1/3})$ fm, μ is the reduced mass of the system. The quantities corresponding to the projectile and target

nuclei are marked by indices 1 and 2, respectively. The potential parameters V_0 , r_0 , and d are taken from [2].

In the classical optical model the CNF cross section is calculated by using the imaginary part of the nucleus-nucleus potential. In [2] the empirical systematics of the ratio σ_{CN}/σ_R as a functions of Z_1Z_2 have been established. This systematics is the result of comparison of the experimental values of σ_{CN} and σ_R , obtained in dozens of reactions. The values of σ_{CN} in the reactions $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$ have been found by calculating σ_R and using the empirical systematics of the ratio σ_{CN}/σ_R presented in [2]. In the calculations of σ_{CN} for a heavy system with $Z_1Z_2 > 1500$, the energy displacement between the fusion barrier and interaction barrier should be taken into account. In the present paper this is done according to [2].

B. Calculation of the capture cross section in the framework of the surface friction model

Our calculations are based on one of the recent variants of the surface friction model which takes into account the dynamic deformation of both the colliding nuclei [4]. It should be noted that in the framework of this model the capture cross section can be described satisfactorily for such a massive ion as ^{86}Kr .

In this model the capture (fusion) is assumed to take place if the projectile after full dissipation of the kinetic energy finds oneself in the potential pocket of $V(R, \alpha)$ (α_i are the deformation parameters). By introducing the nuclear friction and solving the system of classical equations of motion, we can obtain the critical angular momentum l_c . All the trajectories with $l < l_c$ lead to the capture or fusion. In the sharp cut off approximation the capture (fusion) cross section is determined by the expression:

$$\sigma_c = \pi \lambda_0^2 \sum_{l=0}^{l_c} (2l+1) = \pi \lambda_0^2 (l_c + 1)^2. \quad (6)$$

In the reactions with relatively light projectiles σ_c is equal to σ_{CN} . However, in the reactions between massive nuclei where the quasi-fission (fast fission) takes place the

surface friction model can give only σ_c . Nevertheless, we have made the calculations of σ_{ER} for the reactions $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$ assuming that $\sigma_{CN} = \sigma_c$. The comparison of the calculation results with the experimental σ_{ER} values give the scale of the quasi-fission in these reactions.

C. Calculation of the compound nucleus formation cross section in the framework of the macroscopic dynamic model

A variant of the macroscopic dynamic model [7] has been used to calculate the CNF cross section in the reaction $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$. In this model σ_{CN} is determined by the following expression:

$$\sigma_{CN}(E_{c.m.}) = \frac{\pi r_c^2}{E_{c.m.}} \left[\sqrt{\left(\frac{c_1 c_2 + 0.5}{c_2^2} \right)^2 - \left(\frac{c_1^2 + E_B - E_{c.m.}}{c_2^2} \right)} - \left(\frac{c_1 c_2 + 0.5}{c_2^2} \right) \right], \quad (7)$$

where

$$c_1 = k^{1/2} \left[(Z^2/A)_{i,f} - (Z^2/A)_{cf}^{thr} \right], \quad c_2 = \frac{k^{1/2}}{c^2/r_0} \frac{8f^2}{(A_1 A_2)^{1/3}}$$

$$k = 2025 \cdot (A_1 A_2)^{1/3} (A_1^{1/3} + A_2^{1/3})^2 32 \left(\frac{3}{\pi} \right)^{2/3} m c^2 a^2 / (A_1 + A_2)$$

$$r_c = D_1 + D_2 + 1.44 \text{ fm}, \quad D_i = R_i - 1/R_i,$$

$$R_i = 1.28 A_i^{1/3} - 0.76 + 0.8 A_i^{-1/3}, \quad m c^2 = 931 \text{ MeV}, \quad r_0 = 1.224 \text{ fm}.$$

In [7] the following values for parameters are recommended for a better description of the experimental data:

$$f = 3/4, \quad a = 12, \quad (Z^2/A)_{cf}^{thr} = 33.$$

In the model [7] the result of massive nuclei collision depends on a relationship between the kinetic energy of the collision, $E_{c.m.}$, the Coulomb barrier, B_C , and the extra-extra push energy, E_{xx} . If $E_{c.m.} > B_C + E_{xx}$ during collision the nuclear system takes a more compact shape than the saddle-point one of the compound nucleus, thus the complete fusion occurs. In the case of $E_{c.m.} < B_C + E_{xx}$ the nuclei cannot fuse and the system

decays via quasi-fission or deep inelastic transfer channel. In the reaction $^{100}\text{Mo}+^{100}\text{Mo}$ the E_{xx} value is equal to 1 MeV, in the reaction $^{110}\text{Pd}+^{110}\text{Pd}$ the E_{xx} value is equal to 60 MeV [7]. Since in the second reaction the Bass barrier equals to 228 MeV, at $E_{c.m.} > 288$ MeV the macroscopic dynamic model was expected to be capable of describing σ_{CN} .

D. De-excitation of compound nucleus

In the reactions $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$ the compound nuclei ^{200}Po and ^{220}U are formed with the excitation energy of dozens of MeV and with a large set of angular momenta. The competition between fission and emission of a light particle determines the part of compound nuclei surviving as evaporation residues.

To describe the decay of the excited nuclei ^{200}Po and ^{220}U , a statistical model based on the Monte-Carlo method has been used [11,12]. The angular momenta of compound nuclei formed in the complete fusion reaction have a respective distribution of values of I . The vector \mathbf{I} is transversal to the ion beam. By means of two random numbers the drawing of the momentum value and its orientation in space are performed. Then for different decay channels of the compound nucleus the maximum of residual energy is defined in the following way

$$E_{\nu}^{\text{res}} = E^* - E_r - E_{\nu} - V_{\nu}; \quad E_f^{\text{res}} = E^* - E_r - B_f.$$

Here E^* is the excitation energy of the compound nucleus, E_r is its rotational energy, V_{ν} is the exit Coulomb barrier for a particle of the kind ν ($\nu = \text{n, p, d, t, } ^3\text{He, } \alpha$), E_{ν} is the kinetic energy of the particle and B_f is the fission barrier. For all $E_{\nu}^{\text{res}} > 0$ the kind of the emitted particle or γ -ray is drawn. For partial widths of the particle ν emission, for the fission and of the γ -quanta emission the following expressions have been used [13]:

$$\Gamma_{\nu}(E_m^*, I_m) \approx \frac{(2s_{\nu} + 1)\mu_{\nu}}{(\pi\hbar)^2\rho_m(U)} \int_{V_{\nu}}^{U-B_{\nu}} \sigma_{\text{inv}}(E_{\nu})\rho_d(U - B_{\nu} - E_{\nu})E_{\nu}dE_{\nu}, \quad (8)$$

$$\Gamma_f(E_m^*, I_m) \approx (2\pi\rho_m(U))^{-1} \int_0^{U-B_f} \rho_s(U_s - B_f - \varepsilon)d\varepsilon, \quad (9)$$

$$\Gamma_{\gamma}(E_m^*, I_m) \approx \frac{3}{(\pi \hbar c)^2 \rho_m(U)} \int_0^U \sigma_{\gamma A}(E_{\gamma}) \rho_d(U - E_{\gamma}) E_{\gamma}^2 dE_{\gamma}. \quad (10)$$

where U is the thermal energy of the mother nucleus, s_{ν} is the spin of emitted particle and μ_{ν} is the reduced mass of the system particle ν plus daughter nucleus. The symbols m and d indicate a mother and a daughter nucleus, respectively. The inverse cross section σ_{inv} is calculated within the model [14]:

$$\sigma_{\text{inv}} = \begin{cases} \sigma_g c_1 (1 + c_2/E_{\nu}), & \nu = n, \\ \sigma_g c_3 (1 - c_4 V_{\nu}/E_{\nu}), & \nu = p, d, t, {}^3\text{He}, \alpha. \end{cases} \quad (11)$$

Here σ_g is the geometrical cross section. r, c_1, c_2, c_3, c_4 are the parameters taken from [14]. In expression (9) the thermal energy U_s and rotational energy E_r^s are connected at the saddle point by the relation $U_s = E^* - E_r^s$. This form of the width Γ_f takes into account the change of the fission barrier of the rotating nucleus so far as $B_f(I) = B_f(0) - (E_r - E_r^s)$ (see details in [15]). In expression (10) for the partial width of the electric dipole radiation $\sigma_{\gamma A}$ is the photoabsorption cross section (the dipole electric γ -transitions dominate in the statistical γ -cascade at $U > 1.5 \div 2.0$ MeV; at lower values of U the quadrupole γ -transitions plays the crucial role).

To describe the level density as a function of the excitation energy, the well known expression from [16]

$$\rho(E^*) = \frac{\sqrt{\pi}}{12} \frac{1}{a^{1/4} (E^*)^{5/4}} \exp[S(E^*)], \quad (12)$$

has been used. In (12) the dependence of the nucleus entropy S on the excitation energy E^* is determined by the relation

$$S = 2at \quad (13)$$

by using the connection of the nucleus temperature with its excitation energy:

$$E^* = at^2 \quad (14)$$

The parameter of the level density $a = \pi^2 g_0/6$ is expressed through the density of single particle states near the Fermi energy $g_0 = f(E_f) = \text{const.}$ The decrease of the influence

of shell effects on the level density with increasing excitation energy is taken into account by the phenomenological expression [16]:

$$a(E^*) = \bar{a}[1 + f(E^*)\delta W/E^*]. \quad (15)$$

Here $f(x) = 1 - \exp(-\gamma x)$, δW is the shell correction in the nucleus mass formula, $\bar{a} = A(\alpha + \beta A)$ is the Fermi-gas value of the level density parameter, A is the mass number of nucleus. The empirical values of the parameters $\alpha = 0.134 \text{ MeV}^{-1}$, $\beta = -1.21 \cdot 10^{-4} \text{ MeV}^{-1}$, $\gamma = 6.1 \cdot 10^{-2} \text{ MeV}^{-1}$ have been obtained from the analysis of the data on the level density with taking into account the contribution of the collective states to the total level density:

$$\rho_{\text{tot}}(E^*) = K_{\text{rot}}K_{\text{vib}}\rho(E^*) \quad (16)$$

(see details in [16]).

After the determination of the de-excitation mode (if fission does not occur) the characteristics of the emitted particles or γ -ray, namely their kinetic energy, orbital momentum and emission angle were drawn. For a given kind of the particle the simultaneous selection of E_ν , i (i is the particle orbital momentum) and $\cos(\Theta)$ (Θ is the angle between \mathbf{I} and \mathbf{i}) has been performed by using three random numbers. Then by using the fourth random number they are rejected according to the three-dimensional probability density

$$\rho'(E_\nu, i, \cos(\Theta)) \propto i \exp \left[2\sqrt{a(E_k^* - E_\nu - \hbar^2(I^2 + i^2)/2j + \hbar^2 I i \cos(\Theta)/j)} \right]. \quad (17)$$

Here j is the moment of inertia of the compound nucleus. The azimuthal angle of the evaporated particle is drawn in the coordinate system with the axis z parallel to \mathbf{I} . The fission process is taken into account by the weight factor

$$F = \prod_{\nu=1}^x [1 - \Gamma_f/\Gamma_{\text{tot}}], \quad (18)$$

where Γ_{tot} is the sum of all partial widths and x is the number of steps in the evaporation cascade. This is convenient, in particular, for strongly fissionable nuclei. All the quantities are transformed to the centre-of-mass system of interacting nuclei and the characteristics




Fig.1. Evaporation residue cross sections for the reactions $^{100}\text{Mo} + ^{100}\text{Mo}$ (a) and $^{110}\text{Pd} + ^{110}\text{Pd}$ (b) as functions of $E_{c.m.}$. The results of calculation in the framework of the optical model, surface friction model, macroscopic dynamic model and our model are presented by dotted line, short dashed line, long dashed line and solid line, respectively. The experimental data are presented by solid squares.

of the residual nucleus are calculated. Then the maximum residual energies of all emission processes and fission channel are calculated for this nucleus. Among the allowed values of E_{ν}^{res} and $E_f^{\text{res}} > 0$ the next drawing of the de-excitation type is performed. This is done while the condition $E^{\text{res}} > 0$ is satisfied. The gathering of the required statistics for the calculation of different reaction characteristics has provided about 5% calculation accuracy.

The computation of the compound nucleus de-excitation on the basis of the Monte-Carlo method is performed for all the considered variants of the CNF cross section calculation (see the subsections A, B and D). The ratio of the level density parameters $a_f/a_n = 1$ is used. The calculated evaporation residue cross sections for the reactions $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$ are compared in Fig. 1 with the experimental data [9]. Strong discrepancy is observed between the calculated and experimental results. The discrepancy is particularly large for the reaction $^{110}\text{Pd}+^{110}\text{Pd}$ when the CNF cross section is calculated within the optical model and the surface friction model. The macroscopic dynamic model gives lower values of σ_{CN} , but the discrepancy between the calculated results and the experimental data is several orders of magnitude. The observed discrepancy cannot be explained in the framework of the standard concepts of the fusion of complex nuclei. Therefore, to investigate the reason of strong decrease of the evaporation residue cross sections in these reactions, the DNS approach was used.

III. PECULIARITY OF THE COMPLETE FUSION OF MASSIVE NUCLEI. FUSION BARRIER OF A NEW TYPE

According to the DNS-approach the first stage of the complete fusion of nuclei ends with the formation of the DNS. The DNS evolution is defined by the potential energy of the system $V(Z, I)$ as a function of the charge asymmetry and the angular momentum. The calculated potential energies of the DNS for the reactions $^{100}\text{Mo}+^{100}\text{Mo}$ and $^{110}\text{Pd}+^{110}\text{Pd}$ are presented in Ref. [10]. The liquid-drop mass of nuclei [17] and nucleus-

nucleus potential $V(R) = V_N(R) + V_C(R) + V_r(R)$ have been used to calculate $V(Z, l)$. The calculation has been performed under a simple assumption about the DNS form. The DNS is considered as two spherical overlap nuclei. The distance R between their centres corresponds to the minimum of potential pocket of $V(R)$. The overlap of nuclei in the DNS is small and we can use the frozen density assumption to calculate the nucleus-nucleus potential. This assumption corresponds to the retaining individuality of the DNS nuclei used in the DNS-approach which is used successfully in the description of deep inelastic transfer reactions [24]. The energy scales are normalized to the total energy of the corresponding spherical compound nucleus (see Fig. 1 in Ref. [10]). The isotopic composition of the nuclei forming the DNS is chosen from the condition of N/Z equilibrium in the system.

The nuclear interaction $V_N(R)$ is taken in two variants: “proximity” and “double folding”. According to [18] the expression for $V_N(R)$ in the “proximity” variant looks like:

$$V_N(R) = -2\pi(\gamma_1 + \gamma_2)R s_0 \begin{cases} \frac{5}{3}(1 + s/s_0)\exp[-1.6s/s_0], & s \geq 0, \\ \frac{5}{3} - s/s_0 - \frac{1}{2}(s/s_0)^2, & s < 0, \end{cases} \quad (19)$$

where $\gamma_i = 0.9517(1 - 1.7826(1 - 2Z_i/A_i)^2)$, $s = R - R_{1p} - R_{2p}$, $R_{ip} = 1.17A_i^{1/3}$ fm, $R = R_{1p}R_{2p}/(R_{1p} + R_{2p})$, $s_0 = 1$ fm. Here s is the distance between the surfaces of interacting spherical nuclei. The expression for $V_N(R)$ in the “double folding” form

$$V_N(R) = \int \omega_1(\mathbf{r}_1)\omega_2(\mathbf{R} - \mathbf{r}_2)\mathcal{F}(\mathbf{r}_1 - \mathbf{r}_2)d\mathbf{r}_1d\mathbf{r}_2 \quad (20)$$

is taken from [19]. Here $\omega_i(\mathbf{r}_i)$ are the densities of interacting nuclei, $\mathcal{F}(\mathbf{r}_1 - \mathbf{r}_2)$ is the nucleon-nucleon interaction potential. To take into account the repulsive part of the nucleus-nucleus potential, the density-dependent nucleon-nucleon interaction is used [20].

The final expression has the form [19]:

$$V_N(R) = C_0 \left\{ \frac{F_{in} - F_{ex}}{\omega_{00}} \left(\int \omega_1^2(\mathbf{r})\omega_2(\mathbf{r} - \mathbf{R})d\mathbf{r} + \int \omega_1(\mathbf{r})\omega_2^2(\mathbf{r} - \mathbf{R})d\mathbf{r} \right) + F_{ex} \int \omega_1(\mathbf{r})\omega_2(\mathbf{r} - \mathbf{R})d\mathbf{r} \right\}, \quad (21)$$

$$F_{in,ex} = f_{in,ex} + f'_{in,ex}(N_1 - Z_1)/A_1(N_2 - Z_2)/A_2,$$

where N_i are neutron numbers. The values of the dimensionless parameters f , f' are known from the fit of a great set of the experimental data within the theory of the finite Fermi-systems [20]: $C_0 = 300\text{MeV fm}^3$, $f_{in} = 0.09$, $f_{ex} = -2.59$, $f'_{in} = 0.42$, $f'_{ex} = 0.54$. For massive nuclei the following expression

$$\omega_i(\mathbf{r}) = \frac{\omega_{00}}{1 + \exp((r - R_{i0})/a_0)} \quad (22)$$

can be used with parameters $\omega_{00} = 0.17 \text{ fm}^{-3}$, $R_{i0} = r_0 A_i^{1/3}$. The position and height of the barrier for many reactions are well described by the values of parameters $r_0 = (1.08 \div 1.17) \text{ fm}$ and $a_0 = (0.50 \div 0.55) \text{ fm}$ [19]. The same set of parameters has been used to calculate the DNS potential energies for all the considered reactions. For the reactions $^{100}\text{Mo} + ^{100}\text{Mo}$ and $^{110}\text{Pd} + ^{110}\text{Pd}$ the nucleus-nucleus potentials are presented in Ref. [10]. The double folding form of $V_N(R)$ has been used there.

A partial overlap of the volumes of interacting nuclei [21] is taken into account in the Coulomb potential. A complete sticking takes place for the DNS evolving to the compound nucleus, therefore the centrifugal potential $V_r(R)$ has the form:

$$V_r(R) = \frac{\hbar^2 l(l+1)}{2(j_1 + j_2 + \mu R^2)}, \quad (23)$$

where $j_i = 2m A_i R_i^2 / 5$ are the rigid-body moments of inertia of the DNS nuclei.

The potential energy of the DNS $V(Z, l)$ as a function of charge asymmetry and the nucleus-nucleus potential $V(R)$ as a function of R are presented in Fig. 2 schematically. It is seen that for the reactions $^{100}\text{Mo} + ^{100}\text{Mo}$ and $^{110}\text{Pd} + ^{110}\text{Pd}$ the initial DNS seems to be at the minimum of the potential energy. The DNS is similar to a giant nuclear molecule. As it has been emphasized in [8], the existence of the shell structure gives significant stability for nuclei of DNS.

It is clear from Fig. 2 that on its way to the compound nuclei the DNS has to overcome the potential barrier that is equal to the difference of the potential energy $V(Z, l)$ at the Businaro-Gallone (BG) point and symmetric configuration. This difference can be called the fusion barrier B_{fus}^* . Even at a considerable surplus of the kinetic energy over the

entrance barrier the fusion barrier appears on the way to the compound nucleus. This is a specific feature of the complete fusion of massive nuclei which can be revealed only within the DNS-approach. In further calculations the double folding potential is used as $V_N(R)$. If we replace it by the proximity potential, the values of B_{fus}^* lower and σ_{CN} increases (Fig. 3). Therefore, the total cross section of the fusion of massive nuclei seems to be very sensitive to the potential $V_N(R)$. This fact can be used for a more precise definition of the potential $V_N(R)$.




Fig.2. Schematic presentation of the process of competition between complete fusion and quasi-fission. Potential energy of DNS $V(Z, l)$ as a function of charge asymmetry and nucleus-nucleus potential $V(R)$ as a function of R are presented.




Fig.3. Potential energy of DNS as a function of charge asymmetry at $l = 0$ and $40\hbar$ in the case of reaction $^{110}\text{Pd} + ^{110}\text{Pd}$. The nuclear interaction is taken in proximity (dashed lines) and double folding (solid lines) forms. The energy scales are normalized to the total energy of the corresponding compound nucleus. The sequence $l = 0, 40, 0, 40, 0\hbar$ is assigned to curves from top to bottom.

The physical nature of this fusion barrier drastically differs from the extra-extra push of the macroscopic dynamic model [5-7]. The extra-extra push is an additional kinetic energy over the entrance potential barrier which should provide the compact form of fusing nuclei, i.e. a more compact form than the form of the fissile compound nucleus to be reached at the saddle point. Unlike the extra-extra push, the source of energy for getting over the fusion barrier B_{fus}^* is the DNS excitation energy. Namely, the excitation energy allows the system to realize such an endoergic redistribution of the nucleons between the

DNS nuclei after which the system turns out to be at the top of the fusion barrier. These changes of A , and Z , can be considered as a large fluctuation in the initial DNS. After reaching the fusion barrier the DNS potential energy begins to decrease with increasing charge asymmetry and the driving forces lead the DNS to a compound nucleus.

At the same time if we look at the nucleus-nucleus potential in the entrance channel $V(R)$ (Fig. 2) we can see the capture of the projectile by the target nucleus into a rather shallow potential pocket. The shallow potential pocket and slight overlap of the two massive nuclei, as a result of strong Coulomb repulsion, can lead to the disintegration of the DNS to two fragments having close masses, i.e. the initial DNS can easily undergo quasi-fission. The small values of $\sigma_{ER}(E)$, especially, in the reaction $^{110}\text{Pd}+^{110}\text{Pd}$ indicate the predominance of the quasi-fission channel over the complete fusion one. To determine $\sigma_{ER}(E)$, it is necessary to calculate the competition between the channels of complete fusion and quasi-fission. The authors propose a simple model of this competition in the initial DNS.

IV. MODEL OF THE COMPETITION BETWEEN COMPLETE FUSION AND QUASI-FISSION IN MASSIVE SYMMETRIC DINUCLEAR SYSTEM

In accordance with [8] the fusion process starts after the formation of the DNS. There is competition between the complete fusion and quasi-fission channels in the initial DNS formed by massive nuclei. Therefore, this competition should be taken into account in the calculation of cross section $\sigma_{CN}(E)$. Unfortunately, the existing models don't allow to calculate the competition between the complete fusion and quasi-fission in the DNS.

The thermal equilibrium is established in the DNS rather fast, for several units of 10^{-22} s. As the quasi-fission time is one order of magnitude longer [22] one can try to use the statistical approach to analyse the competition between complete fusion and quasi-fission. The possibility of using the statistical approach to the DNS decay is indicated by the Q_{gg} -systematics of cross sections of deep inelastic transfer reaction products [23].

We assume that the probability for the initial DNS to evolve via the complete fusion or to decay by the quasi-fission channel is determined by the DNS level densities at the maxima of the fusion and quasi fission barriers. A similar approach was used in [24] to describe the charge distribution of the deep inelastic transfer reactions. The fusion barrier B_{fus}^* can be calculated from the DNS potential energy $V(Z, l)$ (see Fig. 2). How to estimate the quasi-fission barriers? Usually quasi-fission in asymmetric nuclear systems is considered [25]. These systems evolve towards a symmetric shape with a subsequent decay into two nuclear fragments with close masses. In the reactions $^{100}\text{Mo} + ^{100}\text{Mo}$ and $^{110}\text{Pd} + ^{110}\text{Pd}$ the initial DNS has a symmetric shape already at the moment of formation. This shape is favourable for the decay because of the maximum value of the Coulomb repulsion. The assumption about the retaining individuality of the DNS nuclei [8], its small overlap and the system position in the minimum potential energy are favourable to use in the analysis of the decay of a massive symmetric DNS in a sudden approximation. In the process of quasi-fission the DNS should overcome the potential barrier (B_{qf}) which coincides with the depth of the pocket of interaction potential $V(R)$ (see Fig. 2).

To describe the DNS level density, one has to use the expression proposed in Ref. [26]

$$\rho_i(E_i^*) = \left[\frac{g^2}{g_1 g_2} \right]^{1/2} \frac{g}{6^{3/4} (2g E_i^*)^{5/4}} \exp[2(a E_i^*)^{1/2}] \quad (24)$$

where i denotes B_{fus}^* or B_{qf} , g_1 and g_2 are densities of single particle states near the Fermi surface for the two nuclei incorporated in DNS, $2g = g_1 + g_2$ and $a = \pi^2 g/3$. The values of g_1 and g_2 are taken according to the systematics [27]. The excitation energy E_i^* is the difference between the DNS excitation energy in the symmetric configuration $E^* = E_{c.m.} - V(R^*)$ and the value of the corresponding barrier. R^* is the R for the bottom of a pocket in the $V(R)$. Taking into account the above assumption we can get for the probability of complete fusion W_{fus}

$$W_{fus} = \frac{\rho_{B_{fus}^*}}{\rho_{B_{fus}^*} + \rho_{B_{qf}}} \quad (25)$$

The ratio $\rho_{B_{qf}}/(\rho_{B_{fus}^*} + \rho_{qf})$ determines the quasi-fission probability. Therefore, in the general case the fusion cross section can be written in the following form

$$\sigma_{CN}(E_{c.m.}) = \pi\lambda_0^2 \sum_{l=0}^{l_f} (2l+1)T(l, E_{c.m.})W_{fus}(l, E_{c.m.}), \quad (26)$$

where l_f is the angular momentum corresponding to the vanishing fission barrier. In our case the last multiplier takes into account the competition between the fusion and quasi-fission-like process. It is supposed that $W_{fus}=1$ in the optical model and in the surface friction model. Usually the fusion of nuclei is considered there when the quasi-fission-like channel is negligible. However, in the case of symmetric combination of the massive colliding nuclei the quasi-fission dominates.

The DNS excitation energy is defined by $E_{c.m.}$ and the DNS potential energy. We used the data of Ref. [22] where it was shown that in contrast to fission in the process of quasi-fission light particles do not carry away a considerable portion of the excitation energy of the system.

In the classical quasi-fission models [25] the initial asymmetric DNS evolves to the symmetric form of which it decays into two fragments with close masses. A possibility of the system decay from asymmetric configuration is ignored. Here we followed this traditional concept about quasi-fission process. At the same time there is a definite probability of the DNS decay from the intermediate configurations on its way to the Businaro-Gallone point. The question is whether this probability is comparable with the quasi-fission probability from symmetric configuration. To estimate the quasi-fission probability from the intermediate configurations, we replaced the δ -functional distribution of Z in the initial DNS by the function proportional to $\exp(-V(Z, l)/t)$ assuming the thermal equilibrium in the DNS. For each Z between the symmetric and BG point configurations (Fig. 2) the quasi-fission probability has been calculated as given above taking into account the weight factors proportional to $\exp(-V(Z, l)/t)$. The DNS configurations close to the symmetric one seem to give the main contribution ($\approx 90\%$) to the quasi-fission. This allows one to use in the first approximation the traditional statistical model for the analysis of the competition between the complete fusion and quasi-fission in the massive symmetric DNS ignoring quasi-fission from the intermediate configurations of DNS.




Fig.4. Compound nucleus formation cross sections for the reaction $^{100}\text{Mo} + ^{100}\text{Mo}$ (a) and $^{110}\text{Pd} + ^{110}\text{Pd}$ (b), as functions of $E_{c.m.}$. The results of calculation in the framework of the optical model, surface friction model, macroscopic dynamic model and our model are presented by dotted line, short dashed line, long dashed line and solid line, respectively.




Fig.5. Probabilities of CNF defined only by the penetration factor of the entrance parabolic barrier are presented by dotted line and dashed line for the reactions $^{40}\text{Ar} + ^{180}\text{Hf}$ and $^{124}\text{Sn} + ^{96}\text{Zr}$, respectively. For the reaction $^{124}\text{Sn} + ^{96}\text{Zr}$ the result of calculation in the framework of the DNS-approach is presented by solid line. Probabilities of CNF obtained from the experimental data are presented by solid squares and solid circles for the reactions $^{40}\text{Ar} + ^{180}\text{Hf}$ and $^{124}\text{Sn} + ^{96}\text{Zr}$, respectively. The abscissa for the ^{40}Ar -induced reaction (upper scale) is shifted by the difference of the Coulomb barriers of both reactions.

The results of $\sigma_{CN}(E)$ calculation are presented in Fig. 4. Our data for $\sigma_{CN}(E)$ calculated by the standard complete fusion models, namely, the optical [2], surface friction [4] and macroscopic dynamical [7] ones, are also given in these figures for comparison. As one can see, the formation of the potential barrier B_{fus}^* on the way of DNS to the compound nucleus and the competition between complete fusion and quasi-fission in the initial DNS lead to a sharp decrease of $\sigma_{CN}(E)$ values in reactions between massive nuclei. The position of B_{fus}^* slightly shifts towards larger asymmetry of the DNS with increasing l . To simplify our calculation, we fix the place of B_{fus}^* in the configuration with ^{48}Ca as a light nucleus.

The proposed model has been used also to analyse the competition between the complete fusion and quasi-fission for almost symmetric massive dinuclear systems. In Ref. [28] the cross sections of the reaction channels (HI,xn) have been measured for two reactions leading to the same compound nucleus ^{220}Th : $^{40}\text{Ar}+^{180}\text{Hf}$ and $^{124}\text{Sn}+^{96}\text{Zr}$. The standard model of complete fusion was able to describe the experimental data for the first reaction. At the same time, there was a large discrepancy between the calculated and experimental data for the second reaction. From our point of view, the reason of this is the quasi-fission which is a predominant reaction channel in the second reaction. Indeed, the probability for the CNF, which has been determined from the experimental data by using some model assumptions [28], seems to be larger for the reaction $^{40}\text{Ar}+^{180}\text{Hf}$ than for the reaction $^{124}\text{Sn}+^{96}\text{Zr}$. The initial DNS formed in the reaction $^{40}\text{Ar}+^{180}\text{Hf}$ corresponds approximately to the BG point configuration. In this case $W_{fus} \approx 1$ in (26) and the values of the probability for the CNF are described rather well (above barrier) by the penetration factor T of the entrance parabolic barrier (Fig. 5). On the contrary, for the almost symmetric system $^{124}\text{Sn}+^{96}\text{Zr}$ the values of W_{fus} , which can be much smaller than unity, decrease the probability for the CNF. A good description of the experimental data in the framework of the DNS-approach is demonstrated for this reaction in Fig. 5. Thus, our model can be applied to almost symmetric systems.

V. CALCULATION OF EVAPORATION RESIDUE CROSS SECTIONS IN THE REACTIONS $^{100}\text{Mo} + ^{100}\text{Mo}$ AND $^{110}\text{Pd} + ^{110}\text{Pd}$

Three factors taken into account in calculating the $\sigma_{ER}(E)$ for the reactions $^{100}\text{Mo} + ^{100}\text{Mo}$ and $^{110}\text{Pd} + ^{110}\text{Pd}$ are as follows: (i) the capture cross section $\sigma_c(E)$, (ii) the competition between complete fusion and quasi-fission in the initial DNS, and (iii) the competition between fission and the emission of light particles and γ -rays in the compound nucleus de-excitation. The capture cross section $\sigma_c(E)$ has been calculated using an optical model [2]. The competition between complete fusion and quasi-fission has been calculated in the framework of the proposed model (Sec. IV). The compound-nucleus de-excitation has been analyzed in the framework of a statistical model by the Monte Carlo method [11,12].

As is shown in Ref. [29], a considerable part of the excitation energy of the massive compound nucleus is carried away by neutrons before the nucleus reaches the scission point. The independence of pre-scission neutron multiplicities on the total kinetic energy of the fission fragments indicates that the neutron emission just before and just after scission is not very important in fusion-fission reactions [22]. The survival of the evaporation residues of heavy nuclei at a high excitation energy [29] allows one to assume that the pre-scission neutron emission takes place mainly before the compound nucleus reaches the saddle point. Therefore, based on the results of Refs. [22,29] we can assume that the excitation energy at the saddle point is about 30–40 MeV. Taking into account the fission of compound nucleus only at $E^* \leq 35$ MeV we obtained a better agreement between the calculated and experimental data. The calculated results seem to be not too different if instead of 35 MeV we take 50 MeV. The fact that in the reaction $^{110}\text{Pd} + ^{110}\text{Pd}$ the intermediate system, on the way to a compound nucleus, emits an α particle [9] has also been taken into account. Due to the dynamic coupling of the DNS modes of motion [19] this emission takes place mainly on the route from the Businaro-Gallone point to a compound nucleus. So, it does not influence the competition between the fusion and quasi-fission.

The probability of the pre-compound α emission is about 10^{-3} [9] in the reaction $^{110}\text{Pd} + ^{110}\text{Pd}$ and has a negligible influence the quasi-fission that is the dominant channel of the initial DNS decay. In the reaction $^{100}\text{Mo} + ^{100}\text{Mo}$ the probability of the pre-compound α emission is considerably smaller than the probability of the compound nucleus formation and we can neglect it.

The results of $\sigma_{ER}(E)$ calculations by the developed model are presented in Fig. 1. The optical model used by us to calculate $\sigma_c(E)$ does not take into account the coupling of various fusion channels at $E_{c.m.}$ near the Coulomb barrier. This leads to the disagreement between the calculated and experimental data at low $E_{c.m.}$. Therefore, our model is applicable to describe the experimental values of $\sigma_{ER}(E)$ at the collision energy exceeding the Coulomb barrier.

The drastic disagreement between the experimental data and the results of calculations by the optical and surface friction models is due to the fact that these models do not take into account quasi-fission process following the massive DNS formation. We presented these calculated results in Figs. 1 and 4 to demonstrate the quasi-fission role in the fusion of massive nuclei.

The macroscopic dynamic model [7] takes into account many nuclear processes that occur in the entrance channel of the reactions. At $E_{c.m.} > B_c + E_{xx}$ the macroscopic dynamic model was expected to be capable of describing $\sigma_{CN}(E)$. However, the $\sigma_{ER}(E)$ value calculated by this model is about three orders of magnitude larger than the experimental one. From our point of view, this large difference is the result of the absence of the competition between complete fusion and quasi-fission in the macroscopic dynamic model. Indeed, at $E_{c.m.} > B_c + E_{xx}$ the CNF takes place and the quasi-fission is absent. At $E_{c.m.} < B_c + E_{xx}$ the quasi-fission is realised but the CNF is not possible. Therefore, the competition between complete fusion and quasi-fission is not taken into account for given $E_{c.m.}$.

According to Ref. [7], at energies below $B_c + E_{xx}$ no compound nucleus can be formed at all. However as one can see from experimental data (Fig. 1), $\sigma_{ER}(E)$ goes smoothly to

energies several dozens of MeV below $B_C + E_{ex}$. In the used variant of the macroscopic dynamical model the fusion barrier fluctuations are not considered. Introduction of these fluctuations [31] allows one to spread $\sigma_{ER}(E)$ to the region $E_{c.m.} < B_C + E_{ex}$. However, the considerable disagreement between the calculated and experimental data remains for a high collision energy. Apparently the reason for the discrepancy lies in the macroscopic approach itself, in which real nuclei possessing shell structure are replaced by drops of homogeneous nuclear liquid.




Fig.6. (a) Dependence of the fusion barrier, B_{fus}^* on the angular momentum for the system $^{100}\text{Mo}+^{100}\text{Mo}$ (long dashed line) and for the system $^{110}\text{Pd}+^{110}\text{Pd}$ (solid line). (b) Dependence of the quasi-fission barrier, B_{qf} on the angular momentum for the system $^{100}\text{Mo}+^{100}\text{Mo}$ (long dashed line) and for the system $^{110}\text{Pd}+^{110}\text{Pd}$ (solid line). Fission barrier for the compound nucleus ^{220}U according [32] is presented by short dashed line.

A satisfactory description of $\sigma_{ER}(E)$ by the model of competition between complete fusion and quasi-fission can be considered as indicative of the realistic interpretation of mechanism of the compound-nucleus formation proposed in the DNS approach [8].

Calculations of the evaporation residue cross sections on the basis of the DNS-approach allow us to understand reasons of the sharp decrease of cross sections in the transition from the reaction $^{100}\text{Mo}+^{100}\text{Mo}$ to $^{110}\text{Pd}+^{110}\text{Pd}$. In these reactions the mass and charge of compound nuclei are different only by 10% but the cross sections differ by several orders of magnitude. The dependence of the fusion barrier and quasi-fission barrier on the angular momentum for both reactions is shown in Figs. 6a and 6b. One can see that the quasi-fission and fusion barriers change in opposite directions: the quasi-fission barriers decrease in the transition from the reaction $^{100}\text{Mo}+^{100}\text{Mo}$ to the reaction $^{110}\text{Pd}+^{110}\text{Pd}$, and on the contrary the fusion barriers considerably increases.

The exchange with valence nucleons between the nuclei of the DNS increases the nuclear attraction [33]. This leads usually to an increase of the capture cross section but the fusion barrier B_{fus}^* is changed slightly. Therefore, all the peculiarities of the fusion process of massive nuclei seem to be conserved.

The deformation of the DNS nuclei under the influence of the Coulomb forces slightly changes B_{fus}^* as well. This influence of the deformation has been estimated. It was supposed that in the symmetric DNS the nuclei look like rotating prolate ellipsoids with collinear located large axis. In both the reactions on the fusion barriers the light nucleus of the DNS was ^{48}Ca while the heavy nuclei were ^{152}Er and ^{172}Hf (if the particle evaporation is neglected). For this configuration the deformation is introduced only for heavy nuclei of the DNS. The surface of the deformed nucleus is described by the expression:

$$R_{id} = fR_1(1 + \beta Y_{20}(\theta, \varphi)), \quad (27)$$

$$f = \left[(1 + \beta \sqrt{\frac{5}{4\pi}})(1 - \beta \sqrt{\frac{5}{16\pi}})^2 \right]^{-1/3},$$

The factor f is introduced for the volume conservation. Calculations of B_{fus}^* were performed with $V_N(R)$ in the ‘‘proximity’’ form. In (19) \bar{R} has been modified according to

[34]. The potential of the Coulomb interaction $V_C(R)$ was calculated following [34], as well. In $V(Z, l)$ the corresponding changes have been done in the components describing the surface and the Coulomb energies. For the symmetric configuration of the DNS the value $\beta = 0.2$ was taken; and for the BG point, $\beta = 0.4$, which is in accordance with the data on the deformation in symmetric and asymmetric fission [35]. By our estimations the deformation of the DNS nuclei increases B_{fus}^* nearly by 2 MeV for the reaction $^{110}\text{Pd}+^{110}\text{Pd}$ and by 1 MeV for the reaction $^{100}\text{Mo}+^{100}\text{Mo}$.

VI. CONCLUSION

In the paper the cross sections of the evaporation residues in the reactions between massive nuclei are analysed in the framework of different models of complete fusion. This analysis allows us to estimate the validity of the CNF mechanism suggested in different models.

The determination of the evaporation residue cross sections includes the calculation of the CNF cross section and the competition between fission and emission of light particles, γ -rays at the compound nucleus de-excitation. The CNF cross sections have been calculated in the framework of the standard models of complete fusion, and the de-excitation of a compound nucleus has been calculated by the Monte-Carlo method.

The obtained results were calculated by the optical model, the model with surface friction and the macroscopic dynamic model seem to be in sharp contradiction with the experimental data. Especially, for the reaction $^{110}\text{Pd}+^{110}\text{Pd}$ the calculated evaporation residue cross sections exceed the experimental value by several orders of magnitude.

The analysis of the complete fusion in the massive symmetric and almost symmetric systems on the basis of the DNS-approach revealed an important specific feature of this process, the appearance of the fusion barrier after the capture on the way of the DNS evolution to a compound nucleus. This barrier is, in principle, different from the extra-extra push energy of the macroscopic dynamical model. As a result, the competition

between the channels of quasi-fission and complete fusion arises and strongly reduces the CNF cross section.

On the basis of the DNS-approach a model of the competition between complete fusion and quasi-fission in massive symmetric DNS has been developed which includes the fusion and quasi-fission barriers as main elements.

The calculations of the evaporation residues cross sections in the reactions $^{100}\text{Mo} + ^{100}\text{Mo}$ and $^{110}\text{Pd} + ^{110}\text{Pd}$ on the basis of this model give satisfactory agreement with the experimental data, which can be considered as an evidence of the validity of the concept of the CNF mechanism developed in the DNS-approach.

ACKNOWLEDGMENTS

The authors (A.N.V. and N.A.K.) are grateful to the International Science Foundation for financial support under Grant N RFJ-000.

- [1] J. Galin, D. Gurreau, U. Lefort and X. Tarrago, *Phys. Rev. C* **9**, 1081 (1974).
- [2] A.S. Iljinov and E.A. Cherepanov, *JINR Communication P7-84-68* (Dubna, 1984); A.S. Iljinov, Yu. Ts. Oganessian and E.A. Cherepanov, *Sov. J. Nucl. Phys.* **36**, 118 (1982).
- [3] D.H.E. Gross, R.C. Nayak and L. Satpathy, *Z. Phys. A* **299**, 63 (1981).
- [4] P. Fröbrich, *Phys. Rep.* **116**, 337 (1984).
- [5] W.J. Swiatecki, *Phys. Scripta* **24**, 113 (1981).
- [6] S. Bjornholm and W.J. Swiatecki, *Nucl. Phys. A* **391**, 471 (1982).
- [7] J.P. Blocki, H. Feldmeier and W.J. Swiatecki, *Nucl. Phys. A* **459**, 145 (1986).
- [8] V.V. Volkov, *Contributed Papers of Nucleus-Nucleus Collision II*, Visby, 1985, v.1 (North-Holland, 1985), p.54; V.V. Volkov, *Izv. AN SSSR ser. fiz.* **50**, 1879 (1986); V.V. Volkov, *Proceedings of Intern. School-Seminar on Heavy Ion Physics*, Dubna, 1986, D7-87-68 (Dubna, 1987), p.528; V.V. Volkov, *Proceedings of the 6th Intern.*

- Conf. on Nuclear Reaction Mechanisms*, Varenna, 1991, edited by E.Gadioli (Ricerca Scientifica ed Educazione Permanente Supplemento n.84, 1991), p.39.
- [9] W. Morawek, T. Ackermann, T. Brohn, H.G. Clerc, U. Gollerthan, E. Hanlet, M. Horz, W. Schwab, B. Voss, K.H. Schmidt, J.J. Gaimard and F.P. Hessberger, GSI Scientific Report p.38 (1988); K.H. Schmidt and W. Morawek, Rep. Prog. Phys. **54**, 949 (1991); W. Morawek, T. Ackermann, T. Brohn, H.G. Clerc, U. Gollerthan, E. Hanlet, M. Horz, W. Schwab, B. Voss, K.H. Schmidt and F.P. Hessberger, Z. Phys. A **341**, 75 (1991).
- [10] N.V. Antonenko, E.A. Cherepanov, A.K. Nasirov, V.P. Permjakov and V.V. Volkov, Phys. Lett. B **319**, 425 (1993).
- [11] A.S. Iljinov, Yu.Ts. Oganessian and E.A. Cherepanov, Sov. J. Nucl. Phys. **33**, 526 (1981).
- [12] E.A. Cherepanov, *Proceedings of Intern. Symp. on In-Beam Nuclear Spectroscopy*, Debrecen, Hungary, 1984, edited by Zs. Dombrad and T.Fenyés (Akademiai Kiado, Budapest, 1984), p.499.
- [13] R. Bass, *Nuclear Reactions with Heavy Ions* (Springer, Berlin, Heidelberg, New York, 1980).
- [14] I. Dostrovsky, Phys. Rev. **111**, 1659 (1958).
- [15] M. Blann and F. Plasil, Phys. Rev. Lett. **29**, 303 (1972).
- [16] A.V. Ignatyuk, *Statistical properties of excited nuclei* (Energoizdat, Moscow, 1983).
- [17] W.D. Myers, *Droplet model of atomic nucleus* (IFI/Plenum Press, New York, 1977).
- [18] R. Schmidt and J. Teichert, *JINR Communication E4-80-527* (Dubna, 1980).
- [19] N.V. Antonenko and R.V. Jolos, Z. Phys. A **339**, 153 (1991).
- [20] A.B. Migdal, *Theory of finite Fermi systems and applications to atomic nuclei* (Nauka, Moscow, 1988).
- [21] J.P. Bondorf, M.I. Sobel and D. Sperber, Phys. Rep. **15**, 83 (1974).
- [22] D.J. Hinde, Nucl. Phys. A **553**, 255c (1993).
- [23] V.V. Volkov, Phys. Rep. **44**, 93 (1978).

- [24] L.G. Moretto and J.S. Sventek, Phys. Lett. B **58**, 26 (1975).
- [25] Ch. Ngo, Progr. Part. Nucl. Phys. **16**, 139 (1986).
- [26] S. Ayik, B. Schürman and W. Nörenberg, Z. Phys. A **277**, 299 (1976).
- [27] E.A. Cherepanov and A.S. Iljinov, Nucleonica **25**, 611 (1980).
- [28] D. Vermeulen, H.-G. Clerc, C.-C. Sahn, K.-H. Schmidt, J.G. Keller, G. Münzenberg and W. Reisdorf, Z. Phys. A **318**, 157 (1984); C.-C. Sahn, H.-G. Clerc, K.-H. Schmidt, W. Reisdorf, P. Armbruster, F.P. Hessberger, J.G. Keller, G. Münzenberg and D. Vermeulen, Z. Phys. A **319**, 113 (1984); C.-C. Sahn, H.-G. Clerc, K.-H. Schmidt, W. Reisdorf, P. Armbruster, F.P. Hessberger, J.G. Keller, G. Münzenberg and D. Vermeulen, Nucl. Phys. A **441**, 316 (1985).
- [29] D.J. Hinde, D. Hilsher, H. Rossner, B. Gebauer, M. Lehmann and M. Wilpert, Phys. Rev. C **45**, 1229 (1992).
- [30] S. Beiersdorf, R.A. Esterlund, M. Knaack, W. Westmeier, P. Patzelt, F.P. Heßberger, V. Ninov and A. Lüttgen, Phys. Lett. B **286**, 225 (1992).
- [31] P. Armbruster, *Proceedings of Intern. School-Seminar on Heavy Ion Physics*, Dubna, 1986, D7-87-68 (Dubna, 1987), p.82.
- [32] A.J. Sierk, Phys. Rev. C **33**, 2039 (1986).
- [33] R.V. Jolos and A.K. Nasirov, Sov. J. Nucl. Phys. **45**, 1298 (1987).
- [34] N. Malhotra and R.K. Gupta, Phys. Rev. C **31**, 1179 (1985).
- [35] R.V. Hasse, Sov. J. Question of Atomic Science and Technology, ser. Nuclear Constant. **1**, 3 (1988).

Received by Publishing Department
on February 27, 1995.

Антоненко Н.В. и др.

E4-95-80

Образование составных ядер в реакциях между массивными ядрами. Барьер слияния

Проанализированы сечения испарительных остатков (σ_{ER}) в реакциях между массивными ядрами в рамках различных моделей полного слияния. Использование оптической модели, модели поверхностного трения и макроскопической динамической модели может привести к результатам, которые на несколько порядков отличаются от экспериментальных. Это вызвано пренебрежением конкуренции между полным слиянием и квазиделением. Предложен возможный механизм формирования составного ядра с образованием двойной ядерной системы в реакциях с тяжелыми ионами и разработан метод расчета этой конкуренции в слиянии симметричных массивных ядер. Метод может быть применен для энергий столкновения выше кулоновского барьера. Значения величин σ_{ER} , полученных в рамках предложенной модели, оказались близкими к экспериментальным данным. Для иллюстрации рассмотрены реакции $^{100}\text{Mo}+^{100}\text{Mo}$, $^{110}\text{Pd}+^{110}\text{Pd}$ и $^{124}\text{Sn}+^{96}\text{Zn}$.

Работа выполнена в Лаборатории теоретической физики им. Н.Н.Боголюбова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1995

Antonenko N.V. et al.

E4-95-80

Compound Nucleus Formation in Reactions Between Massive Nuclei. Fusion Barrier

The evaporation residue cross sections σ_{ER} in reactions between massive nuclei have been analysed within different models of complete fusion. The calculations in the framework of the optical model, the surface friction model and the macroscopic dynamic model can give the results which are by few orders of magnitude different from experimental data. This takes place due to neglect of the competition between complete fusion and quasifission. A possible mechanism of compound nucleus formation in heavy ion-induced reaction has been suggested. A model is proposed for calculation of this competition in a massive symmetric dinuclear system. This model is applied for collision energies above the Coulomb barrier. The σ_{ER} values calculated in the framework of the approach suggested seem to be close to the experimental data. For illustration the reactions $^{100}\text{Mo}+^{100}\text{Mo}$, $^{110}\text{Pd}+^{110}\text{Pd}$ and $^{124}\text{Sn}+^{96}\text{Zn}$ have been considered.

The investigation has been performed at the Bogoliubov Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 1995

Макет Т.Е.Попеко

Подписано в печать 27.03.95
Формат 60×90/16. Офсетная печать. Уч.-изд.листов 2,27
Тираж 370. Заказ 48076. Цена 1362 р.

Издательский отдел Объединенного института ядерных исследований
Дубна Московской области