

Computational Models of Neuromodulation

Jean-Marc Fellous

Brandeis University, Volen Center for Complex Systems, Waltham, MA 02254-9110, U.S.A.

Christiane Linster

Harvard University, Department of Psychology, Cambridge, MA 02138, U.S.A.

Computational modeling of neural substrates provides an excellent theoretical framework for the understanding of the computational roles of neuromodulation. In this review, we illustrate, with a large number of modeling studies, the specific computations performed by neuromodulation in the context of various neural models of invertebrate and vertebrate preparations. We base our characterization of neuromodulations on their computational and functional roles rather than on anatomical or chemical criteria. We review the main framework in which neuromodulation has been studied theoretically (central pattern generation and oscillations, sensory processing, memory and information integration). Finally, we present a detailed mathematical overview of how neuromodulation has been implemented at the single cell and network levels in modeling studies. Overall, neuromodulation is found to increase and control computational complexity.

1 Introduction ---

Organisms, from invertebrates to mammals, exhibit diverse behaviors when coping with their environments. Correspondingly, the nervous systems of these organisms can differ significantly in their organization and cellular components. Despite such cross-species variability, computational models of nervous systems have shown that complex computations can emerge from the interaction of relatively simple circuits of neurons. A typical connectionist model, for example, involves a transfer function computing the output of the neuron given the sum of its inputs and a synaptic learning rule determining how the strength of synaptic connections is updated. With this type of simple model, a variety of behavioral functions have been modeled, providing insights into how complex phenomena, such as perception, memory, and motor control, can be explained in terms of simple neural mechanisms. Simple models, however, often fail to capture important aspects of neural processing such as neuromodulation (Cooper, Bloom, &

Roth, 1991; Harris-Warrick & Marder, 1991; Hasselmo, 1995; Kaczmarek & Levitan, 1987).

In addition to the classic excitatory and inhibitory neurotransmission, such as those mediated by glutamate or GABA, a large number of biophysical processes serve to modify the response of a neuron to a given input signal or to alter the input signals before their arrival. These modulatory effects often involve substances such as acetylcholine (ACh), norepinephrine (NE), histamine, serotonin (5-HT), dopamine (DA), and a variety of neuropeptides. Although these substances are known to act at different types of receptors, originate from different structures, and have different spatial distributions and time courses of action, they have at least one of the following three functional effects: modulation of intrinsic neural properties (such as input-output function, or threshold), modulation of afferent properties (such as strengthening some neural inputs rather than others), or modulation of efferent properties (such as presynaptic modulation of release). At the behavioral level, such modulations can profoundly affect the function of the nervous tissue involved.

Much is now known about the detailed action of neuromodulatory substances and their agonists and antagonists at the level of small circuits, single neurons, single synapses, or single channels. On the other hand, psychopharmacologists have examined the effects of many drugs that affect various neuromodulatory systems on behaviors such as perception, learning and memory, and motor control. Because of the wider use of modeling techniques and growing interest in systems neuroscience, the computational role of neuromodulation in information processing is receiving increased attention in both the modeling and experimental communities. As we will suggest, the study of neuromodulation may help bridge the gap between elementary neural principles and behavior.

Computational models provide a formal framework in which the function of a neuron or a group of neurons can be expressed rigorously. In general, neural dynamics is represented as a set of equations with variables and parameters. Variables are determined by both the level of description of the model (concentration, membrane potential, firing rate, etc.) and the function under study. Parameters are potential neuromodulatory factors. They are diffuse (nonspecific to each neuron) and are assumed to change more slowly than variables, so that keeping them constant (or very slowly varying) will not perturb the function of the network. In this formalism, neuromodulation appears as a means of changing the way the function is achieved, without changing the function itself. However, not all parameters have a biological meaning. Some are abstract place holders used to make up for the lack of knowledge about the details of a particular phenomenon (the learning rate, for example), others ensure the ad hoc goodness of fit ("tuning") of the model to certain experimental data that are not the primary targets of the model ("time constants" of synaptic alpha functions for example). Moreover, not all the parameters that have putative biological correlates have

identified neuromodulatory roles in situ. For example, some chemical time constants might characterize complex biophysical mechanisms that are normally severely constrained (“regulated”), and consequently have no neuromodulatory function. Furthermore, not all neuromodulatory phenomena can be represented by simple parameter changes.

The purpose of this article is to highlight, through a targeted review of the modeling literature, some of the basic computational roles assigned to neuromodulation and present their possible neural implementation. Due to the diversity and ubiquity of neuromodulatory phenomena, we will not provide a comprehensive review of all neuromodulatory systems in terms of their anatomical loci, detailed biochemical pathways, and individual physiological effects. Nor will we attempt to define it; rather, we will review neuromodulation according to the computational framework provided by a chosen set of modeling studies. Our intent is not to be exhaustive. Many models not mentioned here have discussed how specific neuromodulations can be implemented and how they affect particular aspects of the neural system they consider. We include here a selection of studies that have dealt explicitly with neuromodulation and will help readers understand a specific computational role of neuromodulation.

In the first section, we characterize neuromodulation on the basis of its spatial origin (extrinsic or intrinsic), its functional coupling with neural computation (tuning versus regulation), and its time course. We then review in more detail the computational role of neuromodulation in three important classes of models that address issues pertaining to oscillations and synchrony in small and large networks, sensory processing, and memory function. Finally, in the appendix, we give a detailed mathematical account of the way neuromodulation has been implemented in the various modeling frameworks reviewed.

2 Characterizing Neuromodulation

Neuromodulations can be described by their spatial and temporal characteristics, and in the computational framework chosen here, they can also be characterized by their level of coupling with the specific neural computations under consideration.

2.1 Extrinsic and Intrinsic Neuromodulation. A first class of neuromodulatory signals may originate from an area extrinsic to the neural substrate whose computation is under study, so that lesioning the neuromodulatory center does not usually perturb the function itself, but only modifies its quality. The computational functions of such extrinsic neuromodulation are expected to be somewhat global, because they usually influence many functionally different sites simultaneously. A second class consists of neuromodulations that originate in the relevant substrate itself or in a distant site but are controlled locally within the substrate. In such systems, neuromod-

ulation is an integral part of the computation. Cotransmission (Brezina, Orekhova, & Weiss, 1996; Chan-Palay & Palay, 1984; Kupfermann, 1991; Marder, Christie, & Kilman, 1995), presynaptic receptions (Marder, 1996; Starke, Gothert, & Kilbinger, 1989), glial modulation (Hansson & Ronnback, 1994), and volume transmission (Fuxe & Agnati, 1991; Ridet, Rajaofetra, Teilhac, Geffard, & Privat, 1993) are examples of such phenomena. The functions of such intrinsic modulations are more specific to the substrate under consideration (Katz & Frost, 1996).

2.1.1 Extrinsic Neuromodulation. In many models, the origin of the modulation is known but does not depend in general on the computation of the substrate being modulated. Rather, it depends on the parallel activity of functionally distinct systems, extrinsic to the substrate. Such is the case of most neuromodulatory centers releasing specific neuroactive substances that modify the cellular and synaptic properties of their targets. Most of the actions of dopamine (Cooper, 1991) and norepinephrine (van Dongen, 1981) enter in this category. Here, we illustrate this point with a recent model of sequence learning in hippocampal region CA3 showing that computations may crucially depend on the extrinsic modulation by GABAergic and cholinergic inputs from the septum (Wallenstein & Hasselmo, 1997b).

In this large multicompartmental model, CA3 interneurons receive external periodic (4–10 Hz) inhibitory GABAergic signals from the septum (itself not modeled), while pyramidal cell and interneuron excitability is increased by steadily lowering their leak potassium conductance, simulating the cholinergic influences of the septum. In this modulatory regime, interneurons spontaneously fire gamma (30–100 Hz) bursts of action potentials at the theta (4–10 Hz) frequency externally imposed by the septum (Wallenstein & Hasselmo, 1997a). This pattern of firing in turn entrains the pyramidal cell network at theta frequency, yielding an overall network behavior compatible with much *in vivo* and *in vitro* experimental data. The emerging theta-gamma pattern of interneuronal GABAergic activation results in a periodic activation of GABA_B receptors on pyramidal cells: GABA_B activation is greatest at the start of each theta cycle and decreases smoothly until the end of each cycle. Because GABA_B receptors primarily control synaptic activation at intrinsic (CA3 recurrent collaterals) rather than extrinsic (sensory) pyramidal inputs, their net effect is to modify periodically the balance between internal and external information processing. Sensory inputs dominate at early phases of the theta cycle; intrinsic inputs dominate at later phases. This pattern of modulation is shown to be crucial to the computations of CA3 in that it allows for the development of place fields and for the learning and recall of sequence information modeled as a path learned by a rat running on a linear track. Without GABA_B modulation, the network still functions, but it is qualitatively impaired, yielding place fields that do not develop and making significant errors during the recall of learned sequences.

In this example, extrinsic neuromodulation acts as a separate clocking device whose net effect is to improve the nature of information processing in the CA3 region of the hippocampus. Both the timing of the modulatory signal (theta frequency) and its pharmacological consequences (GABA_B receptor activation) are important and generate testable predictions as to what might happen if either is modified. Other models have viewed extrinsic modulation as a signal influencing synaptic mechanisms. Such is the case of the reward signal entering the weight modification rule, between the ventral tegmental area (VTA) and cortex (Montague, Dayan, & Sejnowski, 1996), discussed in the following sections, or the direct change of synaptic efficacy triggered by an external center (Linster & Gervais, 1996; Linster & Masson, 1996; Raymond, Baxter, Buonomano, & Byrne, 1992).

2.1.2 Intrinsic Neuromodulation. In some interesting instances, it is not possible to isolate the neuromodulatory phenomenon from the system it modulates. In such cases, neuromodulation is intrinsic to the network whose computation is under study. Unfortunately, to our knowledge, there are no direct modeling studies of such phenomena. Some experimental evidence for intrinsic neuromodulation is reviewed elsewhere (Katz & Frost 1996). We briefly mention two examples.

In the stomatogastric ganglion (STG) of the lobster, an afferent axon (SNAX1) has been characterized as both a participant in the rhythmicity of the gastric mill network and as a conveyor of modulatory information (Nusbaum, Weimann, Golowasch, & Marder, 1992). SNAX1 receives (inhibitory) synaptic inputs from the STG and is capable of initiating action potentials (intrinsically, within the STG and not near the cell body, a few centimeters away), which generate excitatory postsynaptic potentials (EPSPs) on the STG elements, therefore participating in the generation of the rhythm. However, because SNAX1 is also electrically coupled with key neurons of the central pattern generator, its level of depolarization (whether or not action potentials are present) modulates the activity of the network.

Similarly, in the tritonia, dorsal swim interneuron DSI (a serotonergic central pattern generator (CPG) neuron), is known to enhance synaptic transmission presynaptically at synapses made by a key CPG neuron (Katz & Frost, 1995b; Katz & Frost, 1996; Katz, Getting, & Frost, 1994). DSI elicits both a fast, neurotransmitter-like EPSP, and a slow neuromodulatory-like EPSP (Katz & Frost, 1995a), both pharmacologically separable. DSI therefore modulates the oscillatory pattern it is contributing to.

It is, of course, possible to envision dual extrinsic and intrinsic neuromodulations, whereby the former would express state or stimulus dependency and the latter would be activity dependent. In the computational framework of modeling studies, extrinsic neuromodulations can be easily implemented by choosing appropriate sets of parameters (tuning), whereas intrinsic neuromodulations require that the neuromodulatory mechanisms be regulated by the computations under consideration.

2.2 Regulation and Tuning. Choosing a computational framework to study neuromodulation inherently places it within a larger continuum.

2.2.1 Regulation. At one extreme, when neuromodulation is tightly coupled with neural computations, it becomes regulatory, an integral part of the computations. Such is the case of second messenger systems described in a Markovian kinetics formalism (Destexhe, Mainen, & Sejnowski, 1994b) or of activity-dependent regulation of maximal conductances (LeMasson, Marder, & Abbott, 1993), which we briefly discuss next.

Using a single-compartment model of the lateral pyloric neuron of the stomatogastric ganglion of the crab (Buchholtz, Golowasch, Epstein, & Marder, 1992), LeMasson et al. (1993) elegantly illustrate how neurons can maintain a given firing behavior in the face of perturbations such as changes in extracellular K^+ concentrations or sudden shifts in certain membrane current maximal conductances. This is achieved by making the intrinsic properties of the neuron (maximal conductances) dependent on the intracellular calcium concentration, and hence indirectly on previous activity. This feedback regulation ensures that conductances are stable and that the firing pattern of the cell (silent, bursting, or tonically firing) is preserved. The authors propose that this regulation, because it happens on a relatively slow timescale, could correspond physiologically to calcium regulation of channel synthesis, insertion, or degradation. Interestingly, in this particular model, the same mechanism that regulates the firing pattern in the face of certain perturbations may also change it in the face of other perturbations, such as external patterns of stimulation, therefore increasing the complexity of the input-output relationship of the cell.

2.2.2 Tuning. At the other extreme, when neuromodulation is entirely decoupled from the network under study, its actual implementation becomes a matter of parameter tuning. Such is the case of the choice of particular parameter sets that yield different bursting modes in invertebrate pattern generators (Epstein & Marder, 1990) or different cell frequency adaptation characteristics in piriform cortex (Barkai & Hasselmo, 1994), as we discuss next.

In slices, piriform cortex pyramidal cells can generally be classified into strongly adapting or weakly adapting cells, depending on their response to long constant depolarizing current pulses (Barkai & Hasselmo, 1994). This difference in firing frequency adaptation may influence the computations at hand. Carbachol, a muscarinic receptor agonist, has been found to decrease the spike frequency adaptation of pyramidal cells and, in effect, switches strongly adapting cells into weakly adapting ones. On the basis of the experimental finding that carbachol primarily modulates two membrane potassium currents, IK (AHP) and IK (M) (Madison, Lancaster, & Nicoll, 1987), Barkai and Hasselmo used a compartmental model and found that different values of the maximal conductances of these two cur-

rents, as well as different values for membrane resistance, would reproduce the range of spiking adaptation behaviors identified experimentally. These authors therefore found two distinct parameter tunings that characterize strongly and weakly adapting cells, and they used the weakly adapting tuning to model cells undergoing cholinergic modulation. The computational role of cholinergic neuromodulation is then illustrated in the context of associative memory (Barkai, Bergman, Horwitz, & Hasselmo, 1994) and will be discussed in more details in a later section.

Regardless of whether neuromodulation is extrinsic or intrinsic, it is possible, if not desirable, to build a model that implements the effects of neuromodulation using external parameter tuning first. As more data about the neuromodulatory processes are made available, the model can be modified to include mechanisms that could trigger the parameter changes in an internal manner (e.g., as a function of network activity). Such is the case, for example, of the regulation of acetylcholine modulation by overall network activity (Hasselmo & Schnell, 1994) rather than by parameter set switching (Barkai et al., 1994).

2.3 Time Course of Neuromodulation. Depending on the function implemented, neural computation may follow different time courses, from a few milliseconds to several minutes or hours. For the neuromodulation of such circuits to be relevant to the computation, it must be adapted to its timing.

2.3.1 Fast Computations, Slow Modulation. Most computational studies that feature neuromodulation implement it as a slow and diffuse process, tonically changing some aspects of membrane or synaptic properties. Such is the case for acetylcholine and its steady depolarizing effects on pyramidal cells and interneurons in CA3 (Wallenstein & Hasselmo, 1997a) or for its slow activity-dependent modulation of CA1 between learning and recall modes (Hasselmo & Schnell, 1994). We discuss these models further in a later section.

2.3.2 Slow Computations, Fast Modulation. Some types of slow computations may be influenced by fast modulations. In a series of experiments in monkeys, Schultz, Apicella, and Ljungberg (1993) showed that during learning, VTA dopaminergic neurons (A9–A10) exhibit transient increases in activity lasting less than 150 ms in response to behaviorally relevant signals, such as reward delivery, or conditioned stimulus presentation. Monkeys were tested in several behavioral paradigms. Here, we focus on the delayed response task in which monkeys are taught to memorize one of two spatially distant targets for a variable (2.5–3.5 sec) amount of time and to respond by arm movement at the onset of a trigger signal. Correct responses are rewarded.

During learning, VTA activity increases transiently after the presentation of the target signal and shortly after the delivery of the reward, regardless

of the delay introduced. This firing behavior is in marked contrast to some of their neural targets, such as the striatum or prefrontal cortex, whose neural activity may be tonically increased during the whole delay period (Goldman-Rakic, Lidow, Smiley, & Williams, 1992). Moreover, the increase of VTA activity after reward appears to be present only during learning, and not once the animal has acquired the task. These data suggest that the transient actions of DA after reward delivery may be specifically involved in learning. The precise duration of the postsynaptic effects of the release of dopamine in the prefrontal cortex during such a task is not known, but it might be as short as 100–200 ms (Jay, Glowinski, & Thierry, 1995). Insofar as one considers that the performance of the delayed response task is a slow process (lasting up to 4 sec), a 150-ms phasic involvement of the dopaminergic system appears as a fast modulatory process influencing a slow sequence of neural computations.

In a model of VTA activity, Montague et al. (1996) propose a way in which DA neuron may transiently affect learning. Their model suggests that DA signals $\delta(t)$ carry a composite information about external reward $r(t)$, and internal fluctuations between present $V(t)$ and immediately past $V(t - 1)$ sensory cortical signals. This DA signal is used to modulate the rate of change of the synaptic weights, which link cortical signals $x(t)$ to dopaminergic neurons. Mathematically, this modulation is expressed as a transient change of learning rate, which tends to reduce the amount of excitation forwarded to the dopaminergic neurons, as learning develops, compatible with experimental data (Schultz et al., 1993), and following the general idea of temporal difference learning (Sutton & Barto, 1990):

$$\Delta w_i^\tau = \eta x_i \delta(t) \text{ if } t = \tau. \text{ 0 if not.}$$

with

$$\delta(t) = r(t) + V(t) - V(t - 1)$$

and

$$V_i(t) = x_i w_i^\tau(t), \quad V(t) = \sum_i V_i(t).$$

Interestingly, this model chooses to label weights explicitly with space (i , origin of cortical activity) and time (τ , relative to the start of each trial). In this paradigm, each weight codes for the occurrence of a particular cortical signal, at a particular time within the experiment.

The spatial diffusion of the DA signal is expressed by the fact that the same $\delta(t)$ affects all synaptic weights equally (it is not indexed by i) and by the fact that it is built on the basis of the sum of all cortical inputs, rather than specialized cortical inputs only (V rather than V_i).

During the initial stages of learning, when weights are uniformly distributed, DA activity closely follows the temporal patterns of reward. During learning, if the time of reward is fixed (such as in the instructed spatial task; Schultz et al., 1993), the weights that code for the particular time (τ_r) of the reward will be strengthened, so that $\delta(\tau_r)$ eventually vanishes. If the

time of reward is variable, as in a delayed spatial task, the activity of DA neurons will become small, but nonzero, around the mean of the times when reward was delivered. After learning, in both cases, DA activity becomes particularly significant when the initial target sensory cue is presented. DA neurons therefore learn to respond to the target sensory cue predictive of the reward rather than to the reward itself.

In this model, the timing characteristics of the modulation are crucial. Its short duration is directly related to the precision with which the prediction of reward is made. Phasic modulation is also important in other modeling studied involving GABA_B receptors (Wallenstein & Hasselmo, 1997b) and in other experimental systems involving norepinephrine and locus coeruleus response to attentional signals and novelty (Aston-Jones, Rajkowski, Kubiak, & Alexinsky, 1994; Rajkowski, Kubiak, & Aston-Jones, 1994; Sara, Vankov, & Herve, 1994).

3 Computational Aspects of Neuromodulation

3.1 Modulation of Oscillation and Synchrony. Neural computation is dynamic and modular and requires that functionally distinct structures communicate in a coordinated fashion. Experimental and theoretical evidence suggests that the generation and synchronization of oscillatory activity may be used to this effect (Gray, 1994). Invertebrate studies have been crucial in furthering our understanding of how both intrinsic membrane properties and synaptic interactions may contribute to the creation and modulation of rhythmic firing (Calabrese & De Schutter, 1992; Harris-Warrick & Marder, 1991). Vertebrate studies of the cortex have built on these results and have proposed ways in which oscillations may synchronize across functionally distinct structures (Gray, 1994). In this context, the neuromodulation of the generation and synchronization of oscillations is bound to play an important computational role.

3.1.1 Central Pattern Generators: Creating and Modulating Rhythmicity. A long tradition of experimental work in invertebrates has led to a detailed knowledge of the effects of various substances on the behavior of individual neurons and small networks of neurons (see for reviews, Calabrese & De Schutter, 1992; Marder, 1996; Marder & Selverston, 1992). Most of these effects can be modeled by changes in the maximal conductance of one or more membrane currents. In these systems, attention is given to neurons whose putative function is to provide, through their rhythmic firing, timing signals necessary for one or several rhythmic motor behaviors (Pearson, 1993), such as chewing in the crustaceans or hormone release during egg-laying behavior in *Aplysia*. These cells are often referred to as conditional bursters because of their ability to fire rhythmically, either intrinsically or under the influence of a small network of connected cells. Two examples can be found in the pyloric network of the crustacean stomatogastric ganglion and in the *Aplysia* bursting neuron R15.

In the STG of the lobster, various modulatory substances such as dopamine, pilocarpine, serotonin, or proctolin can elicit rhythmic burst firing. The mechanisms involved, even though they result in similar bursting behaviors, are by no means simple and depend on the particular substance applied. For example, tetrodotoxin (TTX) may block the effects of serotonin and octopamine but have no effects on the bursting evoked by dopamine and pilocarpine. One possibility is that each of these neuromodulatory effects is mediated by a particular change in the mix of membrane conductances of the cells (Harris-Warrick & Flamm, 1987), which may be studied theoretically using the Hodgkin-Huxley (HH) formalism (Hodgkin & Huxley, 1952; Rinzel & Lee, 1987).

Epstein and Marder (1990) provide a model for the conditional bursting of the anterior burster (AB) neuron of the lobster STG and investigate the effects of the change of a selected set of maximal conductance on the oscillatory properties of the model. They are able to show that two different mixes of fast sodium, leakage, and voltage-dependent calcium maximal conductances were able to model the bursting behaviors of the AB neuron under various neuromodulatory conditions and show why TTX has a different effect on two of these oscillatory modes. Kepler, Marder, and Abbott (1990) showed that, in addition to being intrinsically modulated, the frequency of the modeled AB cell might also depend on the state of follower neurons, provided that both neurons are coupled via gap junctions. Unfortunately, the effects of isolated membrane conductances are often not accessible experimentally. To study the putative effect of pharmacological agents (expressed as continuous maximal conductance changes) on the oscillatory properties of this cell, researchers may then use different modeling techniques, such as exhaustive parameter searches (Bhalla & Bower, 1993; Foster, Ungar, & Schwaber, 1993) or dynamical systems theory (Guckenheimer, Gueron, & Harris-Warrick, 1993; Guckenheimer, Harris-Warrick, Peck, & Willms, 1997). Further experimental and theoretical studies focused on other STG neurons (Golowasch, Buchholtz, Epstein, & Marder, 1992). These models essentially consider neuromodulation to be extrinsic to the oscillatory circuit, and implement it using parameter tuning. Interestingly, further work has attempted to show how maximal conductances may also be changed by intrinsic phenomena. LeMasson et al. (1993), for example, show how intracellular calcium concentrations can be used to implement the activity-dependent modulation of certain maximal conductances (Turigiano, Abbott, & Marder, 1994). Their model shows that depending on the nature of the perturbations imposed onto the cells, this modulation can be regulatory (maintaining the behavior of the cells when extracellular $[K^+]$ is modified) or truly modulatory, by enriching the behavioral repertoire of the cell in response to external patterns of stimulations.

Most modeling studies of the extrinsic effects of neuromodulatory substances have addressed the problem at the level of maximal conductances by tuning them to different values. Very few have actually modeled the

explicit effect of these substances on the conductances (Brezina et al., 1996; Butera, Clark, Canavier, Baxter, & Byrne, 1995). A different line of research in *Aplysia*, however, has achieved this.

Burster neuron R15 in *Aplysia* has been studied in much detail, and its electrophysiological and biochemical properties have been investigated intensively (reviewed in Adams & Benson, 1985; Lechner, Baxter, Clark, & Byrne, 1996). Numerous mathematical models have been developed to explain the cellular basis of single-cell oscillatory activity and bursting. Some models went further and studied the extrinsic modulation of oscillatory dynamics by substances such as DA and 5-HT (Bertram, 1993, 1994; Butera et al., 1995), while others focused on the role of intrinsic modulation by calcium-dependent processes in conditioning (Gingrich & Byrne, 1987; Raymond et al., 1992).

A first series of studies used a simplified HH framework to model the effect of 5-HT as modifications in the conductance of a subthreshold K^+ current (Bertram, 1993, 1994). As for the models of the AB neuron, these models show that changes in maximal conductance can modify the firing properties of R15 from silent to bursting and beating and that the sensitivity of the cell to synaptic inputs is increased. In a separate study Butera et al. (1995) show that even though the apparent effects of DA and 5-HT on the firing properties of R15 are similar, its subsequent responses to depolarizing inputs differ. Effects of 5-HT and DA were implemented as a change in the conductance of two opposing currents: an anomalous delayed rectifier current and a slow inward Ca^{2+} current. Unlike the models mentioned above, this change is directly related to the concentration of extrinsic neuromodulators (see the appendix). Their dynamics are such that both 5-HT and DA can hyperpolarize the cell into silence. However, the subsequent response to a brief depolarizing current pulse elicits a burst of spikes if the cell was silenced with 5-HT and occasional single spikes if it was inhibited by DA, as observed experimentally. Because they make the concentration of these neuromodulatory substances explicit, the authors are able to show that although the effects of 5-HT and DA can be modeled as changes in maximal conductances, they cannot be understood without taking into account the indirect effects of other currents and second messenger systems (such as Ca^{2+} or cAMP). In turn, these indirect effects lead to further modeling that shows their functional importance.

Indeed, the roles of intracellular cAMP and Ca^{2+} are known to be important in activity-dependent neuromodulation in the context of associative classical conditioning in *aplysia* (reviewed in Abrams & Kandel, 1988; Byrne, 1987). In a study using detailed representations of membrane parameters, Gingrich and Byrne have shown that intrinsic regulation of cAMP by Ca^{2+} in an *aplysia* single sensory neuron can simulate the neural analogues of nonassociative learning and classical conditioning (Gingrich & Byrne, 1987). A subsequent study showed that a circuit of six neuron-like elements (including central pattern generators), some of which have synapses mod-

ifiable according to an activity-dependent neuromodulation learning rule, can account for simple features of operant conditioning as well (Raymond et al., 1992).

3.1.2 Modulation of Rhythmicity in the Cortex: Toward Information Processing. Current research in the vertebrate cortex has indicated the functional importance of oscillation and synchronization (Gray, 1994; Singer, 1993). Experimental and theoretical evidence suggest their role in odor coding in the olfactory bulb, in feature integration in the visual cortex, in synaptic plasticity in hippocampus, in attentive behaviors in somatomotor cortex (Gray, 1994), and in the gating of sensory information during awake and sleep states in the thalamocortical circuit (McCormick, 1992). Unfortunately, the computational role of neuromodulation in the generation and synchronization of these rhythms has rarely been studied from a modeling point of view. However, an interesting line of research in the thalamocortical loop is setting the stage for modeling work in other systems.

In the past decade, tremendous breakthroughs have been achieved in the understanding of synchronized oscillations in the thalamocortical circuit (see de Carvalho, 1994; McCormick, 1992, for reviews). Their neuromodulation has been studied *in vitro* and *in vivo*, and their cellular mechanisms explored both experimentally and theoretically through computer simulations. The functional significance of the neuromodulation of this system is summarized next. In slow-wave sleep, with low cholinergic, serotonergic, noradrenergic, and histaminergic modulation, the thalamocortical system presents slow, spontaneous basal intrinsic and circuit oscillations (delta waves and spindle waves). During this state, cholinergic inhibition of thalamic interneurons is absent, resulting in massive inhibition of incoming sensory information, which is consequently only poorly transmitted to the cortex. The increase of cholinergic activation (but decrease of noradrenergic, serotonergic, and histaminergic activation) characteristic of rapid eye movement (REM) sleep results in an abolition of oscillatory activity and an increase of endogenous (without sensory inputs) phasic activity (pontogeniculate-occipital [PGO] waves), thought to be at the origin of the pseudosensorial perceptions experienced during dream states. Finally, the tonic activation of all neuromodulatory systems (including cholinergic, noradrenergic, serotonergic, and histaminergic) results in complex patterns of activity and sets the stage for awake attentive cognitive processing. The precise nature of the sensory processing in the thalamus and its modulation by neuromodulatory centers are limited by the lack of understanding of the nature of the sensory codes themselves. However, understanding how oscillations are generated and how they propagate in a synchronized manner across the thalamic networks might help shed some light on the computations achieved by this structure.

A line of experimental and theoretical work shows that the behavioral-dependent rhythmic firing patterns of thalamocortical (TC) relay cells de-

pend on only a small number of membrane currents (McCormick & Huguenard, 1992) and a functionally intact group of inhibitory thalamic reticular (RE) cells. RE cells are capable of oscillating on their own *in vivo*, and a crucial role for their neuromodulations by NE or 5-HT has been proposed on experimental (McCormick & Wang, 1991) and theoretical (Destexhe, Contreras, Sejnowski, & Steriade, 1994a) grounds. By deactivating a potassium leak current, this extrinsic neuromodulation is able to depolarize RE cells so that GABAergic inhibitory postsynaptic potentials (IPSPs) received from other RE cells deactivate the low-threshold Ca membrane current I_T . This current triggers a rebound burst at the single-cell level, which generates network oscillations in the frequency range of spindle waves (Destexhe et al., 1994a). Through their influence on intracellular levels of G-protein (a second messenger), NE or 5-HT has the potential of switching a silent network of RE cells between quiescent and oscillatory states.

Interestingly, the inclusion of TC cells in this network has prompted the study of a form of intrinsic activity-dependent neuromodulation (Destexhe, Bal, McCormick, & Sejnowski, 1996). In a model of synchronized oscillations and propagating waves in thalamic slices Destexhe et al. (1996) show how the activity-dependent modulation (which they term *upregulation*) of a mixed cationic current I_h in TC cells contributes to the waning phase of the characteristic waning and waxing pattern of spindle oscillations. Whereas neuromodulation is often expressed as a change in maximal conductances, previous work on the STG has indicated how serotonin-mediated shifts in the voltage dependence of the activation curve of I_h could also contribute to the pattern of oscillations of an intrinsically oscillating cell (Golowasch et al., 1992; Harris-Warrick, Coniglio, Levini, Gueron, & Guckenheimer, 1995). In the STG model, shifts were artificially introduced and their effects studied. In this model, however, a different formalism is proposed and introduces an activity-dependent shift of the activation of the I_h current:

$$I_h = \bar{G}([O] + K[O_L])(V - E_{rev}) \text{ with}$$

$$C \xrightleftharpoons{\alpha(V)/\beta(V)} O$$

$$P_u + 2Ca^{2+} \longleftrightarrow P_b$$

$$O + P_b \longleftrightarrow O_L,$$

where C , O , and O_L are closed and opened forms of the h channel and P_u and P_b are unbound and bound forms of a slow intracellular regulating factor, which could be cAMP. The kinetics are such that the transition from O_L to a closed state is very improbable, leading effectively to a locking of the O_L fraction of the channels into the open state. This effect is responsible for a bounded shift of the activation curve of I_h toward depolarized values, as the intracellular calcium concentration is increased during bursting activity. Moreover, because K is chosen greater than 1, the binding of calcium

also triggers an increase in conductance. Both effects have been observed experimentally (Hagiwara & Irisawa, 1989).

Because of the dependence of I_h kinetics on Ca, deactivation of I_h occurs only during low-frequency firing when Ca does not accumulate. During bursts, the accumulation of calcium shifts the activation curve of I_h toward more depolarized states and keeps I_h active. During a burst, therefore, O_L (and consequently I_h) increases, leading to a progressive afterdepolarization (ADP). The ADP eventually counteracts the I_T -mediated rebound bursts, and the spindle oscillatory episodes are terminated. The subsequent slow return of I_h to its basal value results in an 8–10 sec refractory period during which further oscillations cannot be initiated. Evoked or spontaneous activity may ultimately restart the spindle episode, after a total waning phase of 15–25 sec, including the refractory period.

In addition to contributing to the waning phase of spindle oscillation, the modulation of I_h also enables the synchronization of several independent colliding spindle waves into a single propagating wave (but see Contreras, Destexhe, Sejnowski, & Steriade, 1997, for *in vivo* data). Other forms of I_h modulations have been proposed elsewhere in the thalamus (Wallenstein, 1996) and in the STG of the lobster (Golowasch et al., 1992; Harris-Warrick et al., 1995).

In the piriform cortex and olfactory bulb, oscillatory dynamics are modulated by noradrenergic and cholinergic afferents (Biedenbach, 1966; Bressler & Freeman, 1980). Liljenstroem and Hasselmo (1995) investigate the effects of cholinergic modulation on piriform cortex oscillatory dynamics. These include cholinergic suppression of neuronal adaptation, cholinergic suppression of intrinsic fiber synaptic transmission, and cholinergic enhancement of interneuron activity. Their model provides a basis for understanding the involvement of acetylcholine modulation in cortical EEG oscillations (Wilson & Bower, 1992). They demonstrate that the suppression of neuronal adaptation could explain the appearance of evoked gamma oscillations after potentials. They also find that such suppression of adaptation, when coupled with the other cholinergic effects mentioned above, was particularly effective in switching the network into spontaneous theta oscillations. These results are related to others in the hippocampus (Traub, Miles, & Buzsaki, 1992; Traub, Whittington, Colling, Buzsaki, & Jefferys, 1996; Wang & Buzsaki, 1996) that do not involve neuromodulation explicitly. The putative functional significance of neuronal adaptation, and its consequence on rhythmicity, is made apparent in later studies on learning and memory in the hippocampus and will be discussed separately.

3.2 Modulation of the Processing of Sensory Signals: Filtering and Signal-to-Noise Ratio. Processing of sensory information often relies on preprocessing functions like filtering, contrast enhancement, and noise reduction. Many of these functions can be modulated, enabling the sensory

system to respond differently to various components of complex incoming sensory streams.

In the visual domain, one example of such a function is the temporal transformation that some lateral geniculate nucleus (LGN) cells perform on their retinal input (Mukherjee & Kaplan, 1995). The experimental data show that the temporal response of these cells is variable and is related to their ability to burst. Such cells can behave as either relays, responding at the same frequency as their retinal inputs by firing tonically (in alert/awake state), or as bandpass filters, responding optimally at frequencies of 2–8 Hz by firing in a bursting mode (in sleep states), presumably failing to transmit sensory information accurately. In a biophysical model, Mukherjee and Kaplan (1995) show that LGN cell responses can vary from low-pass, with no apparent bursting properties, to bandpass, with frequent burst discharges, depending on the value of their resting membrane potential, and provided that the low-threshold calcium T current is kept active. The authors propose that the LGN acts as a temporal filter, which can be dynamically tuned by attentional signals from the brainstem and the visual cortex, through their modulatory effects on LGN cells' resting membrane potential. In a separate connectionist model, Jackson, Marrocco, and Posner (1994) model such modulatory signals by the putative effects of NE release, expressed as a combination of self-feedback excitation and lateral inhibition. The computational role of such modulation is to achieve contrast enhancement, such that small initial differences in the incoming signal are amplified, and consequently direct attention.

In their model of the olfactory bulb, Linster and Gervais (1996) showed that the modulation of two families of interneurons might sensitively improve odorant signal detection. On the one hand, the modulation (increase) of lateral inhibition mediated by the periglomerular interneurons may result in the sparsification of the mitral activation patterns of complex odors, which otherwise would involve a large, undifferentiated population of mitral cells. On the other hand, under conditions when mitral cell responses are close to noise levels, a global modulation (decrease) of the inhibition mediated by glomerular interneurons may result in an enhancement of their responses. In an extension of this model, Linster and Hasselmo (1997) show that such modulation of inhibition could depend on the global activity of the mitral cells. They introduce a modulator neuron (a putative NE or ACh cell) that receives inputs from all mitral cells and that feeds back on periglomerular cells while simultaneously modulating the connection strength between granule cells and mitral cells. The modulation of inhibition in the glomerular layer ensures a constant average number of active mitral cells, irrespective of the complexity of the input patterns, while modulation of granule cells inhibition ensures a constant average mitral cells spiking probability. Together, these modulations decrease the overlap between pairs of output patterns, making discrimination between overlapping input patterns easier and more reliable.

Addressing similar questions in a model of olfactory processing in the honeybee, Linster and Masson (1996) showed that modulation of inhibition in the antennal lobe may serve for feature extraction of complex and fluctuating chemical signals. This modulation is expressed through the synaptic strength of inhibitory synapses, the biological basis of which has yet to be investigated experimentally. Changes of the balance between excitation and inhibition during the presentation of a stimulus allow the network to act as a short-term memory, displaying the neural activity patterns elicited by the stimulus even after its offset, compatible with experimental data (Sun, Fonta, & Masson, 1993). Expanding on this idea, Linster and Smith (1997) constructed a model of reinforcement learning in the honeybee olfactory system. In this model, modulation of lateral inhibition is introduced via an external modulatory neuron that receives reinforcement signals. This neuron makes plastic synapses onto the circuit under consideration. The authors show that such extrinsic modulation accounts for various behavioral phenomena, such as blocking, unblocking, and overshadowing.

Sensory processing may also involve computations aimed at separating a sensory signal from the background noise. When seen at a system level, the modulation of the signal-to-noise ratio appears as a powerful computational tool by selectively enhancing a signal in a specific pathway, while leaving it undifferentiated with noise in others. A line of modeling work has shown that the cellular mechanisms involved in the known effects of catecholamines on signal detection performance (Clark, Geffen, & Geffen, 1987a, 1987b) may be modeled by a modulation of the slope (gain) of the sigmoid function of a network of leaky-integrator neurons (Servan-Schreiber, Printz, & Cohen, 1990). Changes of this gain at the level of an individual neuron do not affect its signal-detecting capabilities, while increases of this gain in a feedforward chain of neurons augment the signal-to-noise ratio of the whole chain. The model accounts for experimental observations pertaining to the cellular effects of norepinephrine, which show that NE-mediated blockade of Iahp may result in the selective diminution of weak EPSPs and the increase of the depolarization associated with trains of EPSPs, thereby increasing signal-to-noise ratio (Madison & Nicoll, 1986). The model is then used in a backpropagation network to model the improvement in signal detection measured experimentally in human subjects performing a continuous performance task. In this task, subjects are submitted to pharmacological challenges that release catecholamines from synaptic terminals or prevent their reuptake. In an extension of this model, Cohen and Servan-Schreiber (1992) simulate several schizophrenic deficits in selective attention and language processing assessed by tasks such as the Stroop task, the continuous performance test, and a lexical disambiguation task. They successfully show that even though these tasks are seemingly different, the deficits exhibited by schizophrenics can be understood as a general disturbance of the internal representation of contextual information. Such disturbances are implemented as a decrease in the gain of the sigmoid func-

tion of modeled prefrontal cortex units, simulating the possible functional effects of the loss of dopaminergic modulation observed in schizophrenic patients. This theoretical work has been followed by experimental work that confirmed and refined the hypothesis advanced (Cohen, Braver, & O'Reilly, 1996).

In a separate experimental and theoretical study in piriform cortex, Hasselmo and coworkers show that noradrenergic enhancement of the signal-to-noise ratio may also be due to a modulation of synaptic transmission rather than a modulation of input-output function, as was first proposed by Servan-Schreiber et al. (1990). They found that NE, like ACh, may suppress excitatory neurotransmission at intrinsic (collateral) fibers and may also depress feedback inhibition. In a model of piriform cortex, they show that these two effects can act synergistically to increase signal-to-noise ratio (Hasselmo, Linster, Patil, Ma, & Cekic, 1997).

Finally, another interesting body of research has pointed to the role of noise itself as a means of modifying the signal-to-noise ratio (Bulsara, Jacobs, Zhou, Moss, & Kiss, 1991; Levin & Miller, 1996; Longtin, 1993; Longtin, Bulsara, Pierson, & Moss, 1994; McNamara & Wiesenfeld, 1989). To our knowledge, no explicit links to neuromodulation have yet been made.

3.3 Modulation of Memory Function. A large class of memory models is based on the assumption that memories are stored as patterns of synaptic strengths mediating the spread of activation in a network. Learning is achieved according to a synaptic modification rule (or equation) that relates synaptic strength and presynaptic and postsynaptic activities (Brown, Kairiss, & Keenan, 1990; Hasselmo, 1995; Zador, Koch, & Brown, 1990). In this framework, memory function is defined by the synaptic learning rule and the dynamics of individual neurons.

3.3.1 Modulation of the Synaptic Learning Rule. In their model of the response of dopamine neurons to reward and conditioned stimuli (Schultz et al., 1993), Montague, Dayan, and Sejnowski (1996) propose a learning rule in which the postsynaptic activity is augmented by an external reward signal of neuromodulatory origin. In addition, plasticity is made sensitive to temporal differences (Sutton & Barto, 1990) in the postsynaptic activity, rather than to the postsynaptic activity itself. This formulation of the Hebbian learning rule makes time explicit in that some synapses represent early events and others represent later ones. The authors show that after learning a delayed matching-to-sample task, dopaminergic neurons act as a temporal predictor of reward, compatible with experimental data. In this context, dopamine centers have the role of computing and sending diffuse modulatory error signals to the cortex, and hence influence its computation of action in the time domain. The same approach has been used elsewhere (Montague, Dayan, Person, & Sejnowski, 1995) to show how an identified interneuron in the honeybee brain, VUMmx1, could predict reward values

of spatial location during foraging. In this model, VUMmx1 cells influence flight in a manner that accounts for the previous learning of the landscape and its rewarding regions. A similar implementation of VUMmx1 modulation can be found elsewhere (Linster & Smith, 1997).

One of the problems with most learning-rule-based neural models of memory function is the fact that learning and recall may interfere in undesirable ways. Unless care is taken to prevent this, the presentation of a new pattern during learning may elicit an erroneous response from the network. This spurious activity perturbs (if not prevents) learning. In a series of experimental and theoretical studies, Hasselmo and coworkers have shown how selective cholinergic modulation of some synapses, but not others, might provide an elegant solution to this problem.

Experimental data from field recordings in the piriform cortex suggest that cholinergic, noradrenergic, and GABAergic modulation might selectively suppress intrinsic but not afferent excitatory synaptic transmission in the piriform cortex (Hasselmo & Bower, 1992; Hasselmo et al., 1997; Tang & Hasselmo, 1994). In a mathematical model of associative memory, Hasselmo (1993) shows that this selective suppression may prevent previously learned patterns from interfering with the storage of new patterns, especially when previous and new patterns are coded by overlapping populations of neurons (Hasselmo, 1993). This modulation is expressed as a decrease in glutamate release in the activation rule, coupled with a rescaling of the learning rate in the learning rule. In further experimental and theoretical studies, Barkai and Hasselmo (1994) present a detailed biophysical model of a single pyramidal cell in piriform cortex. They show that in addition to its effects on synaptic transmission observed with field potentials, intracellular recordings show that cholinergic modulation of single cells also results in the suppression of neuronal adaptation and in marked depolarization from resting potential. Their single-cell model shows these effects as changes in the maximal conductance of two potassium currents. These results lead to a detailed model of autoassociative memory in the piriform cortex, including 240 pyramidal cells as well as feedforward and feedback interneurons (Barkai et al., 1994). Results from intracellular recordings (suppression of neuronal adaptation and depolarization) and field recordings (suppression of intrinsic synaptic transmission) are included in the model. During learning, the overall effects of cholinergic modulation are to enhance pyramidal cell activity, increasing learning performance. After learning, cholinergic modulation is suppressed and sets the stage for recall. ACh therefore ensures that learning and recall do not interfere and controls the computations of the network.

3.3.2 Modulation of Neural Dynamics. In a large associative network of Fitzhugh-Nagumo-like cells, Abbott (1990) shows that a simple modulation (of putative neuromodulatory origin) of the dynamics of the slow variable (see the appendix) may switch the network from implementing a nonselective short-term latching memory to behaving as a long-term associative

memory. This change in mode of operation of the network increases its computational capabilities without changing its learning rule or architecture. Repetitive firing can also be the result of intrinsic cellular properties such as cholinergically or serotonergically induced afterdepolarization. Models of associative memory based on this phenomenon have shown that repetitive firing can be temporally organized into nested theta and gamma oscillations in order to learn and maintain several memory items active in a short-term memory buffer (Jensen, Idiart, & Lisman, 1996; Lisman & Idiart, 1995).

Building on their work in the piriform cortex, Hasselmo and Schnell (1994) show that the dynamics of learning and recall in the hippocampus can also be regulated by overall network activity. In their model of hippocampal layers CA1 and CA3, the total activity of CA1 pyramidal cells feeds back to the cholinergic system (presumably in the septum) and regulates cholinergic neuromodulation. This model involves a closed and autonomous system that has a clear function and in which neuromodulation is regulated by its target. The septum modulates the function of the hippocampus, which in return regulates the septum in a diffuse, activity-dependent manner. These ideas have been incorporated in a model of corticohippocampal classical eye-blink conditioning (Gluck & Myers, 1993) as a septally driven modification of the learning rate of the hippocampus autoassociative module (Myers et al., 1996). In this model, septal neuromodulation controls the relative amount of time spent by the hippocampus in learning new stimuli and the time necessary to transfer information to neocortical regions.

Finally, in a model of hierarchical associative memory, Cartling (1996) shows that different levels of coupling between activity and excitability may change the dynamics of memory recall. In a Hopfield-like architecture, activity may be chaotic (memories fail to be retrieved), oscillatory (memories are retrieved cyclically, one after the other), or tonic (only one memory item is eventually retrieved) as the coupling is decreased. Neuromodulation is implemented as a change in the shape of the sigmoid transfer function linking membrane potential to firing rate. This change is regulated by overall network activity and depends on intracellular calcium concentrations. However, while some experimental and theoretical work shows that a decrease of cholinergic modulation is associated with stable network dynamics (Hasselmo & Schnell, 1994), this model assumes that an increase of cholinergic modulation yields stable states.

3.4 Neuromodulation for Input Selection and Information Integration. In complex neural networks, information flows along many divergent routes. Much experimental and theoretical work has assigned to neuromodulation the role of selecting the input to particular neural systems, thereby controlling the flow of information. Neuromodulation can act as a routing mechanism and control whether synaptic inputs will activate a particular circuit. The general flow of information between functionally distinct circuits is therefore determined by their modulatory state. Neuromodulation

can also act within a circuit to control what subsets of the available information will be processed.

At the single-cell level, the combined actions of different neuromodulatory systems on cellular or synaptic mechanisms may determine whether the cell will be responsive to a given pattern of synaptic stimulation, therefore enabling or disabling processing.

In *Aplysia*, for example, while both DA and 5-HT silence the bursting neuron R15, only the serotonergic modulation will allow brief depolarization to elicit a sustained bursting response. A modeling study of this system has proposed that the underlying mechanism is rooted in the modulation by DA and 5-HT of two distinct currents (Butera et al., 1995). The authors show that DA prevents input signals from eliciting R15 firing, while 5-HT enhances its response, effectively amplifying synaptic inputs. Together these two neuromodulatory systems control when input signals to R15 may be forwarded to later processing stages.

Similarly, at the network level in the vertebrate, experimental and theoretical evidence suggest that ACh levels, together with other neuromodulatory systems, may control the flow of sensory information through the thalamus to the cortex (see sections 3.1.2 and 3.2).

Modeling studies in piriform cortex and hippocampus show that neuromodulation within a circuit may control the nature of the information processed. In a series of experimental and modeling studies (see section 3.2), it was shown that selective cholinergic (Hasselmo & Bower, 1992), noradrenergic (Hasselmo et al., 1997), or GABAergic (Tang & Hasselmo, 1994) suppression of intrinsic (recurrent) but not extrinsic (sensory) inputs promotes learning, while the absence of such suppression allows for memory recall. In this system, the selection of the information that is processed therefore depends on a rich class of neuromodulatory conditions, itself related to the behavioral state of the animal.

Finally, modulation of signal-to-noise ratio (see section 3.2) can also be considered as a form of input selection. By selectively enhancing certain neural inputs (the signal) and decreasing others (the noise), the system makes a de facto selection, which may change with neuromodulatory and behavioral conditions. This observation is at the basis of several models of selective attention involving the noradrenergic locus coeruleus (Aston-Jones et al., 1994; Rajkowski et al., 1994; Usher, Cohen, Servan-Schreiber, Rajkowski, & Aston-Jones, 1995) and of the DA-mediated control of cognitive processing in the prefrontal cortex and its relation to schizophrenia (Cohen et al., 1996).

4 Conclusion: Neuromodulation Increases and Controls Complexity —

Our review has shown that neuromodulation may play a significant computational role in a large spectrum of systems, from invertebrate central pattern generators to vertebrate cortical memory networks. In all cases, neuromodulation appears to be a powerful tool destined to increase and/or control the

computational complexity of a given network, without necessarily increasing the structural or dynamical complexity of the network itself. Spatially diffuse and slow neuromodulations of current conductances may trigger drastic changes of rhythmic patterns in central pattern generators, as well as in the thalamus, probably changing the nature of the downstream computations and increasing the complexity of the computations achieved by the whole circuit. Spatially selective and phasic neuromodulatory controls of specific neuronal input pathways help complex recurrent memory networks function properly.

Our review has also revealed two major limitations to the study of neuromodulation. Overcoming them requires the design of new theoretical and experimental tools, which undoubtedly will be beneficial.

The first stems from the observation that most modeling studies reviewed consider neuromodulation as an enhancing addition to a basic model. Often it is reduced to ad hoc parameter variations. We believe, however, that such an approach will no longer suffice as efforts are made to make computational models more biologically plausible in both their design and their function. Neuromodulation should be an integral part of the models. Only then will comprehensive theories of neuromodulation emerge and new neural computational principles may be discovered.

Second, in actual biological systems, neuromodulation has multiple simultaneous or sequential (cascade) effects on neural information processing. However, their experimental study almost always consists of individual modulations, keeping others constant. Moreover, in most cases, neuromodulation is present or absent and is rarely studied as a continuous phenomenon. It is not generally known whether the effects of different kinds of neuromodulation are truly independent, and if not, how they interact, nor is it known whether various levels of a single neuromodulation may yield drastically different neural behaviors. If some models propose interesting ways in which various modulatory phenomena might coexist, most of the models reviewed here still assume that multiple neuromodulatory effects are independent. As first proposed elsewhere (Harris-Warrick & Marder, 1991; Marder, Hooper, & Eisen, 1987), it is likely that accounting for the simultaneous effects of several neuroactive substances on a single network may increase its computational complexity in relevant and interesting ways, giving further insight into its function in the larger context of behavior.


Overall, computational and experimental models of neuromodulation appear to be powerful tools for the understanding of the computation of single cells as well as large neural networks.

Appendix: The Mathematical Tools _____

Uppercase letters are constants unless otherwise noted; lowercase letters are variables. The appendix is organized by levels of modeling, from more detailed to more abstract.

A.1 Markovian Chemical Kinetics Models.

$$i = \bar{G}s_1^o, \dots, s_i^o, \dots, (v - E)$$



R are rate constants, and s are concentrations (fraction of channels in state S). s_k^o is an open state.

At the most elementary level of modeling, neuronal processes can be described as chemical reactions, provided that their kinetics are quantitatively determined. In this framework, neuromodulatory phenomena are not distinguishable from others.

Destexhe et al. (1994a) expresses intracellular phenomena, membrane mechanisms, synaptic transmission, and neuromodulation with a single set of kinetic equations. In the model proposed, the neuromodulation by second messenger G-protein gated K⁺ channel (GABA_B, 5HT, M2 (ACh), $\alpha 2$ (NE), D2 (DA), histamine, opioid, and somatostatin receptors) is expressed by the appropriate formulation of rate constants of the type $R_{ij} = R_{ij}(v) = A_{ij}e^{-\frac{v}{B_{ij}}}$ for voltage-dependent gating and $R_{ij} = [L]\bar{R}_{ij}$ for ligand-activated gating.

Using a simplified formulation of this model, Destexhe et al. (1994a) model the putative role of NE and 5-HT in modulating rhythmic activity in thalamic reticular cells. G-protein activation is taken as a consequence of both NE and 5-HT neuromodulation. It is implemented as a modulating factor to the activation dynamics of a leak potassium current according to $g_{Kleak} = \bar{G}_{Kleak} \cdot m$, with $\frac{dm}{dt} = K[S]m - K'(1 - m)$, $[S]$ representing the concentration of second messenger present in the cell.

A.2 Hodgkin-Huxley Models. For a multicompartment model (x indexes compartments):

$$C \frac{dv}{dt} = \sum i + \frac{E_{leak} - v}{R} + \sum_x \frac{v_x - v}{R_a} + i_{syn} + I_{inject}$$

$$i = \bar{G}m^A n^B (v - E) \quad \text{with} \quad \frac{dm}{dt} = \frac{L_{\infty}^m(v) - m}{\tau_m(v)} \quad \text{and} \quad \frac{dn}{dt} = \frac{L_{\infty}^n(v) - n}{\tau_n(v)}.$$

m and n are activation and inactivation variables, respectively. Eventual synaptic potentials are modeled by:

$$i_{syn} = g_{syn}(v - E_{syn}) \quad \text{with} \quad g_{syn} = W\bar{G}_{syn} \frac{\tau_1}{\tau_2 - \tau_1} \left(-e^{-\frac{t}{\tau_1}} - -e^{-\frac{t}{\tau_2}} \right),$$

where W is the synaptic weight.

In the Hodgkin-Huxley formalism, neuromodulation is often expressed as a change in the maximal conductance of some particular membrane currents. At this level of modeling, it also may be implemented as a variation of the dynamics of some currents, variations in intracellular concentrations of some substances, and variation in synaptic transmission.

When the actual pharmacology of the channels is known, it is possible to express the conductances as functions of other intracellular quantities, including concentrations of neuromodulatory substances. For example, Butera et al. (1995) propose a scheme of interaction between dopamine and serotonin that yields expressions for conductances of the type:

$$\begin{aligned}\bar{G} &= \bar{G}_1 \left(\frac{K}{[Ca]_i + K} \right) \times \left(\frac{K_{DA}}{[DA] + K_{DA}} \right) \times \left(1 + \frac{K'}{1 + e^{-\frac{[cAMP]-K''}{D'}}} \right) \\ \bar{G} &= \bar{G}_1 \left(\frac{v - E}{1 + e^{\frac{ZF(v-E')}{RT}}} \right) \times \left(1 + \frac{K'}{1 + e^{-\frac{[cAMP]-K''}{D'}}} \right) \\ \frac{d[cAMP]}{dt} &= K \left(1 + K' \frac{[5HT]}{[5HT] + K''} \right) + C \frac{[cAMP]}{[cAMP] + K'''} .\end{aligned}$$

In some cases it is possible to obtain only an experimental curve quantitatively measuring the influence of a modulatory substance on given conductances. Bertram (1993) models two serotonin-sensitive conductances using a fit to their experimental values. The fit chosen takes the form:

$$\bar{G} = \bar{G}(s) = A + \frac{B}{1 + e^{-C(Ds-F)}} \text{ with } s \in [0, 1],$$

where s represents the concentration of serotonin applied. A similar formulation is used to describe the influence of two neuromodulatory substances (small cardioactive peptides, and myomodulins) on several currents in invertebrate neuromuscular circuits (Brezina et al., 1996).

In other cases, maximal conductances can be made dynamically dependent on intracellular quantities such as calcium (LeMasson et al., 1993) with

$$\tau \frac{d\bar{G}}{dt} = f([Ca]) - \bar{G} \text{ and } f([Ca]) = \frac{G_{\max}}{1 + e^{\pm \frac{[Ca]-C_T}{\Delta}}} .$$

Barkai et al. (1994) and Barkai and Hasselmo (1994) have access only to a qualitative experimental description of the effects of two potassium conductances on the firing adaptation of cortical cells. They therefore model these effects by choosing two parameter sets that yield adapting or weakly adapting model cells:

$$(\bar{G}, R) \in \{(\bar{G}_1, R_1), (\bar{G}_2, R_2)\} .$$

In a different system Epstein and Marder (1990) consider intermediate values, extrapolated linearly from the experimental ones following:

$$\bar{G} = \bar{G}(\alpha) = \alpha\bar{G}_1 + (1 - \alpha)\bar{G}_2,$$

where α is a dimensionless parameter.

When values for maximal conductances are not accessible experimentally, a theoretical search might be fruitful. In some cases, the set of conductances under neuromodulatory influence is known or hypothesized, and the dynamics of the network are under investigation. Dynamical systems theory (Guckenheimer et al., 1993) maps conductances values to possible network dynamics. The study of their stability leads them to experimental predictions about conductance values and their effects. In other cases, the dynamics of the network is known, but the set of conductances under neuromodulatory influences is unknown. Exhaustive search (Bhalla & Bower, 1993; Foster et al., 1993) allows for a systematic exploration of the parameter space constituted by all the maximal conductances hypothesized to be functional. Some regions of this space yield the dynamics under study. The location and shape of these regions predict what conductances are likely to be important (i.e., under neuromodulatory control) and what their possible values are.

Neuromodulation can also be expressed as a change in the dynamics (rather than maximal conductance) of some particular membrane currents. Such is the case of a variation in an inactivation time constant (Mukherjee & Kaplan, 1995) such as,

$$\tau_n(v) = \bar{T}_n \bar{\tau}_n(v) \quad \text{with } \bar{T}_n \in [\bar{T}_n^{\min}, \bar{T}_n^{\max}],$$

or of a variation in the voltage dependence of the steady-state activation curve $L_\infty^m(v)$, as for I_h (Destexhe et al., 1996; Golowasch et al., 1992).

Neuromodulation can also be expressed through changes in the intracellular concentration of some substances such as cAMP (Raymond et al., 1992) rather than as changes in maximal conductance of some membrane current.

Finally, neuromodulation can be expressed at the level of synaptic transmission. Such is the case for the presynaptic modulation of synaptic transmission by the activation of GABA_B receptors (Wallenstein & Hasselmo, 1997b).

In this model, the concentration of [GABA]_o in the synaptic cleft is first calculated as a function of the number of local active inhibitory synapses (n_{pre}) and a local diffusion term leading to:

$$\frac{d[\text{GABA}]_o}{dt} = C.n_{pre} - D.[\text{GABA}]_o,$$

where C and D are constants. At any point in time, [GABA]_o is then used to decrease synaptic currents, with $i_{syn} = i_{syn} - A.[\text{GABA}]_o$ where A is a constant.

Other models have viewed modulation as signal influencing synaptic mechanisms. Such is the case of the reward signal entering the weight modification rule, between VTA and cortex (Montague et al., 1996), or the direct change of synaptic efficacy triggered by an external center (Linster & Haselmo, 1997; Linster & Smith, 1997; Raymond et al., 1992).

A.3 Fitzhugh-Nagumo Models.

$$C \frac{dv}{dt} = f(v) - w + I_{inject}$$

$$\tau \frac{dw}{dt} = v - Dw$$

v is the fast (voltage-like, C small) variable; w is the slow (recovery-like) variable.

In this simplified framework (as for BonHoeffer-van der Pol or Morris-Lecar systems), individual concentrations and current conductance are not accessible, and fast Hodgkin-Huxley-type timescales are relaxed to pseudo-steady-state values. Neuronal behavior is assessed macroscopically through overall activity.

In a model of associative learning, Abbott (1990) proposes that neuromodulation may serve as a mechanism for initiating and terminating learning. Using the following formulation for the slow variable,

$$\tau \frac{dw}{dt} = av - (1 - a)w$$

he shows that depending on the value of a and the strength of the external inputs (I_{inject}), single cells may settle in regions of hyperpolarization, depolarization, oscillations, or bistability. At the network level, for values of a yielding oscillation, the network behaves like an associative memory (phase-locked oscillations, patterned according to synaptic coupling). For values of a yielding bistability, a putative consequence of neuromodulation, the network behaves like a nonselective latching short-term memory, maintaining the activity elicited initially by the input pattern, and allowing Hebbian plasticity to take place.

Interesting approaches to neuromodulation have also focused on the role of noise. Longtin (1993) uses stochastic resonance theory to show that the introduction of noise can have modulatory effects on the signal-to-noise ratio of a neuronal system, measured on the basis of the transfer of the oscillatory inputs to the output. The formulation used to illustrate this point introduces noise in v and a periodic forcing on w :

$$C \frac{dv}{dt} = v(v - A)(1 - v) - w + I_{inject} + \xi(t)$$

$$\tau \frac{dw}{dt} = v - Dw - [B + R \sin(\omega t)],$$

where $\xi(t)$ is a white noise (gaussian distributed) function, and $B + R \sin(\omega t)$ is a subthreshold oscillatory forcing. Experimental evidence has recently been found in support of the role of noise in improving information processing (Levin & Miller, 1996).

In this framework, neuromodulation can also be expressed as a change of electrical coupling between two cells, as in the STG (Kepler et al., 1990). It can be expressed as a shared current following:

$$I_{inject} = W(v_f - v) \text{ and } f(v) = Gv - (G + \alpha)\bar{V}$$

$$C_f \frac{dv_f}{dt} = -G_f(v_f - \bar{V}_f) + W(v - v_f).$$

We also should mention attempts at modeling NE-mediated decrease in K^+ current effects on the oscillatory behavior of a small thalamo-cortical model (Wallenstein, 1993). The effects were modeled by current injection in a Bonhoeffer–van der Pol modeling framework.

A.4 Leaky Integrator Models.

$$\tau \frac{dv}{dt} = -v + \sum_i W_i S(v_i) + I_{inject}$$

where $S(\cdot)$ is usually nonlinear (the sigmoid function), and v is the average membrane potential.

This representation allows for qualitative descriptions of the overall effects of average pools of neurons on behavior. Neuromodulation can be expressed by a change in firing threshold or a significant modification of synaptic weights (Linster & Masson, 1996) with:

$$v \leq \theta_{\min} \Rightarrow S(v) = 0 \quad \text{and} \quad v \geq \theta_{\max} \Rightarrow S(v) = 1$$

$$\theta_{\min} \leq v \leq \theta_{\max} \Rightarrow S(v) = \alpha v.$$

Neuromodulation can also be expressed by introducing a multiplicative factor to the upper and lower bounds of the sigmoid function or by decreasing weights by a fraction (Liljenstrom & Hasselmo, 1995):

$$(\theta_{\min}, \theta_{\max}) \Rightarrow (\beta\theta_{\min}, \beta\theta_{\max}) \quad \text{or} \quad W \Rightarrow \frac{W}{\gamma}.$$

It can also be expressed as a dependence of the sigmoid function on other quantities such as the intracellular calcium concentration. Cartling (1996) models neuromodulation as a change in neuronal adaptability (coupling between activity and excitability). It is expressed as a multiplicative factor (a) to the intracellular Ca concentration (c) with activity-dependent second-order dynamics:

$$S(v) = S(v, c) = \text{MAX}(\tanh(Av - ac - \theta), 0)$$

$$\text{with } \frac{dc}{dt} = \frac{K}{K' + c} S(v, c) + \frac{K'' - c}{T},$$

where c is the intracellular calcium concentration and a is the adaptability. Neuromodulation is measured by a , which depends on the total activity of the network:

$$a = A_{\max}(1 - n) \quad \text{with} \quad \frac{dn}{dt} = C(1 - n) \sum_i \alpha_i v_i.$$

α_i is the size of the population having v_i as state variable.

A.5 Connectionist Models.

$$o = \sum_i W_i S(o_i).$$

In case of modifiable synapses:

$$\frac{dw}{dt} = \eta x(t)y(t),$$

where $x(t)$ represents the presynaptic activity, and $y(t)$ represents the post-synaptic activity. $S(\cdot)$ is analogous to the sigmoid function of leaky integrator models.

Neuromodulation in connectionist models has been expressed in two general ways. The first expresses neuromodulation in the sigmoid function, the second in the dynamics of the synaptic weights.

Neuromodulation can be implemented as a modification of the slope (gain) of the sigmoid function (Cohen & Servan-Schreiber, 1992; Servan-Schreiber et al., 1990) in a small network (chain) of connectionist elements, following:

$$S(o) = \frac{1}{1 + e^{-(Go+B)}} \quad \text{with} \quad G \in [G_{\min}, G_{\max}].$$

Other modifications to the sigmoid function can be made to express other neuromodulatory properties, such as suppression of neuronal adaptation (Liljenstrom & Hasselmo, 1995). Using a nonmodulated sigmoid function,

$$S(o_i) = CQ_i \left(1 - e^{-\frac{o_i+1}{Q_i}} \right),$$

activity-dependent neuromodulation is expressed as:

$$S(o_i) = CQ_i \left(1 - e^{-\frac{o_i+1}{Q_i}} \right) \times e^{-\alpha(S(o_i))_{i-T}^2},$$

where T is a fixed time window and t is time.

A second modeling approach consists of introducing neuromodulatory effects to the learning rule. Montague et al. (1996) modeled the activity of

dopamine cells in the VTA by augmenting the postsynaptic activity with an external reward signal:

$$S(o) = o$$

$$\Delta w = \eta x(t)(r(t) + y(t)),$$

where $r(t)$ is the external reward signal.

Hasselmo (1993) selectively modifies the learning rate of certain synapses to include the effects of ACh with

$$\frac{dw}{dt} = \eta(1 - c)x(t)y(t) \quad \text{with } c \in [0, 1],$$

where c measures the amount of cholinergic suppression. Myers et al. (1996) adopt a similar approach in their model of cholinergic influence on cortico-hippocampal interaction during eye-blink conditioning.

Both sigmoid and synaptic modulation can coexist and have been modeled by Hasselmo and Schnell (1994). The synaptic modulation is expressed using:

$$W_i = (1 - cC_{W_i})w_i \quad \text{and} \quad \frac{dw_i}{dt} = \eta(1 + cC_\eta - C_\eta)x(t)y(t)$$

and replacing θ with $(1 - cT_{\max})\theta$ in the normal ramplike sigmoid function:

$$o \leq \theta \Rightarrow S(o) = 0$$

$$o \geq \theta \Rightarrow S(o) = o - \theta.$$

Finally we should mention the modeling of morphological changes in Alzheimer's disease (Horn, Ruppin, Usher, & Herrmann, 1993) expressing modulation of activity by random synaptic deletion, and appropriate compensation with:

$$o = c \sum_{i \in \Delta} W_i S(o_i) \quad \text{with } |\Delta| = (1 - d)N \quad \text{and} \quad S(o) = \text{Step}(o - T).$$

c is the compensation factor, d is the deletion factor, and $c = 1 + \frac{dk}{1-d}$ with $k \in [0, 1]$.

Acknowledgments

We thank Larry Abbott, Mike Hasselmo, John Lisman, Eve Marder, and Akaysha Tang for their critical reading of the manuscript for this article.

References

Abbott, L. F. (1990). Modulation of function and gated learning in a network memory. *Proc Natl Acad Sci USA*, 87(23), 9241–9245.

- Abrams, T. W., & Kandel, E. R. (1988). Is contiguity detection in classical conditioning a system or a cellular property? Learning in *Aplysia* suggests a possible molecular site. *Trends Neurosci*, *11*(4), 128–135.
- Adams, W. B., & Benson, J. A. (1985). The generation and modulation of endogenous rhythmicity in the *Aplysia* bursting pacemaker neurone R15. *Prog Biophys Mol Biol*, *46*(1), 1–49.
- Aston-Jones, G., Rajkowski, J., Kubiak, P., & Alexinsky, T. (1994). Locus coeruleus neurons in monkey are selectively activated by attended cues in a vigilance task. *J Neurosci*, *14*(7), 4467–4480.
- Barkai, E., Bergman, R. E., Horwitz, G., & Hasselmo, M. E. (1994). Modulation of associative memory function in a biophysical simulation of rat piriform cortex. *J Neurophysiol*, *72*(2), 659–677.
- Barkai, E., & Hasselmo, M. E. (1994). Modulation of the input/output function of rat piriform cortex pyramidal cells. *J Neurophysiol*, *72*(2), 644–658.
- Bertram, R. (1993). A computational study of the effects of serotonin on a molluscan burster neuron. *Biological Cybernetics*, *69*, 257–267.
- Bertram, R. (1994). Reduced-system analysis of the effects of serotonin on a molluscan burster neuron. *Biol Cybern*, *70*(4), 359–368.
- Bhalla, U. S., & Bower, J. M. (1993). Exploring parameter space in detailed single neuron models: Simulations of the mitral and granule cells of the olfactory bulb. *J Neurophysiol*, *69*(6), 1948–1965.
- Biedenbach, M. A. (1966). Effects of anesthetics and cholinergic drugs on prepyriform electrical activity in cats. *Exp Neurol*, *16*(4), 464–479.
- Bressler, S. L., & Freeman, W. J. (1980). Frequency analysis of olfactory system EEG in cat, rabbit, and rat. *Electroencephalogr Clin Neurophysiol*, *50*(1–2), 19–24.
- Brezina, V., Orekhova, I. V., & Weiss, K. R. (1996). Functional uncoupling of linked neurotransmitter effects by combinatorial convergence. *Science*, *273*(5276), 806–810.
- Brown, T. H., Kairiss, E. W., & Keenan, C. L. (1990). Hebbian synapses: Biophysical mechanisms and algorithms. *Annu Rev Neurosci*, *13*, 475–511.
- Buchholtz, F., Golowasch, J., Epstein, I. R., & Marder, E. (1992). Mathematical model of an identified stomatogastric ganglion neuron. *J Neurophysiol*, *67*(2), 332–340.
- Bulsara, A., Jacobs, E. W., Zhou, T., Moss, F., & Kiss, L. (1991). Stochastic resonance in a single neuron model: Theory and analog simulation. *J Theor Biol*, *152*(4), 531–555.
- Butera, R. J., Jr., Clark, J. W., Jr., Canavier, C. C., Baxter, D. A., & Byrne, J. H. (1995). Analysis of the effects of modulatory agents on a modeled bursting neuron: Dynamic interactions between voltage and calcium dependent systems [published erratum appears in *J Comput Neurosci* 1996 Sep;3(3):265]. *J Comput Neurosci*, *2*(1), 19–44.
- Byrne, J. H. (1987). Cellular analysis of associative learning. *Physiol Rev*, *67*(2), 329–439.
- Calabrese, R. L., & De Schutter, E. (1992). Motor-pattern-generating networks in invertebrates: Modeling our way toward understanding. *Trends Neurosci*, *15*(11), 439–445.

- Cartling, B. (1996). Dynamics control of semantic processes in a hierarchical associative memory. *Biol Cybern*, 74(1), 63–71.
- Chan-Palay, V., & Palay, S. L. (Eds.). (1984). *Coexistence of neuroactive substances in neurons*. New York: Wiley.
- Clark, C. R., Geffen, G. M., & Geffen, L. B. (1987a). Catecholamines and attention. I: Animal and clinical studies. *Neurosci Biobehav Rev*, 11(4), 341–352.
- Clark, C. R., Geffen, G. M., & Geffen, L. B. (1987b). Catecholamines and attention. II: Pharmacological studies in normal humans. *Neurosci Biobehav Rev*, 11(4), 353–364.
- Cohen, J. D., Braver, T. S., & O'Reilly, R. C. (1996). A computational approach to prefrontal cortex, cognitive control and schizophrenia: Recent developments and current challenges. *Philos Trans R Soc Lond B Biol Sci*, 351(1346), 1515–1527.
- Cohen, J. D., & Servan-Schreiber, D. (1992). Context, cortex, and dopamine: A connectionist approach to behavior and biology in schizophrenia. *Psychol Rev*, 99(1), 45–77.
- Contreras, D., Destexhe, A., Sejnowski, T. J., & Steriade, M. (1997). Spatiotemporal patterns of spindle oscillations in cortex and thalamus. *J Neurosci*, 17(3), 1179–1196.
- Cooper, J. R., Bloom, F. E., & Roth, R. H. (1991). *The biochemical basis of neuropharmacology* (6th ed.). New York: Oxford University Press.
- Cooper, S. J. (1991). Interactions between endogenous opioids and dopamine: Implications for reward and aversion. In P. Willner & J. Scheel-Kruger (Eds.), *The mesolimbic dopamine system: From motivation to action* (pp. 331–366). New York: Wiley.
- de Carvalho, L. A. (1994). Modeling the thalamocortical loop. *Int J Biomed Comput*, 35(4), 267–296.
- Destexhe, A., Bal, T., McCormick, D. A., & Sejnowski, T. J. (1996). Ionic mechanisms underlying synchronized oscillations and propagating waves in a model of ferret thalamic slices. *J Neurophysiol*, 76(3), 2049–2070.
- Destexhe, A., Contreras, D., Sejnowski, T. J., & Steriade, M. (1994a). Modeling the control of reticular thalamic oscillations by neuromodulators. *Neuroreport*, 5(17), 2217–2220.
- Destexhe, A., Mainen, Z. F., & Sejnowski, T. J. (1994b). Synthesis of models for excitable membranes, synaptic transmission and neuromodulation using a common kinetic formalism. *J Comput Neurosci*, 1(3), 195–230.
- Epstein, I. R., & Marder, E. (1990). Multiple modes of a conditional neural oscillator. *Biol Cybern*, 63(1), 25–34.
- Foster, W. R., Ungar, L. H., & Schwaber, J. S. (1993). Significance of conductances in Hodgkin-Huxley models. *J Neurophysiol*, 70(6), 2502–2518.
- Fuxe, K., & Agnati, L. F. (1991). Volume transmission in the brain: Novel mechanism for neural transmission. *Advances in Neuroscience*, 1, 1–9.
- Gingrich, K. J., & Byrne, J. H. (1987). Single-cell neuronal model for associative learning. *J Neurophysiol*, 57(6), 1705–1715.
- Gluck, M. A., & Myers, C. E. (1993). Hippocampal mediation of stimulus representation: A computational theory. *Hippocampus*, 3(4), 491–516.
- Goldman-Rakic, P. S., Lidow, M. S., Smiley, J. F., & Williams, M. S. (1992). The anatomy of dopamine in monkey and human prefrontal cortex. *J Neural Transm Suppl*, 36, 163–177.

- Golowasch, J., Buchholtz, F., Epstein, I. R., & Marder, E. (1992). Contribution of individual ionic currents to activity of a model stomatogastric ganglion neuron. *J Neurophysiol*, *67*(2), 341–349.
- Gray, C. M. (1994). Synchronous oscillations in neuronal systems: Mechanisms and functions. *J Comput Neurosci*, *1*(1–2), 11–38.
- Guckenheimer, J., Gueron, S., & Harris-Warrick, R. M. (1993). Mapping the dynamics of a bursting neuron. *Philos Trans R Soc Lond B Biol Sci*, *341*(1298), 345–359.
- Guckenheimer, J., Harris-Warrick, R., Peck, J., & Willms, A. (1997). Bifurcation, bursting, and spike frequency adaptation. *J Comput Neurosci*, *4*(3), 257–277.
- Hagiwara, N., & Irisawa, H. (1989). Modulation by intracellular Ca²⁺ of the hyperpolarization-activated inward current in rabbit single sino-atrial node cells. *J Physiol (Lond)*, *409*, 121–141.
- Hansson, E., & Ronnback, L. (1994). Astroglial modulation of synaptic transmission. *Perspect Dev Neurobiol*, *2*(3), 217–223.
- Harris-Warrick, R. M., Coniglio, L. M., Levini, R. M., Gueron, S., & Guckenheimer, J. (1995). Dopamine modulation of two subthreshold currents produces phase shifts in activity of an identified motoneuron. *J Neurophysiol*, *74*(4), 1404–1420.
- Harris-Warrick, R. M., & Flamm, R. E. (1987). Multiple mechanisms of bursting in a conditional bursting neuron. *J Neurosci*, *7*(7), 2113–2128.
- Harris-Warrick, R. M., & Marder, E. (1991). Modulation of neural networks for behavior. *Annual Reviews of Neuroscience*, *14*, 39–57.
- Hasselmo, M. E. (1993). Acetylcholine and learning in a cortical associative memory. *Neural Computation*, *5*, 32–44.
- Hasselmo, M. E. (1995). Neuromodulation and cortical function: Modeling the physiological basis of behavior. *Behav Brain Res*, *67*(1), 1–27.
- Hasselmo, M. E., & Bower, J. M. (1992). Cholinergic suppression specific to intrinsic not afferent fiber synapses in rat piriform (olfactory) cortex. *J Neurophysiol*, *67*(5), 1222–1229.
- Hasselmo, M. E., Linster, C., Patil, M., Ma, D., & Cekić, M. (1997). Noradrenergic suppression of synaptic transmission may influence cortical signal-to-noise ratio. *J Neurophysiol*, *77*(6), 3326–3339.
- Hasselmo, M. E., & Schnell, E. (1994). Laminar selectivity of the cholinergic suppression of synaptic transmission in rat hippocampal region CA1: Computational modeling and brain slice physiology. *J Neurosci*, *14*(6), 3898–3914.
- Hodgkin, A. L., & Huxley, A. F. (1952). A quantitative description of membrane current and its application to conduction and excitation in nerve. *Journal of Physiology*, *117*, 500–544.
- Horn, D., Ruppin, E., Usher, M., & Herrmann, M. (1993). Neuronal network modeling of memory deterioration in Alzheimer's disease. *Neural Computation*, *5*(5), 736–749.
- Jackson, S. R., Marrocco, R., & Posner, M. I. (1994). Networks of anatomical areas controlling visuospatial attention. *Neural Networks*, *7*(6/7), 925–944.
- Jay, T. M., Glowinski, J., & Thierry, A. M. (1995). Inhibition of hippocampoprefrontal cortex excitatory responses by the mesocortical DA system. *Neuroreport*, *6*(14), 1845–1848.

- Jensen, O., Idiart, M. A. P., & Lisman, J. E. (1996). Physiologically realistic formation of autoassociative memory in networks with theta/gamma oscillations: Role of fast NMDA channels. *Learning and Memory*, 3, 243–256.
- Kaczmarek, L. K., & Levitan, I. B. (1987). *Neuromodulation: The biochemical control of neuronal excitability*. New York: Oxford University Press.
- Katz, P. S., & Frost, W. N. (1995a). Intrinsic neuromodulation in the Tritonia swim CPG: Serotonin mediates both neuromodulation and neurotransmission by the dorsal swim interneurons. *J Neurophysiol*, 74(6), 2281–2294.
- Katz, P. S., & Frost, W. N. (1995b). Intrinsic neuromodulation in the Tritonia swim CPG: The serotonergic dorsal swim interneurons act presynaptically to enhance transmitter release from interneuron C2. *J Neurosci*, 15(9), 6035–6045.
- Katz, P. S., & Frost, W. N. (1996). Intrinsic neuromodulation: Altering neuronal circuits from within. *Trends Neurosci*, 19(2), 54–61.
- Katz, P. S., Getting, P. A., & Frost, W. N. (1994). Dynamic neuromodulation of synaptic strength intrinsic to a central pattern generator circuit. *Nature*, 367(6465), 729–731.
- Kepler, T. B., Marder, E., & Abbott, L. F. (1990). The effect of electrical coupling on the frequency of model neuronal oscillators. *Science*, 248(4951), 83–85.
- Kupfermann, I. (1991). Functional studies of cotransmission. *Physiol Rev*, 71(3), 683–732.
- Lechner, H. A., Baxter, D. A., Clark, J. W., & Byrne, J. H. (1996). Bistability and its regulation by serotonin in the endogenously bursting neuron R15 in *Aplysia*. *J Neurophysiol*, 75(2), 957–962.
- LeMasson, G., Marder, E., & Abbott, L. F. (1993). Activity-dependent regulation of conductances in model neurons. *Science*, 259(5103), 1915–1917.
- Levin, J. E., & Miller, J. P. (1996). Broadband neural encoding in the cricket cercal sensory system enhanced by stochastic resonance. *Nature*, 380(6570), 165–168.
- Liljenstrom, H., & Hasselmo, M. E. (1995). Cholinergic modulation of cortical oscillatory dynamics. *J Neurophysiol*, 74(1), 288–297.
- Linster, C., & Gervais, R. (1996). Investigation of the role of interneurons and their modulation by centrifugal fibers in a neural model of the olfactory bulb. *J Comput Neurosci*, 3(3), 225–246.
- Linster, C., & Hasselmo, M. (1997). Modulation of inhibition in a model of olfactory bulb reduces overlap in the neural representation of olfactory stimuli. *Behav Brain Res*, 84(1–2), 117–127.
- Linster, C., & Masson, C. (1996). A neural model of olfactory sensory memory in the honeybee's antennal lobe. *Neural Computation*, 8, 94–114.
- Linster, C., & Smith, B. H. (1997). A computational model of the response of honey bee antennal lobe circuitry to odor mixtures: Overshadowing, blocking and unblocking can arise from lateral inhibition. *Behavioural Brain Research*, 87, 1–14.
- Lisman, J. E., & Idiart, M. A. (1995). Storage of 7 ± 2 short-term memories in oscillatory subcycles. *Science*, 267(5203), 1512–1515.
- Longtin, A. (1993). Stochastic resonance in neuron models. *Journal of Statistical Physics*, 70(1/2), 309–327.
- Longtin, A., Bulsara, A., Pierson, D., & Moss, F. (1994). Bistability and the dy-

- namics of periodically forced sensory neurons. *Biol Cybern*, 70(6), 569–578.
- Madison, D. V., Lancaster, B., & Nicoll, R. A. (1987). Voltage clamp analysis of cholinergic action in the hippocampus. *J Neurosci*, 7(3), 733–741.
- Madison, D. V., & Nicoll, R. A. (1986). Actions of noradrenaline recorded intracellularly in rat hippocampal CA1 pyramidal neurones, in vitro. *J Physiol (Lond)*, 372, 221–244.
- Marder, E. (1996). Neural modulation: Following your own rhythm. *Curr Biol*, 6(2), 119–121.
- Marder, E., Christie, A. E., & Kilman, V. L. (1995). Functional organization of cotransmission systems: Lessons from small nervous systems. *Invertebrate Neuroscience*, 1, 105–112.
- Marder, E., Hooper, S. L., & Eisen, J. S. (1987). Multiple neurotransmitters provide a mechanism for the production of multiple outputs from a single neuronal circuit. In G. M. Edelman, E. W. Gall, & W. M. Cowan (Eds.), *Synaptic function* (pp. 305–327). New York: Wiley.
- Marder, E., & Selverston, A. I. (1992). Modeling the stomatogastric nervous system. In R. M. Harris-Warrick, E. Marder, A. I. Selverston, & M. Moulins (Eds.), *Dynamic biological networks: The stomatogastric nervous system*. Cambridge, MA: MIT Press.
- McCormick, D. A. (1992). Neurotransmitter actions in the thalamus and cerebral cortex and their role in neuromodulation of thalamocortical activity. *Prog Neurobiol*, 39(4), 337–388.
- McCormick, D. A., & Huguenard, J. R. (1992). A model of the electrophysiological properties of thalamocortical relay neurons. *J Neurophysiol*, 68(4), 1384–1400.
- McCormick, D. A., & Wang, Z. (1991). Serotonin and noradrenaline excite GABAergic neurones of the guinea-pig and cat nucleus reticularis thalami. *J Physiol (Lond)*, 442, 235–255.
- McNamara, B., & Wiesenfeld, K. (1989). Theory of stochastic resonance. *Physical Review A*, 39(9), 4854–4869.
- Montague, P. R., Dayan, P., Person, C., & Sejnowski, T. J. (1995). Bee foraging in uncertain environments using predictive Hebbian learning. *Nature*, 377(6551), 725–728.
- Montague, P. R., Dayan, P., & Sejnowski, T. J. (1996). A framework for mesencephalic dopamine systems based on predictive Hebbian learning. *J Neurosci*, 16(5), 1936–1947.
- Mukherjee, P., & Kaplan, E. (1995). Dynamics of neurons in the cat lateral geniculate nucleus: In vivo electrophysiology and computational modeling. *J Neurophysiol*, 74(3), 1222–1243.
- Myers, C. E., Ermita, B. R., Harris, K., Hasselmo, M., Solomon, P., & Gluck, M. A. (1996). A computational model of cholinergic disruption of septohippocampal activity in classical eyeblink conditioning. *Neurobiol Learn Mem*, 66(1), 51–66.
- Nusbaum, M. P., Weimann, J. M., Golowasch, J., & Marder, E. (1992). Presynaptic control of modulatory fibers by their neural network targets. *J Neurosci*, 12(7), 2706–2714.
- Pearson, K. G. (1993). Common principles of motor control in vertebrates and

- invertebrates. *Annu Rev Neurosci*, 16, 265–297.
- Rajkowski, J., Kubiak, P., & Aston-Jones, G. (1994). Locus coeruleus activity in monkey: Phasic and tonic changes are associated with altered vigilance. *Brain Res Bull*, 35(5–6), 607–616.
- Raymond, J. L., Baxter, D. A., Buonomano, D. V., & Byrne, J. H. (1992). A learning rule based on empirically-derived activity-dependent neuromodulation supports operant conditioning in small network. *Neural Networks*, 5, 789–803.
- Ridet, J. L., Rajaofetra, N., Teilhac, J. R., Geffard, M., & Privat, A. (1993). Evidence for nonsynaptic serotonergic and noradrenergic innervation of the rat dorsal horn and possible involvement of neuron-glia interactions. *Neuroscience*, 52(1), 143–157.
- Rinzel, J., & Lee, Y. S. (1987). Dissection of a model for neuronal parabolic bursting. *J Math Biol*, 25(6), 653–675.
- Sara, S. J., Vankov, A., & Herve, A. (1994). Locus coeruleus-evoked responses in behaving rats: A clue to the role of noradrenaline in memory. *Brain Res Bull*, 35(5–6), 457–465.
- Schultz, W., Apicella, P., & Ljungberg, T. (1993). Responses of monkey dopamine neurons to reward and conditioned stimuli during successive steps of learning a delayed response task. *J Neurosci*, 13(3), 900–913.
- Servan-Schreiber, D., Printz, H., & Cohen, J. D. (1990). A network model of catecholamine effects: Gain, signal-to-noise ratio, and behavior. *Science*, 249(4971), 892–895.
- Singer, W. (1993). Synchronization of cortical activity and its putative role in information processing and learning. *Annu Rev Physiol*, 55, 349–374.
- Starke, K., Gothert, M., & Kilbinger, H. (1989). Modulation of neurotransmitter release by presynaptic autoreceptors. *Physiol Rev*, 69(3), 864–989.
- Sun, X., Fonta, C., & Masson, C. (1993). Odour quality processing by bee antennal lobe neurons. *Chemical Senses*, 18, 355–377.
- Sutton, R. S., & Barto, A. G. (1990). Time-derivative models of Pavlovian reinforcement. In M. Gabriel & J. Moore (Eds.), *Learning and computational neuroscience*. Cambridge, MA: MIT Press.
- Tang, A. C., & Hasselmo, M. E. (1994). Selective suppression of intrinsic but not afferent fiber synaptic transmission by baclofen in the piriform (olfactory) cortex. *Brain Res*, 659(1-2), 75–81.
- Traub, R. D., Miles, R., & Buzsaki, G. (1992). Computer simulation of carbachol-driven Rhythmic population oscillations in the CA3 region of the in vitro rat hippocampus. *J Physiol (Lond)*, 451, 653–672.
- Traub, R. D., Whittington, M. A., Colling, S. B., Buzsaki, G., & Jefferys, J. G. (1996). Analysis of gamma rhythms in the rat hippocampus in vitro and in vivo. *J Physiol (Lond)*, 493(Pt 2), 471–484.
- Turrigiano, G., Abbott, L. F., & Marder, E. (1994). Activity-dependent changes in the intrinsic properties of cultured neurons. *Science*, 264(5161), 974–977.
- Usher, M., Cohen, J., Servan-Schreiber, D., Rajkowski, J., Kubiak, P., & Aston-Jones, G. (1995). *A computational model of locus coeruleus function and its influence on behavioral performance in primate* (Tech. rep. No. PDP.CNS 95.1). Pittsburgh: Carnegie Mellon University, University of Pittsburgh.
- van Dongen, P. A. (1981). The central noradrenergic transmission and the locus

- coeruleus: A review of the data, and their implications for neurotransmission and neuromodulation. *Prog Neurobiol*, 16(2), 117–143.
- Wallenstein, G. V. (1993). *Spatial mode dynamics of a thalamo-cortical network*. Paper presented at the SPIE, Chaos in Biology and Medicine.
- Wallenstein, G. V. (1996). Adenosinic modulation of 7-14 Hz spindle rhythms in interconnected thalamic relay and nucleus reticularis neurons. *Neuroscience*, 73(1), 93–98.
- Wallenstein, G. V., & Hasselmo, M. E. (1997a). Functional transitions between epileptiform-like activity and associative memory in hippocampal region CA3. *Brain Res Bull*, 43(5), 485–493.
- Wallenstein, G. V., & Hasselmo, M. E. (1997b). GABAergic modulation of hippocampal population activity: Sequence learning, place field development, and the phase precession effect. *J Neurophysiol*, 78(1), 393–408.
- Wang, X. J., & Buzsaki, G. (1996). Gamma oscillation by synaptic inhibition in a hippocampal interneuronal network model. *J Neurosci*, 16(20), 6402–6413.
- Wilson, M., & Bower, J. M. (1992). Cortical oscillations and temporal interactions in a computer simulation of piriform cortex. *J Neurophysiol*, 67(4), 981–995.
- Zador, A., Koch, C., & Brown, T. H. (1990). Biophysical model of a Hebbian synapse. *Proc Natl Acad Sci U S A*, 87(17), 6718–6722.

Received January 2, 1997; accepted October 23, 1997.