

**Computing the Minimum Eigenvalue
of a Symmetric Positive
Definite Toeplitz Matrix**

George Cybenko*
Charles Van Loan†

TR 82-527
April 1984

*Department of Mathematics
Tufts University
Medford, Massachusetts 02155

and
The Statistics Center
Massachusetts Institute of Technology
Cambridge, Massachusetts 02135

†Department of Computer Science
Cornell University
Ithaca, New York 14853

*Partially supported by contract NSF MCS 8003364

†Partially supported by contracts NSF-MCS-8004106 and ONR N00014-83-K-0640.

COMPUTING THE MINIMUM EIGENVALUE OF A SYMMETRIC POSITIVE
DEFINITE TOEPLITZ MATRIX

George Cybenko^{*}
Department of Mathematics
Tufts University
Medford, Mass. 02155

and

The Statistics Center
MIT
Cambridge, Mass. 02135

Charles Van Loan^{**}
Department of Computer Science
Cornell University
Ithaca, NY 14853

Abstract

A method for computing the smallest eigenvalue of a symmetric positive definite Toeplitz matrix is given. It relies solely upon the Levinson-Durbin algorithm. The procedure involves a combination of bisection and Newton's method. Good starting values are also shown to be obtainable from the Levinson-Durbin algorithm.

^{*}Partially supported by contract NSF MCS 8003364

^{**}Partially supported by contracts NSF-MCS-8004106 and ONR N00014-83-K-0640

1. Introduction

Recent progress in signal processing and estimation has generated considerable interest in the problem of computing the minimal eigenvalue of a Toeplitz matrix. The fundamental modelling and solution that led to this are due to Pisarenko [P73], while more recently numerous authors have discussed the computational aspects of the problem [F83,H80,H83a,H83c].

In this paper we shall not discuss the underlying assumptions, merits, or potential applications of the model-- instead pointing the interested reader to the literature concerned with these issues [H83a,P73]. We hasten to add that the quantities of ultimate interest in applications are the roots of the polynomial whose coefficients are given by the eigenvector associated with the minimal eigenvalue of the Toeplitz matrix. We shall only discuss the computation of the minimal eigenvalue noting that the associated eigenvector can be obtained as a by-product. Furthermore, methods exist for computing the roots that altogether avoid the explicit formation of the eigenvector [C84b].

The essence of our minimum eigenvalue procedure involves solving systems of shifted Yule-Walker (YW) systems. Initially, the solutions to these systems are used in a bisection scheme that repeatedly halves a bracketing subinterval. Subsequently, a Newton iteration takes over that quadratically converges to the desired eigenvalue. We stress the fact that only YW systems are involved--an important point since extremely efficient methods for YW systems exist. (They require half the computational resources needed by general symmetric Toeplitz system solvers.)

In an absolute sense, only modest use is made of Toeplitz structure. Indeed, this is true of all currently known Toeplitz eigenvalue solvers. The

study of the eigenstructure of finite Toeplitz matrices is proceeding rather slowly. Recent developments include [C84a,C84b,D83]. An indication of the collective ignorance about Toeplitz eigenstructure is that the inverse eigenvalue problem for real symmetric Toeplitz matrices is currently unsolved. We suspect that the process of designing efficient algorithms for this problem will go hand in hand with the uncovering of Toeplitz eigenstructure properties.

Our paper is organized as follows. Section 2 describes a rational function intimately related to the eigenvalue problem for Hermitian matrices. Section 3 specializes the discussion to real symmetric Toeplitz matrices and then develops our bisection/Newton scheme. In the last section we discuss the numerical behavior of our procedure.

2. A Rational Eigenvalue Equation

In this section we derive a rational function from a given Hermitian matrix that has the property that its zeros are eigenvalues of the original matrix. A feature of this rational function is that both it and its derivatives are easily evaluated thereby making Newton-type schemes feasible. Strictly speaking, parts of our derivation are not new and can be found in [W65], but we present the details for the sake of completeness.

Let T be an $n \times n$ Hermitian matrix partitioned as follows:

$$T = \begin{array}{cc|c} \left[\begin{array}{cc} \gamma & r^* \\ r & G \end{array} \right] & & \begin{array}{c} 1 \\ n-1 \end{array} \\ \hline & \begin{array}{c} 1 \\ n-1 \end{array} & \end{array} .$$

Here, r^* denotes the conjugate transpose of r . It is well known [G83] that the eigenvalues of T and G are real and satisfy an interlacing property. In particular, if $\lambda_i(T)$ and $\lambda_i(G)$ are the i -th largest eigenvalues of T and G

respectively then

$$\lambda_n(T) \leq \lambda_{n-1}(G) \leq \lambda_{n-1}(T) \leq \dots \leq \lambda_2(T) \leq \lambda_1(G) \leq \lambda_1(T)$$

Note that if T has a repeated eigenvalue then the repeated value is also an eigenvalue of G . Adopting the notation

$$\lambda_{\min} \equiv \lambda_n(T)$$

we shall assume throughout this paper that

$$(2.1) \quad d = \lambda_{n-1}(G) - \lambda_{\min} > 0 .$$

The strict separation of $\lambda_{n-1}(G)$ and λ_{\min} guarantees that the eigenvector associated with λ_{\min} is unique up to scalar premultiplication. It is a realistic assumption in many important problems such as the estimation of Pisarenko frequencies. See [C84b].

Suppose

$$(2.2) \quad \begin{bmatrix} \gamma & r^* \\ r & G \end{bmatrix} \begin{bmatrix} \alpha \\ y \end{bmatrix} = \lambda_{\min} \begin{bmatrix} \alpha \\ y \end{bmatrix}$$

where it is assumed that α and y are not both zero. From this equation we obtain

$$(2.3) \quad \begin{aligned} \gamma\alpha + r^*y &= \lambda_{\min}\alpha \\ \alpha r + Gy &= \lambda_{\min}y \end{aligned}$$

We must have $\alpha \neq 0$ for otherwise $Gy = \lambda_{\min}y$ contradicting (2.1). Noting that

$G - \lambda_{\min} I$ is positive definite we obtain the following rational equation for λ_{\min} :

$$(2.4) \quad \gamma - r^*(G - \lambda_{\min} I)^{-1}r - \lambda_{\min} = 0$$

Thus, the smallest eigenvalue of T is the smallest root of the rational function

$$(2.5) \quad f(\lambda) = \gamma - \lambda - r^*(G - \lambda I)^{-1}r .$$

In addition, $f(\lambda)$ has the following important properties whenever $\lambda < \lambda_{n-1}(G)$:

$$(2.6) \quad f'(\lambda) = -1 - \|(G - \lambda I)^{-1}r\|_2^2 \leq -1$$

$$(2.7) \quad f''(\lambda) = -2r^*(G - \lambda I)^{-3}r \leq 0$$

Now consider the following Newton iteration:

Algorithm 2.1

Let $\lambda \in [\lambda_{\min}, \lambda_{n-1}(G))$ be given along with a tolerance $\delta > 0$.

Do Until ($|f(\lambda)| < \delta / (1 + \|w\|_2^2)^{1/2}$)

$$\left[\begin{array}{l} \text{Solve } (G - \lambda I)w = -r \text{ for } w . \\ \lambda := \lambda + \frac{\gamma + r^*w - \lambda}{1 + w^*w} = \lambda - \frac{f(\lambda)}{f'(\lambda)} \end{array} \right.$$

Properties (2.6) and (2.7) ensure that the iteration converges to λ_{\min} . To see this assume that $\lambda \in (\lambda_{\min}, \lambda_{n-1}(G))$ and set

$$\lambda_+ = \lambda - \frac{f(\lambda)}{f'(\lambda)} .$$

Since f is monotone decreasing in this interval, it follows that both $f(\lambda)$ and $f'(\lambda)$ are negative. Thus, $\lambda_+ < \lambda$. On the other hand, from truncated Taylor series we have

$$0 = f(\lambda_{\min}) = f(\lambda) + f'(\lambda)(\lambda_{\min} - \lambda) + \frac{f''(\zeta)}{2} (\lambda_{\min} - \lambda)^2$$

with $\zeta \in [\lambda_{\min}, \lambda]$. It follows that

$$(2.8) \quad \lambda_+ - \lambda_{\min} = \frac{f''(\zeta)}{2f'(\lambda)} (\lambda_{\min} - \lambda)^2 > 0.$$

Thus, the iterates in the algorithm converge monotonically to λ_{\min} from the right and at a rate that is ultimately quadratic. Note from (2.8) that in the limit we have

$$(\text{error in new } \lambda) \approx C \cdot (\text{error in old } \lambda)^2$$

where

$$C = \frac{f''(\lambda_{\min})}{2f'(\lambda_{\min})} = \frac{w^*(G - \lambda_{\min} I)^{-1} w}{1 + w^* w}$$

and $w = -(G - \lambda_{\min} I)^{-1} r$. Since $\| (G - \lambda_{\min} I)^{-1} \|_2 = 1/d$ it is easy to show that $C \leq 1/d$. It follows that Algorithm 2.1 may converge slowly in problems where the separation d is small. We return to this point later.

The termination criteria in Algorithm 2.1 gives good absolute error in the final λ provided the tolerance δ is small enough. This follows from

$$\left\| \begin{bmatrix} \gamma & r^* \\ r & G \end{bmatrix} \begin{bmatrix} 1 \\ w \end{bmatrix} - \lambda \begin{bmatrix} 1 \\ w \end{bmatrix} \right\|_2 = \left\| \begin{bmatrix} f(\lambda) \\ 0 \end{bmatrix} \right\|_2 = |f(\lambda)|.$$

Applying standard Hermitian matrix perturbation theory (see [G83]) we may conclude that there exists an exact eigenvalue λ_e of T that satisfies

$$|\lambda - \lambda_e| < |f(\lambda)| (1 + \|w\|_2^2)^{\frac{1}{2}} < \delta$$

If δ is sufficiently small compared to the separation d , then one can ensure that $\lambda_e = \lambda_{\min}$.

Despite the nice mathematical properties of Algorithm 2.1, its practical value hinges on two critical factors: how is the starting value determined and how is the linear system $(G - \lambda I)w = -r$ to be solved? We address these questions in the next section for the case when T is symmetric, positive definite, and Toeplitz.

3. The Symmetric Positive Definite Toeplitz Case

Let $(t_0, t_1, \dots, t_{n-1})$ be the first row of a symmetric positive definite Toeplitz matrix $T = (t_{ij})$, i.e., $t_{ij} = t_{|i-j|}$. Assume that T is normalized so that $t_0 = 1$ and partition it as follows:

$$T = \begin{bmatrix} 1 & r^T \\ r & G \end{bmatrix} \quad r^T = (t_1, \dots, t_{n-1}) .$$

Recall that in order to apply Algorithm 2.1 we must find a starting value λ that belongs to the interval $[\lambda_n(T), \lambda_{n-1}(G)]$. This requirement can be couched in the language of signatures. The signature $\text{sig}(A)$ of a symmetric matrix A is a triplet of integers $(\text{neg}, z, \text{pos})$ where neg , z , and pos are the number of negative, zero and positive eigenvalues of A . Our starting value problem is to find λ such that $\text{sig}(G - \lambda I) = (0, 0, n-1)$ while $\text{sig}(T - \lambda I) = (1, 0, n-1)$ or $(0, 1, n-1)$.

This problem can be solved by exploiting the well-known Levinson-Durbin algorithm:

Algorithm 3.1

$$E_0 = 1$$

For $i = 1$ to $n-1$

$$\left[\begin{array}{l} k_i = -(t_i + \sum_{j=1}^{i-1} a_{i-1,j} t_j) / E_{i-1} \\ \text{For } j = 1 \text{ to } i-1 \\ \quad \left[\begin{array}{l} a_{ij} = a_{i-1,j} + k_i a_{i-1,i-j} \\ a_{ii} = k_i \\ E_i = E_{i-1} (1 - k_i^2) \end{array} \right. \end{array} \right.$$

The a_{ij} satisfy the Yule-Walker (YW) systems

$$\begin{bmatrix} 1 & t_1 & \cdot & \cdot & \cdot & t_{i-1} \\ t_1 & 1 & & & & \cdot \\ \cdot & & \cdot & & & \cdot \\ \cdot & & & \cdot & & \cdot \\ \cdot & & & & \cdot & \\ t_{i-1} & \cdot & \cdot & \cdot & & 1 \end{bmatrix} \begin{bmatrix} a_{i1} \\ \cdot \\ \cdot \\ \cdot \\ a_{ii} \end{bmatrix} = - \begin{bmatrix} t_1 \\ \cdot \\ \cdot \\ \cdot \\ t_i \end{bmatrix}$$

for $i = 1, \dots, n-1$. The quantities k_i and E_i are referred to as the i -th partial correlation coefficient and the i -th prediction error respectively. (k_i is also known as the i -th reflection coefficient.) See [G83] for a discussion of Algorithm 3.1.

In [C80] it is shown that if

Algorithm 3.2

```

i = 0
E0 = 1
Do While ( Ei > 0 & i < n-1 )
  i := i+1
  ki = -(ti + ∑j=1i-1 ajtj)/[(1-λ)Ei-1]
  [ a1 ]   [ a1 ]   [ ai-1 ]
  [ ⋮ ]     [ ⋮ ]     [ ⋮ ]
  [ ai-1 ] := [ ai-1 ] + ki [ a1 ]
  [ ai ]     [ 0 ]     [ 1 ]
  Ei = Ei-1(1 - ki2)

```

We have dropped the double subscripting of the a's since we need only be in possession of the most recent YW solution at any one time.

Note that if the loop terminates because $i = n-1$, then we have

$$(G - \lambda I) \begin{bmatrix} a_1 \\ \vdots \\ a_{n-1} \end{bmatrix} = - \begin{bmatrix} t_1 \\ \vdots \\ t_{n-1} \end{bmatrix} .$$

Recall that being able to solve this shifted YW system is critical to Algorithm 2.1, the Newton iteration for $f(\lambda)$.

Equally important, the final value of i in Algorithm 3.2 enables us to determine the position of λ with respect to λ_{\min} and $\lambda_{n-1}(G)$:

- (a) If $i = n-1$ and $E_{n-1} > 0$, then $\lambda < \lambda_{\min}$.
- (b) If $i = n-1$ and $E_{n-1} \leq 0$, then $\lambda_{\min} \leq \lambda < \lambda_{n-1}(G)$
- (c) If $i < n-1$ then $\lambda_{n-1}(G) \leq \lambda$.

Hence, Algorithm 3.2 can be used in a bisection scheme to eventually position λ in the interval $[\lambda_{\min}, \lambda_{n-1}(G)]$. Thereafter, it can be used to carry out the Newton iteration. All we need is an initial interval $[\alpha, \beta]$ with the property that

$$(3.2) \quad \alpha \leq \lambda_{\min} \leq \beta \cdot$$

Algorithm 3.3

Compute α and β satisfying (3.2) and let $\delta > 0$ be a given tolerance.

$k = 0$

$$\lambda^{(0)} = (\alpha + \beta) / 2$$

Do Until ($|\lambda^{(k)} - \lambda^{(k-1)}| \leq \delta |\lambda^{(k-1)}|$)

 Apply Algorithm 3.2 with $\lambda = \lambda^{(k)}$ to generate i and a_1, \dots, a_i .

$k := k + 1$

 If ($i < n - 1$)

 then

$$\beta = \lambda \ ; \ \lambda^{(k)} = (\alpha + \beta) / 2$$

 else

 If ($E_{n-1} > 0$)

 then

$$\alpha = \lambda \ ; \ \lambda^{(k)} = (\alpha + \beta) / 2$$

 else

$$\lambda^{(k)} = \lambda + \frac{1 - \lambda + t_1 a_1 + \dots + t_{n-1} a_{n-1}}{1 + a_1^2 + \dots + a_{n-1}^2}$$

The last expression for $\lambda^{(k)}$ above is same as the λ update expression in Algorithm 2.1 with $\gamma = 1$, $r^T = (t_1, \dots, t_{n-1})^T$ and $w^T = (a_1, \dots, a_{n-1})^T$.

There are several possible ways to choose the initial bracketing interval $[\alpha, \beta]$.

Method 1.

Set $[\alpha, \beta] = [0, 1]$. The choice for β follows from the inequality $\lambda_{\min} \leq e_1^T T e_1 = 1$ where $e_1 = (1, 0, \dots, 0)^T$.

Method 2.

Set $[\alpha, \beta] = [0, 1 - |t_1|]$. Since the smallest eigenvalue of

$$T_1 = \begin{bmatrix} 1 & t_1 \\ t_1 & 1 \end{bmatrix}$$

is given by $1 - |t_1|$, we have from separation theory that $1 - |t_1| = \lambda_2(T_1) \geq \lambda_n(T)$

Method 3.

Set $[\alpha, \beta] = [0, \min_i \{1 - |t_i|\}]$. The reasoning is the same as for Method 2 with t_1 replaced by t_i . Note that $\begin{bmatrix} 1 & t_i \\ t_i & 1 \end{bmatrix}$ is a principal submatrix of T .

Method 4.

Set $[\alpha, \beta] = [0, E_{n-2}(1 - |k_{n-1}|)]$ where E_{n-2} and k_{n-1} are generated by Algorithm 3.3 with $\lambda = 0$. To understand the choice for β , consider the effect of replacing t_{n-1} with $\tilde{t}_{n-1} = t_{n-1} + \epsilon$ in Algorithm 3.3 and that we set $\lambda = 0$. Nothing changes except during the last pass through the loop when we compute

$$\tilde{k}_{n-1} = -(\tilde{t}_{n-1} + \sum_{j=1}^{n-2} a_j t_j) / E_{n-2} = -\frac{\epsilon}{E_{n-2}} + k_{n-1}.$$

Note that if we choose ϵ so that $1 - \tilde{k}_{n-1}^2 = 0$, then the resulting \tilde{E}_{n-1} will be zero. Thus, a perturbation of size ϵ transforms T into a singular matrix. It follows that $\lambda_n(T) \leq \epsilon$. The choice for β is the smaller of the two ϵ values that render $\tilde{k}_{n-1}^2 = 1$

Method 5.

$$\text{Set } [\alpha, \beta] = \left[\frac{1}{\sqrt{n} \|T^{-1}\|_\infty}, \frac{\sqrt{n-1}}{\|G^{-1}\|_\infty} \right]. \text{ The value for } \alpha \text{ follows}$$

from the inequality

$$\frac{1}{\lambda_n(T)} = \|T^{-1}\|_2 \leq \sqrt{n} \|T^{-1}\|_\infty$$

Here, $\|\cdot\|_\infty$ denotes maximum row sum. The value for β follows from

$$\frac{1}{\lambda_n(T)} > \frac{1}{\lambda_{n-1}(G)} = \|G^{-1}\|_2 \geq \sqrt{n-1} \|G^{-1}\|_\infty$$

The quantities $\|T^{-1}\|_\infty$ and $\|G^{-1}\|_\infty$ can be calculated in $O(n^2)$ operations and $O(n)$ storage using the Trench algorithm [T64].

4. Analysis, Discussion, and Numerical Experiments.

Another method for finding λ_{\min} via the Levinson-Durbin algorithm is presented in [H80]. They propose solving $f(\lambda) = 0$ ($E_n(\mu) = 0$ in their notation) using a linear interpolation scheme. A key aspect of our work and what distinguishes it from [H80] is the recognition that one can apply Newton's method using by-products from the Levinson-Durbin algorithm. In addition, we have attempted to handle the problem of starting values more rigorously than [H80].

The importance of only having to solve Yule-Walker systems should be stressed. Methods based on inverse iteration, for example, require the solution of general Toeplitz systems. This doubles the amount of work per step. Moreover, there currently exist highly concurrent algorithms and VLSI architectures for solving Yule-Walker systems in $O(n)$ time. See [K83].

The total amount of work required by Algorithm 3.3 is determined by the number of YW systems that must be solved. The number N_B of bisection steps is bounded above by

$$N_B \leq -\log_2[d / (\beta - \alpha)] + 1 .$$

Note that during this phase of the algorithm, calls to Algorithm 3.2 do not require a full $n-1$ steps so it is a little hard to quantify the overall work. As a function of n , N_B appears to grow as $\log(n)$. A simple explanation of this is possible if we assume that the eigenvalues of T are uniformly distributed. In this case, the distance $\lambda_{n-1}(G) - \lambda_{\min}$ is roughly $1/n^2$. Hence the worst case limit on N_B is proportional to $\log(n)$.

The number of Newton steps N_N tends to be around 5 or 6 based on our experience with numerous examples a subset of which we now describe. For each $n = 11, 21, \dots, 91$ we generated 25 random positive definite symmetric Toeplitz matrices. These matrices had the form

$$T = \frac{1}{m} \sum_{k=1}^n w_k T_{2\pi\theta_k}$$

where n is the dimension, m is chosen so that T is normalized,

$$T_{\theta} = (t_{ij}) = (\cos(\theta(i-j))),$$

and the w_k and θ_k are uniformly distributed random numbers taken from $[0,1]$. It can be shown that T_θ is rank two, symmetric, semidefinite, and Toeplitz.

Table 1 summarizes the results of these experiments. Only initial interval Methods 2 and 4 were considered. Method 1 is unnecessarily crude while Methods 3 and 5 were too similar in performance to Methods 2 and 4 respectively for us to report. (Note: in tabulating the work associated with Method 4, the single call to Algorithm 3.2 necessary to compute β is accounted for in the Table.)

Order	Starting Values Via Method 2		Starting Values Via Method 4	
	Bisection Steps	Newton Steps	Bisection Steps	Newton Steps
11	4.8	5.0	5.7	4.8
21	8.5	5.7	4.8	5.4
31	9.7	5.3	6.6	5.0
41	10.0	4.6	6.7	5.2
51	11.7	5.8	8.1	5.5
61	12.1	5.0	9.2	5.3
71	12.0	5.3	9.2	5.2
81	13.6	5.4	9.8	5.0
91	11.6	5.0	8.4	5.7

Table 1. Behavior of Algorithm 3.3 ($\delta = 10^{-6}$) based on 25 random examples per dimension.

The matrices were generated by a Fortran program and the eigenvalues λ_{\min} and $\lambda_{n-1}(G)$ were computed by the EISPACK routine `rs` [S76]. Although our generation technique was guaranteed to generate at least a semi-definite matrix (definite with probability one) rounding errors led to the generation of some isolated slightly indefinite cases. Although indefinite matrices (due to finite arithmetic) ought to be expected in practice, they provide no realistic test for our procedure. In fact, if we know that our data has t significant bits (floating point) then more than t calls to the bisection step is useless. Given the quality of our data, after t steps of bisection we must conclude the the matrix is either not definite or that the condition $|\lambda_{n-1}(G) - \lambda_{\min}| < 2^{-t}$ holds and so to the precision of our data, $\lambda_n(G) = \lambda_{\min}$. This follows from standard eigenvalue perturbation arguments [P80].

While on the subject of small separations, it is important to point out that $1/d$ is the condition of λ_{\min} 's eigenvector. This is often the quantity of ultimate interest and so it is good to know that slow convergence in Algorithm 3.3 goes hand in hand with ill-conditioning in the minimum eigenvector.

The actual procedure was implemented in C on a DEC-10. Computations were done in the "double" data type. The tolerance δ in Algorithm 3.3 was set to 10^{-6} . The Newton iteration terminated successfully on all strictly separated trials and gave six significant digit agreement with EISPACK generated solutions.

REFERENCES

- [B80] R. Bitmead and B.D.O. Anderson, "Asymptotically fast solutions of Toeplitz and related systems of equations," *Lin.Alg.&Applic.*, 34 (1980), 103-116.
- [B82] R. Brent and F. Luk, "A systolic array for linear-time solution of Toeplitz systems of equations," Cornell Computer Science Technical Report TR82-526, 1982, Ithaca, New York 14853.
- [C80] G. Cybenko, "The numerical stability of the Levinson-Durbin algorithm for Toeplitz systems of equations," *SIAM J. Sci.&Stat. Comp.*, 1 (1980), 303-319.
- [C84a] G. Cybenko, "On the eigenstructure of Toeplitz matrices," *IEEE Trans. ASSP*, to appear.
- [C84b] G. Cybenko, "Computing Pisarenko Frequency estimates," Proceedings of the 1984 Princeton Conference on Information Systems & Sciences, March 14-16, 1984, to appear.
- [D82] T.S. Durrani and K.C. Sharman, "Extraction of an eigenvector oriented spectrum from the MESA coefficients," *IEEE Trans. ASSP*, 30 (1982), 649-651.
- [D83] P. Delsarte and Y. Genin, "Spectral properties of finite Toeplitz matrices," Proceedings of the 1983 mathematical theory of networks and systems, Beer-Sheva, Israel, 1983.
- [F83] D. Fuhrman and Bede Liu, "Approximating the eigenvectors of a symmetric Toeplitz matrix," 1983, preprint.
- [G83] G.H. Golub and C. Van Loan, Matrix Computations, Johns Hopkins University Press, Baltimore, Md., 1983.
- [H80] Y.H. Hu and S.Y. Kung, "Computation of the minimum eigenvalue of a Toeplitz matrix by the Levinson algorithm," Proceedings SPIE 25th International Conference (Real Time Signal Processing), San Diego, August 1980, 40-45.
- [H83a] Y.H. Hu and S.Y. Kung, "A Toeplitz eigensystem solver," 1983, preprint.
- [H83b] Y.H. Hu, "Resolution enhanced spectrum estimation via Toeplitz eigensystem solvers," Proceedings 22nd IEEE Conference on Decision and Control, San Antonio, TX, Dec. 14-16, 1983, pp. 1345-46.
- [H83c] Y.H. Hu and S.Y. Kung, "Highly concurrent Toeplitz eigensystem solver for high resolution spectral estimation," Proceedings of IEEE 1983 ICASSP, Boston, Mass., 1983, pp. 1422-1425.
- [K83] S.Y. Kung and Y.H. Hu, "A highly concurrent algorithm and architecture for solving Toeplitz systems," *IEEE Trans. on ASSP*, 31 (1983), pp.
- [P73] V.F. Pisarenko, "The retrieval of harmonics from a covariance function," *Geophysical J. Royal Astronomical Soc.*, 33 (1973), 347-366.

- [S76] B.T. Smith et al, Matrix Eigensystem Routines - EISPACK Guide, Springer-Verlag, New York, 1976.
- [T64] W.J. Trench, "An algorithm for the inversion of finite Toeplitz matrices," J. SIAM, 12, 1964, 515-522.
- [W65] J. Wilkinson, The algebraic eigenvalue problem, Oxford, New York, 1965.