



**Peer Reviewed**

**Title:**

A Conceptual Model of Sedimentation in the Sacramento–San Joaquin Delta

**Journal Issue:**

[San Francisco Estuary and Watershed Science, 10\(3\)](#)

**Author:**

[Schoellhamer, David H.](#), U.S. Geological Survey  
[Wright, Scott A.](#), U.S. Geological Survey  
[Drexler, Judy](#), U.S. Geological Survey

**Publication Date:**

2012

**Publication Info:**

San Francisco Estuary and Watershed Science, John Muir Institute of the Environment, UC Davis

**Permalink:**

<http://www.escholarship.org/uc/item/2652z8sq>

**Acknowledgements:**

We thank David Fryberg, Denise Reed, Stuart Siegel, Mark Stacey, the DRERIP review panel, and the other DRERIP model authors for their assistance in developing this model. Justin Ferris, Mick van der Wegen, and two anonymous reviewers provided helpful comments on initial drafts of this manuscript.

**Keywords:**

Sediment, sediment transport, sedimentation, Sacramento – San Joaquin River Delta, conceptual model, feedback, open water, tidal marsh, watershed, vegetation, sea level rise, Central Valley

**Local Identifier:**

jmie\_sfews\_11152

**Abstract:**

Sedimentation in the Sacramento–San Joaquin River Delta builds the Delta landscape, creates benthic and pelagic habitat, and transports sediment-associated contaminants. Here we present a conceptual model of sedimentation that includes submodels for river supply from the watershed to the Delta, regional transport within the Delta and seaward exchange, and local sedimentation in open water and marsh habitats. The model demonstrates feedback loops that affect the Delta ecosystem. Submerged and emergent marsh vegetation act as ecosystem engineers that can create a positive feedback loop by decreasing suspended sediment, increasing water column



light, which in turn enables more vegetation. Sea-level rise in open water is partially countered by a negative feedback loop that increases deposition if there is a net decrease in hydrodynamic energy. Manipulation of regional sediment transport is probably the most feasible method to control suspended sediment and thus turbidity. The conceptual model is used to identify information gaps that need to be filled to develop an accurate sediment transport model.

**Copyright Information:**



Copyright 2012 by the article author(s). This work is made available under the terms of the Creative Commons Attribution 3.0 license, <http://creativecommons.org/licenses/by/3.0/>



**eScholarship**  
University of California

eScholarship provides open access, scholarly publishing services to the University of California and delivers a dynamic research platform to scholars worldwide.

# Conceptual Model of Sedimentation in the Sacramento–San Joaquin River Delta

David H. Schoellhamer<sup>1†</sup>, Scott A. Wright<sup>1</sup>, and Judith Z. Drexler<sup>1</sup>

## ABSTRACT

Sedimentation in the Sacramento–San Joaquin River Delta builds the Delta landscape, creates benthic and pelagic habitat, and transports sediment-associated contaminants. Here we present a conceptual model of sedimentation that includes submodels for river supply from the watershed to the Delta, regional transport within the Delta and seaward exchange, and local sedimentation in open water and marsh habitats. The model demonstrates feedback loops that affect the Delta ecosystem. Submerged and emergent marsh vegetation act as ecosystem engineers that can create a positive feedback loop by decreasing suspended sediment, increasing water column light, which in turn enables more vegetation. Sea-level rise in open water is partially countered by a negative feedback loop that increases deposition if there is a net decrease in hydrodynamic energy. Manipulation of regional sediment transport is probably the most feasible method to control suspended sediment and thus turbidity. The conceptual model is used to identify information gaps that need to be filled to develop an accurate sediment transport model.

## KEY WORDS

Sediment, sediment transport, sedimentation, Sacramento–San Joaquin River Delta, conceptual model, feedback, open water, tidal marsh, watershed, vegetation, sea-level rise, Central Valley, Delta Regional Ecosystem Restoration Implementation Plan (DRERIP).

## INTRODUCTION


The Sacramento–San Joaquin River Delta (Delta) is where the rivers of the Central Valley of California merge to become the San Francisco Estuary (Figure 1). The rivers deliver sediment from the Central Valley watershed to the Delta. A sedimentation model was included in the Delta Regional Ecosystem Restoration Implementation Plan (DRERIP) suite of conceptual models because sediment deposition creates and sustains the Delta landscape, pelagic habitat depends on suspended sediment, and sediment transports adsorbed nutrients and contaminants.

Deposited sediment creates and sustains the Delta landscape, including habitats such as tidal marsh, floodplain, open channels, and flooded islands. Massive sediment supply during the period of hydraulic mining in the late 1800s caused deposition in Sacramento Valley rivers, the Delta, and San Francisco Bay (Gilbert 1917). Today, a key manage-

<sup>1</sup> California Water Science Center, U.S. Geological Survey, Placer Hall, 6000 J Street, Sacramento, CA 95819 USA.

† Corresponding author: [dschoell@usgs.gov](mailto:dschoell@usgs.gov).

**SAN FRANCISCO ESTUARY & WATERSHED SCIENCE**



**Figure 1** Sacramento-San Joaquin River Delta. Locations in red correspond to the sediment fluxes and budget presented in Figures 3 and 6.

ment question is whether the existing Delta landscape can be sustained as sea level rises. Sea-level rise and floodplain, marshplain, and channel-form changes are habitat stressors, which can be counteracted by artificial and natural movement of sediment. The preferred option is natural sediment movement because it is usually less costly and more sustainable. Sediment is the raw material for habitat restoration projects and levee construction. Sediment also deposits in ports, marinas, and shipping channels, which sometimes require dredging to maintain navigation.

Suspended sediment affects habitat for pelagic organisms. Suspended sediment is the primary attenuator of sunlight in the water column of the Delta which, in turn, limits photosynthesis and primary photosynthetic carbon production (Cloern 1987; Jassby and others 2002). Abundance of some fish species increases in more turbid waters (Nobriga and others 2005; Feyrer and others 2007). A decline in fish abundance in the 2000s, locally called the pelagic organism decline, has had the most serious consequences for delta smelt (*Hypomesus transpacificus*), for which larval feeding sharply decreases when turbidity is less than 18 nephelometric turbidity units (NTU) (Baskerville–Bridges and others 2004), which is roughly equivalent to a suspended-sediment concentration (SSC) of 24 mg L<sup>-1</sup> (Ganju and others 2007). Water exports from the Delta, which provide drinking and irrigation water for much of California, have been limited because of the pelagic organism decline.

Many contaminants are associated with sediment (Turner and Millward 2002; Schoellhamer and others 2007a; Luengen and Flegal 2009). Suspended sediment moving into, within, and out of estuaries provides a pathway for the transport of sediment-associated contaminants (Turner and others 1999; Turner and Millward 2000; Bergamaschi and others 2001; Le Roux and others 2001). Thus, the fate of these substances is largely determined by the fate of sediment. The San Francisco Estuary is an impaired water body because of several sediment-associated contaminants (Schoellhamer and others 2007a).

In this paper, we summarize a conceptual model of sedimentation in the Delta (Schoellhamer and others 2007b). We discuss critical feedback loops, habitat restoration, uncertainty, and numerical modeling.

## Study Area

The Sacramento River drains the northern part of the Central Valley, an area of approximately 60,900 km<sup>2</sup>. The San Joaquin River drains approximately 35,060 km<sup>2</sup> in the southern Central Valley. The Cosumnes and Mokelumne rivers enter the Delta directly from the east, draining areas of approximately 1900 and 1700 km<sup>2</sup>, respectively. River discharge is greatest during winter and spring, and smallest during the dry summer and early autumn. Flood bypasses were constructed for Central Valley rivers in the early 20th century (Singer and others 2008). Dams were built in most Central Valley rivers in the mid-20th century (Singer 2007). The Sacramento River was stabilized with rip rap mostly during the latter half of the 20th century (USFWS 2000; Florshiem and others 2008). Tides propagate into most of the Delta when river discharge is small. Suisun Bay is the sub-embayment of San Francisco Bay that is seaward of the Delta. Tides in Suisun Bay are mixed diurnal and semidiurnal, and the tidal range varies from about 0.6 m during the weakest neap tides to 1.8 m during the strongest spring tides. Most of the waters of the Delta are fresh and during the dry season flow from reservoirs is managed to try to maintain a salinity of 2 parts per thousand (ppt) in Suisun Bay.

At the confluence of the four rivers, a complex network of natural and man-made channels has developed (Figure 1). The *Sacramento–San Joaquin Delta Atlas* (DWR 1995) contains detailed information on the history of the Delta; a brief summary is provided here. Levee construction and draining of marshlands began in late 1850. As a result, the Delta today consists of a network of slough channels that surround former marshlands commonly termed “islands,” which are primarily used for agriculture. Because of this channelization, only 0.02 km<sup>2</sup> of non-vegetated tidal flats exist in the Delta today (DFG 1997). Before channelization, the top soil in the Delta was a peat layer 2 to 15 m thick (Atwater and Belnap 1980). The organic material in peat contains plant litter, root biomass, as well as allochthonous material from the watershed. Carbon sequestration in the Delta ranges from 0.38 to 0.79 Mg ha<sup>-1</sup> yr<sup>-1</sup> (Drexler 2011). Because of the high organic content of Delta soils,



draining of marshes has resulted in significant land subsidence, such that most of the islands are currently below mean sea level, some by as much as 4 m. During the latter half of the 20th century there was a large increase in water exports from the Delta for urban and agricultural use primarily south of the Delta.

### Delta Sediment Properties

Important surficial sediment properties in the Delta are size, density, and organic content. Sediment in channels that convey relatively large flows, such as the Sacramento River, San Joaquin River, and Threemile Slough, tend to be sandy and contain bed forms (Dinehart 2002). The relative quantity of fine sediment generally increases seaward. From 1993 to 2003, 18 surficial bed samples collected by the Regional Monitoring Program (RMP) in the lower Sacramento River had a mean of 19% fines and a range of 8% to 50% (RMP [cited 2007]). Eighteen samples from the lower San Joaquin River had a mean of 48% fines and a range of 16% to 79%. Larger river floods that winnow fines from the bed and larger sand supply in the Sacramento River probably account for the difference between the two rivers. At Rio Vista, Thompson and others (2000) observed that large floods increased the percent of sand on the bed (up to nearly 100% from nearly 0%), and benthic assemblages shifted from species that favor muddy sediment to sandy sediment. During the intervals between floods, finer sediment deposited, the bed sediment became finer, and benthos that prefer muddy sediment returned. At two other sites in the Sacramento and San Joaquin rivers, however, the fraction of sand varied over a similar range but appeared unrelated to flow, perhaps because of spatial heterogeneity. Fine sediments tend to deposit on marshes, which typically have large expanses of emergent macrophytes (Byrne and others 2001). Grain size affects substrate stability and establishment and growth of submerged aquatic vegetation.


The dry density of bed sediment is the mass of dry sediment per unit volume and it generally increases as grain size increases and organic content decreases. A survey of studies shows that dry density of bed

sediment can vary by a factor of two. Porterfield (1980) estimated that the dry density of estuarine sediments was  $851 \text{ kg m}^{-3}$ . Ogden Beeman & Associates and Krone & Associates (1992) estimated a value of  $529 \text{ kg m}^{-3}$ . Caffrey (1995) measured a dry density of  $1,144 \text{ kg m}^{-3}$  for the top centimeter of bed sediment in the channel at Rio Vista.

Organic content of bed sediment is greatest in marshes. Reed (2002) found organic contents of wetland soils ranged from 7.80 to 39.38%. From 1993 to 2003, 17 bed samples from the lower Sacramento River had a mean of 0.55% total organic carbon (TOC) and a range of 0.14% to 2.10% (RMP [cited 2007]). Seventeen samples from the lower San Joaquin River had a mean of 0.68% TOC and a range of 0.26% to 1.38%. At both sites, TOC generally increased as the fraction of fine sediment increased.

In this paper, we use SSC to quantify the mass of sediment in the water column, expressed as mass of suspended sediment per unit volume. Typical SSC in the Delta ranges from 10 to  $50 \text{ mg L}^{-1}$ , except during large river discharge when SSC can exceed  $200 \text{ mg L}^{-1}$  (Wright and Schoellhamer 2005; McKee and others 2006). In the San Francisco Estuary, concentrations reported as total suspended solids or suspended particulate matter are equivalent to SSC (Gray and others 2000). Turbidity and Secchi depth are other measures of the light characteristics of Delta waters. Turbidity is an optical measure of light scattering in water with units of nephelometric turbidity (NTU). As the number of fine particles in suspension increases, SSC and turbidity generally increase. Because a given mass of fine sediment is a more effective scatterer than the same mass of coarse sediment, turbidity and SSC are not necessarily correlated. In the San Francisco Estuary, however, suspended sediment is predominantly fine sediment and floc sizes are spatially homogeneous, so turbidity and SSC are well correlated (Ganju and others 2007; Figure 2). Lowering a Secchi disk into water until it reaches a depth at which it is no longer visible provides another common measurement of light penetration in the water column.

Other properties of suspended sediment that are of interest include particle diameter, floc size, density,



**Figure 2** Comparison of turbidity and SSC measurements, from four locations in San Francisco Bay. Source: Ganju and others 2007.

and organic content. Suspended sediment is primarily fine sediment less than  $63\ \mu\text{m}$  in diameter. At Freeport on the Sacramento River the median percent of fine suspended sediment was 85% from July 1998 to September 2001, and the range was 46% to 98% (Schoellhamer and Wright 2003). Fine suspended sediment is cohesive, and small primary particles combine to form larger flocs of suspended sediment. In other systems, fine sediment in rivers is flocculated (Droppo and others 1997), and limited measurements from the Sacramento River at Freeport (Ganju, unpublished data) confirm this. Ganju and others (2007) estimate that the primary particle diameter is  $2.5\ \mu\text{m}$ . During a slack after ebb tide at Rio Vista, the median volumetric floc diameter ( $D_{50}$ ) increased from  $20\ \mu\text{m}$  near the water surface to  $80\ \mu\text{m}$  near the bed as a result of the settling of larger flocs in the water column (Ganju and others 2007). During maximum flood tide,  $D_{50}$  was 45 to  $65\ \mu\text{m}$  and more uniform in the water column. At slack after flood tide,  $D_{50}$  ranged from 40 to  $110\ \mu\text{m}$  increasing with depth. Salinity varied from 0 to 1.6 psu.

Schemel and others (1996) found that the particulate organic carbon (POC) content of suspended sediment was inversely related to flow. During floods in 1983 and 1984, POC was 1% to 2% of the suspended sediment at Rio Vista. During periods of low flow,

POC was higher, mostly 2% to 4% of the suspended sediment at Rio Vista. Algal production probably accounted for higher POC in summer when flows were low.

## MODEL DESCRIPTION

The conceptual model is qualitative and is composed of drivers, linkages, and outcomes. First, the scope of the model is described, specifically the treatment of suspended and bed load and time scales. The model includes submodels for river supply, regional transport, and local sedimentation in open water and marsh habitats.

### Suspended and Bed Load

Sand and coarser sediment (diameter larger than  $63\ \mu\text{m}$ ) not only move in suspension but also can move along the bed by rolling, sliding, and jumping, which is called bed load. Based on measurements in the late 1950s, Porterfield (1980) estimated that the bed load was 109 metric tons per day in the Sacramento River at Sacramento. This bed load was only 1.4% of the total sediment load. From 1998 to 2000, Dinehart (2002) collected several pairs of bedform measurements about one week apart to estimate bedload transport rates of 15 to 73 metric tons per day at Garcia Bend downstream from Sacramento. Wright and Schoellhamer (2005) report that the daily suspended-sediment load at Freeport, about 6 kilometers downstream from Garcia Bend, averaged about 3,000 metric tons per day during water years 1999 to 2002. Although not temporally aligned, these more recent measurements, when compared, appear to confirm as still valid, Porterfield's finding that suspended load is about 2 orders of magnitude greater than bed load. Because bedload is a small fraction of suspended load, this conceptual model neglects the mass of sediment transported as bed load. Here, we consider how bed load affects the size and properties of bed sediment, however, because it can affect erosion. Suspended load in the model includes sand and finer particles.

## Time Scale

The time scale of our conceptual model is tidally-averaged. Sediment deposits and re-suspends during flood and ebb tides at the semidiurnal tidal time scale. For example, suspended-sediment deposits at slack tide when water velocity and turbulence are small, and bottom sediment is re-suspended when tidal currents and thus shear stresses increase. In addition, tidal currents are stronger and resuspension is more likely during fortnightly spring tides compared to neap tides. By integrating deposition and erosion over many tidal cycles, a tidally-averaged rate of deposition or erosion can be calculated. Thus, for the purposes of this conceptual model, while some of the drivers of sediment transport occur at tidal time scales, a tidally-averaged outcome results. Geomorphic change takes place over years and decades, and is estimated by summing deposition and erosion of the tidally-averaged model over time. While we are not attempting to resolve the tidal time scale, the model should, nonetheless, be applicable to tidal time scales.

Nontidal sedimentation processes are largely episodic. For example, rivers supply most sediment to the Delta during large floods over only a few days per year. This episodic nature is driven largely by sediment pulses from the Sacramento River that deposit sediment in the Delta and move into Suisun Bay (Wright and Schoellhamer 2005; [Figure 3](#)). During water years 1999 to 2002, 82% of the sediment was delivered during the wet period (31% of the time) (Wright and Schoellhamer 2005). McKee and others (2006) found that for sediment supplied to Suisun Bay from the Delta: (1) a large flood in January 1997 transported 11% of the sediment supplied from 1995 to 2003; (2) 88% of the annual sediment supply occurred during the wet season; and (3) 43% of the annual sediment supply occurred during the wettest 30-day period. Another episodic forcing is wind waves and associated sediment resuspension in shallow water generated by storms (Ruhl and Schoellhamer 2004).

## Spatial Sub-Models

Our conceptual model includes linked submodels at three different spatial scales. First, a conceptual


submodel of riverine sediment supply considers how relevant processes in the watershed affect sediment supply to the Delta. Second, a regional transport submodel transports sediment from the rivers to the Delta, within the Delta itself, and between the Delta and Suisun Bay. Advection and dispersion move suspended sediment horizontally, and deposition and erosion occur along the transport pathway. Finally, we describe submodels of local erosion and deposition for open water, marsh, floodplain, and riparian habitats. For brevity, only the open water and marsh submodels are presented in this paper. Floodplain and riparian models are available from Schoellhamer and others (2007b).

## River Supply Sub-Model

The primary outcome of the river supply submodel ([Figure 4](#)) is the amount of suspended sediment that enters the Delta from river sources. Several studies (e.g. Porterfield 1980; Wright and Schoellhamer 2005) have documented how the Sacramento River dominates in contributing sediment to the Delta. The structure of this conceptual submodel, however, is such that it could be used to describe the sediment supply from any of the Central Valley watersheds. This river supply submodel is not intended to be a complete watershed submodel.

The amount of water and sediment delivered to a watershed outlet, under natural conditions, is a complex function of the watershed's climate, geology, topography, and vegetation. For the watersheds that drain into the Sacramento–San Joaquin Delta, precipitation comes primarily during winter months as rain and snow. Hydrologic processes acting throughout the watershed (e.g., snowmelt, evapotranspiration, infiltration, etc.) then determine the amount and timing of water (i.e., flow regime) that reaches the watershed outlet. The flow regime of the river (which sets the transport capacity, i.e., the amount of sediment that could be transported if the supply were unlimited) and the supply of sediment available from the landscape determines the amount of sediment that reaches the watershed outlet. For the size of sediment that dominates the yield to the Delta (finer than 63  $\mu\text{m}$ , or silt and clay), it is likely that the






**Figure 3** Tidally-averaged suspended sediment flux in the Delta, 1998–2002. Arrows indicate downstream/down estuary flow paths for the Sacramento (left) and San Joaquin (right) rivers. The vertical scale for the Sacramento River flow path is larger than that for the San Joaquin River flow path and Threemile Slough (TMS). Sacramento River at Freeport (FPT) and Rio Vista (RVS), San Joaquin River at Vernalis (VNS), Stockton (STN), and Jersey Point (JPT), and Mallard Island (MAL) are also shown. Source: Wright and Schoellhamer 2005.

available supply limits sediment transport volumes. Observations of seasonal variability in the relationship between sediment concentration and flow, where sediment concentrations are higher for the same flow during “first flush” events and lower during spring snowmelt events (Goodwin and Denton 1991; Schemel and others 1996; Curtis and others 2006) support the conclusion that fine sediment is supply limited. Thus, human activities that alter watershed sediment supply are likely to have a greater effect on river supply to the Delta than those that modify the flow regime (most activities influence both). In accord with this philosophy, we have not included

two anthropogenic drivers that primarily affect flow regime: climate change and consumptive water use.

Since the discovery of gold in 1848, the watersheds that drain to the Sacramento–San Joaquin Delta have been heavily affected by human activities (Gilbert 1917; James 1991; Singer and others 2008). In general, human activities tend to increase the amount of sediment transported in rivers through soil erosion, but this increase can be offset by sediment retention in reservoirs, leading Syvitski and others (2005) to conclude that the worldwide flux of terrestrial sediment to the oceans has decreased from prehuman



**Figure 4** River Supply Sub-model. Each box is a driver and each arrow is a linkage. River supply is a driver of the regional and local sedimentation submodels.

conditions. A detailed accounting of the changes to the landscape that have occurred in the Central Valley of California is far beyond the scope this conceptual model. Rather, we focus on the more significant changes, with respect to sediment supply to the Delta, and describe conceptually how each of these changes can affect sediment transport. Many of the anthropogenic drivers are interrelated. For example, as climate change alters the volume and timing of water runoff, this may in turn affect how reservoirs are operated.

Probably the most significant anthropogenic driver for river sediment supply is hydraulic mining. During the late 1800s and early 1900s, large deposits were washed into flumes by high-powered water jets in order to separate out the gold. The mine tailings were routed into the rivers, which dramatically increased the sediment supply. Gilbert (1917) estimated a 9-fold increase in sediment supply to San Francisco Bay

during the mining period. Though the primary pulse of mining sediment has moved through the watershed, remnant terrace deposits remain (Meade 1982; James 1991). Also, recent estimates of river sediment supply to the Delta are substantially higher than Gilbert's pre-mining estimate, but have continued to decrease since the mid-1950s potentially indicating continued exhaustion of remnant mining-derived deposits (Wright and Schoellhamer 2004). Thus, although hydraulic mining has stopped and does not seem likely to occur in the future, its legacy may still be affecting river sediment supply and thus should be included in the conceptual model to assess future scenarios.

Two major water supply projects have been constructed in the watersheds that drain to the Delta, the Central Valley Project and the State Water Project, with each project containing multiple large dams and reservoirs. Dams have also been constructed for other purposes, such as trapping hydraulic min-

ing sediments. The U.S. Army Corps of Engineers National Inventory of Dams ([http://geo.usace.army.mil/pgis/f?p=397:3:0::NO::P3\\_STATES:CA](http://geo.usace.army.mil/pgis/f?p=397:3:0::NO::P3_STATES:CA)) contains 1,468 dams in California (see web page for inclusion criteria) and Nilsson and others (2005), in their recent study of flow regulation of the world's large river systems, classified the Sacramento–San Joaquin basin as “strongly affected” by dams. Dams primarily affect the reservoir by retaining sediment in it; the channel immediately downstream from the dam will be sediment-deficient and erode to a new equilibrium (Porterfield and others 1978). While this provides a short-term sediment source, the long-term effect is decreased sediment supply (Williams and Wolman 1984). Dams also affect the flow regime, typically reducing high flows and increasing low flows (Singer 2007), which also reduces downstream sediment supply. It follows that dam removal (compared to keeping the dam in place) would increase downstream sediment supply by making reservoir sediment deposits available (short-term) and by no longer retaining incoming sediment (long-term).

A major system of levees and bypass channels has been constructed in the basin to reduce flood hazards, particularly in the lower reaches of the Sacramento River watershed (Kelley 1998). Before this flood-control system, the Sacramento River would overflow its banks and fill vast flood basins for significant periods of time during wet years. More than half the banks of the lower Sacramento River were ripped during the latter half of the 20th century (USFWS 2000; Florsheim and others 2008). While bank protection would tend to reduce sediment supply from the banks of the river, levees tend to confine flood flows, potentially resulting in erosion of the channel bed and increased sediment supply. The levees also serve to isolate the flood basins, which were likely depositional environments during floods, keeping the flood flows and suspended sediment in the channel. However, flood-control bypasses built in the Sacramento River floodplain during the early 20th century (e.g., Yolo Bypass) provide a similar function as the flood basins because they trap substantial amounts of sediment (Singer and others 2008). Thus, the various flood-control measures that have been implemented affect sediment supply in dif-


ferent ways, and these processes have not been quantified such that a net effect can be discerned.

Much of the Central Valley has been transformed to agricultural and urban land uses. Logging has also taken place in many of the watersheds. As stated previously, human transformation of the landscape typically results in increased soil erosion (Syvitski and others 2005), thus increasing river sediment supply. Though dams and reservoirs may counteract this effect, many of the land-use changes, particularly by agriculture and urbanization, have taken place downstream from the major dams in the system. Thus, these changes are likely to have increased sediment supply to the Delta; even in the early 1900's Gilbert (1917) estimated substantial increases in sediment supply from human activities other than mining. Today, erosion-control practices are often used to minimize these effects.

There has been no comprehensive study of the anthropogenic drivers of sediment supply, so determination of their relative importance is difficult. Wright and Schoellhamer (2004) showed that the amount of sediment trapped annually in just Oroville, Folsom, and Englebright dams in the Sacramento River watershed ( $2.4 \text{ Mm}^3 \text{ yr}^{-1}$ ) is of the same order of magnitude as the annual watershed sediment supply, indicating that dams are significantly affecting the supply. Singer and others (2008) calculated sediment-deposition rates in flood bypasses that total  $1.3 \text{ Mm}^3 \text{ yr}^{-1}$ , about one-half the deposition rate in the three reservoirs. It is more difficult to similarly quantify the effects of the other anthropogenic effects (land-use changes, levee protection), and the studies required to do this have not yet been conducted. However, given the extensive land-use changes that have occurred in the Central Valley, and the understanding that these changes generally result in accelerated erosion (ASCE 1975; Syvitski and others 2005), it seems likely that land-use changes have been an important driver of river sediment supply.

### Regional Transport Submodel

The Delta is where the rivers that drain the Central Valley merge and become an estuary. The regional transport submodel (Figure 5) transports sediment



**Figure 5** Regional submodel of sediment transport in the Delta. Line thickness indicates the approximate importance of external sediment supplies (see Figure 6). The Delta exchanges sediment with Suisun Bay.

from the rivers into the Delta. The regional model also transports sediment from the Delta to Suisun Bay. It is also possible to transport sediment from Suisun Bay landward into the Delta because of the complex hydrodynamics in Suisun Bay and the western Delta (see below).


For the regional transport submodel, the Delta is represented as a triangle (Figure 5). The northeast (upper right) apex is where the Sacramento River enters the Delta, and the southeast (lower right) apex is where the San Joaquin River enters. For convenience, these apexes are considered at the U.S. Geological Survey (USGS) sediment gages at Freeport on the Sacramento River and Vernalis on the San Joaquin River. The Yolo Bypass diverts high Sacramento River flows around the city of Sacramento to the Delta, and is shown as an arrow entering the northwest side. The Mokelumne and Cosumnes rivers enter on the east side. The western (left) apex is the boundary between

Suisun Bay and the Delta located at the USGS continuous suspended-sediment monitoring station at Mallard Island.

Several rivers supply sediment to the Delta, primarily the Sacramento River. During water years 1999 to 2002 the sediment discharge at Freeport (FPT) on the Sacramento River was 5 times greater than the San Joaquin River at Vernalis (Wright and Schoellhamer 2005; Figure 6, VNS). Sediment discharge at Freeport decreased by about one-half from 1957 to 2001 (Wright and Schoellhamer 2004). Reduced sediment supply reduces sediment deposition in the Delta; deposition rates were 4 to 8 times greater from 1944 to 1972 than 1972 to 2005 (Canuel and others 2009). In addition, total suspended solids concentrations in the Delta decreased 50% from 1975 to 1995 (Jassby and others 2002). The second largest source of sediment was the Yolo Bypass (YOL, 28% of the sediment discharge at Freeport). The east side tributaries (EAST)



**Figure 6** Average annual Delta sediment budget based on water years 1999–2002, except for Threemile Slough (TMS), which is based on water years 2001 and 2002 only. Numbers are the annual suspended-sediment flux and the estimated error in thousand metric tons. Arrow thickness indicates relative magnitude of the suspended-sediment flux. Sediment deposition accounts for the decreased sediment fluxes from east to west. Additional sites are Sacramento River at Freeport (FPT), Yolo Bypass (YOL), Delta Cross Channel (DCC), Sacramento River at Rio Vista (RVS), Mallard Island (MAL), Eastside tributaries (EAST), San Joaquin River at Vernalis (VNS), San Joaquin River at Stockton (STN), exports from the State and Federal water projects (EXP), Dutch Slough (DCH), and San Joaquin River at Jersey Point (JPT). Source: Wright and Schoellhamer 2005.



supplied only 3.3% of the sediment discharge at Freeport. Most sediment is supplied during high flows (Schemel and others 1996; Wright and Schoellhamer 2005; McKee and others 2006).

State and federal water projects export water from the southern Delta (Figure 6, EXP). Wright and Schoellhamer (2005) used sediment deposition volumes in Clifton Court Forebay to estimate that exported sediment was about 2% of the sediment discharge at Freeport.

Tidally-averaged sediment transport is usually from the Delta into Suisun Bay. For water years 1999 to 2002, Mallard Island suspended sediment flux was seaward, and 51% of the Freeport sediment discharge (Wright and Schoellhamer 2005). On a daily time scale, which is roughly tidally-averaged, McKee and others (2006) found that sediment transport was landward on 9 of 198 days for which data were available. Occasional net sediment transport from Suisun Bay into the Delta can be caused by small river flows, gravitational circulation, meteorologically-forced tidally-averaged flow into the Delta (Tobin and others 1995), and tidal asymmetries

such as greater bottom shear stress during flood tide (Brennan and others 2002) and higher concentrations in Suisun Bay from wind-wave re-suspension (Ruhl and Schoellhamer 2004) or a turbidity maximum (Schoellhamer 2001). Landward sediment transport is approximately 11% of the seaward sediment transport during high flows and 52% during low flows (McKee and others 2006). On a tidal time scale, flood tides transport sediment from Suisun Bay into the Delta and ebb tides reverse sediment transport. Tides thus exchange and mix suspended sediment between Suisun Bay and the Delta.

Water movement within the Delta transports suspended sediment horizontally, which is represented with gray arrows in the regional submodel (Figure 5). Horizontal transport has two components we will consider: advection is the down current displacement of suspended sediment; dispersion is the mixing caused by turbulence and large eddies. For example, a pulse of sediment delivered from a river will oscillate upstream and downstream because of flood and ebb tides resulting from advection, move closer to the bay after each tidal cycle due to tidal averaging of



the downstream river flow, and spread out as a result of dispersion.

Suspended sediment has an additional complication because the bed acts as a source because of, erosion or sink because of deposition, represented by blue arrows in the regional submodel (Figure 5). For example, to determine the quantity of sediment from the Sacramento River delivered to a point in the Delta, horizontal transport and deposition and erosion along the transport pathway must be considered. On the tidal time scale, deposition is more likely to occur near slack tides when water velocity and turbulence are small, and erosion is more likely to occur during strong tides when water velocity and turbulence are greatest. On a time scale of years, Wright and Schoellhamer (2005) found that two-thirds of the sediment that entered the Delta was deposited in the Delta during water years 1999 to 2002.

Regional spatial variability of suspended sediment within the Delta is dominated by supply from the Sacramento River. The Sacramento River is the primary pathway for sediment transport (Wright and Schoellhamer 2005; Figure 6). At least 82% of the sediment entering the Delta from the Sacramento River watershed either deposits along the Sacramento River or moves past Mallard Island into Suisun Bay. No more than 18% moves into the San Joaquin River portion of the Delta. The suspended-sediment signal of the San Joaquin River attenuates more rapidly than that of the Sacramento River (Wright and Schoellhamer 2005; Figure 3).

### **Local Sedimentation Sub-Models**

Local processes determine how fast sediment erodes from the bed and how fast suspended sediments settle onto the bed. As a parcel of water moves from the river into the Delta, it tidally oscillates within the Delta, and ultimately exits the Delta. Erosion, deposition, and dispersion determine the suspended-sediment concentration (SSC) of the parcel and the properties of the suspended sediment (primarily size, density, organic content, and settling velocity). At a fixed point in the Delta, the time history of erosion and deposition determine the properties of the bed material. Tidally-averaged rates of deposition and


erosion are integrated over decades to determine the geomorphic evolution of a fixed point in the Delta.

Local sedimentation processes will differ in different habitats. In this paper, we present local sedimentation submodels for open water and marsh habitats. Schoellhamer and others (2007b) also present local sedimentation submodels for floodplain and riparian habitats. Each local sedimentation submodel contains the same drivers and outcomes; only the importance, qualitative understanding, and quantitative predictability of the linkages differ.

The open water submodel (Figure 7) includes channels and flooded islands. Delta channels provide the pathways for sediment from the watershed to move through the Delta and enter Suisun Bay (Wright and Schoellhamer 2005).

The marsh submodel for sedimentation (Figure 8) differs from the open water submodel in several ways. Vegetation has a stronger influence on sedimentation in a marsh compared to open water because it dampens hydrodynamic energy, which favors deposition over erosion. Thus, the marsh submodel shows depositional linkages being more important than erosional linkages. Vegetation can also provide shading, so the effect of suspended sediment on water column light is less important than open water. Dredging, filling, and armoring now rarely occur in marshes so we consider them inconsequential. A common goal of restoration projects is to encourage marsh vegetation, so we consider the linkage between restoration and vegetation to be highly important. We do not consider eposition of leaf litter in this submodel.

Most of the drivers are fundamental physical processes that are always present in all habitats and the linkages between these drivers are generally important. Some drivers may or may not be present at a given location (i.e. restoration, dredging, fill, vegetation) and are irrelevant if absent. Because the drivers are linked by fundamental physical laws, the conceptual linkages between drivers are well understood. This includes the direction of change, i.e., increased bed stress will erode, not deposit, bottom sediment. Quantitative prediction of the linkages, however, is generally very difficult. This is especially true for erosion. Marshes tend to be depositional, so our limi-




**Figure 7** Local sedimentation submodel for open water. Each box is a driver and each arrow is a linkage. An increase in a driver either increases (+) or decreases (-) the intermediate outcomes it affects. In addition to being outcomes, purple boxes are also drivers. In addition to being drivers, several green boxes are also intermediate outcomes. The regional submodel is shown in Figure 5.

tations on quantitative prediction of erosion are less daunting.

Water in the Delta moves primarily because of tides, waves, river flows, and pumping. Moving water exerts shear on sediment particles in the bed. Turbulence that causes vertical mixing is also generated. In general, faster water generates more shear stress and turbulence. Larger flows from rivers and stronger tides (i.e. spring tides) increase shear stress and turbulence (Brennan and others 2002). Waves

with relatively large height and period will apply greater shear to the bed and increase erosion (Ruhl and Schoellhamer 2004). Winds are strongest and waves largest during spring and summer. Waves generated by boats can also induce erosion from the bed and bank. Bauer and others (2002) found that levees in Georgiana Slough eroded 0.01 to 0.22 mm per boat passage.

Local erosion and deposition is a key driver that affects all model outcomes (SSC and properties, water



**Figure 8** Local sedimentation submodel for marsh. Each box is a driver and each arrow is a linkage. An increase in a driver either increases (+) or decreases (-) the intermediate outcomes it affects. In addition to being outcomes, purple boxes are also drivers. In addition to being drivers, several green boxes are also intermediate outcomes. The regional submodel is shown in Figure 5.

column light, bed particle size and properties, and geomorphic change). Erosion and deposition alter SSC and properties. Net erosion increases SSC, and net deposition decreases SSC. For non-cohesive sediment, erosion of coarser material increases the average settling velocity of suspended sediment. Coarser material settles faster than finer material, so settling velocity will decrease as deposition occurs.

Model outcomes of SSC and properties and bed particle size and properties are also drivers of local erosion and deposition. Bed shear stress and verti-

cal mixing are additional drivers that affect local erosion and deposition. These hydrodynamic drivers are intermediate outcomes from geographic, anthropogenic, and other hydrodynamic drivers and the geomorphic change outcome/driver.

As erosion and deposition occur at a site, the bed elevation will change, especially over many years. For example, large sediment supply during the period of hydraulic mining in the late 1800s caused deposition in Sacramento Valley rivers, the Delta, and San Francisco Bay (Gilbert 1917). Channels may fill in or



scour. When deposition (including organic accumulation from plant material) on a tidal marsh equals or exceeds sea-level rise, the marsh is sustainable. When deposition is less than sea-level rise, the marsh will eventually become permanently flooded. In this conceptual model, local erosion and deposition are considered on a tidally-averaged time scale. Geomorphic change takes place on a decadal time scale, and is the sum (or integration) of local erosion and deposition. That integration includes any episodic riverine supply or erosion events.

Erosion and deposition also alter bed particle size and properties. If bed shear stress exceeds a critical value, particle motion will occur (ASCE 1975), which decreases the stability of the bed and the potential for submerged aquatic vegetation to establish and grow (see DRERIP aquatic plant model, [http://www.dfg.ca.gov/ERP/conceptual\\_models.asp](http://www.dfg.ca.gov/ERP/conceptual_models.asp)). Erosion tends to winnow finer sediment from the bed, making the bed coarser. Deposition supplies new sediment to the bed, changing organic content, and usually decreasing bed particle size. For cohesive sediment, newly deposited material will have a low density and be relatively erodible (Brennan and others 2002). Consolidation and biostabilization decrease the erodibility of cohesive bed sediment with time (Mehta 1989; Widdows and others 2000). Sediment beds that contain both cohesive and non-cohesive sediment generally become less erodible as the fraction of fine cohesive sediment increases (Le Hir and others 2008).

If present, anthropogenic factors can greatly affect local suspended sediment. To prevent erosion, large boulders (rip-rap) are often placed on levees and the sides of channels. Armoring increases bed particle size and decreases erosion. Dredging deepens channels and can suspend bottom sediment. Fill or dredged material disposal decreases water depths and suspends sediment. Filling estuaries and diking tidal wetlands can reduce tidal prism in an estuary, which decreases tidal velocities and increases deposition (Hood 2004). Wetland restoration projects are designed to increase vegetation and to provide low hydrodynamic energy at the restoration site to encourage deposition. Restoration of tidal flooding to diked lands can increase tidal prism and tidal veloci-

ties in channels adjacent to the restoration site, which in turn can erode the channels (Kirby 1990).

## DISCUSSION

The model includes vegetation and sea-level rise feedback loops, and demonstrates that managing regional transport can affect the distribution of suspended sediment in the Delta.

### Vegetation Feedback Loops

The ability of macrophytes to slow water velocity is well established (Carpenter and Lodge 1986; Sand-Jensen and Mebus 1996). Macrophytes can choke streams and slow flow, sometimes forming dams that alter channel morphology and width (Wilcox and others 1999). The effect of a particular macrophyte on flow is strongly related to its areal extent, density, canopy height, and phenology. There is a threshold velocity (i.e., extreme events) over which macrophytes can no longer reduce flow but simply bend out of the way (Wilcox and others 1999).

By reducing velocity, macrophytes attenuate waves and reduce vertical mixing and bed shear stress, which in turn leads to deposition of sediment (Yang 1998; Braskerud 2001). Sedimentation in wetlands and near-shore environments is a function of sediment supply and retention. Whereas sediment supply depends on watershed processes (Wright and Schoellhamer 2004), sediment retention is greatly influenced by local-scale factors such as plant community composition and particular plant species characteristics (Eisma and Dijkema 1997; Pasternack and Brush 1998, 2001; Alizai and McManus 1980). Sedimentation rate is strongly related to plant architecture, canopy height, and plant density (Yang 1998; Alizai and McManus 1980; Leonard and others 2002).

Our open water submodel assumes that vegetation is not an important factor (Figure 7). An exception is the plant *Egeria densa*, which invaded some of the open waters of the Delta beginning in the 1960s (Jassby and Cloern 2000) and is a likely contributor to the decline in Delta turbidity (Nobriga and others 2008). A positive feedback loop in our con-

ceptual model demonstrates how *Egeria densa* could successfully invade the Delta and reduce turbidity. Where submerged aquatic vegetation successfully colonizes, hydrodynamic energy and bed shear stress are reduced (Carpenter and Lodge 1986; Sand-Jensen and Mebus 1996). This increases local deposition, which decreases local SSC (Yang 1998; Braskerud 2001). Tidal currents transport this clearer water within the Delta (Figure 7). Thus, *Egeria densa* filters suspended sediment out of the water that passes through it, and that clearer water moves elsewhere in the Delta and mixes with more turbid water to reduce turbidity elsewhere. Increased light in the water column at the colonization site and downcurrent (landward and seaward because of bidirectional flow) increases vegetation (via the DRERIP aquatic plant model). This increase in vegetation further decreases hydrodynamic energy in this positive feedback loop.

Inundation of the marsh plain via sea-level rise also establishes another positive feedback loop. As sea level rises, plants at lower elevations are the first to be lost because they drown. This results in less overall organic accumulation on the marsh plain and less trapping of sediment, which ultimately accelerates drowning in other areas of the marsh (Kirwan and others 2008; D'Alpaos 2011). In the worst-case scenario, as sea level continues to rise, the entire marsh drowns and converts to open water habitat. The rate at which this process unfolds strongly depends on the rate of sea-level rise, sediment availability from the watershed and wind-wave re-suspension, and above-ground and below-ground productivity. Other factors that also may be important in particular marsh settings include bioturbation, tidal meandering, and plant species composition of the marsh (Morris 2006; D'Alpaos 2011).

Water depth also affects vegetation colonization. Only where the land surface elevation is greater than mean tide level, can brackish emergent vegetation colonize the site (Orr and others 2003). In salt marshes in the South Bay, vegetation was shown to colonize in the range between 0.98 and 2.94 m above mean low water (Orlando and others 2005). Freshwater emergent vegetation colonizes down to 0.2 m below mean lower low water (Simenstad and others 2000).

### Sea-Level Rise Feedback Loop

Water depth affects hydrodynamics. As cross-sectionally averaged water depth decreases due to deposition or fill, water velocity will increase if the landward tidal prism and water flow passing through that point is unchanged or increases. As cross-sectionally averaged water depth increases because of erosion, dredging, or sea-level rise, water velocity will decrease if the landward tidal prism and the water flow that passes through that point is unchanged or decreases. Wave bottom orbital velocity and the resulting bottom shear stress increases as water depth decreases (Dean and Dalrymple 1984). Because of water depth, geomorphic change is both an outcome and driver of this model.

Unlike the vegetation positive feedback loops, sea-level rise in open waters creates a negative feedback loop. Sea-level rise increases water depth, which decreases hydrodynamic energy and bed shear stress, thus increasing the rate of deposition. The net result is that water depth will increase at a slower rate than sea-level rise. Ganju and Schoellhamer (2010) conducted 30-year scenarios of sea-level rise and sedimentation in Suisun Bay, and found that deposition increased but did not keep up with an assumed sea-level rise of 2 mm yr<sup>-1</sup>. Thus, increased deposition can reduce the effect of sea-level rise on water depth. This negative feedback loop, however, may still not be enough to prevent marsh drowning. Pfeffer and others (2008) estimated that sea level will rise 800 to 2,000 mm during the 21st century, a much greater rate than Ganju and Schoellhamer (2010) assumed. If sea-level changes are large enough to increase hydrodynamic energy, by increasing tidal prism or fetch for example, then feedback may be positive rather than negative.

### Managing Regional Transport

Regional transport is a driver of SSC and suspended-sediment properties at a specific site in the Delta. Regional drivers of local suspended sediment and properties are shown in the dashed box in Figures 7 and 8. River supply is probably the most important regional driver, and it is discussed in the river supply submodel. Exchange with Suisun Bay was discussed




in the regional submodel. Bay and river boundary conditions communicate with any point in the Delta via advective and dispersive transport and local erosion and deposition along the transport pathway. An example of regional influence is that the dominant sediment transport pathway in the Delta is the Sacramento River (Figure 3, Wright and Schoellhamer 2005) and wetland deposition is greatest along the Sacramento River (Reed 2002).

Suspended-sediment concentration and the local properties of SSC and bed sediment are controlled by the interaction between river supply and the hydrodynamics in the Delta. Thus, manipulations of Delta hydrodynamics through reservoir releases, pumping rates, or gate operations may affect the distribution of suspended-sediment concentration and bed-particle size throughout the Delta, and are probably the most feasible techniques for influencing sedimentation in the Delta. For example, during flooding on the San Joaquin and Sacramento rivers when sediment delivery to the Delta is high, the SSC distribution in the Delta could potentially be manipulated through pump and gate operations. Predicting these outcomes is beyond the capabilities of a conceptual model and requires numerical modeling (discussed below).

## Restoration

Marshes that are restored via levee breaches start their development as open water habitats. Depositing sediment increases the elevation of the site until vegetation eventually colonizes the site and creates a marsh. Deposition rate at a restoration site increases as SSC of the water that inundates the site increases. Williams and Orr (2002) used a numerical model to show the effect of SSC on vertical accretion in salt marshes, assuming no erosion of bottom sediment (Figure 9). Vertical accretion of inorganic sediment when SSC is  $100 \text{ mg L}^{-1}$  is about  $0.025 \text{ m yr}^{-1}$ . SSC in the Delta is usually less than  $100 \text{ mg L}^{-1}$ , the smallest SSC considered in Williams and Orr's study, so it would take at least several decades for one meter of sediment to deposit at a Delta restoration site.

In tidal freshwater marshes such as the Delta, however, organic accumulation is a much larger con-



**Figure 9** Effect of SSC on marshplain evolution over time for a site sheltered from wind-wave action. Shaded bar identifies the approximate *Spartina* colonization elevation. Prediction is based on tides at the San Francisco Presidio, no sea-level rise, and  $550 \text{ kg m}^{-3}$  dry density of inorganics typical for San Francisco Bay. National Geodetic Vertical Datum (NGVD) is a vertical datum fixed at the mean sea level of 1929. Source: Williams and Orr (2002).

tributor to marsh formation than it is in saltwater marshes. Organic accumulation includes root biomass, plant litter, and any allochthonous inputs from the watershed (Schlesinger 1997). Organic accumulation in tidal freshwater marshes has been shown to be of greater importance than inorganic sediment for vertical accretion (Neubauer 2008; Drexler 2011). Therefore, although inorganic sedimentation may be relatively low in Delta marshes, the overall vertical accretion rate may be considerably greater than Williams and Orr (2002) estimated because of organic accumulation.

## Uncertainty

Here we discuss some key uncertainties of the conceptual model, but this is not intended as a complete discussion of uncertainties about sedimentation in the Delta. The sediment budget for the Delta (Figure 6) is quantitatively uncertain. Most of the values are based on only 4 years of data and may not represent future conditions. The second-largest source of sediment is the Yolo Bypass which provides 28% of the sediment discharge of the Sacramento River at Freeport

(Figure 6, YOL). Wright and Schoellhamer (2005) used a rating curve based on data from 1957 to 1961 and 1980 to determine Yolo Bypass sediment flux, and a large uncertainty (42%) is associated with that value. In addition, the effect of decreased sediment yield from the Sacramento Valley is unknown, because of lack of data. Uncertainty in sediment supply leads to uncertainty in answering basic questions about the Delta, such as what the deposition rate is, the quantity of sediment available for restoration, and whether the Delta will be sustainable given sea-level rise. The east side tributaries (Figure 6, EAST) supplied only 3.3% of the sediment discharge at Freeport using similarly outdated rating curves. Current data on sediment supply are needed to reduce uncertainty.

Sediment discharge at Freeport decreased by about one-half from 1957 to 2001 (Wright and Schoellhamer 2004). Whether or not that trend will continue is a key uncertainty in predicting future geomorphic changes to the Delta and northern San Francisco Bay (Ganju and Schoellhamer 2010).

Overall, within the conceptual model, physical processes are qualitatively understood but difficult to quantify. Erosion is more difficult to quantify than deposition. Biological processes that affect sedimentation are not as well understood qualitatively, and are very difficult to predict quantitatively. Thus, studies to quantify sediment supply and physical processes and studies of the interaction of biota and sedimentation are needed.

### From Conceptual to Numerical Modeling

The conceptual model described herein is a useful tool for describing the fundamentals of sediment transport in the Delta as well as some of the important feedback loops. It also provides the basis for the development of numerical modeling tools that would provide more predictive capabilities. The conceptual framework remains the same, while the drivers become boundary conditions, the linkages become the numerical algorithms, and the outcomes are the numerical model results. Thus, a numerical model of the Delta would use the river supply as boundary conditions, along with a numerical model of Delta hydrodynamics, to predict local SSC, bed sediment

properties, and geomorphic change. The primary linkages are relations for erosion and deposition rates, which in turn depend on local hydrodynamics, SSC, and bed size and properties (i.e. bed erodibility).

Information gaps that need to be filled to develop an accurate numerical sediment transport model for the Delta can be identified with the conceptual model presented in this paper. The key information gaps are boundary conditions of the regional transport sub-model (Figure 5), initial condition of the bed, data on important but poorly quantified linkages (Figures 7 and 8), and suspended-sediment flux and temporal bed sediment data to calibrate and validate the numerical model.

Numerical models simulate a fixed domain and are driven by external boundary conditions. For a numerical model of sedimentation in the Delta, the regional transport model (Figure 5) shows the external boundary conditions that need to be measured. Suspended-sediment flux in the Sacramento River at Freeport and San Joaquin River at Vernalis has been measured for many years by the USGS. The Yolo Bypass is the second largest source of sediment to the Delta (Figure 6) and sediment discharge measurements were restarted in WY2011. Sediment discharge from the eastside tributaries is smaller and measurements were restarted in WY2011. The Yolo Bypass is the second largest source of sediment to the Delta (Figure 6) and sediment discharge measurements were restarted in WY2011. Sediment discharge from the eastside tributaries is smaller and measurements were restarted in WY2011. McKee and others (2006) estimated net sediment discharge from the Delta to Suisun Bay but it is not measured directly. Measurements at these unmeasured locations would provide the boundary condition data needed to accurately drive a numerical model.

Initial conditions must also be supplied to a numerical model. The bed property that has the greatest effect on sediment transport is the size of surficial bed sediment. While periodic sampling has occurred at a range of sites in the Delta, no comprehensive compilation and analysis is currently available. Delta-wide bed-sediment sampling the USGS initiated

in WY 2011 should provide the most comprehensive assessment to date.

Important but poorly quantified linkages are shown by thick dotted lines in Figures 7 and 8. These represent the physical processes that would limit accuracy of a numerical sedimentation model and are primarily associated with erosion and deposition. To select an optimal erosion algorithm and coefficients, erosion experiments can be conducted in situ (Amos and others 2010) or with undisturbed sediment cores (McNeil and others 1996; Law and others 2008; Le Hir and others 2008). To better quantify deposition, especially of fine cohesive sediment, images of particles collected by video cameras can be analyzed (Manning and Dyer 2002). An additional important but poorly quantified linkage in the marsh submodel is the effect of vegetation on hydrodynamics, which, in turn, affects sediment deposition. It has long been known that sediment deposition decreases along a transect from a tidal channel into a marsh (e.g., Hatton and others 1983), however, it is unclear how much deposition decreases with various types of emergent and submersed vegetation. Therefore, factors such as the structure, areal extent, density, and phenology of dominant marsh plants, which have all been shown to influence sediment deposition and flow velocity (e.g., Carpenter and Lodge 1986; Wilcox and others 1999), need to be better understood to construct an accurate numerical sedimentation model.

To calibrate model coefficients to data, and to test the model by validating it with an independent data set, measurements of suspended-sediment flux (mass per unit time) within the Delta are preferred to measurements of concentration (mass per unit volume) (Ganju and Schoellhamer 2009). To accurately simulate geomorphic change or the load of sediment-associated contaminants, the model must accurately simulate sediment flux. For example, to accurately simulate whether a segment of the Delta is sustainable given sea-level rise, the model must deliver and carry away the correct masses of sediment to and from the segment. During WY 2011 the U.S. Geological Survey began to continuously measure sediment flux at key locations in the Delta, similar to those used by Wright and Schoellhamer (2005) to develop the sediment budget in Figure 6. Velocity-weighted, cross-

sectionally averaged samples of SSC are collected and used to calibrate a continuous point measurement of turbidity. To calculate suspended-sediment flux at a site, the resulting continuous SSC time series is multiplied by continuously measured water discharge. The measurements have to be continuous because of tidal variability and episodic floods. Measurements of how surficial bed sediment size varies in time also provide calibration and validation data for a numerical model.

## CONCLUSIONS

The sedimentation conceptual model demonstrates two key feedback loops. Emergent macrophytes and *Egeria densa* act as ecosystem engineers that can create a positive feedback loop for sedimentation. This feedback loop helps explain the successful invasion of the Delta by *Egeria densa* since the 1960s and its effect on Delta turbidity. It also highlights the importance of maintaining existing marsh vegetation to prevent extensive marsh drowning if sea-level rise accelerates greatly. Sea-level rise in open water is partially countered by a negative feedback loop that increases deposition if there is a net decrease in hydrodynamic energy. Manipulation of regional sediment transport is probably the most feasible method to manipulate SSC to create or control turbidity and thus fish habitat.

Other conclusions include:

1. **Sediment supply:** Sediment supply is decreasing as a result of trapping behind dams, deposition in flood bypasses, protection of river banks, and diminishment of the hydraulic mining sediment pulse. About two-thirds of the sediment that enters the Delta from the rivers deposits in the Delta.
2. **Sustainability:** The Delta is depositional, and adequate sediment deposition is required for wetlands to keep up with sea-level rise. If sediment supply continues to decrease, and if sea-level rise accelerates, it will become more difficult to sustain the existing Delta landscapes and habitats.
3. **Restoration:** Typical SSCs in the Delta range from 10 to 50 mg L<sup>-1</sup> except during river floods when the SSC can exceed 200 mg L<sup>-1</sup>. Natural inorganic

wetland sedimentation when SSC is  $100 \text{ mg L}^{-1}$  is about  $0.025 \text{ m yr}^{-1}$ , so inorganic sediment deposition rates in the Delta typically are smaller than this value. Plants provide organic accumulation that can exceed inorganic deposition.

4. **Light and biota:** Suspended sediment is the primary attenuator of sunlight in the water column of the Delta. Photosynthesis, primary production, and fish behavior depend on light, and are likely to change if sediment supply and turbidity continue to decrease.
5. **Model uncertainty:** Physical processes are qualitatively understood but difficult to quantify. Erosion is more difficult to quantify than deposition. Biological processes that affect sedimentation are not as well understood qualitatively, and are very difficult to predict quantitatively.
6. **Numerical modeling:** The conceptual model indicates that boundary conditions, bed sediment size, erosion, deposition, biological processes, and suspended-sediment fluxes are key information gaps that need to be filled to develop an accurate numerical model of sedimentation in the Delta.

## ACKNOWLEDGEMENTS

We thank David Fryberg, Denise Reed, Stuart Siegel, Mark Stacey, the DRERIP review panel, and the other DRERIP model authors for their assistance in developing this model. Justin Ferris, Mick van der Wegen, and two anonymous reviewers provided helpful comments on initial drafts of this manuscript.

## REFERENCES

- Alizai SAK, McManus J. 1980. The significance of reed beds on siltation in the Tay Estuary. *Proceedings of the Royal Society Edinburgh*. 78:1–13.
- [ASCE] American Society of Civil Engineers. 1975. *Sedimentation Engineering*: Vanoni VA, editor. *Manuals and Reports on Engineering Practice* 54.
- Amos CL, Umgieser G, Ferrarin C, Thompson CEL, Whitehouse RJS, Sutherland TF, Bergamasco A. 2010. The erosion rates of cohesive sediments in Venice lagoon, Italy: *Continental Shelf Research* 30:859–870.
- Atwater BF, Belknap DF. 1980. Tidal-wetland deposits of the Sacramento–San Joaquin Delta, California. *Pacific Coast Paleogeography Symposium* 4:89–103.
- Bauer BO, Lorang MS, Sherman DJ. 2002. Estimating boat-wake-induced levee erosion using sediment suspension measurements. *Journal of Waterway, Port, Coastal, and Ocean Engineering* 128(4):152–162.
- Bergamaschi BA, Kuivila KM, Fram MS. 2001. Pesticides associated with suspended sediments entering San Francisco Bay following the first major storm of water year 1996. *Estuaries* 24(3):368–380.
- Braskerud BC. 2001. The influence of vegetation on sedimentation and resuspension of soil particles in small constructed wetlands. *Journal of Environmental Quality* 30:1447–1457.
- Brennan ML, Schoellhamer DH, Burau JR, Monismith SG. 2002. Tidal asymmetry and variability of bed shear stress and sediment bed flux at a site in San Francisco Bay, USA. In: Winterwerp JC, Kranenburg C, editors. *Fine sediment dynamics in the marine environment*. Amsterdam, The Netherlands: Elsevier. p 93–108.
- Byrne R, Ingram BL, Starratt S, Malamud–Roam F, Collins JN, Conrad ME. 2001. Carbon-isotope, diatom, and pollen evidence for late Holocene salinity change in a brackish marsh in the San Francisco Estuary. *Quaternary Research* 55:66–76.
- Caffrey JM. 1995. Spatial and seasonal patterns in sediment nitrogen remineralization and ammonium concentrations in San Francisco Bay, California. *Estuaries* 18(1B):219–233.
- [DFG] California Department of Fish and Game. 1997. *California wetland and riparian geographic information system*. Final report. Sacramento (CA): California Department of Fish and Game.



- [DWR] California Department of Water Resources. 1995. Sacramento–San Joaquin Delta atlas. Sacramento (CA): California Department of Water Resources. 121 p.
- Canuel EA, Lerberg EJ, Dickhut RM, Kuehl SA, Bianchi TS, Wakeham SG. 2009. Changes in sediment and organic carbon accumulation in a highly-disturbed ecosystem: the Sacramento–San Joaquin River Delta (California, USA). *Marine Pollution Bulletin* 59(4–7):154–163.
- Carpenter SR, Lodge DM. 1986. Effects of submerged macrophytes on ecosystem processes. *Aquatic Botany* 26:341–370.
- Cloern JE. 1987. Turbidity as a control on phytoplankton biomass and productivity in estuaries. *Continental Shelf Research* 7(11/12):1367–1381.
- Curtis JA, Flint LE, Alpers CN, Wright SA, Snyder NP. 2006. Use of sediment rating curves and optical backscatter data to characterize sediment transport in the upper Yuba River watershed, California, 2001–03. U.S. Geological Survey Scientific Investigations Report 05-5246. 84 p. Available from: <http://pubs.er.usgs.gov/publication/sir20055246>.
- D’Alpaos A. 2011. The mutual influence of biotic and abiotic components on the long-term ecomorphodynamic evolution of salt-marsh ecosystems: *Geomorphology* 126(3–4):269–278. doi:10.1016/j.geomorph.2010.04.027
- Dean RC, Dalrymple RA. 1984. *Water wave mechanics for engineers and scientists*. Englewood Cliffs (NJ): Prentice–Hall. 353 p.
- Dinehart RL. 2002. Bedform measurement recorded by sequential single-beam surveys in tidal rivers. *Journal of Hydrology* 258:25–39.
- Drexler JZ. 2011. Peat formation processes through the millennia in tidal marshes of the Sacramento–San Joaquin Delta, California, USA. *Estuaries and Coasts* 34:900–911.
- Droppo IG, Leppard GG, Flannigan DT, Liss SN. 1997. The freshwater floc: A functional relationship of water and organic and inorganic floc constituents affecting suspended sediment properties. *Water, Air, and Soil Pollution* 99:43–54.
- Eisma D, Kijkema KS. 1997. The influence of salt marsh vegetation on sedimentation. In: Eisma D, editor. *Intertidal deposits*. Boca Raton (FL): CRC Press. p 403–414.
- Feyrer F, Nobriga ML, Sommer TR. 2007. Multidecadal trends for three declining fish species: habitat patterns, and mechanisms in the San Francisco Estuary, California, USA. *Canadian Journal of Fisheries and Aquatic Sciences* 64:723–734.
- Florsheim J, Mount JF, Chin A. 2008. Bank erosion as a desirable attribute of rivers. *Bioscience* 58(6):519–529.
- Ganju NK, Schoellhamer DH. 2010. Decadal-timescale estuarine geomorphic change under future scenarios of climate and sediment supply. *Estuaries and Coasts* 33:15–29.
- Ganju NK, Schoellhamer DH. 2009. Calibration of an estuarine sediment transport model to sediment fluxes as an intermediate step for simulation of geomorphic evolution. *Continental Shelf Research* 29(1):148–158.
- Ganju NK, Schoellhamer DH, Murrell MC, Gartner JW, Wright SA. 2007. Constancy of the relation between floc size and density in San Francisco Bay. In Maa JPY, Sanford LP, Schoellhamer DH, editors. *Estuarine and Coastal Fine Sediments Dynamics*. Elsevier. p. 75–91.
- Gilbert GK. 1917. *Hydraulic-mining debris in the Sierra Nevada*. Washington, DC: U.S. Geological Survey Professional Paper 105. 154 p. Available from: <http://pubs.er.usgs.gov/publication/pp105>.
- Goodwin P, Denton RA. 1991. Seasonal influences on the sediment transport characteristics of the Sacramento River. *Proceedings of the Institution of Civil Engineers* 91(Part 2):163–172.



## SAN FRANCISCO ESTUARY & WATERSHED SCIENCE

- Gray JR, Glysson GD, Turcios LM, Schwarz GE. 2000. Comparability of suspended-sediment concentration and total suspended-solids data. U.S. Geological Survey Water Resources Investigations Report 00-4191. 14 p. Available from: <http://pubs.er.usgs.gov/publication/wri004191>.
- Hatton RS, DeLaune RD, Patrick WH. 1983. Sedimentation, accretion, and subsidence in marshes of Barataria Basin, Louisiana. *Limnology and Oceanography* 28:494–502.
- Hood WG. 2004. Indirect environmental effects of dikes on estuarine tidal channels: thinking outside of the dike for habitat restoration and monitoring: *Estuaries* 27(2):273–282.
- James LA. 1991. Incision and morphologic evolution of an alluvial channel recovering from hydraulic mining sediment. *Geological Society of America Bulletin* 103:723–736.
- Jassby AD, Cloern JE. 2000. Organic matter sources and rehabilitation of the Sacramento–San Joaquin Delta (California, USA). *Aquatic Conservation: Marine and Freshwater Ecosystems* 10:323–352.
- Jassby AD, Cloern JE, Cole BE. 2002. Annual primary production: Patterns and mechanisms of change in a nutrient-rich tidal ecosystem. *Limnology and Oceanography* 47(3):698–712.
- Kelley R. 1998. Battling the inland sea: Floods, Public Policy, and the Sacramento Valley. Berkeley (CA): University of California Press. 420 p.
- Kirby R. 1990. The sediment budget of the erosional intertidal zone of the Medway Estuary, Kent. *Proceedings of the Geological Association* 101(1):63–77.
- Kirwan ML, Murray AB, Boyd WS. 2008. Temporary vegetation disturbance as an explanation for permanent loss of tidal wetlands. *Geophysical Research Letters* 35(L05403):5 p. doi:10.1029/2007GL032681. Available from: <http://www.agu.org/pubs/crossref/2008/2007GL032681.shtml>.
- Law BA, Hill PS, Milligan TG, Curran KJ, Wiberg PL, Wheatcroft RA. 2008. Size sorting of fine-grained sediments during erosion: results from the western Gulf of Lions. *Continental Shelf Research* 28:1935–1946.
- Le Hir P, Cann P, Waeles B, Jestin H, Basslullet P. 2008. Erodibility of natural sediments: experiments on sand/mud mixtures from laboratory and field erosion tests. In: Kusuda T, Yamanishi H, Spearman J, Gailani JZ, editors. *Sediment and ecohydraulics*. Amsterdam, The Netherlands: Elsevier. p 137–153.
- Le Roux SM, Turner A, Millward GE, Ebdon L, Apprion P. 2001. Partitioning of mercury onto suspended sediments in estuaries. *Journal of Environmental Monitoring* 3:37–42.
- Leonard LA, Wren PA, Beavers RL. 2002. Flow dynamics and sedimentation in *Spartina alterniflora* and *Phragmites australis* marshes of the Chesapeake Bay. *Wetlands* 22(2):415–424.
- Luengen AR, Flegal AR. 2009. Role of phytoplankton in mercury cycling in the San Francisco Bay estuary. *Limnology and Oceanography* 54(1):23–40.
- Manning AJ, Dyer KR. 2002. The use of optics for the *in situ* determination of flocculated mud characteristics. *Journal of Optics A: Pure and Applied Optics* 4:S71–S81.
- McKee LJ, Ganju NK, Schoellhamer DH. 2006. Estimates of suspended sediment flux entering San Francisco Bay from the Sacramento and San Joaquin Delta, San Francisco Bay, California. *Journal of Hydrology* 323:335–352.
- McNeil J, Taylor C, Lick W. 1996. Measurements of erosion of undisturbed bottom sediments with depth. *Journal of Hydraulic Engineering* 122(6):316–324.
- Meade RH. 1982. Sources, sinks, and storage of river sediment in the Atlantic drainage of the United States. *Journal of Geology* 90:235–252.
- Mehta AJ. 1989. On estuarine cohesive sediment suspension behavior. *Journal of Geophysical Research* 94(C10):14303–14314.

- Morris J. 2006. Competition among marsh macrophytes by means of geomorphological displacement in the intertidal zone. *Estuarine, Coastal and Shelf Science* 69:395–402.
- Neubauer SC. 2008. Contributions of mineral and organic components to tidal freshwater marsh accretion. *Estuarine, Coastal and Shelf Science* 78:78–88.
- Nilsson C, Reidy CA, Dynesius M, Revenga C. 2005. Fragmentation and flow regulation of the world's large river systems. *Science* 308:405–408.
- Nobriga ML, Feyrer F, Baxter RD, Chotkowski M. 2005. Fish community ecology in an altered river delta: spatial patterns in species composition, life history strategies, and biomass. *Estuaries* 28(5):776–785.
- Nobriga ML, Sommer TR, Feyrer F, Fleming K. 2008. Long-term trends in summertime habitat suitability for delta smelt (*Hypomesus transpacificus*): San Francisco Estuary and Watershed Science [Internet]. [cited 2012 Sept 6]; 6(1). Available from <http://escholarship.org/uc/item/5xd3q8tx>.
- Ogden Beeman & Associates, Ray B. Krone & Associates. 1992. Sediment budget study for San Francisco Bay. Final report prepared for the San Francisco District, U.S. Army Corps of Engineers.
- Orlando JL, Drexler JZ, Dedrick KG. 2005. South San Francisco Bay tidal marsh vegetation and elevation surveys—Corkscrew Marsh, Bird Island, and Palo Alto Baylands, California, 1983. U.S. Geological Survey Data Series Report 134. 51 p. Available from: <http://pubs.er.usgs.gov/publication/ds134>.
- Orr M, Crooks S, Williams PB. 2003. Will restored tidal marshes be sustainable?. *San Francisco Estuary and Watershed Science* [Internet]. [cited 2012 Sept 6]; 1(1). Available from: <http://escholarship.org/uc/item/8hj3d20t>.
- Pasternack GB, Brush GS. 1998. Sedimentation cycles in a river-mouth tidal freshwater marsh. *Estuaries* 21(3):407–415.
- Pasternack GB, Brush GS. 2001. Seasonal variations in sedimentation and organic content in five plant associations on a Chesapeake Bay tidal freshwater delta. *Estuarine, Coastal and Shelf Science* 53:93–106.
- Pfeffer WT, Harper JT, O'Neel S. 2008. Kinematic constraints on glacier contributions to 21st-century sea-level rise. *Science* 321:1340–1343.
- Porterfield G. 1980. Sediment transport of streams tributary to San Francisco, San Pablo, and Suisun Bays, California, 1909–1966. U.S. Geological Survey Water-Resources Investigations Report 80–64. 92 p.
- Porterfield G, Busch RD, Waananen AO. 1978. Sediment transport in the Feather River, Lake Oroville to Yuba City, California. U.S. Geological Survey Water-Resources Investigations Report 78–20. 73 p.
- Reed DJ. 2002. Understanding tidal marsh sedimentation in the Sacramento–San Joaquin Delta, California. *Journal of Coastal Research Special Issue* 36:605–611.
- Regional Monitoring Program (RMP) [Internet]. Data Query Tool; [cited 2007 Feb 21]. Available from: <http://www.sfei.org/rmp/wqt>.
- Ruhl CA, Schoellhamer DH. 2004. Spatial and temporal variability of suspended-sediment concentrations in a shallow estuarine environment. *San Francisco Estuary and Watershed Science* [Internet]. [cited 2012 Sept 6]; 2(2). Available from <http://escholarship.org/uc/item/1g1756dw>.
- Sand-Jensen K, Mebus JR. 1996. Fine-scale patterns of water velocity within macrophyte patches in streams. *Oikos* 76:169–180.
- Schemel LE, Hager SW, Childers D. 1996. The supply and carbon content of suspended sediment from the Sacramento River to San Francisco Bay. In: Hollibaugh JT, editor. *San Francisco Bay: the ecosystem*. Pacific Division, American Association for the Advancement of Science. p 237–260.
- Schlesinger WH. 1997. *Biogeochemistry: an analysis of global change*. New York (NY); Academic Press. 588 p.

## SAN FRANCISCO ESTUARY & WATERSHED SCIENCE

- Schoellhamer DH. 2001. Influence of salinity, bottom topography, and tides on locations of estuarine turbidity maxima in northern San Francisco Bay. In: McAnally WH, Mehta AJ, editors. Coastal and estuarine fine sediment transport processes. Amsterdam, The Netherlands: Elsevier. p 343–357. Available from <http://ca.water.usgs.gov/abstract/sfbay/elsevier0102.pdf>.
- Schoellhamer DH, Wright SA. 2003. Continuous monitoring of suspended sediment discharge in rivers by use of optical backscatterance sensors. In: Bogen J, Fergus T, Walling DE, editors. Erosion and sediment transport measurement: technological and methodological advances. International Association for Hydrological Science Publication 283. p 28–36. Available from [http://iahs.info/redbooks/a283/iahs\\_283\\_028.pdf](http://iahs.info/redbooks/a283/iahs_283_028.pdf).
- Schoellhamer DH, Mumley TE, Leatherbarrow JE. 2007a. Suspended sediment and sediment-associated contaminants in San Francisco Bay. Environmental Research 105:119–131.
- Schoellhamer D, Wright S, Drexler J, Stacey M. 2007b. Sedimentation conceptual model. CALFED Science Program. Delta Regional Ecosystem Restoration Implementation Plan conceptual models. Available from: [http://science.calwater.ca.gov/pdf/drerip/DRERIP\\_sediment\\_conceptual\\_model\\_final\\_111307.pdf](http://science.calwater.ca.gov/pdf/drerip/DRERIP_sediment_conceptual_model_final_111307.pdf).
- Simenstad C, Toft J, Higgins H, Cordell J, Orr M, Williams P, Grimaldo L, Hymanson Z, Reed D. 2000. Sacramento–San Joaquin Delta breached levee wetland study (BREACH) preliminary report. Wetland Ecosystem Team, University of Washington, School of Fisheries.
- Singer MB. 2007. The influence of major dams on hydrology through the drainage network of the Sacramento River basin, California. River Research and Applications 23:55–72.
- Singer MB, Aalto R, James LA. 2008. Status of the lower Sacramento Valley flood-control system within the context of its natural geomorphic setting. Natural Hazards Review 9(3):104–115.
- Syvitski JPM, Vörösmarty CJ, Kettner AJ, Green P. 2005. Impact of humans on the flux of terrestrial sediment to the global coastal ocean. Science 308:376–380.
- Thompson B, Lowe S, Kellog M. 2000. Results of the benthic pilot study, 1994–1997. Part 1–macrobenthic assemblages of the San Francisco Bay–Delta, and their responses to abiotic factors. Regional Monitoring Program Technical Report 39. Oakland (CA): San Francisco Estuary Institute. 39 p.
- Tobin A, Schoellhamer DH, Burau JR. 1995. Suspended-solids flux in Suisun Bay, California. In: Proceedings, First International Conference on Water Resources Engineering; 1995 Aug 14–18; San Antonio (TX). 2:1511–1515.
- Turner A, Hyde TL, Rawling MC. 1999. Transport and retention of hydrophobic organic micropollutants in estuaries: implications of the particle concentration effect. Estuarine, Coastal and Shelf Science 49:733–746.
- Turner A, Millward GE. 2000. Particle dynamics and trace metal reactivity in estuarine plumes: Estuarine, Coastal, and Shelf Science 50:761–774.
- Turner A, Millward GE. 2002. Suspended particles: their role in estuarine biogeochemical cycles, Estuarine, Coastal and Shelf Science 55:857–883.
- [USFWS] U.S. Fish and Wildlife Service. 2000. Impacts of riprapping to ecosystem functioning, Lower Sacramento River, California. Available from: [http://www.fws.gov/sfbaydelta/documents/Riprap\\_Effects\\_2004\\_revision.pdf](http://www.fws.gov/sfbaydelta/documents/Riprap_Effects_2004_revision.pdf).
- Widdows J, Brown S, Brinsley MD, Salkeld PN, Elliott M. 2000. Temporal changes in intertidal sediment erodibility: influence of biological and climatic factors. Continental Shelf Research 20:1275–1290.
- Wilcox RJ, Champion PD, Nagels JW, Croker GF. 1999. The influence of aquatic macrophytes on the hydraulic and physico-chemical properties of a New Zealand lowland stream. Hydrobiologia 416:203–214.

Williams GP, Wolman MG. 1984. Downstream effects of dams on alluvial rivers. U.S. Geological Survey Professional Paper 1286 p. Available from: <http://pubs.er.usgs.gov/usgspubs/pp/pp1286>.

Williams PB, Orr MK. 2002. Physical evolution of restored breached leveed salt marshes in the San Francisco Bay estuary. *Restoration Ecology* 10(3):527–542.

Wright SA, Schoellhamer DH. 2004. Trends in the Sediment yield of the Sacramento River, California, 1957–2001. *San Francisco Estuary and Watershed Science* [Internet]. [cited 2012 Sept 6]; 2(2). Available from: <http://escholarship.org/uc/item/891144f4>.

Wright SA, Schoellhamer DH. 2005. Estimating sediment budgets at the interface between rivers and estuaries with application to the Sacramento–San Joaquin River Delta. *Water Resources Research* 41. Available from: <http://www.agu.org/pubs/crossref/2005/2004WR003753.shtml>. doi:10.1029/2004WR003753.

Yang SL. 1998. The role of a *Scirpus* marsh in attenuation of hydrodynamics and retention of fine sediment in the Yangtze Estuary: *Estuarine, Coastal and Shelf Science* 47:227–233.