

CONDITIONAL EXPECTATIONS AND AN ISOMORPHIC CHARACTERIZATION OF L_1 -SPACES

L. TZAFRIRI¹

ABSTRACT. Conditional expectations can be defined in Banach spaces whose elements can be represented as measurable functions. In the present paper it is shown that such a space (precisely a cyclic space) is isomorphic to an L_1 -space if and only if the conditional expectations act as bounded operators for sufficiently many representations.

Let (Ω, Σ, μ) be a finite measure space and Σ_0 a subring of Σ with maximal element Ω_0 ; then for each Σ -measurable function f which is bounded on Ω one can consider the measure $\mu_0(\sigma) = \int_{\sigma \cap \Omega_0} f(\omega) \mu(d\omega)$; $\sigma \cap \Omega_0 \in \Sigma_0$. Since μ_0 is evidently absolutely continuous with respect to the restriction of μ to the subfield generated by Σ_0 and Ω , due to the Radon-Nikodym theorem, there exists a Σ_0 -measurable function denoted $E(\Sigma_0, \mu)f$ for which

$$\int_{\sigma} f(\omega) \mu(d\omega) = \int_{\sigma} E(\Sigma_0, \mu)f \mu(d\omega); \quad \sigma \in \Sigma_0.$$

Obviously the operator $E(\Sigma_0, \mu): f \rightarrow E(\Sigma_0, \mu)f$ can be extended uniquely to a contractive projection in $L_p(\Omega, \Sigma, \mu)$; $1 \leq p \leq +\infty$, which is called the conditional expectation relative to Σ_0 .

However, if the L_p -norm is replaced by a general monotonic norm ρ in the sense of the theory of Banach function spaces (see for instance W. A. J. Luxemburg and A. C. Zaanen [9, Note I]), usually, $E(\Sigma_0, \mu)$ does not act as a bounded operator in L_ρ —the space of all Σ -measurable functions for which $\rho(f) < +\infty$, even when we assume that $L_1(\Omega, \Sigma, \mu) \supset L_\rho \supset L_\infty(\Omega, \Sigma, \mu)$. Furthermore, a Banach function space L_ρ admits many isometric representations; e.g. to every positive function $\phi \in L_\rho$ whose support is Ω one can define a new norm $\rho_\phi(f) = \rho(\phi f)$, obtaining in this way a new Banach function space L_{ρ_ϕ} which is isometric to L_ρ and satisfies $L_1(\Omega, \Sigma, \phi\mu) \supset L_{\rho_\phi} \supset L_\infty(\Omega, \Sigma, \phi\mu)$.

The main result of this paper states that L_ρ is isomorphic to an L_1 -space over a finite measure space provided for every subring Σ_0 of

Received by the editors September 24, 1969.

AMS 1969 subject classifications. Primary 4606, 4635, 4725; Secondary 4610, 4720.

Key words and phrases. Conditional expectation, Banach function spaces, cyclic spaces, L_1 -spaces, Boolean algebras of projections, isomorphism between Banach spaces.

Supported in part by National Science Foundation, Grant GP 8217.

Σ and positive function $\phi \in L_p$ whose support is Ω the conditional expectation $E(\Sigma_0, \phi\mu)$ is a *bounded* operator in $L_{p\phi}$.

Since a Banach space isomorphic to an L_1 -space is not in general a Banach function space, in order to have a complete characterization we will consider the more natural context (for this purpose) of cyclic spaces introduced by W. G. Bade [3] rather than that of Banach function spaces (for other characterizations using cyclic spaces see [11]).

Conditional expectations in cyclic spaces. We shall start by summarizing some notions and results needed in the sequel. A Boolean algebra of projections \mathfrak{B} in a Banach space X is called σ -complete (cf. W. G. Bade [2]) provided for every sequence $P_n \in \mathfrak{B}; n = 1, 2, \dots$, the projections $\bigvee_{n=1}^{\infty} P_n$ and $\bigwedge_{n=1}^{\infty} P_n$ exist in \mathfrak{B} and satisfy

$$\left(\bigvee_{n=1}^{\infty} P_n\right) X = \text{clm}_n \{P_n X\}; \quad \left(\bigwedge_{n=1}^{\infty} P_n\right) X = \bigcap_{n=1}^{\infty} \{P_n X\}.$$

It is well known that \mathfrak{B} can be regarded as a spectral measure $P(\cdot)$ defined on the Borel sets Σ of its Stone space Ω and it follows from W. G. Bade [2, Theorem 2.2] and N. Dunford [4] that there exists a constant K such that for every Borel bounded function f , the integral $S(f) = \int_{\Omega} f(\omega) P(d\omega)$ exists in the uniform operator topology and satisfies the inequality: $\|S(f)\| \leq K \sup_{\omega \in \Omega} |f(\omega)|$. For f unbounded we can consider $S(f)$ as an unbounded operator whose domain is

$$D(S(f)) = \left\{x \mid x \in X, \lim_{m \rightarrow \infty} \int_{e_m} f(\omega) P(d\omega)x \text{ exists}\right\}$$

and $e_m = \{\omega \mid \omega \in \Omega, |f(\omega)| \leq m\}; m = 1, 2, \dots$. According to W. G. Bade [3], X is a *cyclic space* relative to a σ -complete Boolean algebra of projections \mathfrak{B} if there is $x_0 \in X$ such that $X = \mathfrak{M}(x_0) = \text{clm} \{Px_0 \mid P \in \mathfrak{B}\}$. In this case, by W. G. Bade [3, Theorem 4.5], $X = \mathfrak{M}(x_0) = \{S(f)x_0 \mid x_0 \in D(S(f))\}$. Let us also mention that for every cyclic space $X = \mathfrak{M}(x_0)$ there exists a functional $x_0^* \in X^*$, which will be called *Bade functional*, with the following properties:

- (i) $x_0^* Px_0 \geq 0; P \in \mathfrak{B}$;
- (ii) if $x_0^* Px_0 = 0$ for some $P \in \mathfrak{B}$, then $P = 0$ (cf. W. G. Bade [2, Theorem 3.1]).

With this preparation we can state our principal result which is contained in the following theorem.

THEOREM 1. *A Banach space X is isomorphic to an L_1 -space over a finite measure space if and only if:*

(a) X is a cyclic space $\mathfrak{M}(x_0)$; $x_0 \in X$, relative to some σ -complete Boolean algebra of projections \mathfrak{B} , and

(b) there exists a Bade functional $x_0^* \in X^*$ such that the series

$$\sum_{n=1}^{\infty} \frac{x_0^* P(\sigma_n) S(f) x_0}{x_0^* P(\sigma_n) S(\phi) x_0} P(\sigma_n) S(\phi) x_0;$$

$$S(f)x_0, S(\phi)x_0 \in \mathfrak{M}(x_0); \quad \phi(\omega) > 0; \quad \omega \in \Omega,$$

converges strongly in X for every sequence of disjoint sets $\sigma_n \in \Sigma$; $n = 1, 2, \dots$

PROOF. Let τ be an isomorphism between an L_1 -space $L_1(T, \mathfrak{F}, m)$; $m(T) < +\infty$ and X ; $x_0 = \tau(1)$; $x_0^* = (\tau^*)^{-1}(1)$ and $\mathfrak{F} = \{F(e) \mid e \in T; F(e)\tau(f) = \tau(\chi_e f); f \in L_1(T, \mathfrak{F}, m)\}$. Obviously X is a cyclic space relative to the σ -complete Boolean algebra of projections \mathfrak{F} for which the series in condition (b) of the theorem converges and its sum is bounded in norm by $\|\tau\| \cdot \|\tau^{-1}\| \cdot \|S(f)x_0\|$.

Conversely, let us set

$$Q(S(f)x_0) = \sum_{n=1}^{\infty} \frac{x_0^* P(\sigma_n) S(f) x_0}{x_0^* P(\sigma_n) S(\phi) x_0} P(\sigma_n) S(\phi) x_0; \quad S(f)x_0 \in X.$$

One can easily see that Q is a linear projection in X . In order to show that Q is bounded assume there exist $S(f_n)x_0 \in X$; $\|S(f_n)x_0\| = 1$; $n = 1, 2, \dots$, such that $\|Q(S(f_n)x_0)\| \geq n^3$. The inequality

$$\|S(|f_n|)x_0\| = \left\| S\left(\frac{|f_n|}{f_n}\right) S(f_n)x_0 \right\| \leq K \|S(f_n)x_0\|$$

shows that $S(|f_n|)x_0 \in X$ and $\|S(|f_n|)x_0\| \leq K$; $n = 1, 2, \dots$. Set $Q(S(f_n)x_0) = S(g_n)x_0$; $Q(S(|f_n|)x_0) = S(h_n)x_0$; since $|g_n(\omega)| \leq h_n(\omega)$; $\omega \in \Omega$, we obtain

$$\|S(h_n)x_0\| \geq \frac{1}{K} \|S(|g_n|)x_0\| \geq \frac{1}{K^2} \|S(g_n)x_0\| \geq \frac{n^3}{K^2}.$$

Thus for $S(f)x_0 = \sum_{n=1}^{\infty} (S(|f_n|)x_0/n^2) \in X$ we have

$$\|Q(S(f)x_0)\| \geq \frac{1}{K} \left\| Q\left(\frac{S(|f_n|)x_0}{n^2}\right) \right\| = \frac{1}{Kn^2} \|S(h_n)x_0\| \geq \frac{n}{K^3}$$

which shows that Q is not defined in $S(f)x_0 \in X$ i.e. condition (b) of the theorem does not hold. We have to point out that Q depends on the choice of $S(\phi)x_0 \in X$ and $\sigma_n \in \Sigma$; $n = 1, 2, \dots$

Now, let $\{\delta'_n\}$ and $\{\delta''_n\}$ be two sequences of mutually disjoint sets; $\delta'_n, \delta''_n \in \Sigma$; $n = 1, 2, \dots$, $(\bigcup_{n=1}^\infty \delta'_n) \cap (\bigcup_{n=1}^\infty \delta''_n) = \emptyset$ for which $P(\delta'_n)$ as well as $P(\delta''_n)$ are nonzero projections. Set $\mu(\sigma) = x_0^* P(\sigma) x_0$; $\sigma \in \Sigma$; $a_n = \min \{ \mu(\delta'_n), \mu(\delta''_n) \}$ and

$$S(\phi)x_0 = \sum_{n=1}^\infty \frac{a_n}{2^n} \frac{P(\delta'_n)x_0}{\mu(\delta'_n)} + \sum_{n=1}^\infty \frac{a_n}{2^n} \frac{P(\delta''_n)x_0}{\mu(\delta''_n)} + P\left(\Omega - \left(\bigcup_{n=1}^\infty \delta'_n \cup \bigcup_{n=1}^\infty \delta''_n\right)\right)x_0.$$

Since $\phi(\omega) > 0$; $\omega \in \Omega$, we can consider the projection Q corresponding to $S(\phi)x_0$ and the partition $\sigma_n = \delta'_n \cup \delta''_n$; $n = 1, 2, \dots$. Then

$$Q(P(\delta'_n)x_0) = \frac{2^{n-1}}{a_n} \mu(\delta'_n) \left[\frac{a_n}{2^n} \frac{P(\delta'_n)x_0}{\mu(\delta'_n)} + \frac{a_n}{2^n} \frac{P(\delta''_n)x_0}{\mu(\delta''_n)} \right]$$

and further

$$Q\left(\frac{P(\delta'_n)x_0}{\mu(\delta'_n)}\right) = \left(\frac{P(\delta'_n)x_0}{\mu(\delta'_n)} + \frac{P(\delta''_n)x_0}{\mu(\delta''_n)}\right) / 2$$

and similarly:

$$Q\left(\frac{P(\delta''_n)x_0}{\mu(\delta''_n)}\right) = \left(\frac{P(\delta'_n)x_0}{\mu(\delta'_n)} + \frac{P(\delta''_n)x_0}{\mu(\delta''_n)}\right) / 2.$$

This implies that a series $\sum_{n=1}^\infty c_n P(\delta'_n)x_0 / \mu(\delta'_n)$ converges if and only if the series $\sum_{n=1}^\infty c_n P(\delta''_n)x_0 / \mu(\delta''_n)$ does, i.e., the basis $\{P(\delta'_n)x_0 / \mu(\delta'_n)\}$ is equivalent to the basis $\{P(\delta''_n)x_0 / \mu(\delta''_n)\}$. If π is a permutation of the natural numbers, the bases $\{P(\delta'_n)x_0 / \mu(\delta'_n)\}$ and $\{P(\delta'_{\pi(n)})x_0 / \mu(\delta'_{\pi(n)})\}$ will be equivalent since both are equivalent to $\{P(\delta''_n)x_0 / \mu(\delta''_n)\}$. In the terminology of I. Singer [10] (see also M. J. Kadec and A. Pełczyński [7]) this means that both bases $\{P(\delta'_n)x_0 / \mu(\delta'_n)\}$ and $\{P(\delta''_n)x_0 / \mu(\delta''_n)\}$ are symmetric and thus by [7, Theorem 5] there exists M' and M'' such that

$$\frac{1}{\|x_0^*\|} \leq \frac{\|P(\delta'_n)x_0\|}{\mu(\delta'_n)} \leq M'; \quad \frac{1}{\|x_0^*\|} \leq \frac{\|P(\delta''_n)x_0\|}{\mu(\delta''_n)} \leq M''; \quad n = 1, 2, \dots$$

Consequently, for any partition $\{\delta_n\}$ there exists M such that $\|P(\delta_n)x_0\| / \mu(\delta_n) \leq M$; $n = 1, 2, \dots$, (since we can take $\delta'_n = \delta_{2n-1}$ and $\delta''_n = \delta_{2n}$; $n = 1, 2, \dots$).

The next step will be to show that for any partition $\{\delta_n\}$ the basis $\{P(\delta_n)x_0 / \mu(\delta_n)\}$ is equivalent to the natural basis of l_1 . Indeed, if a series $\sum_{n=1}^\infty c_n P(\delta_n)x_0 / \mu(\delta_n)$ is convergent, then it is easy to see that

the series $\sum_{n=1}^{\infty} |c_n| P(\delta_n)x_0/\mu(\delta_n)$ is convergent too. By repeating the previous part of the proof for

$$S(\psi)x_0 = \sum_{n=1}^{\infty} |c_n| \frac{P(\delta_n)x_0}{\mu(\delta_n)} + P\left(\Omega - \bigcup_{n=1}^{\infty} \delta_n\right)x_0$$

instead of x_0 (which is possible since $\psi(\omega) > 0; \omega \in \Omega$) we get $\sum_{n=1}^{\infty} \|P(\delta_n)S(\psi)x_0\| < +\infty$ which implies the convergence of $\sum_{n=1}^{\infty} |c_n|$. Since the converse is obvious the assertion is completely proved.

The crucial point in the proof is to show that

$$\sup_{0 \neq \delta \in \Sigma} \frac{\|P(\delta)x_0\|}{\mu(\delta)} < +\infty.$$

Suppose there exists a sequence $\eta_n \in \Sigma$, for which $\|P(\eta_n)x_0\|/\mu(\eta_n) \geq n; n = 1, 2, \dots$, and $\mu(\eta_n) \neq 0$. We shall construct by induction another sequence $\{\sigma_n\}$ with the properties

- (i) $\sigma_n \cap \sigma_i$ is equal to σ_n or \emptyset ;
- (ii) $\|P(\sigma_n)x_0\|/\mu(\sigma_n) \geq n; 1 \leq i \leq n-1$.

Indeed, set $\sigma_1 = \eta_1$ and assume that $\sigma_1, \dots, \sigma_n$ are already constructed and satisfying the conditions (i) and (ii). The following equality

$$\eta_{n+1} = \left(\eta_{n+1} - \bigcup_{k=1}^n \sigma_k\right) \cup (\eta_{n+1} \cap \sigma_n) \cup \bigcup_{k=1}^{n-1} \left(\eta_{n+1} \cap \left(\sigma_k - \bigcup_{j=k+1}^n \sigma_j\right)\right)$$

splits η_{n+1} into $n+1$ disjoint sets; hence

$$\begin{aligned} & \left\| P\left(\eta_{n+1} - \bigcup_{k=1}^n \sigma_k\right)x_0 \right\| + \|P(\eta_{n+1} \cap \sigma_n)x_0\| \\ & \quad + \sum_{k=1}^{n-1} \left\| P\left(\eta_{n+1} \cap \left(\sigma_k - \bigcup_{j=k+1}^n \sigma_j\right)\right)x_0 \right\| \\ & \geq \|P(\eta_{n+1})x_0\| \geq (n+1)\mu(\eta_{n+1}) \\ & = (n+1) \left[\mu\left(\eta_{n+1} - \bigcup_{k=1}^n \sigma_k\right) + \mu(\eta_{n+1} \cap \sigma_n) \right. \\ & \quad \left. + \sum_{k=1}^{n-1} \mu\left(\eta_{n+1} \cap \left(\sigma_k - \bigcup_{j=k+1}^n \sigma_j\right)\right) \right]. \end{aligned}$$

Thus, at least for one of these disjoint sets, which will be denoted η_{n+1} we have $\|P(\eta_{n+1})x_0\| \geq (n+1)\mu(\sigma_{n+1})$ and in this way all conditions imposed on $\{\sigma_n\}$ will hold.

Condition (i) satisfied by the sequence $\{\sigma_n\}$ shows that it contains either a subsequence of disjoint sets or a nested subsequence $\{\sigma_{n_j}\}$. The first possibility leads immediately to a contradiction of condition (ii) while in the second case we will have the basis

$$\{P(\sigma_{n_j} - \sigma_{n_{j+1}})x_0/\mu(\sigma_{n_j} - \sigma_{n_{j+1}})\}$$

which is equivalent to the natural basis of l_1 . In view of the closed graph theorem this implies the existence of a constant A such that:

$$\left\| \sum_{j=1}^{\infty} \alpha_j \frac{P(\sigma_{n_j} - \sigma_{n_{j+1}})x_0}{\mu(\sigma_{n_j} - \sigma_{n_{j+1}})} \right\| \leq A \sum_{j=1}^{\infty} |\alpha_j|$$

for any sequence $(\alpha_j) \in l_1$. Taking $\alpha_j = \mu(\sigma_{n_j} - \sigma_{n_{j+1}})/\mu(\sigma_{n_k} - \sigma_{n_{m+1}})$; $k \leq j \leq m$, we obtain

$$\left\| \sum_{j=k}^m \frac{P(\sigma_{n_j} - \sigma_{n_{j+1}})x_0}{\mu(\sigma_{n_k} - \sigma_{n_{m+1}})} \right\| \leq A$$

i.e.

$$\left\| \frac{P(\sigma_{n_k})x_0 - P(\sigma_{n_{m+1}})x_0}{\mu(\sigma_{n_k}) - \mu(\sigma_{n_{m+1}})} \right\| \leq A.$$

But $\mu(\sigma_{n_{m+1}}) \leq K\|x_0\|/n_{m+1}$ i.e. $\lim_{m \rightarrow \infty} \mu(\sigma_{n_{m+1}}) = 0$ and therefore $\lim_{m \rightarrow \infty} \|P(\sigma_{n_{m+1}})x_0\| = 0$. Thus $\|P(\sigma_{n_k})\|/\mu(\sigma_{n_k}) \leq A$; $k = 1, 2, \dots$, which contradicts again condition (ii).

In conclusion we have proved the existence of a constant L such that

$$\|P(\delta)x_0\|/\mu(\delta) \leq L; \quad \delta \in \Sigma; \mu(\delta) \neq 0.$$

Finally, let f be a simple function; one can easily see that

$$\begin{aligned} \frac{1}{K\|x_0^*\|} \int_{\Omega} |f(\omega)| \mu(d\omega) &\leq \frac{1}{K} \|S(|f|)x_0\| \leq \|S(f)x_0\| \\ &\leq L \int_{\Omega} |f(\omega)| \mu(d\omega) \end{aligned}$$

which shows that $X = \mathfrak{M}(x_0)$ is isomorphic to $L_1(\Omega, \Sigma, \mu)$.

Let $X = \mathfrak{M}(x_0)$ be a cyclic space relative to a σ -complete Boolean algebra of projections \mathfrak{B} (regarded as a spectral measure $P(\cdot)$ on (Ω, Σ) and x_0^* a Bade functional. It is quite clear that for any $x_{\phi} = S(\phi)x_0 \in X$; $\phi(\omega) > 0$; $\omega \in \Omega$, we have $\mathfrak{M}(x_0) = \mathfrak{M}(x_{\phi})$. Considering the positive measure $\nu_{\phi}(\sigma) = x_0^*P(\sigma)x_{\phi}$ we can define the conditional expectation $E(\Sigma_0, \nu_{\phi})$ relative to a subring Σ_0 of Σ as the operator in $\mathfrak{M}(x_{\phi})$

$= X$ which assigns to $S(f)x_\phi \in \mathfrak{M}(x_\phi)$ the vector $E(\Sigma_0, \nu_\phi)S(f)x_\phi = S(h)x_\phi$ where h would be the Radon-Nikodym derivative of the measure $x_0^*P(\sigma)S(f)x_\phi$; $\sigma \in \Sigma_0$, with respect to the restriction of ν_ϕ to Σ_0 ($S(f)x_\phi$ belongs to the domain of $E(\Sigma_0, \nu_\phi)$ if and only if $x_\phi \in D(S(h))$).

Now the previous theorem can be restated as follows:

THEOREM 2. *A Banach space X is isomorphic to an L_1 -space over a finite measure space if and only if:*

(a) *X is a cyclic space $\mathfrak{M}(x_0)$; $x_0 \in X$, relative to some σ -complete Boolean algebra of projections \mathfrak{B} , and*

(b) *there exists a Bade functional $x_0^* \in X^*$ such that for every subring Σ_0 of Σ and $x_\phi = S(\phi)x_0 \in X$; $\phi(\omega) > 0$; $\omega \in \Omega$, the conditional expectation $E(\Sigma_0, \nu_\phi)$ is a linear bounded projection in $X = \mathfrak{M}(x_\phi)$.*

PROOF. It suffices to observe that the series involved in the statement of Theorem 1 converges and its sum is $E(\Sigma_0, \nu_\phi)S(f\phi^{-1})x_\phi$ where Σ_0 is the subring generated by the sets σ_n ; $n = 1, 2, \dots$. Q.E.D.

REMARKS. 1. Using Banach function spaces instead of cyclic spaces might simplify the statement of Theorem 2 but only a sufficient condition for such spaces to be isomorphic to an L_1 -space can be obtained. The precise assertion appears in the introduction.

2. In defining the measures ν_ϕ we use the same Bade functional x_0^* ; if instead we set $\lambda_\phi(\cdot) = x_\phi^*P(\cdot)x_\phi$, where x_ϕ^* depends on x_ϕ , L_p will be isometric to an L_p -space; $1 \leq p < +\infty$, provided all the conditional expectations $E(\Sigma_0, \lambda_\phi)$ will be contractive projections in L_p (cf. T. Ando [1]).

3. The problems discussed in this paper are related to the so called "leveling property" of a norm ρ in a Banach function space (cf. H. W. Ellis and I. Halperin [5]) and to the property (J) introduced by N. E. Gretskey [6]. It follows from Theorem 1 that unless a weakly sequentially complete Banach function space is isomorphic to an L_1 -space, there exists always an isometric representation L_{ρ_ϕ} of L_ρ in which $\sup_\phi \rho_\phi(f) = +\infty$ (in the notation of [6]), and consequently ρ_ϕ does not admit an equivalent rearrangement-invariant norm with respect to the measure ν_ϕ (cf. W. A. J. Luxemburg [8, Theorem 14.4]).

REFERENCES

1. T. Ando, *Banachverbände und positive Projektionen*, Math. Z. 109 (1969), 121-130.
2. W. G. Bade, *On Boolean algebras of projections and algebras of operators*, Trans. Amer. Math. Soc. 80 (1955), 345-360. MR 17, 513.
3. ———, *A multiplicity theory for Boolean algebras of projections in Banach spaces*, Trans. Amer. Math. Soc. 92 (1959), 508-530. MR 21 #7443.

4. N. Dunford, *Spectral operators*, Pacific J. Math. **4** (1954), 321–354. MR **16**, 142.
5. H. W. Ellis and I. Halperin, *Function spaces determined by a levelling length function*, Canad. J. Math. **5** (1953), 576–592. MR **15**, 439.
6. N. E. Gretskey, *Representation theorems on Banach function spaces*, Bull. Amer. Math. Soc. **74** (1968), 705–709. MR **37** #2009.
7. M. I. Kadec and A. Pelczyński, *Bases, lacunary sequences and complemented subspaces in the spaces L_p* , Studia Math. **21** (1961/62), 161–176. MR **27** #2851.
8. W. A. J. Luxemburg, *Rearrangement-invariant Banach function spaces*, Proc. Sympos. Anal. (Queen's Univ., 1967), Queen's Papers in Pure and Appl. Math., no. 10, 1967, pp. 88–144.
9. W. A. J. Luxemburg and A. C. Zaanen, *Notes on Banach function spaces. I*, Nederl. Akad. Wetensch. Proc. Ser. A **66** = Indag. Math. **25** (1963), 135–147. MR **26** #6723a.
10. I. Singer, *Some characterizations of symmetric bases in Banach spaces*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **10** (1962), 185–192. MR **26** #5393.
11. L. Tzafriri, *An isomorphic characterization of L_p and c_0 -spaces*, Studia Math. **32** (1969), 287–296.

NORTHWESTERN UNIVERSITY, EVANSTON, ILLINOIS 60201

UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON 98105