



# Connected Weak Edge Detour Number of a Graph

J. M. Prabakar\* and S. Athisayanathan†

## Abstract

Certain general properties of the *detour distance*, *weak edge detour set*, *connected weak edge detour set*, *connected weak edge detour number* and *connected weak edge detour basis* of graphs are studied in this paper. Their relationship with the detour diameter is discussed. It is proved that for each pair of integers  $k$  and  $n$  with  $2 \leq k \leq n$ , there exists a connected graph  $G$  of order  $n$  with  $cdn_w(G) = k$ . It is also proved that for any three positive integers  $R, D, k$  such that  $k \geq D$  and  $R < D \leq 2R$ , there exists a connected graph  $G$  with  $rad_D G = R$ ,  $diam_D G = D$  and  $cdn_w(G) = k$ .

**Keywords:** Detour, Detour number, Weak edge detour number, Connected weak edge detour number

Mathematics Subject Classification (2010): 05C12

## 1. Introduction

Graphs are discrete structures that represent objects and their relations among them. For a *graph*  $G = (V, E)$ , with the vertex (object) set  $V$  and edge set, i.e., the set of relations,  $E$ , the order and size of  $G$  are denoted by  $n$  and  $m$  respectively. For basic definitions and terminologies we refer to [4, 1]. Throughout this paper  $G$  denotes a finite undirected connected simple graph with at least two vertices.

For vertices  $u$  and  $v$  in  $G$ , the *distance*  $d(u, v)$  is the length of a shortest  $u-v$  path in  $G$ . A  $u-v$  path of length  $d(u, v)$  is called a  $u-v$  *geodesic*. For a vertex  $v$  of  $G$ , the *eccentricity*  $e(v)$  is the distance between  $v$  and a vertex farthest from  $v$ . The minimum eccentricity among the vertices

---

\*St. Xavier's College (Autonomous), Palayamkottai 627 002; [jmpsoxc@gmail.com](mailto:jmpsoxc@gmail.com)

†St. Xavier's College; [athisxc@gmail.com](mailto:athisxc@gmail.com)

of  $G$  is the *radius*,  $rad G$  and the maximum eccentricity is its *diameter*,  $diam G$  of  $G$ .

The *detour distance*  $D(u, v)$  is the length of a longest  $u - v$  path in  $G$  for vertices  $u$  and  $v$  in  $G$ . A  $u - v$  path of length  $D(u, v)$  is called a  $u - v$  *detour*. For a vertex  $v$  of  $G$ , the *detour eccentricity*  $e_D(v)$  is the detour distance between  $v$  and a vertex farthest from  $v$ . The *detour radius*,  $rad_D G$  of  $G$  is the minimum detour eccentricity among the vertices of  $G$ , while the *detour diameter*,  $diam_D G$  of  $G$  is the maximum detour eccentricity among the vertices of  $G$ . These concepts were studied by Chartrand *et al.* [2]

A vertex  $x$  is said to lie on a  $u - v$  detour  $P$  if  $x$  is a vertex of  $P$  including the vertices  $u$  and  $v$ . A set  $S \subseteq V$  is called a *detour set* if every vertex  $v$  in  $G$  lies on a detour joining a pair of vertices of  $S$ . The *detour number*  $dn(G)$  of  $G$  is the minimum order of a detour sets and any detour set of order  $dn(G)$  is called a *detour basis* of  $G$ . A vertex  $v$  that belongs to every detour basis of  $G$  is a *detour vertex* in  $G$ . If  $G$  has a unique detour basis  $S$ , then every vertex in  $S$  is a detour vertex in  $G$ . [3]

A set  $S \subseteq V$  is called a *weak edge detour set* of  $G$  if every edge in  $G$  has both its ends in  $S$  or it lies on a detour joining a pair of vertices of  $S$ . The *weak edge detour number*  $dn_w(G)$  of  $G$  is the minimum order of its weak edge detour sets and any weak edge set of order  $dn_w(G)$  is called a *weak edge detour basis* of  $G$ . These concepts were studied by Santhakumaran and Athisayanathan. [5]

A set  $S \subseteq V$  is called a *connected detour set* of  $G$  if  $S$  is a detour set of  $G$  and the subgraph  $G \langle S \rangle$  induced by  $S$  is connected. The *connected detour number*  $cdn(G)$  of  $G$  is the minimum order of its connected detour sets and any connected detour set of order  $cdn(G)$  is called *connected detour basis* of  $G$ . [6] This motivated us to introduce and investigate the concepts of *connected weak edge detour set* and *connected weak edge detour number* of a graph  $G$ .

The following theorems are used in this paper for proving the results.

**Theorem 1.1.** [3] *Every end-vertex of a non-trivial connected graph  $G$  belongs to every detour set of  $G$ . Also if the set  $S$  of all end-vertices of  $G$  is a detour set, then  $S$  is the unique detour basis for  $G$ .*

**Theorem 1.2.** [5] *Every end-vertex of a non-trivial connected graph  $G$  belongs to every weak edge detour set of  $G$ . Also if the set  $S$  of all end-vertices of  $G$  is a weak edge detour set, then  $S$  is the unique weak edge detour basis for  $G$ .*

**Theorem 1.3.** [5] *If  $T$  is a non-trivial tree with  $k$  end-vertices, then  $dn(T) = dn_w(T) = k$ .*

## 2. Connected Weak Edge Detour Number of a Graph

**Definition 2.1.** Let  $G = (V, E)$  be a connected graph with at least two vertices. A set  $S \subseteq V$  is a connected weak edge detour set of  $G$  if  $S$  is a weak edge detour set of  $G$  and the subgraph  $\langle S \rangle$  induced by  $S$  is connected. The connected weak edge detour number  $cdn_w(G)$  of  $G$  is the minimum order of its connected weak edge detour sets and any connected weak edge detour set of order  $cdn_w(G)$  is called a connected weak edge detour basis of  $G$ .

**Example 2.2.** For the graph  $G$  given in Figure 2.1, it is clear that no two element subset of  $V$  is a connected weak edge detour set of  $G$ . The set  $S = \{v_1, v_2, v_3\}$  is a connected weak edge detour basis of  $G$  so that  $cdn_w(G) = 3$ . The set  $S_1 = \{v_1, v_2, v_4\}$  and  $S_2 = \{v_1, v_3, v_5\}$  are also connected weak edge detour bases of  $G$ . Thus there can be more than one connected weak edge detour basis for a graph  $G$ .




Figure 2.1:  $G$

**Remark 2.3.** Every connected weak edge detour set is a weak edge detour set but the converse is not true. For the graph  $G$  given in figure 2.1, the set  $U = \{v_1, v_4, v_5\}$  is a weak edge detour set but not a connected weak edge detour set of  $G$ .

**Example 2.4.** For the graph  $G$  given in Figure 2.2, the set  $S_1 = \{v_2, v_3\}$  is a connected weak edge detour basis for  $G$  so that  $cdn_w(G) = dn_w(G) = 2$ .




Figure 2.2:  $G$

**Theorem 2.5.** For any graph  $G$  of order  $n \geq 2$ ,  $2 \leq cdn_w(G) \leq n$ .


*Proof.* A connected weak edge detour set needs at least two vertices so that  $cdn_w(G) \geq 2$  and the set of all vertices of  $G$  is a connected weak edge detour set of  $G$  so that  $cdn_w(G) \leq n$ . Thus  $2 \leq cdn_w(G) \leq n$ .  $\square$

**Remark 2.6.** The bounds in Theorem 2.5 are sharp. For the complete graph  $K_2$ ,  $cdn_w(K_2) = 2$ . The set of all vertices of path  $P_n$  ( $n \geq 2$ ) is


its unique connected weak edge detour set so that  $cdn_w(G) = n$ . Also the inequalities in Theorem 2.5 can be strict. For the graph  $G$  given in Figure 2.1,  $n = 5$ ,  $cdn_w(G) = 3$  so that  $2 < cdn_w(G) < n$ . Thus the complete graph  $K_2$  has the smallest possible connected weak edge detour number 2 and the non-trivial paths have the largest possible connected weak edge detour number  $n$ .

**Definition 2.7.** A vertex  $v$  in a graph  $G$  is a connected weak edge detour vertex if  $v$  belongs to every connected weak edge detour basis of  $G$ . If  $G$  has a unique connected weak edge detour basis  $S$ , then every vertex in  $S$  is a connected weak edge detour vertex of  $G$ .

**Example 2.8.** For the graph  $G$  given in Figure 2.3,  $S = \{u, v, w, x, y\}$  is the unique connected weak edge detour basis so that every vertex of  $S$  is a connected weak edge detour vertex of  $G$ .



**Example 2.9.** For the graph  $G$  given Figure 2.4,  $S_1 = \{u, s, w, t, v\}$ ,  $S_2 = \{u, s, x, t, v\}$  and  $S_3 = \{u, s, y, t, v\}$  are the connected weak edge detour bases of  $G$  so that  $u, s, t$  and  $v$  are the connected weak edge detour vertices of  $G$ .



In the following theorems we show that there are certain vertices in a non-trivial connected graph  $G$  that are connected weak edge detour vertices of  $G$ .

**Theorem 2.10.** Every end-vertex of a non-trivial connected graph  $G$  belongs to every connected weak edge detour set of  $G$ .

*Proof.* Let  $v$  be an end-vertex of  $G$  and  $uv$  an edge in  $G$  incident with  $v$ . Then  $uv$  is either an initial edge or the terminal edge of any detour containing the edge  $uv$ . Hence it follows that  $v$  belongs to every connected weak edge detour set of  $G$ . □

**Theorem 2.11.** Let  $G$  be a connected graph with cut-vertices and  $S$  a connected weak edge detour set of  $G$ . Then for any cut-vertex  $v$  of  $G$ , every component of  $G - v$  contains an element of  $S$ .

*Proof.* Let  $v$  be a cut-vertex of  $G$  such that one of the components, say  $C$  of  $G - v$  contains no vertex of  $S$ . Then by Theorem 2.10,  $C$  does not contain any end-vertex of  $G$ . Hence  $C$  contains at least one edge, say  $uw$ . Since  $S$  is a connected weak edge detour set there exists vertices  $x, y \in S$  such that  $uw$  lies on some  $x - y$  detour  $P : x = u_0, u_1, \dots, u, w, \dots, u_t = y$  in  $G$  or both the ends  $u$  and  $w$  of the edge  $uw$  are in  $S$ . Suppose that  $uw$  lies on the detour  $P$ . Let  $P_1$  be the  $x - u$  subpath of  $P$  and  $P_2$  be the  $u - y$  subpath of  $P$ . Since  $v$  is a cut-vertex of  $G$  both  $P_1$  and  $P_2$  contain  $v$  so that  $P$  is not a detour, which is a contradiction. Suppose that  $u$  and  $w$  are in  $S$ , then  $C$  contains vertices of  $S$ , which is again a contradiction.  $\square$


**Theorem 2.12.** *Let  $G$  be a connected graph with cut-vertices. Then every cut-vertex of  $G$  belongs to every connected weak edge detour set of  $G$ .*

*Proof.* Let  $G$  be a connected graph and  $v$  be a cut-vertex of  $G$ . Let  $G_1, G_2, \dots, G_k$  ( $k \geq 2$ ) be the components of  $G - v$ . Let  $S$  be any connected weak edge detour set of  $G$ . Then by Theorem 2.11,  $S$  contains at least one element from each component  $G_i$  ( $1 \leq i \leq k$ ) of  $G - v$ . Since  $\langle S \rangle$  is connected it follows that  $v \in S$ .  $\square$

**Corollary 2.13.** *All the end-vertices and the cut-vertices of a connected graph  $G$  belong to every connected weak edge detour set of  $G$ .*

*Proof.* Proof is immediate from the Theorems 2.10 and 2.12.  $\square$

**Remark 2.14.** *For the graph  $G$  given in Figure 2.5,  $S_1 = \{u, w, x\}$ ,  $S_2 = \{u, w, y\}$ ,  $S_3 = \{v, w, x\}$  and  $S_4 = \{v, w, y\}$  are the four connected weak edge detour bases. The cut vertex  $w$  belongs to every connected weak edge detour basis so that the cut-vertex  $w$  is the unique connected weak edge detour vertex of  $G$ .*



**Corollary 2.15.** *If  $T$  is a tree of order  $n \geq 2$ , then  $cdn_w(T) = n$ .*

*Proof.* Corollary 2.13 gives the proof.  $\square$

**Corollary 2.16.** *For any connected graph  $G$  with  $k$  end-vertices and  $l$  cut-vertices,  $\max\{2, k + l\} \leq cdn_w(G) \leq n$ .*


*Proof.* The Theorem 2.5 and the corollary 2.13 give the proof.  $\square$

For the graph  $H$  and an integer  $k \geq 1$ , we write  $kH$  for the union of the  $k$  disjoint copies of  $H$ .

**Theorem 2.17.** *Let  $G = (K_{n_1} \cup K_{n_2} \cup \dots \cup K_{n_r} \cup kK_1) + v$  be a block graph of order  $n \geq 4$  such that  $r \geq 1$ , each  $n_i \geq 2$  and  $n_1 + n_2 + \dots + n_r + k = n - 1$ . Then  $cdn_w(G) = r + k + 1$ .*

*Proof.* Let  $u_1, u_2, \dots, u_k$  be the end-vertices of  $G$ . Let  $S$  be any connected weak edge detour set of  $G$ . Then by Corollary 2.13,  $v \in S$  and  $u_i \in S (1 \leq i \leq k)$ . Also by Theorem 2.11,  $S$  contains a vertex from each component  $K_{n_i} (1 \leq i \leq r)$ . Now choose exactly one vertex  $v_i$  from each  $K_{n_i}$  such that  $v_i \in S$ . Then  $|S| \geq r + k + 1$ . Let  $T = \{v, v_1, v_2, \dots, v_r, u_1, u_2, \dots, u_k\}$ . Since every edge in  $G$  has both its ends in  $T$  or it lies on a detour joining a pair of vertices of  $T$ , it follows that  $T$  is a weak edge detour basis of  $G$ . Also, since  $\langle T \rangle$  is connected,  $cdn_w(G) = r + k + 1$ .  $\square$

**Remark 2.18.** *If the blocks of the graph  $G$  in Theorem 2.17 are not complete, then the theorem is not true. For the graph  $G$  given in Figure 2.6 there are two blocks and  $\{v_4, v_9, v_5, v_7\}$  is a connected weak edge detour basis so that  $cdn_w(G) = 4$ .*



**Theorem 2.19.** *Let  $G$  be the complete graph  $K_n (n \geq 2)$ . Then a set  $S \subseteq V$  is a connected weak edge detour basis of  $G$  if and only if  $S$  consists of any two vertices of  $G$ .*

*Proof.* Let  $G$  be the complete graph  $K_n (n \geq 2)$  and  $S = \{u, v\}$  be any set of two vertices of  $G$ . It is clear that  $D(u, v) = n - 1$ . Let  $xy \in E$ . If  $xy = uv$ , then both its ends are in  $S$ . Let  $xy \neq uv$ . If  $x \neq u$  and  $y \neq v$ , then the edge  $xy$  lies on the  $u - v$  detour  $P : u, x, y, \dots, v$  of length  $n - 1$ . If  $x = u$  and  $y \neq v$ , then the edge  $xy$  lies on the  $u - v$  detour  $P : u = x, y, \dots, v$  of length  $n - 1$ . Hence  $S$  is a connected weak edge detour of  $G$ . Since  $|S| = 2$ ,  $S$  is a connected weak edge detour basis of  $G$ .

Conversely, let  $S$  be a connected weak edge detour basis of  $G$ . Let  $S'$  be any set consisting of two vertices of  $G$ . Then as in the first part of this theorem  $S'$  is a connected weak edge detour basis of  $G$ . Hence  $|S| = |S'| = 2$  and it follows that  $S$  consists of any two vertices of  $G$ .  $\square$

**Theorem 2.20.** *Let  $G$  be a cycle of order  $n \geq 3$ . Then a set  $S \subseteq V$  is a connected weak edge detour basis of  $G$  if and only if  $S$  consists of any two adjacent vertices of  $G$ .*

*Proof.* Let  $S = \{u, v\}$  be any set of two adjacent vertices of  $G$ . It is clear that  $D(u, v) = n - 1$ . Then every edge  $e \neq uv$  of  $G$  lies on the  $u-v$  detour and the both ends of the edge  $uv$  belong to  $S$  so that  $S$  is a connected weak edge detour set of  $G$ . Since  $|S| = 2$ ,  $S$  is a connected weak edge detour basis of  $G$ .

Conversely, assume that  $S$  is a connected weak edge detour basis of  $G$ . Let  $S'$  be any set of two adjacent vertices of  $G$ . Then as in the first part of this theorem  $S'$  is a connected weak edge detour basis of  $G$ . Hence  $|S| = |S'| = 2$ . Let  $S = \{u, v\} \subseteq V$ . If  $u$  and  $v$  are not adjacent, it is clear that  $u$  and  $v$  are not connected. Thus  $S$  consists of any two adjacent vertices of  $G$ .  $\square$

**Theorem 2.21.** *Let  $G$  be the complete bipartite graph  $K_{m,n}$  ( $2 \leq m \leq n$ ). Then a set  $S \subseteq V$  is a connected weak edge detour basis of  $G$  if and only if  $S$  consists of any two adjacent vertices of  $G$ .*

*Proof.* Let  $X$  and  $Y$  be the bipartite sets of  $G$  with  $|X| = m$  and  $|Y| = n$ . Let  $S = \{u, v\}$ , where  $u \in X$  and  $v \in Y$  be any two adjacent vertices of  $G$ . It is clear that  $D(u, v) = 2m - 1$ . Then every edge  $e \neq uv$  of  $G$  lies on the  $uv$ -detour and the both ends of the edge  $uv$  belongs to  $S$  so that  $S$  is a connected weak edge detour set of  $G$ . Since  $|S| = 2$ ,  $S$  is a connected weak edge detour basis of  $G$ .

Conversely, assume that  $S$  is a connected weak edge detour basis of  $G$ . Let  $S'$  be any set of two adjacent vertices of  $G$ . Then as in the first part of this theorem  $S'$  is a connected weak edge detour basis of  $G$ . Hence  $|S| = |S'| = 2$ . Let  $S = \{u, v\} \subseteq V$ . If  $u$  and  $v \in X$  or  $Y$  it is clear that  $u$  and  $v$  are not connected. Thus  $S$  consists of any two adjacent vertices of  $G$ .  $\square$

**Corollary 2.22.** (a) *If  $G$  is the complete graph  $K_n$ , then  $cdn_w(G) = 2$ .*

(b) *If  $G$  is the complete bipartite graph  $K_{m,n}$  ( $2 \leq m \leq n$ ), then  $cdn_w(G) = 2$ .*

(c) *If  $G$  is the cycle  $C_n$ , then  $cdn_w(G) = 2$ .*

*Proof.* (a) It follows from Theorem 2.19.

(b) It follows from Theorem 2.21.

(c) It follows from Theorem 2.10.  $\square$

The following theorems give realization results.

**Theorem 2.23.** For each pair of integer  $k$  and  $n$  with  $2 \leq k \leq n$ , there exists a connected graph  $G$  of order  $n$  with  $cdn_w(G) = k$ .

*Proof.* **Case 1.**  $k = n$ . Then any tree of order  $n$  has the desired property by Corollary 2.15.

**Case 2.**  $2 = k < n$ , the cycle  $C_n$  has the desired property by Corollary 2.22 (c).

**Case 3.**  $2 < k < n$ . Let  $G$  be the graph obtained from the cycle  $C_{n-k+2} : u_1, u_2, \dots, u_{n-k+2}, u_1$  of order  $n - k + 2$  by adding  $k - 2$  new vertices  $v_1, v_2, \dots, v_{k-2}$  and joining each vertex  $v_i$  ( $1 \leq i \leq k - 2$ ) to  $u_1$ . The resulting graph  $G$  is connected of order  $n$  and is shown in Figure 2.7. Now we show that  $cdn_w(G) = k$ . Let  $S = \{u_1, v_1, v_2, \dots, v_{k-2}\}$  be the set of all end-vertices together with the cut-vertex  $u_1$  of  $G$ . It is clear that  $S$  is not a connected weak edge detour set of  $G$ . Let  $T = S \cup \{u_2\}$ . Then every edge of  $G$  has both its ends in  $T$  or it lies on a detour joining a pair of vertices of  $T$  and also  $\langle T \rangle$  is a connected so that  $T$  is a connected weak edge detour basis of  $G$ , so that  $cdn_w(G) = k$ .  $\square$




Figure 2.7:  $G$

**Theorem 2.24.** For each positive integer  $k \geq 2$  there exists a connected graph  $G$  and a vertex  $v$  of degree  $k$  in  $G$  such that  $v$  belongs to a connected weak edge detour basis of  $G$  and  $cdn_w(G) = k$ .

*Proof.* **Case 1.**  $k = 2$ , the complete graph  $K_3$  has a desired properties by Corollary 2.22 (a).

**Case 2.**  $k > 2$ , let  $G$  be the graph obtained from the complete graph  $K_3$ , where  $V(K_3) = \{v_1, v_2, v_3\}$  by adding  $k - 2$  new vertices  $u_1, u_2, \dots, u_{k-2}$  and joining  $u_i$  ( $1 \leq i \leq k - 2$ ) to  $v_1$ . The resulting graph  $G$  is connected of order  $n$  and is shown in the Figure 2.8. Then  $deg_G v_1 = k$ . Let  $S = \{u_1, u_2, \dots, u_{k-2}, v_1\}$  be the set of all end-vertices and cut-vertices. However, by Corollary 2.13,  $S$  is not a connected weak edge detour set of  $G$ . Let  $T = S \cup \{v\}$ , where  $v \in \{v_2, v_3\}$  is a vertex in  $K_3$ . Then  $T$  is a connected weak edge detour basis of  $G$  and hence so that  $cdn_w(G) = k$ .  $\square$






Figure 2.8:  $G$

**Theorem 2.25.** For every pair of positive integer  $a, b$  with  $2 \leq a \leq b$ , there exists a connected graph  $G$  such that  $dn_w(G) = a$  and  $cdn_w(G) = b$ .

*Proof.* **Case 1:**  $a = b$ , we have the following two sub cases.

**Sub case (i):**  $a = 2$ , the complete graph  $K_2$  has the desired property.

**Sub case (ii):**  $a > 2$ . Let  $C_3 : u_1, u_2, u_3$  be the cycle of length 3. Now, by adding  $a - 2$  new vertices  $v_1, v_2, \dots, v_{a-2}$  and joining the vertex  $u_2$  as shown in the Figure 2.9. Let  $S = \{v_1, v_2, \dots, v_{a-2}, u_2\}$  be the set of all end vertices and cut-vertices of  $G$ . It is clear that  $S$  is not a weak edge detour set of  $G$ . Let  $T = S \cup \{u\}$ , where  $u \in \{u_1, u_3\}$  is a vertex in  $C_3$ . Then  $T$  is a weak edge detour basis of  $G$  so that  $dn_w(G) = a$ . Also the sub graph  $\langle T \rangle$  induced by  $T$  is connected so that  $cdn_w(G) = a$ .




Figure 2.9:  $G$

**Case 2:**  $a < b$ . Let  $G$  be any tree with  $a$  end-vertices and  $b - a$  cut-vertices. Then by Theorem 1.3,  $dn_w(G) = a$  and by Corollary 2.15,  $cdn_w(G) = b$ . □

### 3. Connected Weak Edge Detour Number and Detour Diameter of a graph

In [3], an upper bound for the detour number, of a graph is given in terms of its order and detour diameter  $D$  as follows:

**Proposition A[3]** If  $G$  is a non-trivial connected graph of order  $n \geq 3$  and detour diameter  $D$ , then  $dn(G) \leq n - D + 1$ .

**Remark 3.1.** In the case of weak edge detour number  $dn_w(G)$  of a graph  $G$  it is show in [5] that, there are graphs  $G$  for which  $dn_w(G) = n - D + 1$ ,

$dn_w(G) > n - D + 1$  and  $dn_w(G) < n - D + 1$ . Similarly, in the case of connected weak edge detour number  $cdn_w(G)$  of the graph  $G$ , we show that there are graphs for which  $cdn_w(G) = n - D + 1$ ,  $cdn_w(G) < n - D + 1$  and  $cdn_w(G) > n - D + 1$ . For the graph  $G$  given in Figure 3.1(a),  $n = 6$ ,  $D = 4$ ,  $cdn_w(G) = 5$  so that  $cdn_w(G) > n - D + 1$ . For the graph  $G$  given in Figure 3.1(b),  $n = 8$ ,  $D = 4$  and  $cdn_w(G) = 5$  so that  $cdn_w(G) = n - D + 1$ . For the graph  $G$  given in Figure 3.1(c),  $n = 6$ ,  $D = 4$  and  $cdn_w(G) = 2$  so that  $cdn_w(G) < n - D + 1$ .






Figure 3.1: G

**Theorem 3.2.** Let  $G$  be a connected graph of order  $n \geq 2$ . If  $D = n - 1$ , then  $cdn_w(G) \geq n - D + 1$ .


*Proof.* For any graph  $G$ ,  $cdn_w(G) \geq 2$ . Since  $D = n - 1$ , we have  $n - D + 1 = 2$  and so  $cdn_w(G) \geq n - D + 1$ . □

**Remark 3.3.** *The converse of the Theorem 3.2 is not true. For the graph  $G$  given in Figure 3.1 (b), as in the Remark 3.1,  $cdn_w(G) = n - D + 1$ , but  $D \neq n - 1$ . Also for the graph  $G$  given in Figure 3.1 (a), as in the Remark 3.1,  $cdn_w(G) > n - D + 1$ , but  $D \neq n - 1$ .*

**Theorem 3.4.** *Let  $R, D, k$  be three positive integers such that  $k > D$  and  $R < D \leq 2R$ . Then there exists a connected graph  $G$  such that  $rad_D G = R$ ,  $diam_D G = D$  and  $cdn_w(G) = k$ .*

*Proof. Case 1:* When  $R = 1$  and  $D = 2$ , let  $G = K_{1,k-1}$ . Clearly  $rad_D G = 1$ ,  $diam_D G = 2$  and by corollary 2.15,  $cdn(G) = k$ .

*Case 2:* When  $R \geq 2$  and  $R < D \leq 2R$ , we construct a graph  $G$  with the desired properties as follows: Let  $C_{R+1} : v_0, v_1, \dots, v_R, v_0$  be a cycle of order  $R + 1$  and let  $P_{D-R+1} : u_0, u_1, \dots, u_{D-R}$  be a path of order  $D - R + 1$ . Let  $H$  be the graph obtained from  $C_{R+1}$  and  $P_{D-R+1}$  by identifying  $v_0$  of  $C_{R+1}$  with  $u_0$  of  $P_{D-R+1}$ . The required graph  $G$  is obtained from  $H$  by adding  $k - D + R - 2$  new vertices  $w_1, w_2, \dots, w_{k-D+R-2}$  to  $H$  and joining each  $w_i (1 \leq i \leq k - D + R - 2)$  to the vertex  $u_{D-R-1}$  and is shown in Figure 3.2. Clearly,  $G$  is connected such that  $rad_D G = R$  and  $diam_D G = D$ . Now, we show that  $cdn_w(G) = k$ . Let  $S = \{u_0, u_1, \dots, u_{D-R-1}, u_{D-R}, w_1, w_2, \dots, w_{k-D+R-2}\}$  be the set of all cut-vertices and end-vertices. However, by Corollary 2.13,  $S$  is not a connected weak edge detour set of  $G$ . Let  $T = S \cup \{v\}$ , where  $v \in \{v_R, v_1\}$  is a vertex in  $C_{R+1}$ . Then  $T$  is a connected weak edge detour basis of  $G$  so that  $cdn_w(G) = k$ . □



**References**

- [1] F. Buckley and F. Harary, *Distance in Graphs*. Reading, MA: Addison-Wesley, 1990.
- [2] G. Chartrand, H. Escudro and P. Zhang, "Detour Distance in Graphs," *J. Combin. Math. Combin. Comput.*, vol. 53, pp. 75-94, 2005.
- [3] G. Chartrand, L. John and P. Zhang, "The Detour Number of a graph," *Util. Math.* vol. 64, pp. 97-113, 2003.
- [4] F. Harary, *Graph Theory*. New Delhi: Narosa, 1997.
- [5] A. P. Santhakumaran and S. Athisayanathan, "Weak edge detour number of a graph," *Ars Combin.*, vol. 98, pp. 33-61, 2011.
- [6] A. P. Santhakumaran and S. Athisayanathan, "The connected detour number of a graph," *J. Combin. Math. Combin. Comput.*, vol. 69, pp. 205-218, 2009.