

Open access • Journal Article • DOI:10.1002/ESP.4434

Connectivity as an emergent property of geomorphic systems — Source link

Ellen Wohl, Gary Brierley, Daniel Cadol, Tom J. Coulthard ...+11 more authors

Institutions: Colorado State University, University of Auckland, New Mexico Institute of Mining and Technology,

University of Hull ...+9 more institutions

Published on: 01 Jan 2019 - Earth Surface Processes and Landforms (John Wiley & Sons, Ltd)

Topics: Landscape connectivity

Related papers:

- Geomorphometric assessment of spatial sediment connectivity in small Alpine catchments
- · Sediment connectivity: a framework for understanding sediment transfer at multiple scales
- · Prolegomena to sediment and flow connectivity in the landscape: A GIS and field numerical assessment
- · Indices of sediment connectivity: opportunities, challenges and limitations
- (Dis)Connectivity in catchment sediment cascades: a fresh look at the sediment delivery problem







Durham Research Online

Deposited in DRO:

15 May 2018

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Wohl, E. and Brierley, G. and Cadol, D. and Coulthard, T. and Covino, T. and Fryirs, K. and Grant, G. and Hilton, R. and Lane, S. and Magilligan, F. and Meitzen, K. and Passalazqua, P. and Poeppj, R. and Rathburn, S. and Sklar, L. (2019) 'Connectivity as an emergent property of geomorphic systems.', Earth surface processes and landforms., 44 (1). pp. 4-26.

Further information on publisher's website:

https://doi.org/10.1002/esp.4434

Publisher's copyright statement:

This is the accepted version of the following article: Wohl, E., Brierley, G., Cadol, D., Coulthard, T., Covino, T., Fryirs, K., Grant, G., Hilton, R., Lane, S., Magilligan, F., Meitzen, K., Passalazqua, P., Poeppj, R., Rathburn, S. Sklar, L. (2019). Connectivity as an Emergent Property of Geomorphic Systems. Earth Surface Processes and Landforms 44(1): 4-26 which has been published in final form at https://doi.org/10.1002/esp.4434. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a link is made to the metadata record in DRO
- $\bullet \,$ the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the full DRO policy for further details.



Connectivity as an Emergent Property of Geomorphic Systems

Journal:	Earth Surface Processes and Landforms
Manuscript ID	ESP-17-0351.R2
Wiley - Manuscript type:	State of Science
Date Submitted by the Author:	n/a
Complete List of Authors:	Wohl, Ellen; Colorado State University, Geosciences Brierley, Gary; University of Auckland, School of Environment Cadol, Daniel; New Mexico Institute of Mining and Technology, Earth and Environmental Science Coulthard, Tom; University of Hull, Geography Covino, Tim; Colorado State University, Ecosystem Science and Sustainability Fryirs, Kirstie; Macquarie University, Department of Environmental Sciences; Grant, Gordon; USDA Forest Service, Pacific Northwest Research Station Hilton, Robert; Durham University, Geography Lane, Stuart; Université de Lausanne, Institute of Earth Surface Dynamics Magilligan, Frank; Dartmouth College, Geography; Meitzen, Kimberly; Texas State University, Geography Passalacqua, Paola; The University of Texas at Austin, Dept. of Civil, Architectural, and Environmental Engineering Poeppl, Ronald; University of Vienna, Department of Geography and Regional Research Rathburn, Sara; Colorado State University, Geosciences Sklar, Leonard; Concordia University, Geography, Planning, and Environment
Keywords:	connectivity, geomorphic systems, sediment, landscape, hydrological

SCHOLARONE™ Manuscripts

Connectivity as an Emergent Property of Geomorphic Systems

Ellen Wohl, Gary Brierley, Daniel Cadol, Tom J. Coulthard, Tim Covino, Kirstie A. Fryirs, Gordon Grant, Robert G. Hilton, Stuart N. Lane, Francis J. Magilligan, Kimberly M. Meitzen, Paola Passalacqua, Ronald E. Poeppl, Sara L. Rathburn, Leonard S. Sklar

Abstract

Connectivity describes the efficiency of material transfer between geomorphic system components such as hillslopes and rivers or longitudinal segments within a river network. Representations of geomorphic systems as networks should recognize that the compartments, links, and nodes exhibit connectivity at differing scales. The historical underpinnings of connectivity in geomorphology involve management of geomorphic systems and observations linking surface processes to landform dynamics. Current work in geomorphic connectivity emphasizes hydrological, sediment, or landscape connectivity. Signatures of connectivity can be detected using diverse indicators that vary from contemporary processes to stratigraphic records or a spatial metric such as sediment yield that encompasses geomorphic processes operate over time and space. One approach to measuring connectivity is to determine the fundamental temporal and spatial scales for the phenomenon of interest and to make measurements at a sufficiently large multiple of the fundamental scales to capture reliably a representative sample. Another approach seeks to characterize how connectivity varies with scale, by applying the same metric over a wide range of scales or using statistical measures that characterize the frequency distributions of connectivity across scales. Identifying and measuring connectivity is useful in basic and applied geomorphic research and we explore the implications of connectivity for river management. Common themes and ideas that merit further research include; increased understanding of the importance of capturing landscape heterogeneity and

connectivity patterns; the potential to use graph and network theory metrics in analyzing connectivity; the need to understand which metrics best represent the physical system and its connectivity pathways, and to apply these metrics to the validation of numerical models; and the need to recognize the importance of low levels of connectivity in some situations. We emphasize the value in evaluating boundaries between components of geomorphic systems as transition zones and examining the fluxes across them to understand landscape functioning.

1. Introduction

Connectivity has become a widely used conceptual framework within geomorphology. Our primary objectives in this paper are to; (i) facilitate careful consideration of how to define and measure connectivity and disconnectivity across diverse spatial and temporal scales; (ii) explore the implications of connectivity, including the situations in which connectivity provides a useful framework or new insight, potential signatures of connectivity in geomorphic systems, and how connectivity can be used in resource management; and (iii) to highlight gaps in current understanding of connectivity and potential pathways for future research. We first introduce some basic characteristics of connectivity as viewed in a geomorphic context, then review both the historical underpinnings of and recent work on connectivity in geomorphology. We then discuss the challenges of identifying and measuring connectivity, use river basins to illustrate the management implications of connectivity, and conclude with a summary of key questions and challenges to understanding and using connectivity in a geomorphic context.

1.1. Connectivity in a geomorphic context

As the scientific study of surface processes and landforms, and as a discipline that has largely developed from geology and physical geography, geomorphology has come to focus upon the fluxes of fluids (air, water) and sediment and the landforms resulting from, and influencing, those fluxes. The term *geomorphic systems* recognizes couplings among seemingly discrete components of Earth's surface and near-surface environments, such as water and sediment fluxes from hillslopes that govern the configuration of river channels or fluxes of eolian dust that influence rates of soil formation in geographically distant locations (e.g., Martignier et al., 2013). Attention to fluxes of material through landscapes dates to the founding of geomorphology as a discipline (e.g., Gilbert, 1880). The term connectivity has become widely used to describe these fluxes within the past two decades.

Several definitions of connectivity have been proposed (Table 1). We define connectivity as the efficiency of transfer of materials between system components. Definition of system components varies between disciplines, such as between geomorphology and ecology, and in relation to the material under consideration (e.g., water versus sediment). Geomorphic systems can be represented as networks with compartments, links, and nodes. Using a drainage basin as an example, hillslopes and valley bottoms are compartments, channel segments are links, and channel junctions are nodes.

Connectivity has value as a conceptual framing for investigating the spatial and temporal variability of fluxes because it directs attention to (i) interactions among geomorphic system components that may appear to be isolated in time and space, such as how relative base level fall triggers river incision and subsequent hillslope adjustments over timespans of 10^3 - 10^4 years (Burbank et al., 1996), (ii) the response of diverse geomorphic systems to varying

inputs, such as how water and sediment fluxes from individual drainage basins respond to extreme storms as a function of characteristics such as basin size, river network structure, and the temporal sequence of extreme storms (Cenderelli and Wohl, 2003), (iii) the specific features of geomorphic systems that govern connectivity, such as the landforms that limit sediment fluxes within a drainage basin (Fryirs et al., 2007a), and (iv) how human alterations of geomorphic systems influence system behavior, such as how flow regulation and associated changes in water and sediment connectivity alter river geometry and biotic communities.

Connectivity also has value as a common framing shared among disciplines (e.g., Tetzlaff et al., 2007; Werner and McNamara, 2007; Larsen et al., 2012; Puttock et al., 2013; Hauer et al., 2016).

Connectivity is not an either/or attribute, but rather a continuum. Consequently, representations of geomorphic systems as networks must recognize that the compartments, links, and nodes exhibit connectivity at differing spatial and temporal scales and include diffuse and concentrated fluxes, and variable rates of flux (e.g., Passalacqua, 2017).

Connectivity is typically limited to some degree through time and across space, so that understanding of one extreme of the continuum, disconnectivity, is equally important (e.g., Faulkner, 2008). Components or processes that are disconnected are those that either are too remote from each other in space or time, so that a change in one component or process does not lead to change in another, or those in which a threshold must be overcome to allow connectivity: a critical shear stress must be exceeded to allow sediment transport, for example, or a flow magnitude must be exceeded to overtop the channel banks and laterally connect the channel and floodplain. The end member of disconnectivity must be treated with caution

because something that is disconnected at a short time-scale may be connected at a longer time-scale. In general, all measures of connectivity are dependent on time and space scales and are relational in the sense of describing transfers between components of a system (Grant et al., 2017).

Figure 1 illustrates the temporal aspect of connectivity in a manner similar to Schumm and Lichty's (1965) conceptualization of variables changing between dependent and independent status over diverse time scales. In this figure, sediment transport is highly connected and continuous over longer time and larger space scales, but disconnected in time and space when considered over periods of years to decades that include substantial periods of lower flow without sediment transport. Analogously, the longitudinal profile may be continuously adjusting to fluctuations in relative base level and thus longitudinally connected over cyclic time scales, but segmented by the presence of knickpoints and thus less longitudinally connected over graded and steady time scales.

Investigations of connectivity and disconnectivity in geomorphic systems can focus on fluxes of different types of materials, such as water (Bracken et al., 2013; Larsen et al., 2017) or sediment (Fryirs et al., 2007a; Bracken et al., 2015; Li et al., 2016). Investigations can emphasize features that enhance or limit connectivity, such as landforms that create physical thresholds which must be exceeded before material can move between compartments (Kondolf et al., 2006; Fryirs et al., 2007a). Alternatively, investigations can emphasize the magnitude, duration, frequency, strength, timing, or spatial extent of connectivity (e.g., Cote et al., 2009; Cavalli et al., 2013). Jaeger and Olden (2012), for example, used electrical resistance sensors to quantify

the longitudinal extent and duration of stream flow in an ephemeral channel network in Arizona, USA.

Framing connectivity in a geomorphic context provides a basis for considering both structural and functional components of the landscape. What has been referred to as structural connectivity is dependent on the position and spacing of landscape units and the extent to which they are in contact or distant from one another (Wainwright et al., 2011). Landscape units can vary from entire mountain ranges or drainage basins down to patches of land cover (e.g., forest versus grassland) or individual grass clumps on a hillslope with spatially discontinuous vegetation cover. Structural connectivity influences the thresholds of magnitude and duration necessary to create fluxes between individual landscape units. Floodplain wetlands adjacent to an active channel and at lower elevations may require a lower magnitude flood to achieve surface hydrologic connectivity with the channel than do floodplain wetlands farther from and/or higher than the channel (Galat et al., 1997; Poole et al., 2002). The occurrence of longitudinally continuous flow along intermittent or ephemeral channels in drylands depends partly on the magnitude and duration of precipitation inputs, but also on the structural connectivity governed by valley surface and subsurface geometry as this geometry creates alluvial reservoirs that must be saturated before surface flow occurs (Falke et al., 2011; Jaeger and Olden, 2012).

The assemblage and spatial pattern of landforms (i.e., type, size, and adjacency)

produces the structural, physical template from which to examine the extent to which

interactions between landforms at different spatial and temporal scales occur. For example,

Jain and Tandon (2010) and Hooke (2003) describe connectivity patterns in terms of whether

landforms are connected, partially connected or discrete. Fryirs et al. (2007a) describe the position of landforms that act as blockages within the landscape. As water flows over landforms, elements that influence structural connectivity may be modified as the landscape evolves by weathering and erosion processes. The timescale of this evolution can be rapid, such as during large mass wasting events (e.g. Korup et al., 2004), progressive over seasons and decades (e.g. Lane et al., 2017), or acting over long term timescales >10³ years (e.g. Prasicek et al., 2015).

Because we define connectivity as the efficiency of material transfer, we suggest that the structural configuration of geomorphic systems, although strongly influencing connectivity, be described as system configuration rather than structural connectivity. This leaves connectivity as referring specifically to what has been called functional connectivity.

Functional connectivity operates within this structural template. In geomorphic terms functional connectivity refers to the processes associated with the sources and fluxes of water, sediment, and solutes through a landscape and the transfer of those materials between multiple, contiguous structural components or between components of a system that are physically isolated except for relatively brief periods of connectivity (Jain and Tandon, 2010; Wainwright et al., 2011). In analyses of functional connectivity, the strength of connectivity or linkage between different parts of landscapes is considered. These linkages may be strong, weak, or non-existent (i.e., disconnected). Functional connectivity emphasizes the need to think about how the landscape limits the connectivity of the material under consideration, whether water, sediment, or nutrients. Frameworks for assessing hydrological connectivity and sediment connectivity and how these fluxes function in geomorphic terms have been developed and

applied in many different landscape settings in order to understand landscape change through time and to develop strategies for managing landscape processes (e.g., Fryirs et al., 2007b; Lane et al., 2009;). Lane and Milledge (2013), for example, used a catchment-scale model to evaluate the effect of shallow upland drains on flow hydrographs. Lisenby and Fryirs (2017a) compare the spatial distributions of landforms expected to influence coarse-sediment transport to downstream patterns of bed-sediment fining and evaluate the effects of landform-induced disconnectivity on sediment size distributions.

The configuration and state of the system under consideration strongly influence the expression of connectivity (Gran and Czuba, 2017; Rice, 2017). Increasing landscape morphological complexity can correspond to decreasing connectivity (Baartman et al., 2013), for example, and segments of a river network with wider valley bottoms can produce longitudinal and lateral disconnectivity in fluxes (Fryirs, 2013; Wohl et al., 2017b). The state of the system includes the capacity for adjustment and proximity to thresholds, as well as location within an evolutionary trajectory or spatially within a larger system (Brierley and Fryirs, 2016). Configuration and state are interrelated. Places in a river network with local sediment disconnectivity, for example, can accumulate sediment through time and become sites with higher potential for geomorphic change, or they can be areas that absorb change and limit manifestation of disturbance at off-site locations (Czuba and Foufoula-Georgiou, 2015; Lisenby and Fryirs, 2017a,b).

Structural and functional connectivity are tightly interwoven. Many studies focus on how the spatial template created by structural configuration interacts with variations in available energy to drive spatial and temporal fluctuations in functional connectivity (e.g.,

Jencso et al., 2009, 2010, 2011; Souza et al., 2016; Wohl et al., 2017). Croke et al. (2013) and Thompson et al. (2016), for example, use longitudinal variations in valley-bottom configuration and the measured and modeled extent of floodplain inundation during an extreme flood to infer connectivity between channel and floodplain. Other investigations examine how changes in structural configuration alter functional connectivity (e.g., Puttock et al., 2013; Segurado et al., 2015) or how changes in available energy or material inputs to a geomorphic system create simultaneous changes in structural configuration and functional connectivity (e.g., Wester et al., 2014; Micheletti et al., 2015). Vanacker et al. (2005) provides an example of how changes in structural configuration can alter functional connectivity by relating changes in the spatial distribution of agriculture and forested lands within a catchment in the Ecuadorian Andes to river channel response. Although the overall land use did not change, the changed spatial distribution of land use altered water and sediment connectivity within the catchment, resulting in channel narrowing, incision, and streambed fining. Wester et al. (2014) provides an example of how changes in energy and material inputs can alter structural configuration and connectivity by quantifying changes in morphodynamics and sediment transport on hillslopes following wildfire and rainstorms.

A reliable connectivity framework should allow for analysis of both the static and the dynamic aspects of landscapes, and therefore be flexible enough to consider structural configuration and functional connectivity over varying timeframes. Static frameworks provide a snapshot of how the landscape is structured and functioning at any particular point in time.

Dynamic frameworks recognize three key factors. First, the structure of the landscape can change and therefore the type, position, and pattern of landforms in a landscape can change,

producing alterations in connectivity. Second, the strength of functional connectivity is likely to change in association with changes to structural configuration. Third, structural configuration and functional connectivity may change depending on the magnitude of the disturbances that drive fluxes of water and sediment through landscapes (e.g., rainfall, floods, or mass wasting).

A common theme among investigations of connectivity is the response of a system to some change or lack of change in boundary conditions that may be external to the system (e.g., climate or tectonic inputs) or internal within the system (e.g., fluxes of water and sediment). Boundary conditions can vary depending on the time and space scales of the investigation. How such conditions, and their changes, are transferred to an output depends on the system configuration, which may also vary with time and space scales in response to the changes in flux caused by those boundary conditions (e.g., Romans et al., 2016). Thus, changes in boundary conditions can be modified – either dampened or amplified – by linked sets of processes operating within the system. Resulting outputs may always converge or, more likely, be unique but similar in magnitude and frequency. For example, a mountainous drainage basin considered over the timespan of a century has relatively fixed boundary conditions such as the river network configuration and geometry of individual valley segments. Varying boundary conditions include inputs of water and sediment from adjacent uplands. Outputs fluctuate across space and through time in a manner that reflects river network configuration and valley geometry, but also water and sediment inputs at any point in time, as well as the history of water and sediment inputs and associated changes in the alluvial configuration of the channel and floodplain within valley segments (Figure 2). This conceptualization of connectivity focuses

on how efficiently change is communicated to an output and therefore which processes and links need to be studied to effectively understand a geomorphic system.

Fluxes can also be conceptualized as information propagation, which Ruddell and Kumar (2009) define as the contribution of uncertainty-reducing or predictive information provided by the time lag history of one variable to the future value of another (Figure 3). In this conceptualization, the key questions become whether information can be propagated and how information propagation can be discontinuous in space and time (degree of connectivity), how it is propagated (processes of connectivity), and what is the transfer entropy of information propagation (defined as the asymmetric information flow between two variables, or directionality of connectivity; Schreiber, 2000).

Although structural configuration and functional connectivity are tightly interrelated, functional connectivity is the focus of most studies of connectivity in geomorphic systems.

Consequently, connectivity refers primarily to functional connectivity in the rest of this paper unless stated otherwise.

2. Connectivity research in geomorphology: origins and current focus

2.1. Historical underpinnings of connectivity in geomorphology

The historical underpinnings of connectivity in geomorphic systems emerged from two fundamental perspectives. One involves the cultural or societal management of geomorphic systems, such as river basin management for flood control, irrigation, and water supply (e.g., Kondolf et al., 2006). The other perspective involves basic and applied observations linking Earth surface processes to landform dynamics, such as source to sink connections linking

erosion, transport, and deposition to processes of hillslope and valley formation (e.g., Harvey, 1987; Anthony and Julian, 1999; Warrick et al., 2015). These origins can be traced back thousands of years and are still relevant in a contemporary context for why connectivity in geomorphic systems merits our attention (Table 2).

The earliest known societal actions seeking to understand and to measure connectivity in geomorphic systems can be traced back to at least 5,000 BC when the Sumerians and the Egyptians engineered elaborate projects to manipulate the movement and storage of river water for flood control and irrigation (Newson, 2007). These and subsequent manipulations of geomorphic systems and associated changes in connectivity were sometimes undertaken in ignorance of basic aspects of the hydrologic cycle. Perhaps the most fundamental question faced by early naturalists confronted with a river was the source of continued flow in the absence of precipitation (Tuan, 1968; Duffy, 2017). Although notions regarding the hydrological cycle can be traced back to much earlier (Duffy, 2017), it was not until Bernard Palissy formally elucidated the hydrological cycle in the late 16th century that a comprehensive account of the connectivity between the ocean, atmosphere, precipitation, and river systems was developed (Katerakis et al., 2007).

Italian and French Renaissance scholars, ca. 1400-1800, examined landscape-scale processes of erosion, transport, and deposition, and their role in creating channel networks on the landscape and major river valleys (Hugget, 2007). In the 18th and 19th centuries, Scottish geologists James Hutton and John Playfair made many of the same observations regarding the role of erosion as a dominant force on the landscape, noting that rivers are systematically ordered from smaller headwater streams to progressively larger rivers. The underlying concept

that a river is connected to, and is responsible for forming the landscape -- particularly the valley -- through which it flows, is usually attributed to Playfair (1802). During the 19th century, this perceived connectivity between the river and the landscape that it drains prompted the recognition that geomorphic effects could propagate through the landscape, linking, for example, deforestation on slopes and floods in channels (Marsh, 1864). Indigenous peoples and Asian civilizations may have recognized these forms of connectivity earlier, but the contemporary geomorphic tradition largely derives from western Europe and North America.

Early human modifications of river connectivity in the United States occurred during 19th century artificial river cutoffs and wood removal (Table 2). Human manipulations of connectivity in U.S. rivers continued with hydrologic surveys conducted by John Wesley Powell that led to the commissioning of numerous large dams on western rivers. After the pace of dam building increased to a peak in the mid-20th century in the U.S. and western Europe, subsequent recognition of the detrimental effects of altered connectivity within river corridors drove efforts to remove dams or modify their operating regime (e.g., Bednarek, 2001). Another widespread form of river engineering, the channelization of large meandering or anastomosing rivers such as the Danube (Pisut, 2002) and the Rhine (Diaz-Redondo et al., 2017) in Europe during the latter half of the 19th century also resulted in increased longitudinal connectivity and reduced lateral connectivity for water and sediment.

During this 19th century period of intensified river engineering, geomorphology was being established as a scientific discipline and with that grew a conceptual framework that described landforms relative to systems theories and linkages between process and response dynamics. G.K. Gilbert, a prominent founder of geomorphology in the United States, was the

first to discuss feedbacks among inputs, outputs, and exchanges of material and energy through geomorphic processes (Gilbert, 1877).

In their modern incarnations, the principles of geomorphic connectivity draw heavily on historical antecedent ideas of the watershed as a fundamental unit, the linkages between process and form, and the importance of understanding how materials and disturbances propagate through watersheds. A century after Gilbert's seminal 1877 publication on the geology of the Henry Mountains, for example, Chorley and Kennedy (1971) defined the geomorphic system as a process-response complex consisting of two interacting sub-systems — the morphological system and the cascading system. The morphological system includes the physical, geomorphic landforms on Earth's surface and the cascading system includes the energy and mass/material fluxes interacting with the morphology of the landforms. One sub-system cannot function independently of the other and collectively they function relative to process-response dynamics that vary in space and time, and at varied scales of space (spatial area considerations) and time (temporal period or rate consideration). Within the connectivity framework, the morphological system defines the structural connectivity and the cascading system represents the functional connectivity.

Chorley and Kennedy's (1971) coupled process-response complex, and the transfer of energy and matter between landforms and within the system, represents the first mention of connectivity in geomorphology. This work built on the classic equilibrium theory of cyclic, graded, and steady time (Schumm and Lichty, 1965) and threshold-lag-reaction-recovery equilibriums of dynamic, dynamic meta-stable, and steady time (Chorley and Kennedy 1971; Schumm, 1979; Chorley and Beckinsale, 1980; Graf, 1988).

Brunsden and Thornes (1979) advanced the concepts of process-response coupling by contending that the process of landscape change is driven by the capacity of the landscape to transmit an impulse between system components, and that the capacity is controlled by the landscape connection between components (described as path density) and the strength of the coupling (how (in)directly the impulse is transmitted). The sensitivity of landscape change is then determined by the rate of response. Highly connected and strongly coupled systems respond quickly and are commonly more morphologically complex, whereas less-connected and weakly coupled systems respond slowly and are less complex (Brunsden and Thornes, 1979). These ideas are direct predecessors of the concepts of information propagation (Ruddell and Kumar, 2009) and network-based graph theory (Heckmann et al., 2015).

Other major historical contributions linking landscape connectivity to geomorphic systems involved the development of methods and techniques for quantitatively measuring drainage basin morphometry and surface runoff (Horton, 1945; Strahler, 1952; Strahler 1954). The classic Horton (1945) and Strahler (1954) stream ordering methods also represent one of the few spatial connectivity metrics shared across biological and physical disciplines for communicating structural and functional properties of riverine ecosystems (Stanford and Ward, 1992). The foundations for quantifying stream morphometry using network- and areal-based measurements (Gardiner, 1975) provided a spatial and conceptual framework for organizing river basins relative to dominant processes, such as the production, transport, and, deposition zones described by Schumm (1977). These spatial frameworks led to advances in analyzing source-to-sink sediment budget and sediment yield dynamics (Trimble, 1977, 1983; Walling,

1983) and later to disturbance-driven geomorphic process-domains organized along a river continuum (Montgomery, 1999).

Geomorphic understandings of process-response coupling and its role in landscape sensitivity to change (Brunsden, 1993, 2001; Harvey, 1997, 2002; Nakamura et al., 2000) and sediment budget fluxes (Dietrich and Dunne, 1978; Walling, 1983; Reid and Dunne, 2003, 2016) are increasingly incorporated within broader inter- and multi-disciplinary programs that integrate perspectives from hydrology, biology, ecology, and biogeochemistry with a focus on connectivity relationships (Wohl et al., 2017a). Human manipulations of land cover, topography, and river corridors have strongly altered connectivity across diverse landscapes (Pringle, 2003; Hooke, 2006; Fryirs, 2013). River restoration is now the most widely practiced management action explicitly designed to mitigate some of the negative aspects of past human-induced alterations of connectivity (Buijse et al., 2002; Kondolf et al., 2006; Magilligan et al., 2016). Consequently, it is important to highlight that although recognition of the importance of geomorphic connectivity may seem like a relatively recent development, its roots are deep.

2.2. Current work on connectivity

In the last two decades, research using connectivity as a conceptual framework has experienced a boom in geomorphology, developing new or adapting already existing concepts of connectivity to better understand system complexity and response to change (e.g., Bracken et al., 2013, 2015; Poeppl et al., 2017). In this context, geomorphologists have also begun to assimilate notions of connectivity from other disciplines, especially ecology (Merriam, 1984;

Amoros and Roux, 1988; Ward and Stanford, 1989; Ward, 1997) and hydrology (Pringle, 2001, 2003) (cf., Bracken and Croke, 2007; Poeppl et al., 2017), seeking to better describe water and sediment dynamics in catchment systems (e.g., Croke et al., 2005; Brierley et al., 2006; Fryirs et al., 2007a,b; Turnbull et al., 2008; Wainwright et al., 2011; Fryirs, 2013; Gomez-Velez and Harvey, 2014; Bracken et al., 2015; Lisenby and Fryirs, 2017a,b). Depending on the respective disciplinary basis, three types of connectivity have commonly been differentiated in geomorphic contexts, although all of the types are interdependent: 1) sediment connectivity, which is the potential for sediment to move through geomorphic systems (Hooke, 2003) as governed by the physical coupling of landforms; 2) landscape connectivity, which is the physical coupling of landforms; and 3) hydrological connectivity, which describes the passage of the transporting medium from one part of the landscape to another. Structural configuration and functional connectivity are inherent in each of these types of connectivity.

Considerations of sediment connectivity in geomorphology are generally rooted in; (i) sediment budget approaches, emphasizing how the distribution of sediment stores and sinks reflect and influence the travel distances and pathways of sediment movement in geomorphic systems; or (ii) hillslope-channel connectivity (Harvey, 2012; Li et al., 2016), catchment-scale sediment tracing (Fryirs and Gore, 2013), or continuum-based approaches using the concept of hydrological connectivity (e.g., Lexartza-Artza and Wainwright, 2009; 2011; cf. Bracken et al., 2015). In an ecological context, hydrological connectivity was defined by Pringle (2001) as being the water-mediated transfer of matter, energy, and/or organisms within or between elements of the hydrologic cycle. Prior to that, stream ecology conceptual models including the river continuum concept (Vannote et al., 1980), the flood-pulse model (Junk et al., 1989), and the

serial discontinuity concept (Ward and Stanford, 1983) emphasized the ecological implications of diverse forms of connectivity. Ecological approaches to connectivity have been assimilated by hydrologists, resulting in a novel framework for understanding runoff and run on in catchment systems (e.g., Bracken and Croke, 2007; Ali and Roy, 2009).

Landscape connectivity has been defined in landscape ecology as being the degree to which a landscape facilitates or impedes the movement of individuals (Taylor et al., 1993). Similar notions regarding the role of structural landscape characteristics in a geomorphic context can be found in the coupling concept of Brunsden and Thornes (1979) and, later, in conceptualizations of the four-dimensional nature of lotic ecosystems (Ward, 1989). Brierley et al. (2006) elaborated these ideas and developed a connectivity framework in which they characterized different forms of landscape connectivity based on the position of geomorphic processes in a catchment (i.e., longitudinal, lateral, and vertical connectivity), explaining the efficiency of sediment transfer relationships within catchment systems (see also Fryirs et al., 2007a,b). In bio-geomorphic floodplain systems, the four dimensions of connectivity provide a framework to examine hydrologic-mediated exchanges of organisms, nutrients, carbon, and energy (e.g., Zueg et al., 2005; Opperman et al., 2010; Kupfer et al., 2014; Matella et al., 2015).

Also following ecological literature (e.g., Turner, 1989), geomorphologists drew a distinction between structural connectivity as the extent to which landscape units are physically linked to one another (With et al., 1997; Tischendorf and Fahrig, 2000; Turnbull et al., 2008; Wainwright et al., 2011) and functional connectivity as accounting for the way in which interactions between multiple structural characteristics affect geomorphic processes (Kimberley et al., 1997; With et al., 1997; Turnbull et al., 2008; Wainwright et al., 2011; Bracken

et al., 2015). Recent studies have suggested that geomorphic system response to change can be governed by feedback relationships between structural configuration and functional connectivity (e.g., Turnbull et al., 2008; Wainwright et al., 2011; Bracken et al. 2015; Poeppl et al., 2017). These structural-functional feedback relationships further drive a variety of biogeomorphic interactions in river systems (e.g., exchanges of water, sediment, and propagules) that influence coupled landforms and development of biotic communities (e.g., Hupp and Bornette, 2003; Osterkamp and Hupp, 2010; Meitzen and Kupfer, 2016).

3. Identifying signatures of connectivity in the geomorphic record

One of the challenges of a conceptual framework designed around connectivity is to identify signatures of differing degrees of connectivity in contemporary geomorphic processes and in sedimentary or other records of past processes. The first instinct when looking for a signature of connectivity is to detect changes in a measurement that corresponds to, for example, an input to the system. From a hydrological perspective, this may be looking for a peak in a hydrograph in response to a storm. From a sediment perspective, this could be identifying a pulse of increased eolian dust inputs that affects rate of soil formation. From a geomorphic perspective this may not be quite so straightforward, however, for at least two reasons. First, a peak in, for instance, water or sediment output at a point in space partly reflects what is happening in the basin above that point, but several different combinations of events or circumstances may give rise to this response (equifinality) (Chorley, 1962). Second, the geomorphic response is governed by the availability of transporting mechanisms such as water but also the supply of sediment and the landscape configuration as shaped by the history

of sediment-transporting flows (e.g., Harvey, 1997; Cenderelli and Wohl, 2003). This implies that the geomorphic system has a more effective memory of past events than the hydrological system, in which memory can (literally) evaporate. Therefore, the connectivity signature may be better represented with a spatial metric that encompasses how geomorphic processes operate over time and space. Examples of this include DEMs of difference (DoD) in which topographies from different time periods can be compared to indicate where there have been elevation changes and thus erosion and deposition (Lane et al., 1994; Wheaton et al., 2010). The use of this method has been greatly aided by the recent widespread availability of high resolution lidar topographic data (Jones et al., 2007; Passalacqua et al., 2015; Clubb et al., 2017).

Geomorphic responses that represent changes in connectivity are highly non-linear. Commonly controlled by erosional thresholds such as slope failure angles or entrainment thresholds in bedload transport, the response of a landscape or drainage basin to different magnitude forcings can thus be complex. Evidence of this is widespread throughout geomorphic studies, dating to Schumm's work on complex response (Schumm, 1973) as well as more recent modeling work (Coulthard and Van De Wiel, 2007). Modeling the geomorphic response of basins to climate change, Coulthard et al. (2012) show how increases in rainfall magnitude lead to linear increases in water outputs but exponential increases in sediment delivery. This is driven partly by thresholds in sediment transport but also by spatial and temporal changes in availability of sediment, which in turn are contingent upon the basin's past history of events.

When viewing river bedload transport, Jerolmack and Paola (2010) argue that sediment transport processes can act as a nonlinear filter which can completely erase (or 'shred') the original characteristics of an environmental signal (i.e., its relative magnitude and duration). The degree of this shredding is thought to depend on the ratio between the signal frequency and the timescale of "morphodynamic turbulence" in the system (Jerolmack and Paola, 2010). When signal frequency is shorter than the turnover induced by turbulence, the signal is lost.

This framework can also help explain why some events and/or systems can faithfully record responses to large signals (Romans et al., 2016). One example appears to be the response of suspended sediment in mountain rivers to large-scale landslide sediment inputs triggered by earthquakes (Hovius et al., 2011; Wang et al., 2015). Recent work following the 2008 Mw7.9 Wenchuan earthquake shows immediate (hourly timescale) and multi-annual increases in river suspended sediment concentration and sediment flux following the event (Wang et al., 2015). These observations mirror river suspended sediment data following the 1999 Chi-Chi earthquake in Taiwan (Hovius et al., 2011) and records of sand, silt, and mud accumulation in lakes fed by catchments draining the Alpine Fault in New Zealand over the last ~1000 years (Howarth et al., 2012). These steep mountain catchments have elements of structural connectivity that can greatly enhance the transfer of landslide sediment to river channels (Li et al., 2016). In addition, the erosion and transfer of suspended sediment viewed from the framework of Jerolmack and Paola (2010) may be considered as the morphodynamic equivalent of laminar flows; i.e., the non-linear responses may be less important. Clearly, the internal operation of sediment transport systems needs to be considered when examining records in the context of understanding connectivity.

Choice of metric will also heavily influence the signature. Above, we used the example of sediment output at a point, but other metrics commonly used include slope-area products and hypsometric curves (Sternai et al., 2011; Hancock et al., 2016). Although these provide useful overall indicators of different landscape shapes or form, such metrics can be relatively insensitive to alterations and changes in the landscape that are highly apparent in a visual comparison. For example, Hancock et al. (2016) show how simulated landscapes with very different drainage networks and forms have very similar landscape statistics. Furthermore, landscapes and sedimentary records are palimpsest in that they can be erased and re-written. Therefore, a record of past changes, and indeed a whole landscape, may be incomplete as an indicator of what has driven its final form.

All of the above issues are likely to be affected by the scale of study. For example, simple first-order streams, with limited degrees of freedom to store sediment and then allow this sediment to be re-mobilized, may respond more linearly to forcing than larger second-, third- or higher-order streams (e.g., Trimble, 2013). Here, larger expanses of, for example, floodplain may absorb any stratigraphic change that represents a signature of connectivity or generate false signals through autogenic processes. Temporally, this also affects what type of signature we are looking for. A tectonic signal operating over a long time scale may override the autogenic processes and other factors, muddying or masking the signal. But these processes and factors may be very important when looking for the connectivity signature from a large event (Goodbred, 2003; Jain and Tandon, 2010).

Continual deposition has the potential to serve as an indicator of connectivity, but only at the temporal and spatial scale of the deposit. The depositional record of a limited area does

not indicate how far up channel or gradient the connectivity may have extended into the transport and erosional zones, although sediment fingerprinting (Walling et al., 1999) can be used to infer the extent of longitudinal connectivity. Because of thresholds and complex response, however, a lack of uniformity does not necessarily indicate disconnectivity except at the smallest time scales of depositional processes. Examples include autogenic rhythmites at time scales of 1 to 100 seconds; slip-face bedding planes at hourly to weekly time scales; and repeating sequences resulting from complex basin response at annual to centennial time scales (Schumm, 1981). In all of these cases, the flux driving the deposition could be described as disconnected at time scales smaller than the signal frequency, but connected at greater time scales.

Uniformity and cyclicity can thus be considered indicators of connectivity within the lateral extent of a deposit over the relevant time scale, but their absence does not preclude connectivity. A reach in steady-state equilibrium, which is passing the exact amount of sediment received, is certainly well connected, but it will leave no trace of its role as a connecting part of the landscape. A basin may receive a particular sequence of sediments from its connected source areas, but the lack of repeating cycles does not necessarily negate its connectivity.

Thus, a depositional approach to identifying connectivity is limited to the spatial extent of the deposit or to the linkages that can be inferred from sediment characteristics via techniques such as sediment fingerprinting. Similarly, nothing can be said with certainty about connectivity below the temporal resolution of the stratigraphy. If a daily pulse of sediment slowly builds a delta, then the system would be disconnected at some time scale shorter than a

day, but connected at any scales longer than a day. The bedding resolution is the temporal dividing line. Viewed in the opposite sense, if connectivity is a critical filter in the interpretation of upstream forcing (e.g., climate), inferring changes in that forcing without considering connectivity may be incorrect (Lane et al., 2017).

In summary, issues around spatial and temporal scale of measurements or depositional records, as well as the existence of equifinality and nonlinearity in geomorphic systems, pose fundamental challenges to identifying connectivity. Consequently, the methods used to identify connectivity vary substantially among studies in relation to the specific aspects of connectivity under consideration (Table 3). This is unlikely to change in the future. Most of the methods listed in Table 3 are based on inferred connectivity as reflected in landscape changes through time or as simulated using numerical models calibrated against datasets that span limited time and space scales. Although the list in Table 3 is not exhaustive, the relative proportions of methods relying on direct measurements of fluxes versus inferred fluxes represent the proportions of these approaches in the geomorphic literature.

4. Measuring connectivity

Another basic challenge of a conceptual framework designed around connectivity is to quantify fluxes that reflect connectivity. Although connectivity provides a powerful conceptual framework for understanding geomorphic systems, there is currently a lack of consensus on how to measure and to compare connectivity quantitatively across temporal and spatial scales and between geomorphic systems (Bracken et al., 2013; Wohl, 2017; Table 1). This may be unavoidable given the issues discussed above that arise from the interest in diverse aspects of

connectivity. At some level, however, the lack of consensus on connectivity metrics gives rise to

many challenging questions we are currently unable to answer quantitatively. Examples include questions of broad scope: Is there a spatial scale at which landscape connectivity is most sensitive to human influences (Vanacker et al., 2005)?; where and when is restoring connectivity an appropriate strategy (Kondolf et al., 2006)?' or, under what conditions are Eulerian versus Lagrangian frameworks more appropriate to developing insights into a particular geomorphic system (Doyle and Ensign, 2009)? Examples of more specific questions include: are deltas inherently more connected than dendritic drainage networks (Passalacqua, 2017)?; or can factors that determine thresholds governing longitudinal connectivity of mobile large wood in a river network be quantitatively predicted (Kramer and Wohl, 2017)? Although many studies have quantified connectivity, most have used approaches developed for the specific question at hand (Bracken et al., 2013, 2015; Wohl, 2017). Few studies have sought to develop connectivity metrics that are intended to be general and widely applicable. In this section, we review the wide variety of published methods for quantifying connectivity in geomorphic systems, and explore the opportunities and challenges for developing a general approach to measuring this elusive but vital attribute of landscapes.

Many questions arise in considering how best to measure connectivity, starting with whether connectivity is the state of a geomorphic system (structural connectivity) or a measurable flux (process connectivity)? Is it possible to quantify the essential aspects of connectivity in a single general metric, or are the dominant controls and manifestations of connectivity so varied that site-specific or process-specific metrics will always be needed (Blue and Brierley, 2016)? Are structural connectivity and functional connectivity more or less

amenable to a standardized measurement approach? How sensitive are connectivity metrics to the methods and tools of data collection and the temporal and spatial scales of analysis? Should connectivity be measured directly or is it sufficient to quantify it indirectly, by measuring the factors that influence connectivity or its effects on landforms and material fluxes? Do we need a suite of metrics that can capture the cause and effect relationships among the drivers, attributes, and effects of connectivity? Can we as geomorphologists effectively forecast how connectivity relationships will alter geomorphic forms, processes, and fluxes brought about by climate and land use change? To address these questions, we begin by considering previously published connectivity metrics within a cause and effect framework, according to whether they provide a direct measure of connectivity or indirectly quantify the effects or the causes of connectivity.

Most studies of connectivity are motivated primarily by understanding how connectivity affects specific aspects of landscape dynamics, such as the movement of sediment between hillslopes and channels or through a stream network (Fryirs and Brierley, 2001; Fryirs, 2013; Bracken et al., 2015; Gran and Czuba, 2017; Lane et al., 2017). Hence, a straightforward approach is to quantify fluxes directly and to use those measurements to infer the degree of connectivity in the transport system, which represents an Eulerian approach. This can be done at a single point such as a catchment outlet or at many locations distributed through the system. Sediment transport processes, for example, are measured using erosion plots for small-scale measurements of sediment flux (e.g., Cerdà and García-Fayos, 1997; Wainwright et al., 2000; Boix-Fayos et al., 2006) or suspended sediment sampling methods and/or bedload traps in streams and rivers for larger-scale measurements (e.g., Garcia et al., 2000; Bunte and Abt,

2005). In geomorphic connectivity research, functional connectivity is commonly inferred from measured water and sediment fluxes, either on the plot scale (e.g., Turnbull et al., 2010; Wainwright et al., 2011; Puttock et al., 2013) or on the catchment scale (e.g., Duvert et al., 2011; Lane et al., 2017). Sediment tracers have been increasingly utilized to quantify erosion and deposition of sediments and to derive structural and functional connectivity of geomorphic systems (e.g., D'Haen et al., 2013; Fryirs and Gore, 2013; Koiter et al., 2013), which represents more of a Lagrangian approach.

The sediment delivery ratio (SDR) is one of the most widely-used indirect metrics of the effects of connectivity measured at a point along the boundary of the system (Walling, 1983; Brierley et al., 2006; Fryirs, 2013; Baartman et al., 2013). The SDR quantifies the fraction of mass eroded within an upstream catchment that is transported past the catchment outlet. The SDR varies between 0 and 1, thus providing an integrated, non-dimensional measure of the degree of connectivity of sediment sources and transport pathways within the catchment.

Other studies that use output fluxes to infer upstream connectivity include Ali and Roy (2010), which measures stream discharge and infers connectivity of zones of high soil moisture, and recent work on deltaic systems which quantifies the hydrological connectivity of channels and interdistributary islands (Larsen et al., 2012; Hiatt and Passalacqua, 2015; Hiatt and Passalacqua, 2017), and relates sediment output from delta distributary channels to upstream connectivity between geomorphic elements of the delta system (Liang et al., 2016; Passalacqua, 2017).

In contrast to quantifying the bulk system output at a single point in space, metrics for the local effects of connectivity at many locations across the landscape are inherently more

complex. This approach relies on a conceptual model of the internal dynamics of the system that facilitates identifying which sub-systems to measure and the scale and density of measurements. Numerical models can overcome this challenge by predicting outcomes for every point in the landscape. Modeling approaches including cellular automata (Baartman et al., 2013; Masselink et al., 2016a; Coulthard and Van De Wiel, 2017), process-based modeling (Mueller et al., 2007), statistical models (Poeppl et al., 2012), and GIS approaches based on network theory (Lane et al., 2009; Heckmann and Schwanghart, 2013; Masselink et al., 2016b). For example, Coulthard and Van de Wiel (2017) model changes in sediment fluxes due to the cascading impacts of land use change, and infer landscape connectivity across large distances and in both upstream and downstream directions. Czuba and Foufoula-Georgiou (2015) and Gran and Czuba (2017) use a model of sand transport through a natural channel network to indirectly quantify local connectivity by defining a cluster persistence index (CPI). The CPI is calculated from the time integral of sand mass passing a point in the network and identifies locations where discrete packets of sand coalesce and disperse due to longitudinal variations in transport connectivity. Examples of field-based studies that measure the local outcomes of connectivity include: Vanacker et al. (2005), which infers changes in water and sediment connectivity from measured changes in channel geometry and grain size; Croke et al. (2013) and Thompson et al. (2016), which use the measured and modeled extent of floodplain inundation during an extreme flood to infer connectivity between channel and floodplain; and Wester et al. (2014), which measures topographic elevation changes over time in gullies following wildfire to document spatial variation in sediment transport and deposition and infer patterns of local transport connectivity.

Connectivity in geomorphic systems has also been indirectly quantified through measurements of the key drivers that promote or inhibit connections in natural and humandisturbed landscapes (Bracken and Croke, 2007; Poeppl et al., 2017). For example, Borselli et al. (2008) develop a Connectivity Index (IC) that expresses the relative sediment transport efficiency upstream and downstream of any point in the landscape, using topographic attributes such as drainage area, mean slope, and travel distance between elements. Cavalli et al. (2013) adapt the IC for use in mountainous catchments by including the effect of topographic roughness in reducing connectivity. This spatial index has been able to reconcile temporal variability in sediment export from partly glaciated basins (Micheletti and Lane, 2016). Measures of land surface roughness extracted from DEMs have been used in other studies of landscape disconnectivity, including Baartman et al. (2013), which defines a topographic Complexity Index based on local relief and slope variation, and Lane et al. (2017), which uses a pit-filling and flow-routing algorithms to assess the impact of roughness on sediment throughput. In geomorphic terms, this latter study seeks to avoid the limitations of the common hydrological approach to noise in topographic data that leads to artificial pits, or sites of disconnection. Many hydrological analyses of routing begin by filling pits such that flow continuity can be achieved. In hydrology, but particularly in geomorphology, problems can arise with doing this when real pits, sites of reduced connectivity or disconnection, are eliminated by such algorithms. Other studies that quantify geomorphic drivers of connectivity include: Rice (2017), which uses drainage area, Strahler order and other catchment attributes to predict the relative disconnectivity of tributary junctions; Cadol and Wine (2017), which infers differential connectivity between streams and riparian vegetation in various geomorphic settings defined

by measurements of valley width, topographic curvature and slope; and May et al. (2017), which describes reduced coupling between hillslopes and channels due to wider valley bottoms and gentler hillslope gradients in catchments upstream of bedrock-controlled waterfalls. IC are static representations of connectivity that can be very useful for determining areas of high and low structural connectivity within a geomorphic system under study (e.g., Nicoll and Brierley, 2017).

The connectivity of a geomorphic system can also be explicitly quantified using analytical techniques originally developed in other fields, such as network theory (Newman, 2006) and studies of percolation (Grimmett, 1989). The system must first be represented as a network composed of source or storage elements (nodes) that are connected by pathways of potential transport (links). Nodes can be pixels or other polygons in a continuous representation of a landscape, or one- to three-dimensional elements in a graphical representation of the network structure. Links can be formed uniformly with adjacent elements or specified in terms of network structure and other factors representing distance, direction, transport thresholds and transport efficiency. Once the system is defined spatially and dynamically, connectivity can be quantified using a variety of statistics that measure the central tendency or variability of connections between network elements. For example, Western et al. (2001) characterize the degree of hillslope hydrologic connectivity by defining the integral connectivity scale length (ICSL), which represents the average distance separating hillslope elements that are connected by a continuous downslope path of elements with soil moisture above a threshold value.

As networks grow in size and complexity, matrices are needed for network connectivity and flow computation. David et al. (2011) provides an example, using a matrix-based version of the traditional Muskingum method of flow routing, to develop a river network model in which lateral inflow to a river network is calculated by a land-surface model and flow in all reaches of a river network is calculated using the routing equation.

The mathematical model of a network is the graph. Graph theory, which is the study of graphs, has been applied to geomorphic systems (e.g., Haggett and Chorley, 1969; Phillips, 2012; Heckmann and Schwanghart, 2013; Marra et al., 2013; Tejedor et al., 2015) as a means of characterizing network structure and fluxes within networks, as well as simulating propagation of system changes through networks (Heckmann and Schwanghart, 2013). A graph can be formally described as G = (N, E), in which N indicates nodes and E indicates edges. A graph is represented using an adjacency matrix, which is a square matrix with as many rows and columns as there are nodes in G. Such a matrix can provide a mathematic framework for exploring functional connectivity by analyzing nodes, edges, and paths (Heckmann and Schwanghart, 2013). Kupfer et al. (2014) apply network-based graph theory to model spatial and temporal changes in lateral connectivity of a large floodplain under different flood recurrence scenarios. Meitzen and Kupfer (2015) apply this same model to examine how connectivity influences abandoned channel infilling and vegetation development patterns. Tejedor et al. (2015) use spectral graph theory to develop a quantitative framework for channel network connectivity on deltas. Building on studies of neural networks in the human brain, Passalacqua (2017) uses an adjacency matrix to quantify the interactions between channel,

levee, and island components of a delta. This approach permits an integrated evaluation of structural configuration and functional connectivity.

Cote et al. (2009) develops another direct metric of network connectivity to quantify the impact of barriers to fish migration at the catchment scale. Cote et al. (2009) defines the Dendritic Connectivity Index (DCI) based on summing the length of stream reaches linked by passable potential barriers, normalized by the total length of the stream network, which represents the probability that an organism is able to move between any two points within the network. Grill et al. (2014) builds on this work by defining a River Connectivity Index (RCI) that considers other reach attributes such as volume, habitat classification, and usability by a given species.

The preceding review illustrates the wide variety of metrics developed to quantify connectivity, both directly using techniques from network analysis and indirectly through measurements of fluxes and other outcomes of connectivity, and the topographic and other factors that drive variations in connectivity. Structural connectivity/configuration is captured most explicitly in the direct quantification of network properties, but is also implicit in many of the metrics that quantify the drivers of connectivity. On the other hand, metrics based on measuring the outcomes of connectivity primarily quantify functional connectivity. Studies that compare metrics representing different types of connectivity have the potential to quantify the cause and effect relationships at the heart of geomorphic connectivity. For example, Baartman et al. (2013) shows that Sediment Delivery Ratio, in natural and modeled catchments, declines systematically with increasing Complexity Index, thus linking structural drivers of connectivity with functional outcomes. Similarly, Beckman and Wohl (2014) shows that variations in carbon

content in fine sediment deposits, a proxy for sediment residence time and functional disconnectivity, correlate with boundary conditions including valley morphology, log jam spacing, and forest stand age. They also show that increased connectivity, through destruction of wood jams, leads to reductions in sediment deposition and channel roughness, which in turn inhibits the trapping of mobile wood that might otherwise anchor new wood jams. Thus, by linking the causes and effects of (dis)connectivity, Beckman and Wohl (2014) illustrate how feedback loops, with the potential to form multiple alternative states, can arise in connectivity dynamics.

A key question that arises when quantifying connectivity is: compared to what? In any given geomorphic system, is there a maximum or optimum level of connectivity to compare to? Non-dimensional connectivity metrics have the potential to quantify connectivity relative to a reference value. A simple example is the Sediment Delivery Ratio (Walling, 1983), which at its maximum value of 1.0 implies complete connectivity, albeit without directly quantifying any of the upstream connections or considering the timescale over which connectivity is operating. A more spatially explicit non-dimensional metric is the hydrologic connectivity parameter Tau(h) of Western et al. (2001), which is integrated to calculate the Integral Connectivity Scale Length (ICSL) described above. Tau(h) represents the probability that any two pixels separated by a distance *h* are connected by a continuous path of pixels with soil moisture above a threshold value, and thus varies between 0 and 1. Several other connectivity metrics are composed of dimensionless ratios that vary between 0 and 1, including the Dendritic and River Connectivity Indices (Cote et al., 2009; Grill et al., 2014) and the Complexity Index (Baartman et al., 2013).

making them useful for quantifying the effect of changes within the catchment, such as dam construction, but less useful for comparisons between catchments or other geomorphic systems.

Perhaps the most significant challenge in quantifying connectivity is the issue of scale. The frequency and efficiency of connections within any geomorphic system vary systematically with the temporal and spatial scale of analysis. Hence the measures of connectivity produced by any robust metric should also vary with scale. Because most transport processes are intermittent, connectivity should increase when measured over longer characteristic time scales (McGuire and McDonnell, 2010; Bracken et al., 2013, 2015). Conversely, because movement of material occurs over finite distances in any given transport event, measured values of connectivity should decrease as spatial scale increases (Western et al., 2001; McGuire and McDonnell, 2010; Bracken et al., 2013, 2015).

One approach is to determine the fundamental temporal and spatial scales for the phenomenon of interest and to make measurements at a sufficiently large multiple of the fundamental scales to capture reliably a representative sample of transport events. If landslides are a key component of sediment connectivity within a drainage basin, for example, measuring sediment connectivity across a time span that includes more than one landslide and intervening periods with other modes of sediment delivery is important (e.g., Reid et al., 2007; Cavalli et al., 2013; Dethier et al., 2016). Similarly, measurements of longitudinal hydrologic connectivity within a river during floods should incorporate a time scale that includes multiple floods (e.g., Jaeger and Olden, 2012).

Another approach seeks to characterize how connectivity varies with scale, by applying the same metric over a wide range of temporal and spatial scales. Western et al. (2001) provides an example based on multiple soil moisture datasets. Alternatively, statistical measures that characterize the frequency distributions of connectivity across scales can be used, as Ali and Roy (2010) did for soil moisture and stormflow in a humid temperate forested catchment or Sendrowski and Passalacqua (2017) did for hydrological processes on a delta influenced by river discharge, tides, and wind.

Ultimately, the tools and methods available to collect the relevant data will constrain the scales at which connectivity can be analyzed. Technological advances such as terrestrial lidar, structure-from-motion photogrammetry, wireless sensor networks, and new techniques for tracing and tracking sediment fluxes create opportunities to expand the range of scales over which connectivity can be quantified (e.g., Fonstad et al., 2013; Cavalli et al., 2013; Smith and Vericat, 2015).

Several key ideas for future directions in measuring connectivity emerge from this discussion. First is the recognition that distinct metrics are likely needed to characterize structural configuration and functional connectivity and that no single overarching metric for connectivity is likely to emerge. Second, it will be fruitful to explore combinations of metrics that can represent the cause and effect relationships that link the drivers, structures, and outcomes of geomorphic connectivity and give rise to feedbacks and emergent system behavior. Third, non-dimensional measures that characterize connectivity relative to meaningful reference values are needed to compare connectivity across scales and between systems. Finally, quantifying how connectivity varies with temporal and spatial scales of analysis

will both inform future study designs and provide insight into the nature of connectivity in diverse geomorphic systems. A single metric that represents all aspects of connectivity is unlikely, but meaningful progress can be made in the absence of such a universal metric.

5. Using connectivity

Identifying and measuring connectivity provides a useful approach in basic and applied geomorphic research (Wohl, 2017). An explicit focus on connectivity can facilitate identification of spatial and temporal disparities in material fluxes within geomorphic systems, for example, as well as enhancing understanding of mechanisms of retention of materials within a particular system or component of a system. Characterizing connectivity can provide insight into the response of geomorphic systems to disturbance, the nonlinear behavior that may result from those disturbances, and the resistance or resilience of the system to disturbances. Explicit attention to connectivity can also promote transdisciplinary approaches to understanding and communicating geomorphic process and form.

5.1. Effective approaches to the management of landscape connectivity in river basins

In this section, we explore the implications of connectivity for management of rivers.

The implications discussed here also apply to other geomorphic environments, but rivers are particularly the target of environmental management, including restoration and rehabilitation, and a more extensive literature addresses management of rivers relative to management of other geomorphic systems.

Effective river management programs seek to attain the best achievable state for a healthy and responsive river under prevailing and future conditions. Geomorphically informed river management practices incorporate flexibility and future variability in the design and implementation of management practices through articulation of open-ended and dynamic goals (Downs and Gregory, 2004; Brierley and Hooke, 2015; Brierley and Fryirs, 2016). Such planning and design exercises recognize that what has gone before influences our capacity to manage and modify rivers, but altered boundary conditions and evolutionary trajectories constrain the best achievable state and functionality that can be attained under prevailing and likely future conditions. This entails working with river morphodynamics at the reach scale, framed in relation to catchment-scale sediment and other fluxes. Landscape connectivity exerts a critical influence upon these relationships.

Understanding connectivity in relation to the morphodynamics of rivers is critical for making informed decisions in river management practice. Essentially, such understanding is concerned with the management of fluxes. In an era where forecasting river responses to a range of natural and human disturbances is critical to management and planning, understanding connectivity provides a core foundation from which to work. Forecasting where disturbance is likely to be manifest and the extent to which on-site and off-site impacts will result is critical to risk assessment and planning. Forecasting flood hazards requires an understanding of hydrological connectivity. Managing sediment hazards and legacy sediments requires an understanding of not only sediment sources, transport and deposition, but also the extent to which a catchment contains blockages or pathways of conveyance (James, 2010; Wohl, 2015). Managing contaminants or the spread of exotic flora and fauna requires that the

dispersal pathways provided by rivers and floodplains are well understood (e.g., Haycock and Burt, 1993; Coulthard and Macklin, 2003). The connectivity among the component parts of the system and the manner in which these parts fit together set the template of the dynamic physical habitat mosaic in a river corridor.

Concerns for the sediment regime of a river take account of the nature and rate of sediment generation in a particular landscape setting, and controls upon the effectiveness of erosion and transport mechanisms that move materials through river systems (Benda et al., 2004; Czuba and Foufoula-Georgiou, 2014, 2015; Wohl et al., 2015a; Schmitt et al., 2016). In the development and implementation of catchment-scale river management plans (e.g. Sear et al., 1995; Gilvear, 1999; Brierley and Fryirs, 2009; Toone et al., 2014; Wohl et al., 2015a, b), reachscale sensitivity and catchment-scale connectivity are key considerations in determining river recovery potential and the range of potential trajectories of geomorphic river adjustment (Brierley and Fryirs, 2005; Fryirs, 2013, 2017). Connectivity relationships exert a primary control upon the efficiency with which disturbance responses are mediated through catchments, and associated lag times. This may present significant constraints upon what is achievable in managing sediment flux relationships in any given catchment. The (ir)reversibility of geomorphic adjustments to river type, and appraisals of sediment flux at the catchment scale, are important considerations in assessment of likely trajectories of adjustment (Wohl, 2011; Fryirs et al., 2012; Grabowski et al., 2014; Scorpio et al., 2015; Brierley and Fryirs, 2016; Ziliani and Surian, 2016). These insights support the derivation of moving targets for management programs (Brierley and Fryirs, 2016) and help to ensure that management actions are appropriate for a particular site (Brierley and Fryirs, 2009).

Analysis of geomorphic river recovery appraises how a river has adjusted in the past and what the river is adjusting toward. The potential for river recovery following disturbance reflects a river's inherent sensitivity to change and the severity of impacts to which the system is or has been subject (e.g., Hooke, 2015; Fryirs, 2017). Multiple potential trajectories can emerge, dependent on the condition of a reach, likely responses to disturbances, prevailing, system-specific driving factors and time lags, and how connectivity relationships mediate these processes and shape the evolutionary trajectories adopted (Phillips, 2007; Fryirs et al., 2009; Standish et al., 2014; Phillips and Van Dyke, 2016).

Assessing river recovery requires that the history, pathway, and rate of adjustment of each reach in the catchment of interest is known (Kondolf and Larsen, 1995; Surian et al., 2009b; Wohl, 2011; Fryirs et al., 2012; Fryirs and Brierley, 2012, 2016; Grabowski et al., 2014; Rathburn et al., 2013, 2017). Analysis of each reach in its catchment connectivity context provides a basis to evaluate the impact of pressures and limiting factors that may inhibit or enhance river recovery on the likely future trajectories of adjustment (Brierley and Fryirs, 2005, 2009; Ziliani and Surian, 2012, 2016; Standish et al., 2014; Scorpio et al., 2015; Fryirs and Brierley, 2016). When applied effectively, catalytic management activities in certain parts of catchments may trigger recovery processes that can accelerate recovery elsewhere. Appraisals of additional, off-site impacts require that the connectivity dynamics of the system are understood. Alternatively, analysis of connectivity relationship is required to identify where certain measures may have negative off-site impacts that will damage the recovery process. This helps in choosing passive versus active restoration measures; where in a catchment activities are likely to be most successful; and the scale and form of intervention that is

required (Lane et al., 2008; Fryirs and Brierley, 2016). In some instances, connectivity relationships can be used to guide management that maintains the fully functional portions of a catchment. As an example, Lane et al. (2008) show how in an upland river basin, native woodland planting focused on well-connected tributaries could reduce coarse sediment supply rates as an alternative to downstream sediment dredging and engineering. Understanding catchment-scale, spatial and temporal sediment and hydrological connectivity provides foundational knowledge with which to forecast river recovery potential and determine what is realistically achievable at the reach-scale.

The availability of sediment for river recovery, and hence the timeframe of recovery, may vary markedly from catchment to catchment dependent on the connectivity dynamics of that catchment. Assessment of river recovery potential allows managers to assess in which reaches sediment should be retained and stored for river recovery, and where sediments can be released (Fryirs and Brierley, 2001). It is important to ensure that there is neither too much nor too little sediment to facilitate river recovery (Kondolf, 1998; Brooks and Brierley, 2004; Florsheim et al., 2006; Jacobson et al., 2009; Smith et al., 2011; Fryirs and Brierley, 2016). Conceptual models can be used to communicate stages and timeframes of geomorphic adjustment (e.g., Simon, 1989; Brierley and Fryirs, 2005; Fryirs et al., 2012; Stella et al., 2013; Cluer and Thorne, 2014; Fryirs and Brierley, 2016; Phillips and Van Dyke, 2016). These insights can be used to assess whether geomorphic adjustments are likely to be reversible, considering how channel boundary conditions, flow and sediment inputs, and connectivity relationships have changed over time. Process-based modeling applications can be used to quantify

timeframes of adjustment, using confidence limits to express potential uncertainties in future forecasts (e.g., Smith et al., 2011; Small and Doyle, 2012; Ziliani and Surian, 2016).

Given differences in landscape connectivity relationships in differing environmental and landscape settings, there is profound variability in the ways and rates with which responses to disturbances that disrupt the sediment regime are mediated through a catchment (e.g., Fryirs et al., 2007a,b; Lane et al., 2008; Surian et al., 2009a,b; Kuo and Brierley, 2013, 2014; Lisenby and Fryirs, 2017a,b). Fryirs et al. (2009) refer to this as a response gradient. Highly connected systems rapidly convey disturbance responses through the system, whereas responses to disturbances in disconnected landscapes may be absorbed within certain parts of the system (Harvey, 2002; Hooke, 2003; Fryirs et al., 2007a, b, 2009; Jain and Tandon, 2010; Fryirs, 2013).

Catchment-scale conceptual models of process interactions, connectivity and evolutionary traits provide a basis to predict responses to management interventions (Mika et al., 2010). Analysis of threatening processes helps to identify and prioritize what forms of management intervention are required in what parts of the system (Brierley and Fryirs, 2005, 2009, 2016; Czuba et al., 2014; Fryirs and Brierley, 2016; Ziliani and Surian, 2016). Such efforts seek to maximize cumulative benefits while minimizing off-site impacts of interventions (Schmidt et al., 1998). Catchment-framed analysis of connectivity provides the basis for answering questions such as:

- Where should we prioritize our efforts to enhance the recovery of systems?
- Will a treatment reach experience degrading or positive influences from upstream (e.g., sediment slugs, headcuts)?

- From where will the sediment be sourced and dispersed to enhance river recovery in the study reach? Is enhancing sediment connectivity required?
- Where should sediment conveyance be suppressed to protect other reaches and minimize off-site impacts? Is enhancing sediment disconnectivity required?
- How will rehabilitation of the treatment reach affect downstream reaches?

These questions can be answered using conceptual models, qualitative evaluations, numerical simulations, and quantitative metrics: the key point is to characterize levels of connectivity, the processes and forms that promote or retard connectivity, and the response of the geomorphic system to changes in connectivity. Answers to these questions can be used in a range of management situations, including dam construction, removal, and modification of operating regime; incursions of exotic vegetation; mining activities; land use and land cover changes; post-fire treatment priorities (Figure 4); channelized reaches that flush sediments; and inferring whether sediment slugs enhance or inhibit downstream conveyance. Six general points that may assist efforts to manage landscape connectivity in relation to concerns for sediment regime are outlined in Table 4. The most effective technique(s) for addressing each point are likely to be site-specific. Ultimately, river basin management can focus on specific processes and fluxes without regard to connectivity, but measuring and conceptualizing process and flux in a connectivity framework facilitates an understanding of how basin configuration and fluxes of material respond to varying inputs through time and across space.

5.2. Connectivity, flow regulation and river-floodplain infrastructure

Connectivity is fundamental to management plans for river restoration/rehabilitation both as a goal and as a process. Because dams, water diversions, and other infrastructure such as weirs and check dams fragment waterways, their ubiquitous global presence has had a profound effect on hydrologic, sedimentological, and ecological connectivity (Junk et al., 1989; Nilsson et al., 2005), both within the channel and across the broader riparian zone longitudinally, laterally, and vertically (Kondolf et al., 2006). To combat some of these effects, river managers, scientists, and non-governmental organizations have argued for management plans to ameliorate the effects of impoundment on watershed connectivity. In some instances, dam removal has been the preferred option as it provides the more robust opportunity for reestablishing sediment and hydrologic connectivity (Grant and Lewis, 2015; Major et al., 2017; Foley et al., 2017) while also having immediate impacts ecologically by permitting fish passage (Kornis et al., 2014; Pess et al., 2014; Magilligan et al., 2016a) or by providing the necessary sedimentological conditions for enhancing spawning habitat (Magilligan et al., 2016b). In instances where removal is not an option, watershed managers have advocated for environmental flows (Arthington et al., 2006; Bunn and Arthington, 2002) to best mimic the natural flow regime (Poff et al., 1997) with the goal of re-establishing greater hydrologic connectivity especially across the riparian zone to maintain floodplain forest communities (Rood et al., 2005) or to generate longitudinal and lateral sediment connectivity and bar formation (Schmidt et al., 2001; Topping et al., 2005). However, the reduction in connectivity associated with changes in water flow is commonly emphasized at the expense of the effects of such infrastructure and water management on sediment and sediment regime (Wohl et al.,

2015a; Gabbud and Lane, 2016), which runs the risk of introducing environmental remediation that has less than optimal effects.

6. What do we still need to know?

Throughout our discussions and literature review, we identify common themes and ideas that merit further research. First, there is increased understanding of the importance of capturing the heterogeneity of landscapes and their connectivity patterns in space and time (Fryirs and Brierley, 2009). With new technologies, such as high-resolution topographic data and ever-increasing model capabilities, we have the information needed to capture geomorphic features and thus connectivity pathways over a wide range of spatial and temporal scales (e.g., Passalacqua et al., 2015). These mechanisms of mass and information transfer need to be quantified with appropriate metrics. Although bulk measures are helpful and easy to compute, they prevent us from capturing how connectivity patterns may vary spatially and temporally. Quantifying this heterogeneity is particularly important for restoration efforts that work with process and connectivity principles (e.g., Ward et al., 2001; Kondolf et al., 2006).

Several authors have suggested that graph and network theory metrics may be helpful tools to analyze connectivity in landscapes (Heckmann et al., 2015; Cheung et al., 2016; Gran and Czuba, 2017; Lane et al., 2009, 2017; Passalacqua, 2017). These tools have proven useful in a variety of disciplines and for the analysis of many complex systems (e.g., Newman, 2010). There are obvious applications of these metrics in geomorphology. When dealing with river networks, for example, channels and tributary junctions are easily identified as links and nodes (e.g., Marra et al., 2013; Heckmann et al., 2015). In this case, the natural system essentially

maps into the mathematical model, at least in terms of structure. Thinking about network dynamics, however, and thus the fluxes along the system, mapping into a network mathematical model may not be as obvious. For example, there may be leakages in the system (e.g., due to channel-floodplain connectivity) and these losses will have to be represented in the model, either as a distributed loss along the link or by characterizing the structural configuration (e.g., levee channels) through which this transport may occur and the nonlinearities of fluxes (e.g., stage-dependent lateral connectivity). In addition, the sediment and the nutrient networks – the collection of links and nodes along which solids and solutes are transported – may not be as continuous as the water transportation network, depending on the time scale of analysis. This may call for other approaches (e.g., the dynamic tree approach of Zaliapin et al., 2010) able to represent mathematically the superposition of multiple interacting networks of different spatial structure and temporal dynamics.

We have to understand which metrics are most helpful and representative of the physical system and its connectivity pathways. These metrics are also needed for the validation of numerical models. If we can quantify connectivity pathways through a landscape, we can then use those metrics to evaluate similarity of the couplings and transport pathways in numerical results. These validated models can then be used to simulate scenarios of disturbance and change and to predict landscape response in space and time (e.g., Liang et al., 2016a,b).

Another theme that emerged in our discussions in the general tendency to promote connectivity as a desirable landscape characteristic, thus labeling disconnectivity or low degrees of connectivity as a condition to avoid. However, disconnectivity or low levels of connectivity

are present in landscapes as geomorphic features and boundaries between different process domains, not least because without them today's geomorphic processes would not be creating the long term sedimentary record of the future. In many cases, low levels of connectivity create environmental and societal benefits, such as nutrient retention and biotic uptake that improve water quality (Haycock and Burt, 1993; Wegener et al., 2017), sediment retention that increases habitat abundance and diversity (Jacobson et al., 2009), and attenuation of hydrologic fluxes that reduces flood hazards (Lininger and Latrubesse, 2016). Geomorphologists have a critical role to play in communicating to resource managers and the public the benefits that can be derived from maintaining or restoring varying forms of connectivity and disconnectivity.

Finally, instead of imposing boundaries and studying landscapes and processes in compartments, there is value in evaluating boundaries as transition zones and examining the fluxes across them to understand landscape functioning. Our equations and numerical models are commonly built in the same compartmentalized fashion. This calls for the development of equations and models able to capture process transitions and for a critical understanding of where and when connectivity or disconnectivity may be preferable to favor the long-term sustainability of our planet.

To summarize: Connectivity provides a useful conceptual framework for quantifying transfers of materials; examining factors that enhance or limit these transfers; understanding and predicting geomorphic responses to changed input and boundary conditions; and communicating understanding of geomorphic systems to resource managers and stakeholders.

Landscapes and processes that promote and retard connectivity are heterogeneous in time and space. Geomorphic systems include transitions and leakiness rather than just simple

compartments linked by fluxes. Connectivity within geomorphic systems occurs along a continuum in which levels of disconnectivity can be critical to landscape and ecosystem integrity. There is not likely to be any single connectivity metric that adequately characterizes all forms of connectivity, but the absence of a universal connectivity metric does not preclude meaningful progress in quantifying diverse forms of connectivity.

Acknowledgements

We thank the participants of the 2016 Binghamton Symposium on Connectivity in Geomorphic Systems for useful discussions of the concepts summarized in this paper. In this context, we particularly thank Douglas Jerolmack, Katherine Lininger, Daniel Scott, and Nicholas Sutfin. This manuscript benefited from comments by two anonymous reviewers and an associate editor.

References

Ali GA, Roy AG. 2009. Revisiting hydrologic sampling strategies for an accurate assessment of hydrologic connectivity in humid temperate systems. Geogr. Compass 3 (1): 350–374.

Ali GA, Roy AG. 2010. Shopping for hydrologically representative connectivity metrics in a humid temperate forested catchment. Water Resources Research 46: W12544, doi:10.1029/2010WR009442.

Amoros C, Roux AL. 1988. Interactions between water bodies within the floodplain of large rivers: function and development of connectivity. Munsterische Geographische Arbeiten 29: 125-130.

1013	Anthony EJ, Julian M. 1999. Source-to-sink sediment transfers, environmental engineering and
1014	hazard mitigation in the steep Var River catchment, French Riviera, southeastern France.
1015	Geomorphology 31: 337-354.
1016	Appling AP, Bernhardt ES, Stanford JA. 2014. Floodplain biogeochemical mosaics: a
1017	multidimensional view of alluvial soils. Journal of Geophysical Research Biogeosciences 119:
1018	1538-1553.
1019	Arthington AH, Bunn SE, Poff NL, Naiman RJ. 2006. The challenge of providing environmental
1020	flow rules to sustain river ecosystems. Ecological Applications 16: 1311-1318.
1021	Baartman JE, Masselink R, Keesstra SD, Temme AJ. 2013. Linking landscape morphological
1022	complexity and sediment connectivity. Earth Surf. Process. Landf. 38 (12): 1457–1471.
1023	Bednarek AT. 2001. Undamming rivers: a review of the ecological impacts of dam removal.
1024	Environmental Management 27: 803-814.
1025	Benda L, Andras K, Miller D, Bigelow P. 2004. Confluence effects in rivers: interactions of basin
1026	scale, network geometry, and disturbance regimes. Water Resources Research 40(5).
1027	Beven K, Germann P. 1982. Macropores and water flow in soils. Water Resources Research 18:
1028	1311-1325.
1029	Blue B, Brierley G. 2016. 'But what do you measure?' Prospects for a constructive critical

physical geography. Area 48: 190-197.

- Borselli L, Cassi P, Torri D. 2008. Prolegomena to sediment and flow connectivity in the landscape: a GIS and field numerical assessment. Catena 75: 268–277.
- Bracken LJ, Croke J. 2007. The concept of hydrological connectivity and its contribution to understanding runoff-dominated geomorphic systems. Hydrol. Process. 21 (13): 1749–1763.
- Bracken LJ, Wainwright J, Ali GA, Tetzlaff D, Smith MW, Reaney SM, Roy AG. 2013. Concepts of hydrological connectivity: research approaches, pathways and future agendas. Earth-Sci. Rev.
- 1037 119: 17–34.

277: 100-117.

41: 839-848.

- Bracken LJ, Turnbull L, Wainwright J, Bogaart P. 2015. Sediment connectivity: a framework for understanding sediment transfer at multiple scales. Earth Surf. Process. Landf. 40: 177–188.
- Brantley SL, Lebedeva MI, Balashov VN, Singha K, Sullivan PL, Stinchcomb G. 2017. Toward a conceptual model relating chemical reaction fronts to water flow paths in hills. Geomorphology
- Brierley GJ, Cohen T, Fryirs K, Brooks A. 1999. Post-European changes to the fluvial geomorphology of Bega catchment, Australia: implications for river ecology. Freshwater Biology
- Brierley GJ, Fryirs KA. 2005. Geomorphology and river management: applications of the river

styles framework. John Wiley and Sons, Chichester, UK.

Brierley G, Fryirs, K. 2009. Don't fight the site: three geomorphic considerations in catchmentscale river rehabilitation planning. Environmental Management 43: 1201-1218.

1050	Brierley GJ, Fryirs KA. 2016. The use of evolutionary trajectories to guide 'moving targets' in the
1051	management of river futures. River Research and Applications 32: 823-835.
1052	Brierley G, Fryirs K, Jain V. 2006. Landscape connectivity: the geographic basis of geomorphic
1053	applications. Area 38 (2): 165–174.
1054	Brierley G, Hooke J. 2015. Emerging geomorphic approaches to guide river management
1055	practices. Geomorphology 251: 1-5.
1056	Brooks AP, Brierley GJ. 2004. Framing realistic river rehabilitation targets in light of altered
1057	sediment supply and transport relationships: lessons from East Gippsland, Australia.
1058	Geomorphology 58: 107-123.
1059	Brunsden D, Thornes JB. 1979. Landscape sensitivity and change. Trans. Inst. Br. Geogr. NS 4:
1060	463–484.
1061	Buijse AD, Coops H, Staras M, Jans LH, Van Geest GJ, Grifts RE, Ibelings BW, Oosterberg W,
1062	Roozen FCJM. 2002. Restoration strategies for river floodplains along large lowland rivers in
1063	Europe. Freshwater Biology 47: 889-907.Bunn SE, Arthington AH. 2002. Basic principles and
1064	ecological consequences of altered flow regimes for aquatic biodiversity. Environmental
1065	Management 30: 492-507.

Bunte K, Abt SR. 2005. Effect of sampling time on measured gravel bed load transport rates in a coarse-bedded stream. Water Resources Research 41(11): W11405

New York.

Burbank DW, Leland J, Fielding E, Anderson RS, Brozovic N, Reid MR, Duncan C. 1996. Bedrock incision, rock uplift, and threshold hillslopes in the northwestern Himalayas. Nature 379: 505-510. Cadol D, Wine ML. 2017. Geomorphology as a first order control on the connectivity of riparian ecohydrology. Geomorphology 277: 154-170. Cavalli M, Trevisani S, Comiti F, Marchi L. 2013. Geomorphometric assessment of spatial sediment connectivity in small Alpine catchments. Geomorphology 188: 31-41. Cenderelli DA, Wohl EE. 2003. Flow hydraulics and geomorphic effects of glacial-lake outburst floods in the Mount Everest region, Nepal. Earth Surface Processes and Landforms 28: 385-407. Cerdà A, García-Fayos P. 1997. The influence of slope angle on sediment, water and seed losses on badland landscapes. Geomorphology 18: 77–90. Cheung AKL, Brierley G, O'Sullivan D. 2016. Landscape structure and dynamics on the Qinghai-Tibetan Plateau. Ecological Modelling 339: 7-22. Chorley RJ. 1962. Geomorphology and general systems theory. U.S. Geological Survey Professional Paper 500-B. Chorley RJ, Kennedy, BA. 1971. Physical Geography: a systems approach. Prentice Hall. London, UK.

Chorley RJ, Stanley AS, Sugden DE. 1985. Geomorphology. Methuen and Co. Ltd. New York,

Clubb FJ, Mudd SM, Milodowski DT, Valters DA, Slater LJ, Hurst MD, Limaye AB. 2017.
Geomorphometric delineation of floodplains and terraces from objectively defined topographic
thresholds. Earth Surface Dynamics Discussions : 1–30. DOI: 10.5194/esurf-2017-21
Coops H, Buijse LL, Buijse AD, Constantinescu A, Covaliov S, Hanganu J, Ibelings BW, Menting G,
Navodaru I, Oosterberg W, Staras M, Török L. 2008. Trophic gradients in a large-river delta:
ecological structure determined by connectivity gradients in the Danube Delta (Romania). River
Research and Applications 24: 698-709.
Cote D, Kehler DG, Bourne C, Wiersma YF. 2009. A new measure of longitudinal connectivity for
stream networks. Landscape Ecology 24: 101-113.
Coulthard TJ, Macklin MG. 2003. Modeling long-term contamination in river systems from
historical metal mining. Geology 31: 451-454.
Coulthard TJ, Ramirez J, Fowler HJ, Glenis V. 2012. Using the UKCP09 probabilistic scenarios to
model the amplified impact of climate change on drainage basin sediment yield. Hydrology and
Earth System Sciences 16: 4401-4416.
Coulthard TJ, Van De Wiel MJ. 2007. Quantifying fluvial non linearity and finding self organized
criticality? Insights from simulations of river basin evolution. Geomorphology 91: 216–235.DOI:
10.1016/j.geomorph.2007.04.011.
Coulthard TJ, Van De Wiel. 2017. Modelling long term basin scale sediment connectivity, driven

by spatial land use changes. Geomorphology 277: 265–281.

- Covino T. 2017. Hydrologic connectivity as a framework for understanding biogeochemical flux through watersheds and along fluvial networks. Geomorphology 277: 133-144.
- 1108 Croke J, Mockler S, Fogarty P, Takken I. 2005. Sediment concentration changes in runoff
 1109 pathways from a forest road network and the resultant spatial pattern of catchment
 1110 connectivity. Geomorphology 68: 257–268.
- 1111 Croke J, Fryirs K, Thompson C. 2013. Channel-floodplain connectivity during an extreme flood
 1112 event: implications for sediment erosion, deposition, and delivery. Earth Surface Processes and
 1113 Landforms 38: 1444-1456.
- Czuba JA, Foufoula-Georgiou E. 2014. A network-based framework for identifying potential synchronizations and amplifications of sediment delivery in river basins. Water Resources

 Research 50: 3826-3851.
- 1117 Czuba JA, Foufoula-Georgiou E. 2015. Dynamic connectivity in a fluvial network for identifying
 1118 hotspots of geomorphic change. Water Resources Research 51: 1401-1421.
- David CH, Maidment DR, Niu G-Y, Yang Z-L, Habets F, Eijkhout V. 2011. River network routing on the NHDPlus dataset. Journal of Hydrometeorology 12: 913-934.
- Dethier E, Magilligan FJ, Renshaw CE, Nislow KH. 2016. The role of chronic and episodic
 disturbances on channel-hillslope coupling: the persistence and legacy of extreme floods. Earth
 Surface Processes and Landforms 41: 1437-1447. Diaz-Redondo M, Egger G, Marchamalo M,
 Hohensinner S, Dister E. 2017. Benchmarking fluvial dynamics for process-based river
 restoration: the Upper Rhine River (1816-2014). River Research and Applications 33: 403-414.

1089.

Dietrich WE, T Dunne. 1978. Sediment budget for a small catchment in mountainous terrain. Zeitschrift fur Geomorphologie 29: 191-206. D'Haen K, Dusar B, Verstraeten G, Degryse P, De Brue H. 2013. A sediment fingerprinting approach to understand the geomorphic coupling in an eastern Mediterranean mountainous river catchment. Geomorphology 197: 64-75. Downs PW, Gregory KJ. 2004. River Channel Management: Towards sustainable catchment hydrosystems. Hodder Arnold, London, United Kingdom. Doyle MW, Ensign SH. 2009. Alternative reference frames in river system science. BioScience 59: 499-510. Duffy CJ. 2017. The terrestrial hydrologic cycle: an historical sense of balance. WIREs Water 4: doi: 10.1002/wat2.1216 Duvert C, Gratiot N, Anguiano-Valencia R, Némery J, Mendoza ME, Carlón-Allende T, Prat C, Esteves M. 2011. Baseflow control on sediment flux connectivity: Insights from a nested catchment study in Central Mexico. Catena 87(1): 129-140. Evrard O, Navratil O, Ayrault S, Ahmadi M, Némery J, Legout C, et al. 2011. Combining suspended sediment monitoring and fingerprinting to determine the spatial origin of fine sediment in a mountainous river catchment. Earth Surface Processes and Landforms 36: 1072-

- Falke JA, Fausch KD, Magelky R, Aldred A, Durnford DS, Riley LK, Oad R. 2011. The role of groundwater pumping and drought in shaping ecological futures for stream fishes in a dryland river basin of the western Great Plains, USA. Ecohydrology 4: 682-697.
- Farnsworth KL, Milliman JD. 2003. Effects of climate and anthropogenic change on small mountainous rivers: the Salinas River example. Global and Planetary Change 39: 53-64.
- Faulkner H. 2008. Connectivity as a crucial determinant of badland morphology and evolution.
- 1150 Geomorphology 100: 91-103.

Water Resources Research 53: 5229-5246.

- Florsheim JL, Mount JF, Constantine CR. 2006. A geomorphic monitoring and adaptive
 assessment framework to assess the effect of lowland floodplain river restoration on channel—
 floodplain sediment continuity. River Research and Applications 22: 353-375.
- Foley MM, Bellmore JR, O'Connor JE, Duda JJ, East AE, Grant GE, Anderson CW, Bountry JA,

 Collins MJ, Connolly PJ, Craig LS, Evans JE, Greene SL, Magilligan FJ, Magirl CS, Major JJ, Pess GR,

 Randle TJ, Shafroth PB, Torgersen CE, Tullos D, Wilcox AC. 2017. Dam removal: listening in.
- Fonstad MA, Dietrich JT, Courville BC, Jensen JL, Carbonneau PE. 2013. Topographic structure from motion: a new development in photogrammetric measurement. Earth Surface Processes and Landforms 38(4): 421-430.
- 1161 Fryirs KA. 2013. (Dis)connectivity in catchment sediment cascades: a fresh look at the sediment delivery problem. Earth Surf. Process. Landf. 38 (1): 30–46.

297-316.

Fryirs KA. 2017. River sensitivity: A lost foundation concept in fluvial geomorphology. Earth Surface Processes and Landforms 42: 55-70. Fryirs K, Brierley GJ. 2001. Variability in sediment delivery and storage along river courses in Bega catchment, NSW, Australia: implications for geomorphic river recovery. Geomorphology 38: 237-265. Fryirs K, Brierley G. 2009. Naturalness and place in river rehabilitation. Ecology and Society 14(1). Fryirs KA, Brierley G J. 2012. Geomorphic analysis of river systems: an approach to reading the landscape. John Wiley and Sons, Chichester, UK. Fryirs KA, Brierley G J. 2016. Assessing the geomorphic recovery potential of rivers: forecasting future trajectories of adjustment for use in management. WIREs Water 3: 727-748. Fryirs K, Brierley GJ, Erskine WD. 2012. Use of ergodic reasoning to reconstruct the historical range of variability and evolutionary trajectory of rivers. Earth Surface Processes and Landforms 37: 763-773. Fryirs KA, Brierley GJ, Preston NJ, Kasai M. 2007a. Buffers, barriers and blankets: the (dis)connectivity of catchment-scale sediment cascades. Catena 70: 49–67. Fryirs KA, Brierley GJ, Preston NJ, Spencer J. 2007b. Catchment-scale (dis) connectivity in

sediment flux in the upper Hunter catchment, New South Wales, Australia. Geomorphology 84:

Office, Washington, DC.

1182	Fryirs K, Gore D. 2013. Sediment tracing in the upper Hunter catchment using elemental and
1183	mineralogical composition: Implications for catchment-scale suspended sediment
1184	(dis)connectivity and management. Geomorphology 193: 128-142.
1185	Fryirs K, Spink A, Brierley G. 2009. Post-European settlement response gradients of river
1186	sensitivity and recovery across the upper Hunter catchment, Australia. Earth Surface Processes
1187	and Landforms 34: 897-918.
1188	Gabbud C, Lane SN. 2016. Ecosystem impacts of Alpine water intakes for hydropower: the
1189	challenge of sediment management. WIREs Water 3: 41–61.
1190	Galat DL, Kubisiak JF, Hooker JB, Sowa LM. 1997. Geomorphology, distribution and connectivity
1191	of lower Missouri River floodplain waterbodies scoured by the flood of 1993. Verhandlungen
1192	Internat. Verein. Limnologie 26: 869-878.
1193	Garcia C, Laronne JB, Sala M. 2000. Continuous monitoring of bedload flux in a mountain gravel-
1194	bed river. Geomorphology 34(1): 23-31.
1195	Gardiner, V. 1975. Drainage Basin Morphometry.
1196	Gilbert, GK. 1877. Report on the Geology of the Henry Mountains. US Government Printing
1197	Office, Washington, DC.
1198	Gilbert GK. 1880. Report on the Geology of the Henry Mountains. US Government Printing

1200	Gilvear DJ. 1999. Fluvial geomorphology and river engineering: future roles utilizing a fluvial
1201	hydrosystems framework. Geomorphology 31: 229-245.
1202	Godfrey AE, Everitt BL, Duque JFM. 2008. Episodic sediment delivery and landscape
1203	connectivity in the Mancos Shale badlands and Fremont River system, Utah, USA.
1204	Geomorphology 102: 242-251.
1205	Gomez-Velez JD, Harvey JW. 2014. A hydrogeomorphic river network model predicts where and
1206	why hyporheic exchange is important in large basins. Geophysical Research Letters 41: 64023-
1207	6412.Goodbred SL. 2003. Response of the Ganges dispersal system to climate change: a source-
1208	to-sink view since the last interstade. Sedimentary Geology 162: 83-104.
1209	Grabowski RC, Surian N, Gurnell AM. 2014. Characterizing geomorphological change to support
1210	sustainable river restoration and management. WIREs Water 1: 483-512.
1211	Gran KB, Czuba JA. 2017. Sediment pulse evolution and the role of network structure.
1212	Geomorphology 277: 17-30.
1213	Grant GE, Lewis SL. 2015. The remains of the dam: what have we learned from 15 years of US
1214	dam removals? In, Engineering geology for society and territory, v. 3. Springer, 31-35.
1215	Grant GE, JE O'Connor, MG Wolman. 2013. A river runs through it: conceptual models in fluvial
1216	geomorphology. In, E Wohl, ed, Fluvial Geomorphology, v. 9 in J Shroder, ed in chief, Treatise

on Geomorphology. Elsevier, Amsterdam, 6-21.

Catena 42: 225-250.

Grant GE, J O'Connor, E Safran. 2017. Excursions in fluvial (dis)continuity. Geomorphology 277: 145-153. Gregory KJ, Lewin J. 2014. The Basics of Geomorphology. Sage, London, UK. Grill G, Dallaire CO, Chouinard EF, et al. 2014. Development of new indicators to evaluate river fragmentation and flow regulation at large scales: A case study for the Mekong River basin. Ecological Indicators 45: 148–159. Grimmett G. 1989. Percolation. Springer-Verlag: New York. Haggett R, Chorley RJ. 1969. Network analysis in geomorphology. Edward Arnold Publishers. Hancock GRR, Coulthard TJJ, Lowry JBCBC. 2016. Predicting uncertainty in sediment transport and landscape evolution – the influence of initial surface conditions. Computers & Geosciences 90: 117-130. Harvey AM. 1987. Seasonality of processes on eroding gullies, a twelve year record of erosion rates. In Goddard A, Rapp A (eds.), Processes et Mesure de l'erosion. CNRS, Paris, 439-454. Harvey AM. 1997. Coupling between hillslope gully systems and stream channels in the Howgill Fells, northwest England: temporal implications. Geomorphologie 3: 3-19. Harvey AM. 2001. Coupling between hillslopes and channels in upland fluvial systems:

implications for landscape sensitivity, illustrated from the Howgill Fells, northwest England.

doi:10.1061/(ASCE)HY.1943-7900.0001309.

Harvey AM. 2002. Effective timescales of coupling within fluvial systems. Geomorphology 44: 175-201. Harvey AM. 2012. The coupling status of alluvial fans and debris cones: a review and synthesis. Earth Surface Processes Landforms 37: 64–76. Hauer FR, Locke H, Dreitz VJ, Hebblewhite M, Lowe WH, Muhlfeld CC, Nelson CR, Proctor MF, Rood SB. 2016. Gravel-bed river floodplains are the ecological nexus of glaciated mountain landscapes. Science Advances 2:e16000026. Haycock NE, Burt TP. 1993. Role of floodplain sediments in reducing the nitrate concentration of subsurface run-off: A case study in the Cotswolds, UK. Hydrological Processes 7: 287–295. Heckmann T, Schwanghart W. 2013. Geomorphic coupling and sediment connectivity in an alpine catchment — exploring sediment cascades using graph theory. Geomorphology 182: 89– 103. Heckmann T, Schwanghart W, Phillips JD. 2015. Graph theory — recent developments of its application in geomorphology. Geomorphology 243: 130–146. Hiatt, M. and P. Passalacqua (2015), Hydrological connectivity in river deltas: The first-order importance of channel-island exchange, Water Resources Research, 51, 2264-2282, doi:10.1002/2014WR016149. Hiatt, M. and P. Passalacqua (2017), What controls the transition from confined to unconfined flow? Analysis of hydraulics in a coastal river delta, Journal of Hydraulic Engineering, 143, 6,

Hooke JM. 2003. Coarse sediment connectivity in river channel systems: a conceptual framework and methodology. Geomorphology 56: 79–94. Hooke JM. 2006. Human impacts on fluvial systems in the Mediterranean region. Geomorphology 79: 311-335. Hooke JM. 2015. Variations in flood magnitude-effect relations and the implications for flood risk assessment and river management. Geomorphology 251: 91-107. Horton RE. 1954. Erosional development of streams and their drainage basins: Hydro-physical approach to quantitative morphology, Geological Society of America Bulletin 56 (3): 275-370. Hovius N, Meunier P, Lin CW, Chen H, Chen YG, Dadson S, Horng MJ, Lines M. 2011. Prolonged seismically induced erosion and the mass balance of a large earthquake. Earth and Planetary Science Letters 304: 347–355, doi:10.1016/j.epsl.2011.02.005. Howarth JD, Fitzsimons SJ, Norris RJ, Jacobsen GE. 2012. Lake sediments record cycles of sediment flux driven by large earthquakes on the Alpine fault, New Zealand. Geology: 40, 1091-1094, doi:10.1130 /G33486.1. Huggett R. 2007. Fundamentals of Geomorphology 2nd Ed. Routledge, New York, NY. Hupp CR, Bornette G. 2003. Vegetation as Tool in the Interpretation of fluvial geomorphic

processes and landforms in Humid Temperate Areas. In. In Kondolf, G.M. and Piegay, H. (Eds.),

Tools in Fluvial Geomorphology (270-288). England: Wiley.

Jacobson RB, Blevins DW, Bitner CJ. 2009. Sediment regime constraints on river restoration—An
example from the Lower Missouri River. Geological Society of America Special Papers 451: 1-22.
Jaeger KL, Olden JD. 2012. Electrical resistance sensor arrays as a means to quantify longitudinal
connectivity of rivers. River Research and Applications 28: 1843-1852.
Jaeger KL, Olden JD, Pelland NA. 2014. Climate change poised to threaten hydrologic
connectivity and endemic fishes in dryland streams. Proceedings National Academy of Sciences
111: 13,894-13,899.
Jain V, Tandon SK. 2010. Conceptual assessment of (dis) connectivity and its application to the
Ganga River dispersal system. Geomorphology 118: 349-358.
James LA. 2010. Secular sediment waves, channel bed waves, and legacy sediment. Geography
Compass 4: 576-598.
Jencso KG, McGlynn BL, Gooseff MN, Wondzell SM, Bencala KE, Marshall LA. 2009. Hydrologic
connectivity between landscapes and streams: transferring reach- and plot-scale understanding
to the catchment scale. Water Resources Research 45: W04428.
Jencso KG, McGlynn BL, Gooseff MN, Bencala KE, Wondzell SM. 2010. Hillslope hydrologic
connectivity controls riparian groundwater turnover: implications of catchment structure for

riparian buffering and stream water sources. Water Resources Research 46: W10524.

31.

1290	Jencso KG, McGlynn BL. 2011. Hierarchical controls on runoff generation: topographically
1291	driven hydrologic connectivity, geology, and vegetation. Water Resources Research 47:
1292	W11527.
1293	Jerolmack DJ, Paola C. 2010. Shredding of environmental signals by sediment transport,
1294	Geophys. Res. Lett., 37, L19401, doi:10.1029/2010GL044638.
1295	Jones AF, Brewer PA, Johnstone E, Macklin MG. 2007. High-resolution interpretative
1296	geomorphological mapping of river valley environments using airborne LiDAR data. Earth
1297	Surface Processes and Landforms 32 : 1574–1592.
1298	Junk W, Bayley P, Sparks R. 1989. The flood pulse concept in river-floodplain systems. In: Dodge
1299	D, Dodge DP (Eds.), Proceedings of the International Large River Symposium. Canadian Special
1300	Publication of Fisheries and Aquatic Sciences 106, 110–127.
1301	Kampf SK, Mirus BB. 2013. Subsurface and surface flow leading to channel initiation. In, Wohl E,
1302	ed, Treatise on fluvial geomorphology. Elsevier, Amsterdam, 23-42.
1303	Kao SJ, Milliman JD. 2008. Water and sediment discharge from small mountainous rivers,
1304	Taiwan: the roles of lithology, episodic events, and human activities. Journal of Geology 116:
1305	431-448.
1306	Karterakis SM, BW Karney, B Singh, A Guergachi. 2007. The hydrologic cycle: a complex history

with continuing pedagogical implications. Water Science and Technology: Water Supply 7: 23-

1309	Kimberly AW, Gardner RH, Turner MG. 1997. Landscape connectivity and population
1310	distributions in heterogeneous environments. Oikos 78: 151 - 169.
1311	Koiter AJ, Lobb DA, Owens PN, Petticrew EL, Tiessen KH, Li S. 2013. Investigating the role of
1312	connectivity and scale in assessing the sources of sediment in an agricultural watershed in the
1313	Canadian prairies using sediment source fingerprinting. Journal of Soils and Sediments 13(10):
1314	1676-1691.
1315	Kondolf GM. 1998. Lessons learned from river restoration projects in California. Aquatic
1316	Conservation: Marine and Freshwater Ecosystems 8: 39-52.
1317	Kondolf GM, Boulton AJ, O'Daniel S, Poole GC, Rahel FJ, Stanley EH, Wohl E, et al. 2006.
1318	Process-based ecological river restoration: visualizing three-dimensional connectivity and
1319	dynamic vectors to recover lost linkages. Ecology and Society 11: 5.
1320	http://www.ecologyandsociety.org/vol11/iss2/art5.
1321	Kondolf GM, Larson M. 1995. Historical channel analysis and its application to riparian and
1322	aquatic habitat restoration. Aquatic Conservation: Marine and Freshwater Ecosystems 5: 109-
1323	126.
1324	Kornis MS, Weidel BC, Powers SM, Diebel MW, Cline TJ, Fox JM, Kitchell JF. 2014. Fish
1325	community dynamics following dam removal in a fragmented agricultural stream. Aquatic
1326	Sciences 77: 465-480.
1327	Korup, O, McSaveney MJ, Davies TRH. 2004. Sediment generation and delivery from large
1000	

historic landslides in the Southern Alps, New Zealand. Geomorphology 61: 189–207.

W08423.

Kramer N, Wohl E. 2017. Rules of the road: a qualitative and quantitative synthesis of large
 wood transport through drainage networks. Geomorphology 279: 74-97.
 Kuo CW, Brierley GJ. 2013. The influence of landscape configuration upon patterns of sediment

Kuo CW, Brierley G. 2014. The influence of landscape connectivity and landslide dynamics upon channel adjustments and sediment flux in the Liwu Basin, Taiwan. Earth Surface Processes and Landforms 39: 2038-2055.

Kupfer JA, Meitzen KM, Gao P. 2014. Flooding and surface connectivity of Taxodium-Nyssa

storage in a highly connected river system. Geomorphology 180: 255-266.

stands in a southern floodplain forest ecosystem. River Research and Applications. DOI:

1338 10.1002/rra.2828

Lane SN, Richards KS, Chandler JH. 1994. Developments in monitoring and modelling small-scale river bed topography. Earth Surf. Process. Landforms, 19: 349–368.

Lane SN, Milledge DG. 2013. Impacts of upland open drains upon runoff generation: a numerical assessment of catchment-scale impacts. Hydrological Processes, 27: 1701–1726

Lane SN, Reaney SM, Heathwaite AL. 2009. Representation of landscape hydrological

1344 connectivity using a topographically driven surface flow index. Water Resources Research, 45:

Lane SN, Reid SC, Tayefi V, Yu D, Hardy RJ. 2008. Reconceptualising coarse sediment delivery

problems in rivers as catchment-scale and diffuse. Geomorphology 98: 227-249.

Lane SN, Bakker M, Gabbud C, Micheletti N, Saugy JN. 2017. Sediment export, transient
landscape response and catchment-scale connectivity following rapid climate warming and
Alpine glacier recession. Geomorphology 277: 210-227.
Larsen LG, Choi J, Nungesser MK, Harvey JW. 2012. Directional connectivity in hydrology and
ecology. Ecological Applications 22: 2204-2220.
Larsen LG, Harvey JW, Maglio MM. 2015. Mechanisms of nutrient retention and its relation to
flow connectivity in river-floodplain corridors. Freshwater Science 34: 187-205.
Larsen LG, Ma J, Kaplan D. 2017. How important is connectivity for surface water fluxes? A
generalized expression for flow through heterogeneous landscapes. Geophysical Research
Letters 44:
Lesschen JP, Schoorl JM, Cammeraat LH. 2009. Modelling runoff and erosion for a semi-arid
catchment using a multi-scale approach based on hydrologic connectivity. Geomorphology 109:
174-183.
Lexartza-Artza I, Wainwright J. 2009. Hydrological connectivity: linking concepts with practical
implications. Catena 79 (2): 146–152.
Lexartza-Artza I, Wainwright J. 2011. Making connections: changing sediment sources and sinks
Lexartza-Artza I, Wainwright J. 2011. Making connections: changing sediment sources and sinks in an upland catchment. Earth Surf. Process. Landf. 36: 1090–1104.

of earthquake-triggered landslides with the fluvial network: implications for landslide sediment

- transport after the 2008 Wenchuan earthquake. Journal of Geophysical Research: Earth Surface 121: 703-724. Liang, M., W. Kim, P. Passalacqua (2016b), How much subsidence is enough to change the morphology of river deltas?, Geophysical Research Letters, 43, 19, 10266-10276, doi:10.1002/2016GL070519. Liang, M., C. Van Dyk, P. Passalacqua (2016a), Quantifying the Patterns and Dynamics of River Deltas Under Conditions of Steady Forcing and Relative Sea-Level Rise, Journal of Geophysical Research Earth Surface, 121, 2, 465-496, doi:10.1002/2015JF003653. Lininger KB, Latrubesse EM. 2016. Flooding hydrology and peak discharge attenuation along the middle Araguaia River in central Brazil. Catena 143: 90-101. Lisenby P, Fryirs K. 2017a. Sedimentologically significant tributaries: Catchment-scale controls
- on sediment (dis)connectivity in the Lockyer Valley, SEQ, Australia. Earth Surface Processes and Landforms 42: 1493-1504.

 Lisenby P, Fryirs K. 2017b. 'Out with the Old?' Why coarse spatial datasets are still useful for
- catchment-scale investigations of sediment (dis)connectivity. Earth Surface Processes and
 Landforms. DOI: 10.1002/esp.4131
 - Lisle TE. 1982. Effects of aggradation and degradation on riffle-pool morphology in natural gravel channels, northwestern California. Water Resources Research 18: 1643-1651.

331-345

Macklin MG, Lewin J. 1989. Sediment transfer and transformation of an alluvial valley floor: the
River South Tyne, Northumbria, U.K. Earth Surface Processes and Landforms 14: 233-246.
Magilligan FJ, Graber BE, Nislow KH, Chipman JW, Sneddon CS, Fox CA. 2016a. River restoration
by dam removal: enhancing connectivity at watershed scales. Elementa: Science of the
Anthropocene 4: 000108. doi:10.12952/journal.elementa.000108.
Magilligan FJ, Nislow KH, Kynard BE, Hackman AM. 2016b. Immediate changes in stream
channel geomorphology, aquatic habitat, and fish assemblages following dam removal in a
small upland catchment. Geomorphology 252: 158-170.
Major JJ, East AE, O'Connor JE, Grant GE, Wilcox AC, Magirl CS, Collins MJ, Tullos DD. 2017.
Geomorphic responses to dam removal in the United States – a two-decade perspective. In,
Tsutsumi D, Laronne JB, eds, Gravel-bed rivers. John Wiley and Sons, Chichester, UK, 255-383.
Marra WA, Kleinhans MG, Addink EA. 2013. Network concepts to describe channel importance
and change in multichannel systems: test results for the Jamuna River, Bangladesh. Earth
Surface Processes and Landforms 39: 766-778.
Marsh GP. 1965 (1864). Man and Nature: or Physical Geography as Modified by Human Action.
Martignier L, Adatte T, and Verrecchia EP. 2013. Bedrock versus superficial deposits in the Swiss
Jura Mountains: what is the legitimate soil parent material?. Earth Surf. Process. Landforms, 38:

doi:10.1029/2010WR009341.

Masselink RJH, Keesstra SD, Temme AJAM, Seeger M, Giménez R, Casalí J. 2016a. Modelling Discharge and Sediment Yield at Catchment Scale Using Connectivity Components. Land Degradation & Development 27: 933-945 Masselink RJH, Heckmann T, Temme AJAM, Anders NS, Gooren HPA, Keesstra SD. 2016b. A network theory approach for a better understanding of overland flow connectivity. Hydrological Processes 31: 207-220. Matella MK, AM Merenlender (2015) Scenarios for restoring floodplain ecology given changes to river flows under climate change: Case from the San Joaquin River, California. River Res Appl 31: 280-290. May C, Roering J, Snow K, Griswold K, Gresswell R. 2017. The waterfall paradox: how knickpoints disconnect hillslope and channel processes, isolating salmonid populations in ideal habitats. Geomorphology 277: 228-236. McClain WE, Boyer EW, Dent CL, Gergel SE, Grimm NB, Groffman PM, Hart SC, Harvey JW, Johnston CA, Mayorga E, McDowell WH, Pinay G. 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic ecosystems. Ecosystems 6: 301-312. McDonnell JJ. 2003. Where does water go when it rains? Moving beyond the variable source area concept of rainfall-runoff response. Hydrological Processes 17: 1869-1875. McGuire KJ, McDonnell JJ. 2010. Hydrological connectivity of hillslopes and streams: characteristic time scales and nonlinearities. Water Resources Research 46: W10543,

Water Resources Association 35, 397-410.

Meade RH. 2007. Transcontinental moving and storage: the Orinoco and Amazon Rivers
transfer the Andes to the Atlantic. In, A Gupta, ed, Large Rivers: Geomorphology and
Management. John Wiley and Sons, Chichester, UK, 45-63.
Meade RH, Dunne T, Richey JE, Santos UM, Salati E. 1985. Storage and remobilization of
suspended sediment in the lower Amazon River of Brazil. Science 228: 488-490.
Meitzen, KM and Kupfer JA. 2015. Abandoned meander forest development in a large alluvial
southeastern floodplain, South Carolina, USA. River Systems 21(2-3): 141-162.
Merriam G. 1984. Connectivity: a fundamental ecological characteristic of landscape pattern.
Proceedings of the International Association for Landscape Ecology 1: 5-15.
Micheletti N, Lane SN. 2016. Water yield and sediment export in small, partially glaciated Alpine
watersheds in a warming climate. Water Resour. Research 52: 4924–4943.
Micheletti N, Lambiel C, Lane SN. 2015. Investigating decadal-scale geomorphic dynamics in an
alpine mountain setting. Journal of Geophysical Research: Earth Surface 120: 2155-2175.
Mika S, Hoyle J, Kyle G, Howell T, Wolfenden B, Ryder D, et al. 2010. Inside the "black box" of
river restoration: using catchment history to identify disturbance and response mechanisms to
set targets for process-based restoration. Ecology and Society 15(4).
Montgomery, DR. 1999. Process domains and the river continuum. Journal of the American

Mueller EN, Wainwright J, Parsons AJ. 2007. Impact of connectivity on the modeling of overland flow within semiarid shrubland environments. Water Resources Research 43: DOI:10.1029/2006WR005006 Nakamura F, Swanson FJ, Wondzell SM. 2000. Disturbance regimes of stream and riparian ecosystems – a disturbance-cascade perspective. Hydrological Processes 14: 2849-2860. Newman MEJ. 2006. Finding community structure in networks using the eigenvectors of matrices. Physical Review E 74: 036104. Newman MEJ. 2010. Networks: An Introduction. Oxford University Press Inc., NewYork, NY. Newman MEJ, Barabasi AL, Watts DJ. 2006. The Structure and Dynamics of Networks. Princeton University Press, Princeton, New Jersey. Newson MD. 1997. Land, water, and development: Sustainable management of river basin systems 2nd Ed. Routledge, New York, NY. Nicoll T, Brierley G. 2017. Within-catchment variability in landscape connectivity measures in the Garang catchment, upper Yellow River. Geomorphology 277: 197-209. Nilsson C, Reidy CA, Dynesius M, Revenga C. 2005. Fragmentation and flow regulation of the world's largest river systems. Science 308: 405-408. Opperman JJ, Luster R, McKenney BA, Roberts M, Meadows AW. 2010. Ecologically functional

floodplains: Connectivity, flow regime, and scale. J Am Water Res Assoc 46: 211-226.

Orlando J, Comas X, Hynek SA, Buss HL, Brantley SL. 2016. Architecture of the deep critical zone in the Rio Icacos watershed (Luquillo Critical Zone Observatory, Puerto Rico) inferred from drilling and ground penetrating radar (GPR). Earth Surface Processes and Landforms 41: 1826-1840. Osterkamp WR, Hupp CR. 2010. Fluvial Processes and vegetation – Glimpses of the past, the present, and perhaps the future. Geomorphology. 116:274-285. Passalacqua P Belmont P, Staley DM, Simley JD, Arrowsmith JR, Bode CA, Crosby C, DeLong SB, Glenn NF, Kelly SA, Lague D, Sangireddy H, Schaffrath K, Tarboton DG, Wasklewicz T, Wheaton JM. 2015. Analyzing high resolution topography for advancing the understanding of mass and energy transfer through landscapes: A review. Earth-Science Reviews 148: 174-193, doi:10.1016/j.earscirev.2015.05.012. Passalacqua P. 2017. The Delta Connectome: A network-based framework for studying connectivity in river deltas. Geomorphology 277: 50-62. Pavelsky TM, Smith LC. 2008. Remote sensing of hydrologic recharge in the Peace-Athabasca Delta, Canada. Geophysical Research Letters 35: L08403, doi:10.1029/2008GL033268. Pess GR, Quinn TP, Gephard SR, Saunders R. 2014. Re-colonization of Atlantic and Pacific rivers by anadromous fishes: linkages between life history and the benefits of barrier removal. Reviews in Fish Biology and Fisheries 24: 881-900.

Phillips JD. 2007. The perfect landscape. Geomorphology 84: 159-169.

Phillips JD. 2012. Synchronization and scale in geomorphic systems. Geomorphology 137: 150-158. Phillips JD. 2013. Hydrological connectivity of abandoned channel water bodies on a coastal plain river. River Research and Applications 29: 149-160. Phillips JD, Van Dyke C. 2016. Principles of geomorphic disturbance and recovery in response to storms. Earth Surface Processes and Landforms 41: 971-979. Pisut P. 2002. Channel evolution of the pre-channelized Danube River in Bratislava, Slovakia (1712-1886). Earth Surface Processes and Landforms 27: 369-390. Playfair J. 1802. Illustrations of the Huttonian Theory of the Earth. Cadell and Davies and William Creech, London and Edinburgh. Poeppl RE, Keiler M, Von Elverfeldt K, Zweimueller I, Glade T. 2012. The influence of riparian vegetation cover on diffuse lateral sediment connectivity and biogeomorphic processes in a medium-sized agricultural catchment, Austria. Geografiska Annaler: Series A, Physical Geography 94: 511–529. Poeppl RE, Keesstra SD, Maroulis J. 2017. A conceptual connectivity framework for understanding geomorphic change in human-impacted fluvial systems. Geomorphology 277: 238-251. Poff NL, Allan JD, Bain MB, Karr JR, Prestegaard KL, Richter BD, Sparks RE, Stromberg JC. 1997.

The natural flow regime. BioScience 47: 769-784.

1497	Poole GC, Stanford JA, Frissell CA, Running SW. 2002. Three-dimensional mapping of	
1498	geomorphic controls on flood-plain hydrology and connectivity from aerial photos.	
1499	Geomorphology 48(4): 329-347.	
1500	Prasicek G, Larsen IJ, Montgomery DR. 2015. Tectonic control on the persistence of glacially	
1501	sculpted topography. Nature Communications 6: 8028, DOI: 10.1038/ncomms9028	
1502	Pringle CM. 2001. Hydrologic connectivity and the management of biological reserves: a global	
1503	perspective. Ecol. Appl. 11: 981–998.	
1504	Pringle CM. 2003. The need for a more predictive understanding of hydrologic connectivity.	
1505	Aquatic Conservation: Marine and Freshwater Ecosystems 13: 467-471.	
1506	Puttock A, Macleod CJA, Bol R, Sessford P, Dugait J, Brazier RE. 2013. Changes in ecosystem	
1507	structure, function and hydrological connectivity control water, soil and carbon losses in semi-	
1508	arid grass to woody vegetation transition. Earth Surface Processes and Landforms 38: 1602–	
1509	1611.	
1510	Rathburn SL, Rubin ZK, Wohl EE. 2013. Evaluating channel response to an extreme	
1511	sedimentation event in the context of historical range of variability: Upper Colorado River, USA.	
1512	Earth Surface Processes and Landforms 38: 391-406.	
1513	Rathburn SL, Shahverdian SM, Ryan SE. 2017. Post-disturbance sediment recovery and	

geomorphic resilience. Geomorphology.

1515	Reid HE, Brierley GJ. 2015. Assessing geomorphic sensitivity in relation to river capacity for
1516	adjustment. Geomorphology 251: 108-121.
1517	Reid LM, T Dunne. 2003. Sediment budgets as an organizing framework in fluvial
1518	geomorphology, in GM Kondolf, H Piegay, eds, Tools in Fluvial Geomorphology, John Wiley and
1519	Sons, Chichester, UK, 463-500.
1520	Reid LM and Dunne T. 2016. Sediment budgets as an organizing framework in fluvial
1521	geomorphology. In Tools in Fluvial Geomorphology. Eds. Kondolf GM and Piegay H. pp 357-375
1522	Wiley Blackwell.
1523	Reid SC, Lane SN, Montgomery DR, Brookes CJ. 2007. Does hydrological connectivity improve
1524	modelling of coarse sediment delivery in upland environments? Geomorphology 90: 263-282.
1525	Rice SP. 2017. Tributary connectivity, confluence aggradation, and network biodiversity.
1526	Geomorphology 277: 6-16.
1527	Romans BW, Castelltort S, Covault JA, Fildani A, Walsh JP. 2016. Environmental signal
1528	propagation in sedimentary systems across timescales. Earth Science Reviews 153: 7–29.
1529	Rood SB, Samuelson GM, Braatne JH, Gourley CR, Hughes FMR, Mahoney JM. 2005. Managing
1530	river flows to restore floodplain forests. Frontiers in Ecology and the Environment 3: 193-201.
1531	Ruddell BL, Kumar P. 2009. Ecohydrologic process networks: 2. Analysis and characterization.

Water Resources Research 45: W03420. doi:10.1029/2008WR007280.

1533	Schmidt JC, Parnell RA, Grams PE, Hazel JE, Kaplinski MA, Stevens LE, Hoffnagle TL. 2001. The
1534	1996 controlled flood in Grand Canyon: flow, sediment transport, and geomorphic change.
1535	Ecological Applications 11: 657-671.
1536	Schmidt JC, Webb RH, Valdez RA, Marzolf GR, Stevens LE. 1998. Science and values in river
1537	restoration in the Grand Canyon. BioScience 48: 735-747.
1538	Schmitt RJ, Bizzi S, Castelletti A. 2016. Tracking multiple sediment cascades at the river network
1539	scale identifies controls and emerging patterns of sediment connectivity. Water Resources
1540	Research 52: 3941-3965.
1541	Schreiber T. 2000. Measuring information transfer. Physics Review Letters 85: 464-464.
1542	Schumm SA. 1973. Geomorphic thresholds and complex response of drainage systems. In,
1543	Fluvial Geomorphology, M Morisawa ed, State University of New York, Binghamton, NY, 299-
1544	310.
1545	Schumm S a. (Colorado S. 1973. Geomorphic Thresholds and Complex Response of Drainage
1546	Systems. Fluvial Geomorphology DOI: Doi 10.2307/622211
1547	Schumm, SA. 1977a. Drainage Basin Morphology. Benchmark Papers in Geology, Volume 41.
1548	Dowden, Hutchinson, and Ross. Stroudsburg, Pennsylvania.
1549	Schumm SA. 1977b. The Fluvial System. Wiley-Interscience, NY.
1550	Schumm SA. 1981. Evolution and response of the fluvial system, sedimentologic implications.

Society of Economic Paleontologists and Mineralogists Special Publication No. 31, 19-29.

Academy of Sciences 110: 18436-18441.

1552	Schumm SA, Lichty RW. 1965. Time, space, and causality in geomorphology. American Journal	
1553	of Science 263: 110-119.	
1554	Scorpio V, Aucelli PP, Giano SI, Pisano L, Robustelli G, Rosskopf CM, Schiattarella M. 2015. River	
1555	channel adjustments in Southern Italy over the past 150years and implications for channel	
1556	recovery. Geomorphology 251: 77-90.	
1557	Sear DA, Newson MD, Brookes A. 1995. Sediment-related river maintenance: The role of fluvial	
1558	geomorphology. Earth Surface Processes and Landforms 20: 629-647.	
1559	Segurado P, Branco P, Avelar AP, Ferreira MT. 2015. Historical changes in the functional	
1560	connectivity of rivers based on spatial network analysis and the past occurrences of diadromous	
1561	species in Portugal. Aquatic Sciences 77: 427-440.	
1562	Sendrowski, A. and P. Passalacqua (2017), Process connectivity in a naturally prograding river	
1563	delta, Water Resources Research, 53, doi:10.1002/2016WR019768.	
1564	Simons DB, Richardson EV. 1966. Resistance to flow in alluvial channels. U.S. Geological Survey	
1565	Professional Paper 422J.	
1566	Singer MB, Aalto R, James LA, Kilham NE, Higson JL, Ghoshal S. 2013. Enduring legacy of a toxic	
1567	fan via episodic redistribution of California gold mining debris. Proceedings of the National	

1569	Smith MW, Vericat D. 2015. From experimental plots to experimental landscapes: topography,
1570	erosion and deposition in sub-humid badlands from Structure-from-Motion photogrammetry.
1571	Earth Surface Processes and Landforms 40(12): 1656-1671.
1572	Smith S, Belmont P, Wilcock PR. 2011. Closing the gap between watershed modeling, sediment
1573	budgeting, and stream restoration. Stream Restoration in Dynamic Fluvial Systems, 293-
1574	317. Sokal MA, Hall RI, Wolfe BB. 2008. Relationships between hydrological and limnological
1575	conditions in lakes of the Slave River Delta (NWT, Canada) and quantification of their roles on
1576	sedimentary diatom assemblages. Journal of Paleolimnology 39: 533-550.
1577	Souza JOP, Correa ACB, Brierley GJ. 2016. An approach to assess the impact of landscape
1578	connectivity and effective catchment area upon bedload sediment flux in Saco Creek
1579	Watershed, semiarid Brazil. Catena 138: 13-29.
1580	Standish RJ, Hobbs RJ, Mayfield MM, Bestelmeyer BT, Suding KN, Battaglia LL, et al. 2014.
1581	Resilience in ecology: Abstraction, distraction, or where the action is?. Biological Conservation
1582	177: 43-51.
1583	Strahler, AN. 1952. Hypsometric (area-altitude) analysis of erosional topology. Geological
1584	Society of America Bulletin 63 (11) 117-1142.
1585	Strahler, AN. 1954. Dimensional analysis in quantitative terrain descriptions. Annals of the

Association of American Geographers 44, 282.

Stanford JA and Ward JV. 1992. Management of aquatic resources in large catchments: recognizing interactions between ecosystem connectivity and environmental disturbance. In Watershed Management, Ed. R. J. Naiman. Pp.91-124. Springer-Verlag. New York. Sternai P, Herman F, Fox MR, Catelltort S. 2011. Hypsometric analysis to identify spatially variable glacial erosion. Journal of Geophysical Research: Earth Surface 116: F03001. doi:10.1029/2010JF001823. Surian N, Rinaldi M, Pellegrini L, Audisio C, Maraga F, Teruggi L, et al. 2009a. Channel adjustments in northern and central Italy over the last 200 years. Geological Society of America Special Papers 451: 83-95. Surian N, Ziliani L, Comiti F, Lenzi MA, Mao L. 2009b. Channel adjustments and alteration of sediment fluxes in gravel-bed rivers of North-Eastern Italy: potentials and limitations for channel recovery. River Research and Applications 25: 551-567. Tank SE, Lesack LFW, Hesslein RH. 2009. Northern delta lakes as summertime CO₂ absorbers within the Arctic landscape. Ecosystems 12: 144-157. Taylor PD, Fahrig L, Henein K, Merriam G. 1993. Connectivity is a vital element of landscape structure. Oikos 68(3): 571-573. Taylor S, Feng X, Kirchner JW, Osterhuber R, Klaue B, Renshaw CE. 2001. Isotope evolution of a

seasonal snowpack and its melt. Water Resources Research 37: 759-769.

1605	Tejedor A, Longjas A, Zaliapin I, Foufoula-Georgiou E. 2015. Delta channel networks: 1. A graph-
1606	theoretic approach for studying connectivity and steady state transport on deltaic surfaces.
1607	Water Resources Research 51: 3998-4018.
1608	Tetzlaff D, Soulsby C, Bacon PJ, Youngson AF, Gibbins C, Malcolm IA. 2007. Connectivity
1609	between landscapes and riverscapes – a unifying theme in integrating hydrology and ecology in
1610	catchment science? Hydrological Processes 21: 1385-1389.
1611	Thompson C, Fryirs K, Croke J. 2016. The disconnected sediment conveyor belt: Patterns of
1612	longitudinal and lateral erosion and deposition during a catastrophic flood in the Lockyer Valley,
1613	southeast Queensland, Australia. River Research and Applications 32: 540-551.
1614	Tischendorf L, Fahrig L. 2000. How should we measure landscape connectivity? Landscape
1615	Ecology 15: 633 – 641.
1616	Topping DJ, Rubin DM, Schmidt JC. 2005. Regulation of sand transport in the Colorado River by
1617	changes in the surface grain size of eddy sandbars over multi-year timescales. Sedimentology
1618	52: 1133-1153.
1619	Tranter M, Brimblecombe P, Davies TD, Vincent CE, Abrahams PW, Blackwood I. 1986. The
1620	composition of snowfall, snowpack and meltwater in the Scottish Highlands – evidence for
1621	preferential elution. Atmosphere and Environment 20: 517-525.

Trimble SW. 1983. A sediment budget for Coon Creek in the Driftless Area, Wisconsin, 1853-

1977. American Journal of Science 283: 454-474.

Trimble SW. 2013. Historical agriculture and soil erosion in the Upper Mississippi Valley Hill Country. CRC Press, Boca Raton, FL. Toone J, Rice SP, Piégay H. 2014. Spatial discontinuity and temporal evolution of channel morphology along a mixed bedrock-alluvial river, upper Drôme River, southeast France: Contingent responses to external and internal controls. Geomorphology 205: 5-16. Tuan YF. 1968. The Hydrologic Cycle and the Wisdom of God: A Theme in Geoteleology. University of Toronto Press. Turnbull L, Wainwright J, Brazier RE. 2008. A conceptual framework for understanding semi-arid land degradation: ecohydrological interactions across multiple-space and time scales. Ecohydrology 1: 23 – 34. Turnbull L, Wainwright J, Brazier RE. 2010. Changes in hydrology and erosion over a transition from grassland to shrubland. Hydrological Processes 24(4): 393-414. Turner MG. 1989. Landscape ecology: the effect of pattern on process. Annual Review of Ecology and Systematics 20: 171–197. Trimble, SW 1977. The fallacy of stream equilibrium in contemporary denudation studies. American Journal of Science. 277: 876-887. Trimble, SW 1983. A sediment budget for Coon Creek basin in the driftless area, Wisconsin,

1853-1977. American Journal of Science 283: 454-474.

1642	Vanacker V, Molina A, Govers G, Poesen J, Dercon G, Deckers S. 2005. River channel response
1643	to short-term human-induced change in landscape connectivity in Andean ecosystems.
1644	Geomorphology 72: 340-353.
1645	Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE. 1980. River continuum
1646	concept. Canadian Journal of Fisheries and Aquatic Sciences 37: 130–137.
1647	Wainwright J, Parsons AJ, Abrahams AD. 2000. Plot-scale studies of vegetation, overland flow
1648	and erosion interactions: case studies from Arizona and New Mexico. Hydrological Processes
1649	14: 2921– 2943.
1650	Wainwright J, Turnbull L, Ibrahim TG, Lexartza-Artza I, Thornton SF, Brazier RE. 2011. Linking
1651	environmental regimes, space and time: Interpretations of structural and functional
1652	connectivity. Geomorphology 126: 387 – 404.
1653	Walling DE. 1983. The sediment delivery problem. Journal of Hydrology 65: 209-237.
1654	Walling DE, Owens PN, Leeks GJL. 1999. Fingerprinting suspended sediment sources in the
1655	catchment of the River Ouse, Yorkshire, UK. Hydrological Processes 13: 955-975.
1656	Wang J, Jin Z, Hilton RG, Zhang F, Densmore AL, Li G, West AJ. 2015. Controls on fluvial
1657	evacuation of sediment from earthquake-triggered landslides. Geology 43: 115-118.
1658	Ward JV. 1997. An expansive perspective of riverine landscapes: pattern and process across

scales. River Ecosystems 6: 52-60.

- Ward JV, Stanford JA. 1983. The serial discontinuity concept of lotic ecosystems. In Fontaine TD, Bartell SM, eds, Dynamics of lotic ecosystems. Ann Arbor Science, Ann Arbor, MI, 29-42. Ward JV, Stanford JA. 1989. The four-dimensional nature of lotic ecosystems. Journal of the North American Benthological Society 8: 2-8. Ward JV, Tockner K, Schiemer F. 1999. Biodiversity of floodplain river ecosystems: ecotones and connectivity. Regulated Rivers: Research & Management 15: 125-139. Ward JV, Tockner K, Uehlinger U, Malard F. 2001. Understanding natural patterns and processes in river corridors as the basis for effective river restoration. River Research and Applications 17: 311-323. Warrick JA, Mertes LAK. 2009. Sediment yield from tectonically active semiarid Western Transverse Ranges of California. Geological Society of America Bulletin 121: 1054-1070. Warrick JA, Bountry JA, East AE, Magirl CS, Randle TJ, Gelfenbaum G, Ritchie AC, Pess GR, Leung V, Duda JJ. 2015. Large-scale dam removal on the Elwha River, Washington, USA: source-to-sink sediment budget and synthesis. Geomorphology 246: 729-750. Wegener P, Covino T, Wohl E. 2017. Beaver-mediated lateral hydrologic connectivity, fluvial carbon and nutrient flux, and aquatic ecosystem metabolism. Water Resources Research 53: 4606-4623.
- 1677 Werner BT, McNamara DE. 2007. Dynamics of coupled human-landscape systems.
- 1678 Geomorphology 91: 393-407.

Resources Research 51: 5974-597.

1679	Western AW, Blöschl G, Grayson RB. 2001. Toward capturing hydrologically significant	
1680	connectivity in spatial patterns. Water Resources Research 37: 83-97.	
1681	Wester T, Wasklewicz T, Staley D. 2014. Functional and structural connectivity within a recently	
1682	burned drainage basin. Geomorphology 206: 362-373.	
1683	Wheaton JM, Brasington J, Darby SE, Sear DA. 2010. Accounting for uncertainty in DEMs from	
1684	repeat topographic surveys: improved sediment budgets. Earth Surface Processes and	
1685	Landforms 35: 136-156.	
1686	With KA, Gardner RH, Turner MG. 1997. Landscape connectivity and population distributions in	
1687	heterogeneous environments. Oikos 78: 151-169.	
1688	Wohl E. 2011. What should these rivers look like? Historical range of variability and human	
1689	impacts in the Colorado Front Range, USA. Earth Surface Processes and Landforms 36: 1378-	
1690	1390.	
1691	Wohl E. 2015. Legacy effects on sediments in river corridors. Earth-Science Reviews 147: 30-53.	
1692	Wohl E, Bledsoe BP, Jacobson RB, Poff NL, Rathburn SL, Walters DM, Wilcox AC. 2015a. The	
1693	natural sediment regime in rivers: broadening the foundation for ecosystem management.	
1694	BioScience 65: 358-371.	
1695	Wohl E. 2017. Connectivity in rivers. Progress in Physical Geography 41: 345-362.	

Wohl E, Lane SN, Wilcox AC. 2015b. The science and practice of river restoration. Water

- Wohl, E, Magilligan FJ, Rathburn SL. 2017a. Introduction to the special issue: Connectivity in Geomorphology. Geomorphology 277: 1-5.
- Wohl E, Rathburn S, Chignell S, Garrett K, Laurel D, Livers B, Patton A, Records R, Richards M, Schook DM, Sutfin NA, Wegener P. 2017b. Mapping longitudinal stream connectivity in the North St. Vrain Creek watershed of Colorado. Geomorphology 277: 172–182.
- Yue TX, Xu B, Liu JY. 2004. A patch connectivity index and its change in relation to new wetland at the Yellow River Delta. International Journal of Remote Sensing 25: 4617-4628.
- Zaliapin, I., Foufoula-Georgiou, E., Ghil, M., 2010. Transport on river networks: a dynamic tree
 approach, J. Geophys. Res. Earth Surf., 115 (F2), F00A15,
- 1707 http://dx.doi.org/10.1029/2009JF001281.
- Zarnetske JP, Haggerty R, Wondzell SM, Baker MA. 2011. Dynamics of nitrate production and
 removal as a function of residence time in the hyporheic zone. Journal of Geophysical Research
 Biogeosciences 116: G01025. Doi:10.1029/2010JG001356.
- Zeug SC, Winemiller KO, Tarim S (2005). Response of Brazos River oxbow fish assemblages to
 patterns of hydrologic connectivity and environmental variability. Trans. American Fishery
 Society 134: 1389-1399.
- Ziliani L, Surian N. 2012. Evolutionary trajectory of channel morphology and controlling factors
 in a large gravel-bed river. Geomorphology 173: 104-117.

- 1716 Ziliani L, Surian N. 2016. Reconstructing temporal changes and prediction of channel evolution
- in a large Alpine river: the Tagliamento River, Italy. Aquatic Sciences 78: 83-94.



Table 1. Definitions and quantitative metrics of connectivity (After Wohl, 2017, Tables 1 and 2) A)

A)	
Definition	Reference
Connectivity in the context of landscape dynamics describes the	(Harvey, 1987, 1997, 2001,
transmission of matter and energy among system components	2002; Godfrey et al., 2008)
Hydrological connectivity as the exchange of matter, energy, and biota between different elements of the riverine landscape via the aqueous medium	Amoros and Roux, 1988
Hydrological connectivity can be defined as the physical linkage of water and sediment through the fluvial system.	Hooke, 2003; Lesschen et al., 2009
Hydrologic connectivity refers to the water-mediated transfer of matter, energy, and/or organisms within or between elements of the hydrologic cycle	Pringle, 2003
River hydrologic connectivity refers to the water-mediated fluxes of material, energy, and organisms within and among components, e.g., the channel, floodplain, alluvial aquifer, etc., of the ecosystem	Kondolf et al., 2006
Static/structural connectivity: static elements of hydrological connectivity are spatial patterns, such as hydrological runoff units, that can be categorized, classified, and estimated; spatial patterns in the landscape (Turnbull et al., 2008)	Bracken and Croke, 2007
Dynamic/functional connectivity: describes both the longer term landscape developments, such as changes following abandonment of agriculture, and short-term variation in antecedent conditions and rainfall inputs to systems that result in nonlinearities in hillslope and catchment response to rainfall; how spatial patterns interact with catchment processes to produce water transfer in catchments (Turnbull et al., 2008)	Bracken and Croke, 2007
Process connectivity: the evolutionary dynamics of how systems operate; also defined as flow of information among a system's drivers, where information is a reduction of the uncertainty in a variable's state	Bracken and Croke, 2007; Passalacqua, 2017; Ruddell and Kumar, 2009
Three stages of landscape connectivity: coupled linkage when there is free transmission between landscape units; partial coupling when a discontinuity between units results in pulses of sediment movement; partly connected stage when there is a decrease of transmission due to impediments, but some material can pass the impediment during an effective event; buffers hinder lateral connectivity, barriers hinder longitudinal connectivity, and blankets hinder vertical connectivity	Fryirs et al., 2007; Jain and Tandon, 2010
Initiation of a shallow groundwater table across hillslope, riparian, and stream zones	Jencso and McGlynn, 2011
Hydrologic connectivity describes connection, via the subsurface flow system, between the riparian zone and the upland zone, which occurs when the water table at the upland-riparian zone interface is above the confining layer (Also presents 10 other definitions from the literature, categorized with respect to water cycle or landscape features at the watershed scale, and landscape features, spatial patterns, and flow processes at the hillslope scale)	Bracken et al., 2013

Sediment connectivity: the degree of linkage that controls sediment Cavalli et al., 2013 fluxes throughout landscapes and in particular between sediment sources and downstream areas Sediment connectivity is the water-mediated transfer of sediment Fryirs, 2013 between two different compartments of the catchment sediment cascade; catchment disconnectivity can be expressed as the degree to which any limiting factor constrains the efficiency of sediment transfer relationships Connectivity defined as the transfer of matter between two different Wester et al., 2014 landscape compartments Connectivity describes the integrated transfer of sediment across all Bracken et al., 2015 possible sources to all potential sinks in a system over the continuum of detachment, transport, and deposition, which is controlled by how the sediment moves between all geomorphic zones; on hillslopes, between hillslopes and channels, and within channels. Describe two fluxes as connected if they are in close spatial proximity Czuba and Foufoulaalong the river network; refer to connectivity as the state of two or more Georgiou, fluxes being connected; dynamic connectivity refers to how the connectivity of fluxes changes in time. Defines five layers of hydrologic connectivity as hillslope, hyporheic, Covino, 2017 stream-groundwater, riparian/floodplain, and longitudinal within

В

channels

Description	Metric	Reference
Primarily hydrologic metrics		
Integral connectivity scale	Average distance over which wet locations are	Western et al.,
lengths (ICSL)	connected using either Euclidean distances or topographically defined hydrologic distances; 1 of 15 indices of hillslope hydrologic connectivity in Bracken et al. (2013: Table 4)	2001
Attenuated imperviousness (I)	Weighted impervious area as a percentage of	Walsh and
$I = \left(\frac{\sum_{j} (A_{j} W_{j})}{A_{c}}\right)$	catchment area; Aj is the area of the j th impervious surface; Wj is the weighting applied to Aj; Ac is catchment area	Kunapo, 2009
River Connectivity Index (RCI) $DCI_P = \sum_{i=1}^{n} \frac{v_i^2}{V^2} * 100$	The size of disconnected river fragments between dams in relation to the total size of the original river network, based on Cote et al. (2009) DCI; size can be described in terms of volume (example at left), length, or other variables	Grill et al., 2014
Primarily sediment metrics		
Sediment delivery ratio (SDR) $SDR = \frac{net\ erosion}{total\ erosion}$	Measure of sediment connectivity	Brierley et al., 2006
Connectivity Index (IC)	D_{up} and D_{dn} are the upslope and downslope components of connectivity, respectively, with	Cavalli et al., 2013

$$IC = log_{10} \left(\frac{D_{up}}{D_{dn}} \right)$$

$$D_{up} = \overline{WS}\sqrt{A}$$

$$D_{dn} = \sum_{i} \frac{d_i}{W_i S_i}$$

$$W = 1 - \left(\frac{RI}{RI_{MAX}}\right)$$

Roughness Index (RI)

$$RI = \sqrt{\frac{\sum_{i=1}^{25} (x_i - x_m)^2}{25}}$$

Complexity index based on overall relief Dh_{max}

 $Dh_{max} = E_{max} - E_{min}$ and slope variability SV $SV = S_{max} - S_{min}$

Cluster Persistence Index (CPI)

$$CPI_{i} = \int_{\substack{\text{over all} \\ \text{times } t}} M_{j}^{(i)}(t)dt$$

connectivity increasing as IC increases; \overline{W} is the average weighting factor of the upslope contributing area, \overline{S} is the average slope gradient of the upslope contributing area, and A is the upslope contributing area; d_i is the length of the flow path along the i^{th} cell according to the steepest downslope direction, W_i and S_i are the weighting factor and the slope gradient of the i^{th} cell, respectively; RI_{MAX} is the maximum value of RI in the study area; 25 is the number of processing cells within a 5 X 5 moving window, x_i is the value of one specific cell of the residual topography within the moving window, and x_m is the mean of the 25 cell values.

Where E_{max} and E_{min} are the maximum and minimum elevations, respectively, in the catchment; S_{max} and S_{min} are the maximum and minimum, respectively, % slope within the area of analysis (moving window)

Defines clusters within a river network where mass (sediment) coalesces into a connected extent of the network; the superscript (i) denotes all clusters $M_j^{(i)}$ that occupy link i at time t

Baartman et al., 2013

Czuba and Foufoula-Georgiou, 2015

Yue et al., 2004

Metrics for diverse fluxes

$$C(t) = \sum_{i=1}^{m(t)} \sum_{j=1}^{n_i(t)} p_{ij}(t) S_{ij}(t)$$

 $= \frac{\sum_{i=1}^{\nu} \sum_{j=r+1}^{R} w_{ij} \frac{dx(j-r)}{d_{ij}}}{\sum_{i=1}^{\nu} \sum_{j=r+1}^{R} w_{ij}}$

Patch connectivity, along with line, vertex, and network connectivity, can be used to characterize landscape connectivity; patch connectivity is the average movement efficiency between patches; C is patch connectivity, $p_{ij}(t)$ is the area proportion of the j^{th} patch in the i^{th} land cover type to the total area under investigation at time t; S is movement efficiency; $0 \le C(t) \le 1.1$.

Directional connectivity index (DCI); i is a node index, j is a row index, r is the row containing the node i, R is the total number of rows in the direction of interest, dx is the relative pixel length along that direction, d_{ij} is the shortest connected structural or functional distance between node i and any node in row j, w_{ij} is a weighting function

Applies a connectivity analysis to a delta by identifying a set of objects (e.g., locations or variables) arranged in a network such that

Larsen et al., 2012

Newman et al., 2006; Heckmann et al., 2015;

Adjacency matrix

DCI

objects are nodes and connections or physical dependencies are links; evaluate connections between nodes using the mathematical technique of an adjacency matrix, which captures whether two nodes are connected, as well as link directionality and the strength of the connection

Passalacqua, 2017



Table 2. Historical (5,000 BC – early 1900s) contributions to connectivity in geomorphic systems from a fluvial perspective. Selections from Hugget (2007), Newsom (1997), Gregory and Lewin (2014) and authors' discretion.

Mesopotamia, Sumerians (5,000 -3,000 BC): Hydraulic-based irrigation and flood control projects of the Tigris and Euphrates via canals and drainage of floodplains and marshlands.

Egypt, Egyptians (5,000 -2,000 BC): River and society connections involving water supply, irrigation, and flood storage projects. Collection of river levels using the Roda 'nilometer' for predicting lateral river-floodplain connections for irrigation and flood control.

Emporer Yu the Great, China (2200-2101 BC): River network and basin mapping, and engineering of flood control using dikes, dams, dredging, and irrigation canal systems.

Lucius Anneaus Seneca (4 BC-AD 65): Roman philosopher recognition that rivers erode and create their valleys.

Claudius Ptolemy (100 AD – 168 AD): Greek-Egyptian scientists depicted the first river basin map of the Nile connecting the 'Mountains of the Moon' headwaters to the 'Upper, Middle, and Lower Lands' and eventually with the Mediterranean Sea.

Leonardo DaVinci (1452-1519): Italian renaissance scholar illustrated how rivers carved valleys and moved materials from one place and deposited them in another, and painted the first slope-contoured, shaded relief drainage map of the Arno River in Italy (1502-1503), complete with headwaters, tributaries, and main stem river connections.

Giovanni Targioni-Tozzetti (1712-1784): Italian scholar observed that river patterns and the courses they took in their valleys were a function of the lithology and processes of differential erosion.

Jean-Étienne Guettard (1715-1786): French naturalist recognized mountain to sea connections, i.e. sediment eroded from mountains was deposited as floodplains or carried to the sea.

James Hutton (1726-1797): Scottish geologist who recognized erosion was the dominant forces carving large river valleys. Hutton also engineered hydrologic connections on the landscape by building canals for navigation and water supply.

John Playfair (1748-1819): Scottish professor who expanded on Hutton's ideas, and showed that channels form in systematic order, whereby small rivers drain into larger rivers and so forth, until you have a mainstem river and valley complex, and that these river networks are organized into drainage basins.

Captain Henry M. Shreve (1785-1851): American soldier, artificially cut a neck through "Turnbull's Bend" on the Mississippi River disconnecting the river from its preferred path down the Atchafalaya in 1831. This lead to the construction of the Old River Control Structure in 1963, which has permanently controlled the course of the river ever since.

George Perkins Marsh (1801-1882): American environmentalists pioneered early understanding of human-induced land cover and land use changes and their connections to impacts to land and water processes.

Charles Darwin (1809-1882): English explorer observed rivers as agents of erosion, attributed anthropomorphic terms youth, middle age, old age, and rejuvenation to cycles of landscape evolution.

John Newberry (1822-1892): American geologist recognized that rivers carved tremendous canyons, i.e. Grand Canyon, through the terrain of the American West

John Wesley Powell (1834-1902): American soldier, professor, and head of USGS (1870-1892), first scientists to descend the Colorado River system from its headwaters. Established hydrologic surveys for commissioning of western US dams.

Grove Karl Gilbert (1843-1918): American geologists contributed substantially to our understanding of geomorphology as an open system of inputs, outputs, and fluxes of energy and material exchanges from his work in the western US and most notable his regional geologic-geomorphic descriptions of the Henry Mountains, Utah. Described landscape evolution through processes of erosion, incision, transport, and deposition.

Robert Horton (1875-1945): Textbook father of network-based stream order patterns from topographic analysis, and applications of stream order to quantifying drainage basin sizes, accumulation through a network, hill-slope erosion, and runoff processes.

John T. Hack (1913-1991): Established early theories of dynamic equilibrium and steady state models that described geomorphic processes and forms changing relative to a balanced steady state of inputs and outputs.

Luna B. Leopold (1915-2006): Established field methods for quantifying fluvial forms and processes, by understanding connections among sediment sources, transport, and deposition.

Arthur N. Strahler (1918-2002): Advanced Horton's stream order concepts into the format commonly used today, and contributed to quantitative methods for measuring other morphometric indices and hillslope erosion processes.

M. Gordan "Reds" Wolman (1924-2010): Established field methods for quantifying fluvial forms and processes, with a focus on floodplain depositional styles and the importance of drainage basin and local scale controls.

Richard Chorley (1927-2002): Introduced complex system theory to geomorphology through subsystem classification of morphological, cascading, process-response, and control systems.

Stanley Schumm (1927-2011): Developed concept of sediment budgets as a method for quantifying sediment sources, transport, and sinks through a drainage basin and for measuring sediment yield.

James C. Knox (1941- 2012): Provided significant evidence that human-induced changes have substantially more sedimentation impacts to rivers and floodplains than natural, climate-driven changes. Knox's work underscores the importance of why understanding river-landscape connectivity dynamics is critical to how we interpret geomorphic form, process, and management practices.

Table 3. Examples of methods used to identify signatures of geomorphic connectivity

Description	Sample references
Measured fluxes	Jampie references
Used 40 years of erosion-pin data, along with sediment trap data and sequential aerial photos and floodplain surveys to measure and infer sediment fluxes from hillslopes to pediments, floodplains, and channels	Godfrey et al., 2008
Measured precipitation, riparian water table, and stream flow and used these data as input to model hydrological connectivity	Jencso et al., 2009
Used arrays of electrical resistance sensors to quantify longitudinal connectivity of flow through time in drylands rivers	Jaeger and Olden, 2012
Used piezometers and subsurface samplers to measure vertical hydraulic gradients and specific discharge as indices of vertical hydrological	Wainwright et al., 2011
connectivity between a channel and hyporheic zone	
Inferred fluxes Catchment-scale sediment flow diagrams that identify spatial variability in patterns of sediment inputs, outputs, and storage based on direct measurements or, more commonly, spatial configuration of landscape units and relative volumes of stored sediment	Trimble, 1983; Fryirs et al., 2007
Visual or morphologic assessments of characteristics (size, spatial distribution, function) of landscape units in relation to facilitating or retarding sediment connectivity	Brierley et al., 2006
Modeled the delivery of landslide-generated sediment to channel networks; sediment generation from hillslopes and channel banks and its delivery to the channel network modeled using a modified form of SHALSTAB coupled to a network index version of TOPMODEL	Reid et al., 2007
Because alkalinity of stream waters reflects relative influence of groundwater and unsaturated zone runoff, used alkalinity as index of hydrologic connectivity at catchment scale	Tetzlaff et al., 2007
Used 1D hydrological modeling to infer hydrological connectivity among lakes and channels in the Danube River delta	Coops et al., 2008
Used in situ water level and MODIS satellite data to relate mainstem river level fluctuations to delta inundation on Canada's Peace-Athabasca delta; temporal covariance between the two datasets allows inference of hydrologic connectivity processes, as well as inundation extent	Pavelsky and Smith, 2008
Used diatom sedimentary assemblages to discriminate between three categories of delta lakes with differing types of hydrological connectivity to the Slave River of Canada	Sokal et al., 2008
Simulated runoff and sediment dynamics at the catchment scale with a dynamic landscape evolution model that can simulate erosion and sedimentation based on a limited number of input parameters	Lesschen et al., 2009
Assessed hydrologic connectivity of the Mackenzie River and lakes on its delta from duration of 'connection time' based on elevation of sill height for a lake and daily river water levels from stream gage records	Tank et al., 2009
Visual evaluation of location of sediment sources, degree of coupling to stream network, channel morphology, and magnitude of erosion & deposition following a rainstorm	Cavalli et al., 2013

Used 2D simulation and river corridor topography to numerically model Croke et al., 2013 flood inundation extent for varying discharges and from this inferred lateral connectivity between channel and floodplain Field observations of surface-flow connectivity combined with topography Phillips, 2013 of river corridor to infer relative degrees of hydrological connectivity among active channel and abandoned channel water bodies on the floodplain Numerically simulated coarse sediment transport via diverse geomorphic Heckmann and processes (rockfall, debris flows, slope wash, fluvial transport) and used Schwanghart, 2013 these data in graph-based network analysis Data from ground surveys used with digital terrain model differencing Wester et al., 2014 techniques & morphological sediment budgets to infer sediment connectivity Measured rates of channel migration from sequential aerial photos used to Czuba and Foufoulaidentify locations of enhanced geomorphic change, which is inferred to Georgiou, 2015 reflect spatial variation in sand transport Used archival digital photogrammetry to reconstruct history of topographic Micheletti et al., 2015 change and inferred sediment fluxes in a catchment Represented sediment transport from each source in a watershed as a Schmitt et al., 2016 suite of individual cascading processes that are incorporated into an

integrated modeling framework of sediment cascades that is used to infer

patterns of connectivity and locations of disconnectivity

Policy.

Table 4. Aspects of landscape connectivity important in managing sediment regime

Identify expectations and realistic targets: The key issue in managing landscape connectivity is determination of 'what are we measuring against' (i.e., what is expected in any given system)? Recognizing explicitly that human disturbance has modified natural process linkages in a given catchment, what attributes of the prevailing sediment regime are manageable (i.e., what is realistically possible)? Inevitably, these are context- and catchment-specific situation.

Identify relevant components of sediment dynamics: Develop an understanding of forms and rates of sediment generation and patterns of sediment stores and their ease/frequency of reworking in a given system. How have human activities modified natural connectivity relationships in that system? How have these changes impacted upon the evolutionary trajectory of the system and over what timeframe? Is it possible, or desirable, for human activities to manage or reverse these traits?

Identify rates of sediment movement and geomorphic recovery: Quantify time frames of sediment movement through a system as a basis to evaluate whether geomorphic river recovery is possible. This entails analysis of the extent to which human disturbance has modified natural patterns and trends of sediment sources, transfer and deposition (Fryirs and Brierley, 2009). In some cases, rates of movement have been accelerated (e.g., deforestation), elsewhere they have been suppressed (e.g., dams). In some instances, excess sediments are available to be reworked such that aggradation may ensue in downstream reaches (e.g., legacy effects of human impacts such as mining activities or abandoned water mills), elsewhere limited upstream availability of sediment (sediment exhaustion) may inhibit prospects for geomorphic recovery in downstream reaches where channel are over-enlarged (e.g., Fryirs and Brierley, 2001; Hooke, 2003; Brooks and Brierley, 2004).

Identify the catchment context: The sediment regime and associated process morphodynamics in any given reach must be viewed in their catchment context, assessing how upstream and downstream reaches influence the reach of interest. Any given reach is subjected to changes in boundary conditions. Most reaches are adjusting to legacy effects (e.g., Coulthard and Macklin, 2003; James, 2010; Evrard et al., 2011; Wohl, 2015). Longitudinal connectivity relationships determine the nature, extent and rate with which changes to boundary conditions in one part of a system impact upon morphodynamic interactions elsewhere in that system. Pulses of sediment movement through river systems operate over different timescales and with variable impacts on a reach-by-reach basis. Resulting aggradational-degradational trends exert a key control upon channel adjustments over a range of timescales.


Identify the historical range of variability: Caution must be applied in the use of theoretical regime principles to predict rates of sediment movement and associated forms and rates of channel adjustment, as these framings assume continuity and uniformity in sediment inputs. However, sediment inputs vary and we need to know when and how they are likely to change if we are to make these assessments. In light of this issue, analysis of the historical range of variability of a river reach provides a critical basis to inform management applications pertaining to the range of channel sizes and configuration that are appropriate or expected for a given setting (Wohl, 2011; Rathburn et al., 2013; Reid and Brierley, 2015).

Identify natural levels of connectivity: If working in a largely disconnected landscape, maintain

disconnectivity of longitudinal process in interactions whenever possible. For example, if wetlands associated with discontinuous watercourses are present, these features exert important controls on downstream fluxes and create unique habitats and nutrient storage that are important to preserve (Brierley et al., 1999).



Geomorphic Process	Cyclic	Graded	Steady
Sediment transport	↑ connected	↓ connected	↓ connected
bedload	↑ connected	↓ connected	↓ connected
suspended load	↑ connected	↓ connected	↑ connected
Knickpoint retreat	↑ connected	↑ connected	<u>+</u> connected
Planform adjustments	↑ connected	↓ connected	<u>+</u> connected
Longitudinal profile evolution	↑ connected	↓ connected	<u>+</u> connected
Drainage network development	<u>+</u> connected	个 connected	个 connected






Figure 2 254x190mm (300 x 300 DPI)




Figure 3 254x190mm (300 x 300 DPI)

