

Connes–Landi Deformation of Spectral Triples

Makoto Yamashita

Cardiff University, School of Mathematics

EU-NCG 4th Annual Meeting Bucharest 2011

Based on:

M. Yamashita. Connes–Landi deformation of spectral triples, *Lett. Math. Phys.* **94**(3):263–291, 2010

- 1 θ -deformation of spectral triples
- 2 Cyclic cohomology of one-parameter crossed product
- 3 Cyclic cohomology of Connes-Landi deformation

Connes-Landi deformation

\mathcal{A} : Fréchet, $\sigma: \mathbb{T}^2 \curvearrowright \mathcal{A}$ smooth

Connes-Landi deformation

\mathcal{A} : Fréchet, $\sigma: \mathbb{T}^2 \curvearrowright \mathcal{A}$ smooth

$\theta\mathbb{R}/\mathbb{Z}$: deformed product $*_{\theta}$ on \mathcal{A}

$$f *_{\theta} g = e^{\pi i \theta (mn' - m'n)} fg$$

when f is a \mathbb{T}^2 -eigenvector of weight $(m, n) \in \mathbb{Z}^2$, g is a one of weight (m', n') .

$$\mathcal{A}_{\theta} = (\mathcal{A}, *_{\theta})$$

Connes-Landi deformation

\mathcal{A} : Fréchet, $\sigma: \mathbb{T}^2 \curvearrowright \mathcal{A}$ smooth

$\theta\mathbb{R}/\mathbb{Z}$: deformed product $*_{\theta}$ on \mathcal{A}

$$f *_{\theta} g = e^{\pi i \theta (mn' - m'n)} fg$$

when f is a \mathbb{T}^2 -eigenvector of weight $(m, n) \in \mathbb{Z}^2$, g is a one of weight (m', n') .

$$\mathcal{A}_{\theta} = (\mathcal{A}, *_{\theta})$$

$A_{\theta} = \overline{\mathcal{A}_{\theta}}$: deformation of $A = \overline{\mathcal{A}}$ by Rieffel (1993)

CL-deformation of spectral triples

(\mathcal{A}, H, D) spectral triple, $U: \mathbb{T}^2 \curvearrowright H$

$$\sigma_t(a) = \text{Ad}_{U_t}(a),$$

$$\text{Ad}_{U_t}(D) = D$$

CL-deformation of spectral triples

(\mathcal{A}, H, D) spectral triple, $U: \mathbb{T}^2 \curvearrowright H$

$$\sigma_t(a) = \text{Ad}_{U_t}(a),$$

$$\text{Ad}_{U_t}(D) = D$$

$$\Rightarrow \exists \mathcal{A}_\theta \curvearrowright H, \quad a^{(\theta)}\xi = e^{\pi i \theta(mn' - m'n)} a\xi$$

when $a \in \mathcal{A}$ is a \mathbb{T}^2 -eigenvector of weight (m, n) and $\xi \in H$ is a one of weight (m', n')

CL-deformation of spectral triples

(\mathcal{A}, H, D) spectral triple, $U: \mathbb{T}^2 \curvearrowright H$

$$\sigma_t(a) = \text{Ad}_{U_t}(a),$$

$$\text{Ad}_{U_t}(D) = D$$

$$\Rightarrow \exists \mathcal{A}_\theta \curvearrowright H, \quad a^{(\theta)}\xi = e^{\pi i \theta(mn' - m'n)} a\xi$$

when $a \in \mathcal{A}$ is a \mathbb{T}^2 -eigenvector of weight (m, n) and $\xi \in H$ is a one of weight (m', n')

\Rightarrow New spectral triple $(\mathcal{A}_\theta, H, D)$

Crossed product presentation of CL-deformation

Strong Morita equivalence

$$A_\theta \simeq (A \otimes C(\mathbb{T}_\theta^2))^{\sigma \otimes \gamma} \simeq_{\text{KK}} \mathbb{T}^2 \ltimes_{\sigma \otimes \gamma} (A \otimes C(\mathbb{T}_\theta^2))$$

Crossed product presentation of CL-deformation

Strong Morita equivalence

$$A_\theta \simeq (A \otimes C(\mathbb{T}_\theta^2))^{\sigma \otimes \gamma} \simeq_{\text{KK}} \mathbb{T}^2 \ltimes_{\sigma \otimes \gamma} (A \otimes C(\mathbb{T}_\theta^2))$$

Automorphisms of $\mathbb{Z} \ltimes_{\widehat{\sigma^{(2)}}} \mathbb{T}^2 \ltimes_\sigma A$:

$$\widehat{\sigma^{(1)}}\left(\sum_{k \in \mathbb{Z}} f_k v^k\right) = \sum_{k \in \mathbb{Z}} \widehat{\sigma^{(1)}}(f_k) v^k,$$

Crossed product presentation of CL-deformation

Strong Morita equivalence

$$A_\theta \simeq (A \otimes C(\mathbb{T}_\theta^2))^{\sigma \otimes \gamma} \simeq_{\text{KK}} \mathbb{T}^2 \ltimes_{\sigma \otimes \gamma} (A \otimes C(\mathbb{T}_\theta^2))$$

Automorphisms of $\mathbb{Z} \ltimes_{\widehat{\sigma^{(2)}}} \mathbb{T}^2 \ltimes_\sigma A$:

$$\widehat{\sigma^{(1)}}\left(\sum_{k \in \mathbb{Z}} f_k v^k\right) = \sum_{k \in \mathbb{Z}} \widehat{\sigma^{(1)}}(f_k) v^k,$$

 $\mathbb{R} \ltimes_{\sigma^{(1)}} A$ is identified with the mapping torus of $\widehat{\sigma^{(1)}}: \mathbb{Z} \curvearrowright \mathbb{T} \ltimes_{\sigma^{(1)}} A$

Crossed product presentation of CL-deformation

Strong Morita equivalence

$$A_\theta \simeq (A \otimes C(\mathbb{T}_\theta^2))^{\sigma \otimes \gamma} \simeq_{\text{KK}} \mathbb{T}^2 \ltimes_{\sigma \otimes \gamma} (A \otimes C(\mathbb{T}_\theta^2))$$

Automorphisms of $\mathbb{Z} \ltimes_{\widehat{\sigma^{(2)}}} \mathbb{T}^2 \ltimes_\sigma A$:

$$\widehat{\sigma^{(1)}}\left(\sum_{k \in \mathbb{Z}} f_k v^k\right) = \sum_{k \in \mathbb{Z}} \widehat{\sigma^{(1)}}(f_k) v^k,$$

$\mathbb{R} \ltimes_{\sigma^{(1)}} A$ is identified with the mapping torus of $\widehat{\sigma^{(1)}}: \mathbb{Z} \curvearrowright \mathbb{T} \ltimes_{\sigma^{(1)}} A$

$$\Rightarrow \mathbb{Z} \ltimes_{\sigma_\theta^{(2)} \widehat{\sigma^{(1)}}} \mathbb{T} \ltimes_{\sigma^{(1)}} A \simeq_{\text{KK}} \mathbb{R} \ltimes_{\sigma_{\theta t}^{(2)} \widehat{\sigma^{(1)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} A$$

an analogue of “noncommutative torus” \Leftrightarrow “Kronecker foliation”

Cyclic cocycles

Differential graded algebra (Ω^*, d) , homomorphism $\mathcal{A} \rightarrow \Omega^0$

Cyclic cocycles

Differential graded algebra (Ω^*, d) , homomorphism $\mathcal{A} \rightarrow \Omega^0$

cyclic n -cocycle \leftrightarrow closed trace on Ω^n

$$\phi(f^0, \dots, f^n) = \tau_\phi(f^0 df^1 \cdots df^n)$$

Cyclic cocycles

Differential graded algebra (Ω^*, d) , homomorphism $\mathcal{A} \rightarrow \Omega^0$

cyclic n -cocycle \leftrightarrow closed trace on Ω^n

$$\phi(f^0, \dots, f^n) = \tau_\phi(f^0 df^1 \cdots df^n)$$

Cyclic cohomology group:

$$\mathrm{HC}^n(\mathcal{A}) = \{\text{cyclic } n\text{-cocycles}\} / \{\text{cyclic } n\text{-coboundary}\}$$

Cyclic cocycles

Differential graded algebra (Ω^*, d) , homomorphism $\mathcal{A} \rightarrow \Omega^0$

cyclic n -cocycle \leftrightarrow closed trace on Ω^n

$$\phi(f^0, \dots, f^n) = \tau_\phi(f^0 df^1 \cdots df^n)$$

Cyclic cohomology group:

$$\mathrm{HC}^n(\mathcal{A}) = \{\text{cyclic } n\text{-cocycles}\} / \{\text{cyclic } n\text{-coboundary}\}$$

$$\mathrm{HC}^n(\mathcal{A}) \times K_n(\mathcal{A}) \rightarrow \mathbb{C},$$

$$\langle [\phi], [e] \rangle = \phi(e, \dots, e) \quad (\phi : \text{cyclic } 2k\text{-cocycle}, e : \text{projection})$$

$$\langle [\phi], [u] \rangle = \phi(u^*, u, u^*, \dots, u) \quad (\phi : \text{cyclic } 2k + 1\text{-cocycle}, u : \text{unitary})$$

Cyclic cocycles

Differential graded algebra (Ω^*, d) , homomorphism $\mathcal{A} \rightarrow \Omega^0$

cyclic n -cocycle \leftrightarrow closed trace on Ω^n

$$\phi(f^0, \dots, f^n) = \tau_\phi(f^0 df^1 \cdots df^n)$$

Cyclic cohomology group:

$$\mathrm{HC}^n(\mathcal{A}) = \{\text{cyclic } n\text{-cocycles}\} / \{\text{cyclic } n\text{-coboundary}\}$$

$$\mathrm{HC}^n(\mathcal{A}) \times K_n(\mathcal{A}) \rightarrow \mathbb{C},$$

$$\langle [\phi], [e] \rangle = \phi(e, \dots, e) \quad (\phi : \text{cyclic } 2k\text{-cocycle}, e : \text{projection})$$

$$\langle [\phi], [u] \rangle = \phi(u^*, u, u^*, \dots, u) \quad (\phi : \text{cyclic } 2k + 1\text{-cocycle}, u : \text{unitary})$$

$$S : \mathrm{HC}^n(\mathcal{A}) \rightarrow \mathrm{HC}^{n+2}(\mathcal{A}), \mathrm{HP}^*(\mathcal{A}) = \lim_{k \rightarrow \infty} \mathrm{HC}^{2k+*}(\mathcal{A})$$

Connes-Thom isomorphism in cyclic cohomology

Theorem (Elliott-Natsume-Nest, 1988)

\mathcal{A} : Fréchet, $\alpha : \mathbb{R} \curvearrowright \mathcal{A}$ smooth

Connes-Thom isomorphism in cyclic cohomology

Theorem (Elliott-Natsume-Nest, 1988)

\mathcal{A} : Fréchet, $\alpha : \mathbb{R} \curvearrowright \mathcal{A}$ smooth

$\Phi : K_*(\overline{\mathcal{A}}) \rightarrow K_{*+1}(\mathbb{R} \rtimes_{\alpha} \overline{\mathcal{A}})$ the Connes-Thom isomorphism

Connes-Thom isomorphism in cyclic cohomology

Theorem (Elliott-Natsume-Nest, 1988)

\mathcal{A} : Fréchet, $\alpha : \mathbb{R} \curvearrowright \mathcal{A}$ smooth

$\Phi : K_*(\overline{\mathcal{A}}) \rightarrow K_{*+1}(\mathbb{R} \rtimes_{\alpha} \overline{\mathcal{A}})$ the Connes-Thom isomorphism

$$\#_{\alpha} : \mathrm{HC}^n(\mathcal{A}) \rightarrow \mathrm{HC}^{n+1}(\mathbb{R} \rtimes_{\alpha}^{\infty} \mathcal{A})$$

$$\langle \phi, x \rangle = \langle \#_{\alpha}(\phi), \Phi(x) \rangle$$

Connes-Thom isomorphism in cyclic cohomology

Theorem (Elliott-Natsume-Nest, 1988)

\mathcal{A} : Fréchet, $\alpha : \mathbb{R} \curvearrowright \mathcal{A}$ smooth

$\Phi : K_*(\overline{\mathcal{A}}) \rightarrow K_{*+1}(\mathbb{R} \rtimes_{\alpha} \overline{\mathcal{A}})$ the Connes-Thom isomorphism

$$\#_{\alpha} : \mathrm{HC}^n(\mathcal{A}) \rightarrow \mathrm{HC}^{n+1}(\mathbb{R} \rtimes_{\alpha}^{\infty} \mathcal{A})$$

$$\langle \phi, x \rangle = \langle \#_{\alpha}(\phi), \Phi(x) \rangle$$

Example

τ : trace on \mathcal{A} , invariant under α

$i_X \tau(a, b) = \tau(ah(b))$: cyclic 1-cocycle on \mathcal{A}

Connes-Thom isomorphism in cyclic cohomology

Theorem (Elliott-Natsume-Nest, 1988)

\mathcal{A} : Fréchet, $\alpha : \mathbb{R} \curvearrowright \mathcal{A}$ smooth

$\Phi : K_*(\overline{\mathcal{A}}) \rightarrow K_{*+1}(\mathbb{R} \rtimes_{\alpha} \overline{\mathcal{A}})$ the Connes-Thom isomorphism

$$\#_{\alpha} : \mathrm{HC}^n(\mathcal{A}) \rightarrow \mathrm{HC}^{n+1}(\mathbb{R} \rtimes_{\alpha}^{\infty} \mathcal{A})$$

$$\langle \phi, x \rangle = \langle \#_{\alpha}(\phi), \Phi(x) \rangle$$

Example

τ : trace on \mathcal{A} , invariant under α

$i_X \tau(a, b) = \tau(ah(b))$: cyclic 1-cocycle on \mathcal{A}

$$\Rightarrow \#_{\hat{\alpha}}(\hat{\tau}) = \mathrm{Tr} \otimes i_X \tau \text{ on } \mathbb{R} \rtimes_{\hat{\alpha}}^{\infty} \mathbb{R} \rtimes_{\alpha}^{\infty} \mathcal{A} \simeq \mathcal{K}^{\infty} \hat{\otimes} \mathcal{A}$$

Invariant cocycles

\mathcal{A} : Fréchet, $\alpha: \mathbb{R} \curvearrowright \mathcal{A}$ smooth

X : generator of α

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under α

Invariant cocycles

\mathcal{A} : Fréchet, $\alpha: \mathbb{R} \curvearrowright \mathcal{A}$ smooth

X : generator of α

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under α

- Dual cocycle $\hat{\phi}$ on $\mathbb{R} \times_{\alpha}^{\infty} \mathcal{A}$

$$\hat{\phi}(f^0, \dots, f^n) = \int_{\sum_{j=0}^n t_j = 0} \phi(f_{t_0}^0, \dots, f_{t_n}^n)$$

Invariant cocycles

\mathcal{A} : Fréchet, $\alpha: \mathbb{R} \curvearrowright \mathcal{A}$ smooth

X : generator of α

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under α

- Dual cocycle $\hat{\phi}$ on $\mathbb{R} \times_{\alpha}^{\infty} \mathcal{A}$

$$\hat{\phi}(f^0, \dots, f^n) = \int_{\sum_{j=0}^n t_j = 0} \phi(f_{t_0}^0, \dots, f_{t_n}^n)$$

- Interior product $i_X \phi$ on \mathcal{A}

$$i_X \phi(a_0, \dots, a_{n+1}) = \sum_{j=1}^{n+1} (-1)^j \tau_{\phi}(a_0 da_1 \cdots X(a_j) \cdots da_{n+1}).$$

Invariant cocycles and ENN-isomorphism

Theorem (Y., 2010)

\mathcal{A} : Fréchet, $\alpha: \mathbb{R} \curvearrowright \mathcal{A}$ smooth

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under α

Invariant cocycles and ENN-isomorphism

Theorem (Y., 2010)

\mathcal{A} : Fréchet, $\alpha: \mathbb{R} \curvearrowright \mathcal{A}$ smooth

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under α

$$[\#_{\hat{\alpha}} \hat{\phi}] = [i_X \phi] \text{ in } \mathrm{HC}^{n+1}(\mathcal{A})$$

Outline: \tilde{X} : generator of $\hat{\sigma} \Rightarrow \hat{\phi} = i_{\tilde{X}} \phi$

Invariant cocycles and ENN-isomorphism

Theorem (Y., 2010)

 \mathcal{A} : Fréchet, $\alpha: \mathbb{R} \curvearrowright \mathcal{A}$ smooth ϕ : cyclic n -cocycle on \mathcal{A} , invariant under α

$$[\#_{\hat{\alpha}} \hat{\phi}] = [i_X \phi] \text{ in } \mathrm{HC}^{n+1}(\mathcal{A})$$

Outline: \tilde{X} : generator of $\hat{\sigma} \Rightarrow \hat{\phi} = i_{\tilde{X}} \phi$ Connes's 2×2 -matrix trick for σ and $\hat{\sigma}$:

$$\mathbb{R} \curvearrowright M_2(\mathbb{R} \rtimes_{\hat{\sigma}} \mathbb{R} \rtimes_{\sigma} \mathcal{A}) \quad (\text{with generator } Y)$$

\Rightarrow Two embeddings $\Psi_1, \Psi_2: \mathbb{R} \rtimes_{\hat{\sigma}} \mathbb{R} \rtimes_{\sigma} \mathcal{A} \rightarrow M_2(\mathbb{R} \rtimes_{\hat{\sigma}} \mathbb{R} \rtimes_{\sigma} \mathcal{A})$

Invariant cocycles and ENN-isomorphism

Theorem (Y., 2010)

 \mathcal{A} : Fréchet, $\alpha: \mathbb{R} \curvearrowright \mathcal{A}$ smooth ϕ : cyclic n -cocycle on \mathcal{A} , invariant under α

$$[\#_{\hat{\alpha}} \hat{\phi}] = [i_X \phi] \text{ in } \mathrm{HC}^{n+1}(\mathcal{A})$$

Outline: \tilde{X} : generator of $\hat{\sigma} \Rightarrow \hat{\phi} = i_{\tilde{X}} \phi$ Connes's 2×2 -matrix trick for σ and $\hat{\sigma}$:

$$\mathbb{R} \curvearrowright M_2(\mathbb{R} \rtimes_{\hat{\sigma}} \mathbb{R} \rtimes_{\sigma} \mathcal{A}) \quad (\text{with generator } Y)$$

\Rightarrow Two embeddings $\Psi_1, \Psi_2: \mathbb{R} \rtimes_{\hat{\sigma}} \mathbb{R} \rtimes_{\sigma} \mathcal{A} \rightarrow M_2(\mathbb{R} \rtimes_{\hat{\sigma}} \mathbb{R} \rtimes_{\sigma} \mathcal{A})$

$$\Psi_1^*(i_Y(\phi \otimes \mathrm{Tr}_{M_2})) = i_{\tilde{X}} \phi \qquad \Psi_2^*(i_Y(\phi \otimes \mathrm{Tr}_{M_2})) = i_X \phi$$

Invariant cocycles and ENN-isomorphism, cont.

Example:

- α : action of \mathbb{Z} on a Fréchet algebra \mathcal{B}

Invariant cocycles and ENN-isomorphism, cont.

Example:

- α : action of \mathbb{Z} on a Fréchet algebra \mathcal{B}
- $\partial: K_{*+1}(\mathbb{Z} \ltimes_{\alpha} \mathcal{B}) \rightarrow K_*(\mathcal{B})$ connecting map in the Pimsner–Voiculescu sequence

Invariant cocycles and ENN-isomorphism, cont.

Example:

- α : action of \mathbb{Z} on a Fréchet algebra \mathcal{B}
- $\partial: K_{*+1}(\mathbb{Z} \ltimes_{\alpha} \mathcal{B}) \rightarrow K_*(\mathcal{B})$ connecting map in the Pimsner–Voiculescu sequence
- D : generator of $\hat{\alpha}$ on $\mathbb{Z} \ltimes_{\alpha} \mathcal{B}$

Invariant cocycles and ENN-isomorphism, cont.

Example:

- α : action of \mathbb{Z} on a Fréchet algebra \mathcal{B}
- $\partial: K_{*+1}(\mathbb{Z} \rtimes_{\alpha} \mathcal{B}) \rightarrow K_*(\mathcal{B})$ connecting map in the Pimsner–Voiculescu sequence
- D : generator of $\hat{\alpha}$ on $\mathbb{Z} \rtimes_{\alpha} \mathcal{B}$
- ϕ : $\hat{\alpha}$ -invariant cyclic n -cocycle on $\mathbb{Z} \rtimes_{\alpha} \mathcal{B}$

Invariant cocycles and ENN-isomorphism, cont.

Example:

- α : action of \mathbb{Z} on a Fréchet algebra \mathcal{B}
- $\partial: K_{*+1}(\mathbb{Z} \rtimes_{\alpha} \mathcal{B}) \rightarrow K_*(\mathcal{B})$ connecting map in the Pimsner–Voiculescu sequence
- D : generator of $\hat{\alpha}$ on $\mathbb{Z} \rtimes_{\alpha} \mathcal{B}$
- ϕ : $\hat{\alpha}$ -invariant cyclic n -cocycle on $\mathbb{Z} \rtimes_{\alpha} \mathcal{B}$
 $\Rightarrow \langle i_D \phi, x \rangle = \langle \phi, \partial(x) \rangle$

Correspondence of invariant cocycles

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under σ

Correspondence of invariant cocycles

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under σ

$\Rightarrow \hat{\phi}$ is a cyclic cocycle over $\mathbb{R} \ltimes_{\widehat{\sigma^{(1)}, \sigma_{\theta t}^{(2)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A}$.

Correspondence of invariant cocycles

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under σ

$\Rightarrow \hat{\phi}$ is a cyclic cocycle over $\mathbb{R} \ltimes_{\widehat{\sigma^{(1)}, \sigma_{\theta t}^{(2)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A}$.

e : minimal projection in \mathcal{K}^∞ , embedding

$$\Psi: \mathcal{A}_\theta \rightarrow \mathcal{A} \hat{\otimes} \mathcal{K}^\infty \simeq \mathbb{R} \ltimes_{\widehat{\sigma^{(1)}, \sigma_{\theta t}^{(2)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A},$$

$$x \mapsto x \otimes e$$

Correspondence of invariant cocycles

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under σ

$\Rightarrow \hat{\phi}$ is a cyclic cocycle over $\mathbb{R} \ltimes_{\widehat{\sigma^{(1)}, \sigma_{\theta t}^{(2)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A}$.

e : minimal projection in \mathcal{K}^∞ , embedding

$$\Psi: \mathcal{A}_\theta \rightarrow \mathcal{A} \hat{\otimes} \mathcal{K}^\infty \simeq \mathbb{R} \ltimes_{\widehat{\sigma^{(1)}, \sigma_{\theta t}^{(2)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A},$$

$$x \mapsto x \otimes e$$

$\phi^{(\theta)} := \Psi^*(\phi)$: cyclic n -cocycle on \mathcal{A}_θ

Correspondence of invariant cocycles

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under σ

$\Rightarrow \hat{\phi}$ is a cyclic cocycle over $\mathbb{R} \ltimes_{\widehat{\sigma^{(1)}, \sigma_{\theta t}^{(2)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A}$.

e : minimal projection in \mathcal{K}^∞ , embedding

$$\Psi: \mathcal{A}_\theta \rightarrow \mathcal{A} \hat{\otimes} \mathcal{K}^\infty \simeq \mathbb{R} \ltimes_{\widehat{\sigma^{(1)}, \sigma_{\theta t}^{(2)}}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A},$$

$$x \mapsto x \otimes e$$

$\phi^{(\theta)} := \Psi^*(\phi)$: cyclic n -cocycle on \mathcal{A}_θ

Rem. ϕ is a trace on $\mathcal{A} \Rightarrow \phi^{(\theta)}(x^{(\theta)}) = \phi(x)$

Comparison of cocycles

$$\#_{\widehat{\sigma^{(1)}}, \sigma_{\theta t}^{(2)}} \circ \#_{\sigma^{(1)}} : \text{HP}(\mathcal{A}) \rightarrow \text{HP}(\mathbb{R} \ltimes_{\widehat{\sigma^{(1)}}, \sigma_{\theta t}^{(2)}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A}) \simeq \text{HP}(\mathcal{A}_\theta)$$

$$\Xi^{(\theta)} := \left(\#_{\widehat{\sigma^{(1)}}, \sigma_{\theta t}^{(2)}} \circ \#_{\sigma^{(1)}} \right)^{-1}$$

Comparison of cocycles

$$\#_{\widehat{\sigma(1)}, \sigma_{\theta t}^{(2)}} \circ \#_{\sigma(1)} : \text{HP}(\mathcal{A}) \rightarrow \text{HP}(\mathbb{R} \ltimes_{\widehat{\sigma(1)}, \sigma_{\theta t}^{(2)}} \mathbb{R} \ltimes_{\sigma(1)} \mathcal{A}) \simeq \text{HP}(\mathcal{A}_\theta)$$

$$\Xi^{(\theta)} := \left(\#_{\widehat{\sigma(1)}, \sigma_{\theta t}^{(2)}} \circ \#_{\sigma(1)} \right)^{-1}$$

Theorem (Y., 2010)

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under σ

Comparison of cocycles

$$\#_{\widehat{\sigma^{(1)}}, \sigma_{\theta t}^{(2)}} \circ \#_{\sigma^{(1)}} : \mathrm{HP}(\mathcal{A}) \rightarrow \mathrm{HP}(\mathbb{R} \ltimes_{\widehat{\sigma^{(1)}}, \sigma_{\theta t}^{(2)}} \mathbb{R} \ltimes_{\sigma^{(1)}} \mathcal{A}) \simeq \mathrm{HP}(\mathcal{A}_\theta)$$

$$\Xi^{(\theta)} := \left(\#_{\widehat{\sigma^{(1)}}, \sigma_{\theta t}^{(2)}} \circ \#_{\sigma^{(1)}} \right)^{-1}$$

Theorem (Y., 2010)

ϕ : cyclic n -cocycle on \mathcal{A} , invariant under σ

$$[\Xi^{(\theta)}(\phi^{(\theta)})] = [\phi] + \theta[i_{X_1} i_{X_2} \phi] \in \mathrm{HC}^{n+2}(\mathcal{A})$$

Invariance of Chern-Connes character

Theorem (Y., 2010)

(\mathcal{A}, H, D) spectral triple, $U: \mathbb{T}^2 \curvearrowright H$

$$\sigma_t(a) = \text{Ad}_{U_t}(a),$$

$$\text{Ad}_{U_t}(D) = D$$

$(\mathcal{A}_\theta, H, D)$ deformed triple

Invariance of Chern-Connes character

Theorem (Y., 2010)

(\mathcal{A}, H, D) spectral triple, $U: \mathbb{T}^2 \curvearrowright H$

$$\sigma_t(a) = \text{Ad}_{U_t}(a),$$

$$\text{Ad}_{U_t}(D) = D$$

$(\mathcal{A}_\theta, H, D)$ deformed triple

$$\langle \text{ch}_{(\mathcal{A}, H, D)}, x \rangle = \langle \Xi^{(\theta)}(\text{ch}_{(\mathcal{A}_\theta, H, D)}), x \rangle,$$

i.e. “ $\Xi^{(\theta)}(\text{ch}_{(H, D)}) = \text{ch}_{(H, D)}$.”

Invariance of Chern-Connes character

Theorem (Y., 2010)

(\mathcal{A}, H, D) spectral triple, $U: \mathbb{T}^2 \curvearrowright H$

$$\sigma_t(a) = \text{Ad}_{U_t}(a), \quad \text{Ad}_{U_t}(D) = D$$

$(\mathcal{A}_\theta, H, D)$ deformed triple

$$\langle \text{ch}_{(\mathcal{A}, H, D)}, x \rangle = \langle \Xi^{(\theta)}(\text{ch}_{(\mathcal{A}_\theta, H, D)}), x \rangle,$$

i.e. “ $\Xi^{(\theta)}(\text{ch}_{(H, D)}) = \text{ch}_{(H, D)}$.”

∴

$$\langle \Xi^{(\theta)}(\text{ch}_{(\mathcal{A}_\theta, H, D)}), x \rangle = \langle \text{ch}_{(\mathcal{A}, H, D)}, x \rangle + \theta \langle i_{X_1} i_{X_2} \text{ch}_{(\mathcal{A}, H, D)}, x \rangle$$

and

$$\forall \theta : \langle \Xi^{(\theta)}(\text{ch}_{(\mathcal{A}_\theta, H, D)}), x \rangle \in \mathbb{Z}.$$