

Review

Connexin-related signaling in cell death: to live or let die?

E Decrock^{1,6,7}, M Vinken^{*2,6,8}, E De Vuyst¹, DV Krysko^{3,4,8}, K D'Herde⁵, T Vanhaecke^{2,8}, P Vandennebee^{3,4}, V Rogiers^{2,9} and L Leybaert^{1,9}

Evidence is accumulating that some forms of cell death, like apoptosis, are not only governed by the complex interplay between extracellular and intracellular signals but are also strongly influenced by intercellular communicative networks. The latter is provided by arrays of channels consisting of connexin proteins, with gap junctions directly connecting the cytoplasm of neighboring cells and hemichannels positioned as pores that link the cytoplasm to the extracellular environment. The role of gap junctions in cell death communication has received considerable interest and recently hemichannels have joined in as potentially toxic pores adding their part to the cell death process. However, despite a large body of existing evidence, especially for gap junctions, the exact contribution of the connexin channel family still remains controversial, as both gap junctions and hemichannels may furnish cell death as well as cell survival signals. An additional layer of complexity is formed by the fact that connexin proteins as such, beyond their channel function, may influence the cell death process. We here review the current knowledge on connexins and their channels in cell death and specifically address the molecular mechanisms that underlie connexin-related signaling. We also briefly focus on pannexins, a novel set of connexin-like proteins that have been implicated in cellular responses to pathological insults.

Cell Death and Differentiation (2009) 16, 524–536; doi:10.1038/cdd.2008.196; published online 6 February 2009

In multicellular organisms, the maintenance of tissue homeostasis ultimately relies on the critical balance between cell growth and cell death. Among the various cell death types, apoptosis has been most extensively characterized. Being the conceptual counterpart of necrosis, apoptosis is a genetically programmed and well-orchestrated process of selective cell deletion that occurs in all tissues as part of the normal cellular turnover.¹ It is also involved in a growing number of pathological conditions, such as in ischemia-related cell injury following stroke.^{2,3} Two major apoptotic pathways can be distinguished, the mitochondria-mediated intrinsic cascade and the death receptor-mediated extrinsic pathway.^{1,4–6} Both rely on the proteolytic activity of an evolutionary conserved family of cysteine proteases – the caspases – which form the

biochemical basis of the apoptotic phenotype.⁷ They are responsible for the cleavage of a large number of cellular proteins including major cytoplasmic and nuclear elements. The extrinsic signaling pathway is triggered by the binding of an extracellular death ligand, such as Fas ligand, tumor necrosis factor α (TNF α) or TNF-related apoptosis-inducing ligand (TRAIL) to their corresponding receptors at the plasma membrane (PM).^{1,4–6} In contrast, the intrinsic signaling pathway is mediated by mitochondria and involves a diverse array of non-receptor-mediated stimuli that produce intracellular signals directly acting on targets within the cell. It is regulated by the B-cell lymphoma-2 (Bcl-2) family of pro- and anti-apoptotic proteins.⁸ Necrotic cell death, on the other hand, has been often considered to be a passive process,

¹Department of Basic Medical Sciences – Physiology group, Faculty of Medicine and Health Sciences, Ghent University, B-9000 Ghent, Belgium; ²Department of Toxicology, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel, B-1090 Brussels, Belgium; ³Molecular Signaling and Cell Death Unit, Department for Molecular Biomedical Research, VIB, B-9052 Ghent, Belgium; ⁴Department of Molecular Biology, Ghent University, B-9052 Ghent, Belgium and ⁵Department of Basic Medical Sciences – Anatomy and Embryology group, Faculty of Medicine and Health Sciences, Ghent University, B-9000 Ghent, Belgium

*Corresponding author: M Vinken, Department of Toxicology, Vrije Universiteit Brussel, Laarbeeklaan 103, B-1090 Brussels, Belgium. Tel: +32 2 477 45 87; Fax: +32 2 477 45 82; E-mail: mvinken@vub.ac.be

⁶These authors contributed equally to this work.

⁷ED is a doctoral research fellow of the Fund for Scientific Research Flanders (FWO-Vlaanderen), Belgium.

⁸MV, TV and DVK are postdoctoral research fellows of the Fund for Scientific Research Flanders (FWO-Vlaanderen), Belgium.

⁹VR and LL share equal seniorship.

Keywords: apoptosis; connexin; hemichannel; gap junction; pannexin; intercellular communication

Abbreviations: ATP, adenosine triphosphate; Bcl-2, B-cell lymphoma-2; cAMP, cyclic adenosine monophosphate; cGMP, cyclic guanosine monophosphate; Cx, connexin; CytC, cytochrome C; ER, endoplasmic reticulum; GCV, ganciclovir; GJ, gap junction; GJIC, gap junctional intercellular communication; HC, hemichannel; HSV-tk, *herpes simplex* virus-thymidine kinase; IP₃, inositol trisphosphate; MEK/ERK, mitogen-activated protein kinase/extracellular signal-regulated kinase; NAD⁺, nicotinamide adenine dinucleotide; NO, nitric oxide; OGD, oxygen/glucose deprivation; P₂X₇R, P₂X₇ receptor; p90^{RSK}, p90 ribosomal S6 kinase; Panx, pannexin; pC/EBP β , phosphorylated CCAAT/enhancer-binding protein β ; PM, plasma membrane; ROS, reactive oxygen species; TNF α , tumor necrosis factor α ; TRAIL, TNF-related apoptosis-inducing ligand

Received 07.11.08; revised 04.12.08; accepted 05.12.08; Edited by JP Medema; published online 06.2.09

lacking underlying signaling events and occurring under extreme physico-chemical conditions, including abrupt anoxia, sudden shortage of nutrients, heat or detergents. Recent evidence, however, suggests that necrotic cell death can also be the morphological manifestation of a molecularly regulated event associated with pathologies such as ischemia-reperfusion injury, neurodegeneration and pathogen infection.^{9,10}

In contrast to the wealth of knowledge concerning the extracellular and intracellular signaling cascades that govern (apoptotic) cell death, our understanding of the role of intercellular (cell-to-cell) communication in this process is still in its infancy. The most direct form of intercellular communication proceeds through gap junction (GJ) channels. These channels arise from the head-to-head interaction of two hemichannels (HCs) (connexons) on adjacent cells, which are hexameric channels composed of connexins (Cxs). The Cx protein has a four membrane-spanning topology with two extracellular loops, one cytoplasmic loop and a cytoplasmic N- and C-terminal region (Figure 1).^{11–13} At present, more than twenty Cx species have been cloned from rodents and human and are named according to their molecular weight.¹⁴ Gap junctional intercellular communication (GJIC) is driven by the passive diffusion of small (<1–1.5 kDa) hydrophilic molecules such as glucose, glutamate, glutathione, cyclic adenosine monophosphate (cAMP), adenosine triphosphate (ATP), inositol trisphosphate (IP₃) and ions (e.g., Ca²⁺, K⁺, Na⁺).^{13,15} The biophysical permeation properties of these substances depend on the nature of the Cx species that form the channel and clear differences in channel permeability have been shown for various ions, reporter dyes and signaling

molecules such as cAMP or ATP.^{16–19} The gating of GJ channels is controlled by a number of factors, with as prominent players the transmembrane voltage (over the PM), transjunctional voltage (over the GJ), Ca²⁺ and the phosphorylation status.^{20–22} With the notable exception of Cx26, all Cxs are phosphoproteins that are targeted, particularly at their C-terminal tail, by a broad panel of kinases. HCs reside as closed channels in the PM but open by a process of ‘loop gating’ when their extracellular loops interact to form a functional GJ channel.^{20,23} Before being incorporated into GJs, HCs can also be opened by various signals or conditions, including membrane depolarization,²⁴ a decrease of extracellular Ca²⁺,²⁵ cytoplasmic Ca²⁺ changes,²⁶ mechanical stimulation,²⁷ changes in phosphorylation status,²⁸ changes in redox status,²⁹ reactive oxygen species (ROS),³⁰ nitrosylation of the Cx protein,³¹ ischemia/hypoxia,^{3,32,33} and also certain Cx mutations.^{34–37} Open HCs allow the entry of below 1 to 1.5 kDa substances (e.g., Ca²⁺, Na⁺)³⁸ or the escape of essential metabolites such as nicotinamide adenine dinucleotide (NAD⁺),³⁹ ATP,⁴⁰ glutamate,⁴¹ prostaglandins⁴² and glutathione.⁴³

Another family of GJ channel-forming proteins, the ‘innexins’, was first reported in invertebrates and later renamed ‘pannexins’ (Panxs) after their orthologs were discovered in vertebrates.⁴⁴ Thus far, three Panxs (Panx1, Panx2 and Panx3) have been characterized in rodents and human. Despite the lack of sequence homology between Cxs and Panxs, they share a similar topology and also display cell-specific expression patterns.⁴⁵ They appear to be endowed with several other Cx-like properties such as the ability to form homomeric Panx1 or heteromeric Panx1/Panx2 HCs⁴⁶ and




Figure 1 Molecular architecture of gap junctions. GJs are grouped in plaques at the membrane surface of two apposed cells, and are composed of twelve Cx proteins, organized as two hexameric HCs. The Cx protein is organized as four membrane-spanning domains (TM1–4), two extracellular loops (EL1 and EL2), one cytoplasmic loop (CL), one cytoplasmic amino tail (NT) and one cytoplasmic carboxy tail (CT) (EC, extracellular; IC, intracellular)

functionally competent GJ channels,^{47,48} although some controversy still exists concerning the latter.^{49,50} Furthermore, Panx channels are also permeable to small molecules,^{51–53} release ATP,^{54,55} open in response to mechanical stimulation,⁵⁴ and intracellular Ca^{2+} ,⁵⁶ and are blocked by certain Cx-based GJ blockers.^{46,53}

Numerous physiological processes are driven by regulatory molecules that pass through GJs and HCs, which are therefore considered as major gatekeepers in the control of tissue homeostasis. Many efforts in this respect have been focused on their roles in the regulation of cellular proliferation and differentiation.^{12,15,57} The exploration of Cx- and Panx-related signaling in cell death, however, has only been initiated in recent years, nevertheless reporting some striking observations in this newly arisen research field.^{12,45,58,59} In this review, we will focus on the role of Cxs and their channels as both positive and negative modulators of cell death, mainly apoptosis. We additionally demonstrate the involvement of the recently discovered Panxs in cellular responses to insults.

Connexin-Based Gap Junction Channels and Cell Death

A substantial body of evidence indicates a positive correlation between GJIC and apoptotic activity (Figure 2). Indirect data come from the observation that chemical GJ inhibitors, such as carbenoxolone and 18-beta-glycyrrhetic acid, prevent apoptosis.^{60–63} *Vice versa*, tumor promoters, including peroxisome proliferators and phenobarbital which are known to counteract apoptosis, also inhibit GJIC.^{64,65} Furthermore, exogenous introduction of Cxs in a plethora of experimental models was found to facilitate apoptotic cell

death (Table 1).^{30,63,66–78} Various cell death models have demonstrated the clustering of dying cells, indicating the spread of death signals to neighboring cells through GJs.^{63,79,80} This phenomenon of ‘bystander death’ (the ‘kiss of death’) has gained a great deal of attention for it opens up the possibility to therapeutically limit the wave of secondary injury in the context of stroke or brain trauma,^{3,80–84} and to amplify the potency of cancer treatment. With respect to the latter, the ‘suicide gene/prodrug therapy’ is a well-known model whereby malignant cells are transfected with the *herpes simplex* virus-thymidine kinase (HSV-tk) gene, followed by treatment with the prodrug ganciclovir (GCV). Following phosphorylation to GCV-triphosphate, this cytotoxic compound competitively inhibits the incorporation of endogenous deoxyguanosine triphosphate into the DNA, resulting in the termination of DNA synthesis and the onset of apoptosis.^{85,86} In several tumor cell models, cells that lack the suicide gene and that surround a HSV-tk⁺ cell are also killed by GCV treatment because of diffusion of GCV-triphosphate through GJs connecting those cells.^{85–87} Another form of bystander killing is mediated by the transfer of viral peptides through Cx-related communication. Neijssen *et al.*⁸⁸ discovered that a cell expressing viral proteins and its closest neighbors are killed by cytotoxic T-cells, because the adjacent cells receive the viral peptides through GJs. Thus, GJ immunological coupling could mediate the elimination of uninfected bystander cells or those in the earliest phases of infection.

In-depth studies have revealed that Cx-related communication is modified during the cell death process (Table 2). Particularly the early phases of apoptosis require




Figure 2 Connexin- and pannexin-related signaling in cell death. Cxs can affect the cell death process through a number of mechanisms, involving GJIC (1), HCs (2–5) and Cx proteins as such (6,7). GJ channels can accommodate direct exchange of cell death and cell survival signals between cells (1). HCs may contribute to cell death by four different mechanisms: by the entry of cell death or the loss of cell survival signals (2), through paracrine signaling of death or survival messengers (3) by HC-mediated transmembrane signal transduction (4) or by affecting mitochondrial functioning (5). Cx proteins as such can associate with cell death regulators (6) or influence the expression of these molecules (7). HCs composed of Panxs may act as a permeabilization pore by itself or as a part of the $\text{P}_2\text{X}_7\text{R}$ death complex (8), allowing ATP to leave the cell or bacterial molecules to make their way into the cell. Although solid scientific data are currently not available, both processes might contribute to cell death. The figure is based on references mentioned in the text. It should be noted that many of the first and second messengers depicted are not cell death or survival messengers *per se*, but rather substances that may lead to cell death or survival under specific conditions that are discussed in the text. (ASK1, apoptosis signal-regulating kinase 1; ATP, adenosine triphosphate; cAMP, cyclic adenosine monophosphate; cGMP, cyclic guanosine monophosphate; ERK, extracellular signal-regulated kinase; GCV, ganciclovir; IP_3 , inositol trisphosphate; pBad, phosphorylated Bad; pC/EBP β , phosphorylated CCAAT/enhancer-binding protein β ; ROS, reactive oxygen species)

Table 1 Interfering with connexin gene expression influences cell death

Model ^a	Cell death	Evaluation method	Mechanism	Reference
<i>Nervous tissue – primary cultures and tissues</i>				
Cx43-deficient mouse astrocytes	↓ ^b	Dextran-LY uptake LDH release		24
Heterozygote Cx43 ^{+/-} mouse astrocytes	↑ ^{b,c}	TUNEL assay	Induced caspase-3 activation	103,116
Cx43-deficient mouse astrocytes		Caspase-3 staining		
Cx43-silenced primary rat astrocytes	↑ ^{b,c}	Hoechst staining		160
Cx43-silenced rat optic nerve segments	↓ ^b	PI staining		83
Rat hippocampal organotypic slice cultures	↓ ^b	PI staining		
Cx26 and Cx32-silenced				
Cx43-silenced				
Cx43 KO mice				
Cx32-deficient mouse oligodendrocyte progenitor cells	↑ ^c	TUNEL assay		111
Cx36-deficient primary rat hypothalamic cultures	↓ ^{b,c}	Calcein-AM staining		61
<i>Nervous tissue – cell lines</i>				
Cx43-transfected mouse N2A neuroblastoma cells	↑ ^{b,c}	Calcein-AM/EB staining	Uptake of ROS	30
Cx43-transfected rat C6 glioma cells (C6Cx43)	↑ ^{b,c}	Caspase staining Annexin V assay		
Cx43-silenced C6Cx43 cells	↓ ^{b,c}	Hoechst 33342 staining EB staining		
Cx43-transfected human U251 glioblastoma cells	↑ ^{b,c,d}	TUNEL assay Hoechst staining Annexin V assay	Reduced Bcl-2 expression No effect on Bax-1, Bad-1, Bcl-x _L , Mcl-1 expressions	68,69
Cx43- and Bcl-2-transfected rat C6 glioma cells in co-culture	↑ ^{b,c}	TUNEL assay Hoechst staining Annexin V assay		
Cx43-transfected rat C6 glioma cells	↓ ^{b,c}	Caspase-3, -8 assay TUNEL assay	Reduced caspase-3 activation No effect on caspase-8 activation	106
Transfected rat C6 glioma cells				
Cx43	↓ ^{b,c,d,e}	TUNEL assay Hoechst staining		110
Cx40	↓ ^{b,c,d,e}	Alamar blue assay		
Cx32	↓ ^{b,c,d,e}			
Cx37-transfected mouse N2A neuroblastoma cells	No effect ^c	XTT assay		74
<i>Other tissues – primary cultures and tissues</i>				
Cx43 dominant negative mutant-transfected chick leg bud mesenchymal cells	↑ ^{b,c}	FACS analysis		108
Cx43-deficient mouse embryonic heart tissue	↑ ^c	TUNEL assay		114
Cx43 antisense oligonucleotide-treated rat neonatal ventricular myocytes	↑ ^c	TUNEL assay Hoechst staining		115
Cx32-mutant-transfected <i>Xenopus laevis</i> oocytes	↑ ^d	Morphological		36
Cx26-mutant-transfected <i>Xenopus laevis</i> oocytes	↑ ^d	Morphological		37
Cx26-inactivated mouse inner ear epithelial cells	↑ ^c	TUNEL assay Caspase-3 staining	Induced caspase-3 activation	105
Cx32- and Cx26-deficient mouse liver tissue	↑ ^c	TUNEL assay XTT assay		
Transfected human umbilical vein endothelial cells				74
Cx37	↑ ^c	TUNEL assay	Induced caspase-3 activation	
Cx40	No effect ^c	PI staining		
Cx43	No effect ^c	Caspase-3 assay Annexin V assay Morphological		
Cx45-deficient mouse vascular tissue	↑ ^c	TUNEL assay		109
<i>Other tissues – cell lines</i>				
Cx43-silenced rat MC lung epithelial cells	↓ ^{b,c}	Yo-Pro staining Hoechst 33342 staining PI staining	Uptake of ROS	30
Cx43-transfected human HeLa cervical carcinoma cells	↑ ^{b,c,f}	AO/EB staining		
Cx43-transfected human LNCaP prostate cancer cells	↑ ^{b,c}	TUNEL assay Annexin V assay PI staining Caspase-8 assay FACS analysis	Induced caspase-8 activation	78
Cx43 dominant negative mutant-transfected rat BC31 bladder carcinoma cells	↓ ^c	EB staining PI staining		
Cx43-transfected HeLa cells	↑ ^{b,c,d,f}	Annexin V assay FACS analysis Hoechst 33258 staining Nucleosomal DNA fragmentation PARP fragmentation Caspase-3 assay		137

Table 1 (Continued)

Model ^a	Cell death	Evaluation method	Mechanism	Reference
Cx32-transfected human Caki-2 renal carcinoma cells	↑ ^c	DNA fragmentation Caspase-3 assay	Induced caspase-3 activation Reduced Bcl-2, Bcl-x _L expression No effect on Bax expression	66,67,177
Cx32-transfected human HeLa cervical carcinoma cells	↑ ^{b,c,f}	AO/EB staining		70
Cx32-transfected human A549 lung adenocarcinoma cells	↑ ^{b,c}	PI staining FACS analysis		72
Cx32-transfected baby hamster kidney cells	↑ ^{b,c}	TUNEL assay Caspase-3, -6, -8, -9 staining Annexin V assay TUNEL assay	Induced activation of caspase-3, -6, -9 No effect on caspase-8 activation	77
Cx26-mutant-transfected human embryonic kidney (HEK) 293 cells	↑ ^{c,d}	TUNEL assay		34
Cx26-transfected human PLC/PRF/5 hepatoma cells	↑ ^c	ssDNA staining		71
Cx26-transfected human T24, UM-UC-3, UM-UC-6, UM-UC-14 bladder cancer cells	↑ ^c	TUNEL assay FACS analysis	Reduced Bcl-2 expression	75
Cx26-transfected human LNCaP, PC-3, DU-145 prostate cancer cells	↑ ^c	FACS analysis	Reduced Bcl-2 expression	76
Cx37-transfected rat NRK kidney epithelial cells	No effect ^c	XTT assay		74

AO, acridine orange; EB, ethidium bromide; FACS, fluorescence-activated cell sorting; LDH, lactate dehydrogenase; LY, lucifer yellow; PARP, poly-ADP-ribose polymerase; PI, propidium iodide; ROS, reactive oxygen species; ssDNA, single-stranded DNA; TUNEL, terminal dUTP nick end labelling. ^aA distinction is made between nervous tissues and tissues/cells from other sources. Studies are further categorized according to the Cx species of which the expression is influenced. ^bExogenously induced cell death. ^cApoptosis-related cell death. ^dEffect on cell death is not related to gap junction function. ^eNecrosis-related cell death. ^fApoptotic necrosis (secondary necrosis).

GJIC,^{62,70,89–91} but with progression of cell death, GJIC drastically declines culminating in the absence of cell coupling between apoptotic bodies. The abolishment of GJIC thus coincides with the typical morphological transformations that occur in apoptosis and results from an increased removal of GJ channels from the PM and not from their functional closure.⁷⁰ In a recent study, Theiss *et al.*⁹² exposed primary bovine lens epithelial cells and mouse NIH-3T3 fibroblasts to a number of well-established apoptosis-inducing agents. The authors noticed a decrease in GJIC that was associated with caspase 3-mediated degradation of Cx43. Similarly, activation of apoptosis in primary cultures of chick lenses resulted in caspase 3-mediated truncation of Cx45.6, a reaction that was inhibited by casein kinase 2-mediated phosphorylation of Cx45.6.⁹³ As a matter of fact, phosphorylation is likely to be a major mechanism responsible for GJIC alterations during apoptosis (Table 2). In rabbit lens epithelial cells, H₂O₂ induced caspase 3-dependent apoptosis that was associated with activation of protein kinase C γ , the phosphorylation of Cx43 and Cx50, and the disassembly of GJ plaques abolishing GJIC.^{94,95} Induction of apoptosis in rat liver epithelial WB-F344 cells by the hydrophobic platinum IV complex LA-12 was also linked to suppression of GJIC and the disappearance of Cx clusters from the cell membrane surface. LA-12 thereby triggered rapid Cx43 hyperphosphorylation mediated by the mitogen-activated protein kinase kinase/extracellular signal-regulated kinase (MEK/ERK) pathway.⁹⁶ Increased Cx phosphorylation does not always go hand in hand with a loss of GJIC. Indeed, in the WB-F344 rat liver epithelial cell model, serum deprivation⁹¹ and exposure to the histone deacetylase inhibitor suberoylanilide hydroxamic acid⁹⁷ enhanced GJIC, particularly during the early phases of apoptosis, and simultaneously increased Cx43 phosphorylation. Inversely, inhibition of GJIC in primary

bovine lens epithelial cells and primary human corneal fibroblasts by staurosporine⁹² and TNF α ⁹⁸ was accompanied by a decrease in the Cx43 phosphorylation status. Of note, alterations in Cx phosphorylation are frequently allied with modifications in cellular localization. The drastic decrease in GJIC that is observed during mitosis, for instance, results from the cytoplasmic redistribution of a unique Cx43 phosphoform called 'Cx43m' or 'Cx43P3'.^{99–101} The latter is generated by the cyclin-dependent kinase 1/cyclin B1 complex, which not only controls the G2/M transition of the cell cycle, but also plays a crucial role in the initiation of apoptosis.^{91,99–101} In fact, induction of apoptosis by choline deficiency increased Cx43 immunoreactivity in the cytoplasm of rat liver epithelial cells. Restoration of the Cx43 membrane localization, and consequently of GJIC, as well as enhanced cell survival was brought about by 8-bromo-cAMP,¹⁰² a well-known inducer of Cx43 phosphorylation.²²

Contradictory to the paradigm that GJIC facilitates apoptosis are a number of investigations showing that GJs may impede the occurrence of cell death (Figure 2).^{103–116} In line with this notion are a number of reports describing induction of apoptosis by chemical GJIC inhibitors^{107,108,117–119} as well as by Cx mimetic peptides,¹²⁰ which are peptides identical to a short Cx sequence that act as GJ blockers.¹²¹ Healthy cells may indeed provide their dying neighbors with rescue signals³ or cells in danger may dilute toxic substances toward healthy neighbors through GJs.^{3,103} This event whereby cell survival is promoted has been designated the 'good Samaritan' effect or the 'kiss of life', as opposed to the 'bystander death' phenomenon.⁸⁵ Thus, there seems to be a clear two-way traffic between (early) apoptotic and non-apoptotic cells⁸⁰ and, depending on the status and environmental context of the cell that receives the signal, two opposite biological outcomes can be achieved, namely cell death or cell survival.³

Table 2 Connexin-related modifications during cell death

Model	Cell death induction	Observation	Reference
Rat hippocampal slices	OGD ^a	Induced GJIC Induced Cx36 expression	60
Rat primary hypothalamic cultures	Different modes of NMDA receptor-regulated cell death ^a	GJIC, but no HC involved	61
Rat MC mammary tumor cells	H ₂ O ₂ ^a	HC activation	30
Rat L2 lung epithelial cells	Cigarette smoke extract ^a		
Cx43-transfected mouse N2A neuroblastoma cells			
Cx43- or Cx32-transfected human HeLa cervical carcinoma cells	Cycloheximide ^a Etoposide ^a Taxol ^a <i>Cis</i> -platinum ^a Camptothecin ^a	GJIC present in early cell death phases Reduced GJIC during cell death progression, associated with gap junction internalization, but not with closure of gap junctions HC functioning decreased during the development of apoptosis	70
Rat BC31 bladder carcinoma cells	Spontaneous ^a	GJIC present between apoptotic and non-apoptotic cells No difference in Cx43 expression between apoptotic and non-apoptotic cells	80
Rat hippocampal organotypic slice cultures	Weight-drop injury	Transient potentiation of GJIC after injury	83
Primary quail granulosa cells	Serum deprivation ^a	Induced GJIC	89
Rat WB-F344 liver epithelial cells	Serum deprivation ^a	Reduced Cx43 expression Induced GJIC in early cell death phases, associated with induced Cx43 expression/phosphorylation	91
Primary bovine lens epithelial cells	Staurosporine ^a Etoposide ^a	Reduced GJIC during cell death progression Reduced GJIC	92
Mouse embryonic NIH 3T3 fibroblasts	Puromycin ^a Cycloheximide ^a	Reduced Cx43 expression/phosphorylation	
Rat WB-F344 liver epithelial cells	LA-12 ^a	Reduced GJIC	96
Rat WB-F344 liver epithelial cells (<i>ras</i> -transformed)	Suberoylanilide hydroxamic acid ^a	Induced Cx43 phosphorylation Induced GJIC	97
Human corneal fibroblasts	TNF α ^a	Induced Cx43 expression/phosphorylation Reduced Cx43 translation and phosphorylation No effect on Cx43 transcription	98
Rat WB liver epithelial cells	Choline depletion ^a	Reduced GJIC, associated with cytosolic Cx43 redistribution	102
Human Hep3B liver epithelial cells		No effect on Cx43 expression	
Chick leg bud mesenchymal cells	TGF- β ₃ ^a	Reduced Cx43 expression	108
Xenopus laevis oocytes	Cytochrome C ^a	Induced GJIC	122
Primary rat astrocytes	Antimycin A and iodoacetic acid-induced metabolic inhibition	HC opening Reduced GJIC Dephosphorylation of Cx43	32
Primary mouse astrocytes	Antimycin A and iodoacetic acid-induced metabolic inhibition	HC opening Increased surface Cx43 expression Dephosphorylation/nitrosylation of Cx43	31
Cx26-mutant-transfected human embryonic kidney (HEK) 293 cells	Spontaneous ^a	Cx26-HC opening	34
Cx32-mutant-transfected Xenopus laevis oocytes	Spontaneous	Cx32-HC opening	36
Cx26-mutant-transfected Xenopus laevis oocytes	Spontaneous	Cx26-HC opening	37
Cx43-transfected human HeLa cervical carcinoma cells	Staurosporine ^a	Cx43-HC opening Increased surface Cx43 expression/dephosphorylation	137
Primary human renal proximal tubule cells	ATP-depletion induced by glycolysis inhibition ^b	Cx43-HC activation, associated with disrupted ATP-flux Dephosphorylation of Cx43 as possible mechanism	141
Rat spinal cord	SCI by cutting into segments	Induced Cx43 expression	159
Rat optic nerve segments	OGD	Increased Cx43 expression	160
Human MG63 osteosarcoma cells	TNF α ^a	Reduced Cx43 translation	189
Human osteoblasts	TRAIL ^a	Induced Cx43 transcription Reduced GJIC	
Human MG63 osteosarcoma cells	Serum deprivation ^a	Reduced Cx43 translation, associated with cytosolic Cx43 redistribution	189
Human osteoblasts		Induced Cx43 transcription Increased GJIC	

ATP, adenosine triphosphate; Cx, connexin; HC, hemichannel; GJIC, gap junctional intercellular communication; JNK, c-Jun N-terminal-kinase; LPS, lipopolysaccharide; MKK7D, mitogen-activated protein kinase kinase 7D; NAD, nicotine adenosine dinucleotide; NMDA, N-methyl D-aspartate; OGD, oxygen/glucose deprivation; SCI, spinal cord injury; TGF- β , transforming growth factor β ; TNF α , tumor necrosis factor α ; TRAIL, TNF-related apoptosis-inducing ligand.
^aApoptosis-related cell death. ^bNecrosis-related cell death.

The biochemical nature of the signals that actually mediate these effects, in particular those that propagate the cell death message, are largely unknown. Ca²⁺ has been proposed as a 'master' cell death messenger (Figure 2). The onset of

apoptosis is indeed frequently accompanied by drastic alterations in cytoplasmic Ca²⁺ concentration, and several crucial apoptosis effectors are known to depend on Ca²⁺.^{3,58,63,80,89,122} However, Ca²⁺ diffusion through GJs

may be limited by its interactions with more slowly mobile Ca^{2+} -binding proteins. The Ca^{2+} messenger IP_3 , produced by phospholipase C activation and triggering Ca^{2+} release from the endoplasmic reticulum (ER), is probably a better suited candidate as it can pass through GJs,¹²³ its ER IP_3 -receptor is modulated by Bcl-2^{124,125} and the ensuing Ca^{2+} changes may trigger cytochrome C (CytC) release from mitochondria.¹²⁶ Released CytC on its turn can interact with IP_3 receptors to potentiate ER Ca^{2+} release and to exacerbate mitochondrial Ca^{2+} -driven CytC release in a vicious circle.¹²⁷ Recent work further suggests that the pattern of Ca^{2+} changes – steady or oscillatory changes – may determine the outcome towards either cell death (sustained Ca^{2+} change) or survival (Ca^{2+} oscillations).¹²⁸ The signature of the Ca^{2+} signal is largely influenced by the level of IP_3 and the sensitivity of its receptor.¹²⁹ As a consequence, the modulation of IP_3 receptors by Bcl-2 family proteins¹²⁵ will determine the Ca^{2+} load of mitochondria and will thus be expected to be decisive in initiating apoptosis through mitochondrial CytC release. CytC has too high a molecular weight (12.4 kDa) to pass as a death messenger through GJs or HCs, so IP_3 remains as a very likely candidate to communicate the cell death message. Other proposed candidate ‘killer messengers’ that can pass through GJs include cAMP and cyclic guanosine monophosphate (cGMP).^{58,74} Potential ‘rescue messengers’ on the other hand include the energy molecules glucose and ATP or the free radical scavengers ascorbic acid and reduced glutathione that are all able to flow through GJs and favor cell survival.³ GJ channels display selective permeability towards the proposed messengers and this typically depends on their Cx composition.^{15–19} Seul *et al.*⁷⁴ found that adenoviral delivery of Cx37, but not of the other vascular Cx40 and Cx43, induces apoptotic cell death in human umbilical vein endothelial cells. This effect was cell- and species-specific as Cx37 did not trigger cell death in mouse N2A neuroblastoma cells and rat NRK kidney epithelial cells. These findings demonstrate that conveying a cell death or survival response through GJ channels is a selective and well-coordinated process, involving specific Cxs and biochemical messengers rather than just the movement of common noxious or vital molecules between cells.

Connexin-Based Hemichannels and Cell Death

For a long time, HCs were classified as mere structural precursors of GJs. A large body of recent evidence, however, suggests that they must be considered as functional channels controlled by various extracellular and intracellular signals, thereby providing a unique pathway for transport and signaling. Of note, many of the signals that trigger HC opening, or responses attributed to HCs, are also involved in the signaling cascades leading to cell death,¹¹ including cytoplasmic Ca^{2+} changes and oxidation or nitrosylation reactions. Being a pore permeable to quite a large range of substances, HC opening may furthermore help in setting off cell death by causing membrane depolarization, the collapse of ionic gradients, loss of small metabolites and elevation of cytoplasmic Ca^{2+} .¹³⁰

As holds for GJs, cell death may have a profound influence on the status of the HCs (Table 2). This has been well exemplified in the case of ischemia-related cell injury where the involvement of HCs as ‘pathogenic pores’ has been demonstrated.^{3,33} Indeed, metabolic inhibition of astrocytes was found to accelerate cell death, which was directly associated with Cx43 HC opening. The molecular mechanism by which it activates HCs is, however, not entirely known, but may involve dephosphorylation and/or oxidation.^{31,32} With respect to the latter, Retamal *et al.*³¹ showed that nitrosylation of intracellular Cx43 cysteine residues by nitric oxide (NO), a protein modification likely to occur under oxidative stress, could be at the heart of HC opening. Along the same line, NO was found to mediate Cx43 HC activation in astrocytes under proinflammatory conditions.¹³¹ The importance of oxidative stress as a key trigger is underscored by the observation that cigarette smoke extract and H_2O_2 may open HCs by depolarizing the cells, thereby facilitating ROS entry into the cell.³⁰ HC function may also be drastically affected by genetic factors. Mutations in Cx32, Cx30 and Cx26 genes result in hereditary peripheral neuropathy, hidrotic ectodermal dysplasia, and nonsyndromic hereditary hearing impairments and skin disease, respectively. In all cases, the phenotype is caused by abnormal HC opening which could adversely affect cell viability.^{34–37,132,133} Gomez-Hernandez *et al.*³⁵ revealed that HC dysregulation in inherited peripheral neuropathy, caused by a Cx32 mutation, is due to alterations in a Ca^{2+} -binding site that interacts with extracellular Ca^{2+} to keep the HCs closed under physiological conditions. A similar mechanism could account for aberrant HC gating and cell death caused by a Cx26 mutation located right next to this putative Ca^{2+} -binding site, a mutation that is known to lead to nonsyndromic hereditary deafness.³⁴

Overall, there are at least four possible mechanisms by which HCs could contribute to cell death or survival (Figure 2). First, the bidirectionally permeable HC pathway may facilitate the uptake of toxic or survival-enhancing substances, and may contribute to the loss of essential metabolites.^{30,38,59,70,134–137} As well-documented ATP release channels, HCs may lower the intracellular ATP content and play as such a decisive role in the balance between cell death through necrosis or apoptosis.^{63,70,138} ATP is also necessary for the phosphorylation of Cxs which, on its turn, influences HC gating.^{28,32,139–141} In Shigella infection, hemichannel ATP release may promote bacterial invasion and dissemination.¹⁴² Another compound released through HCs is glutathione.⁴³ The loss of intracellular glutathione may compromise the antioxidant defense at the cells’ inside but its consequent accumulation outside the cells may as well act beneficially as it can intercept in a more direct manner invading free radicals threatening the cell.⁴³ Another potentially beneficial role of Cx43 HCs is suggested from experiments in astrocytes that demonstrated increased open probability of Cx43 HCs and glucose uptake in response to exposure to proinflammatory cytokines.¹³¹

A second level where HCs can influence the balance between cell death and survival involves a paracrine communication component. The release of metabolites such as ATP, glutamate or prostaglandins through HCs combined with their extracellular diffusion and action (directly or through break-

down products like adenosine)^{143,144} on corresponding receptors on remotely located cells may exert toxic as well as survival effects.^{145–151} Ligand-receptor binding of these messengers is often accompanied by a rise in cytoplasmic Ca^{2+} ,^{126,145,152–154} that on its turn may trigger a new cycle of HC responses or is even influenced by HC activity.^{26,35,38,126,152} In a model of cochlear explant cultures, the local infliction of a cell-damaging stimulus caused a spatiotemporal spread of cell death in hair cells that was related to ERK1/2 activation, dependent on the associated intercellular Ca^{2+} wave and ATP release, and inhibited by carbenoxolone.¹⁵⁵ Once released, ATP is rapidly degraded to adenosine which displays both neuroprotective and -toxic properties.^{143,144} Combined *in vitro* and *in vivo* studies have demonstrated that ischemic pre-conditioning in the brain promotes ATP efflux through HCs and thereby protects against a second exposure to ischemic conditions through adenosine.¹⁵⁶ Another messenger released through HCs is glutamate, a well-documented paracrine messenger of cell death in the brain. The presence of this substance, released by HCs, in conditioned media from microglia and macrophages treated with proinflammatory molecules resulted in marked neuronal cell death when neurons were exposed to it.^{136,157} Conditioned media from activated microglia also affected astrocytes as it reduced their GJIC and induced HC opening.¹³¹ The latter can have a drastic influence on neuronal survival as well, as the capacity of 'spatial buffering', which is provided by coupling of astrocytes through GJ channels and is essential for the control of the composition of the extracellular environment, decreases.^{3,158} This is observed in an *ex vivo* model for spinal cord injury where Cx43 channels appeared to play a dual role; HCs seemed to be involved in cell swelling and the spread of neurotoxins early after injury, whereas GJs were required for spatial buffering between astrocytes and long-term survival of neurons.¹⁵⁹ A paracrine signaling component may not only be of importance for the expansion of cell death or survival but could also assist in the recruitment of both innate and adaptive components of the inflammatory response system.¹⁶⁰ Certain types of dead cells can trigger inflammation by release of danger signals or 'damage-associated molecular patterns' that are normally located inside the cells.^{161,162} At a more speculative level, it is conceivable that these endogenous danger signals (e.g., uric acid, 168 Da) may be shared through GJs with neighboring cells and released through HCs.

A third cell survival/cell death regulatory platform includes HCs as a transmembrane signal transduction pathway. The prototypic example here is the involvement of HCs in the anti-apoptotic actions of alendronate, a drug used to counteract osteoporosis. Alendronate induces the opening of Cx43 HCs in dexamethasone-treated osteoblasts and osteocytes, which activates Src kinase and ERK. Subsequent phosphorylation by the p90 ribosomal S6 kinase (p90^{RSK}) leads to an inactivation of the cytoplasmic pro-apoptotic Bcl-2 protein Bad and binding of pro-caspases by the phosphorylated CCAAT/enhancer-binding protein β (pC/EBP β), both favoring survival of osteoblasts and osteocytes.¹⁶³

A fourth possible link between HCs and cell survival/cell death relates to modifications in the subcellular localization of HCs. Upon ischemic pre-conditioning, a form of cardioprotec-

tion, Cx43 translocates from the PM to the inner mitochondrial membrane through a heat-shock protein 90-dependent pathway.^{164–166} The functional relevance of mitochondrial Cx43, however, is not clear. It has been postulated that Cx43 is part of a multiprotein complex that somehow controls mitochondrial homeostasis.¹¹⁷ Mitochondrial Cx43 could also form HCs that serve as a conduit for ion fluxes,^{59,165–167} a function reminiscent of the apoptosis-regulating Bcl-2 family proteins.¹¹⁷ Mitochondrial Cx43 was recently found to be a major regulator of cardiomyocyte apoptosis.¹¹⁷

Non-Channel Aspects of Connexins in Relation to Cell Death

Several studies have shown that Cx proteins as such can influence tissue homeostasis independent of their channel-forming capabilities. In most cases, the evidence is based on experiments in which Cx expression was artificially modified (Table 1). Cx overexpression was found to trigger cell cycle arrest^{168–173} and cellular differentiation¹⁷⁴ without influencing Cx channel activity in any way. In a similar manner, transfection of C6 rat glioma cells with Cx43, Cx40 or Cx32 genes resulted in an increased resistance to exogenously induced cellular injury, independently of GJIC.¹¹⁰ Forced expression of Cx43 in U251 human glioblastoma cells, on the other hand, did not increase GJ coupling but enhanced the apoptotic response to chemotherapeutics.⁶⁹ Furthermore, research in a model of apoptosis-primed primary quail granulosa cells showed that Cx43 expression was inversely related to both the apoptotic index and the level of GJIC, suggesting distinct roles for GJIC and Cx43 in cell death and cell survival, respectively.⁸⁹

The mechanisms that underlie non-channel Cx protein contributions to cell death remain to be elucidated (Figure 2). It has been suggested that Cxs participate in cell death pathways through direct interaction with apoptotic factors. The cytoplasmic co-localization of Cx26 and Cx43 with the Bcl-2 proteins Bak, Bcl-xL and Bax, in human breast cancer cells and human colorectal cancer cells, respectively, could be in favor of this idea.^{175,176} More recently, Cx43 was reported to directly interact with apoptosis signal-regulating kinase 1 (ASK1).¹⁰⁶ Cxs may also be involved in the control of cell death-related gene expression. A number of papers have reported changes in the expression of individual subsets of apoptotic factors while interfering with Cx expression.^{67–69,72,75,76,168,177} Microarray analysis of heart tissue from Cx43 knockout mice showed altered expression of a wide spectrum of apoptotic genes, including Bok, Bax, Bid, Diablo, caspase 6 and caspase 9.¹¹⁴ Iacobas *et al.*¹⁷⁸ performed a similar in-depth study, by analyzing the transcriptome of Cx43-null astrocytes in mice. Interestingly, they showed that both pro-apoptotic (e.g., Map4k) and anti-apoptotic (e.g., Bcl-xL) genes are affected, a finding that might explain the often contradictory reports, associating both induction and inhibition of apoptotic cell death with the presence of Cxs. The exact nature of the link between Cxs and apoptosis-related gene expression is still a matter of debate. An attractive hypothesis is that Cxs are directly involved in transcriptional control. In this respect, 'Cx response elements' have been proposed to modulate gene expression in rat osteocytes.^{179,180} An active role for Cxs in the

regulation of gene expression is further supported by the observation that (transfected) Cxs, or at least specific parts of these molecules (i.e. the C-terminal region), can reside in the nuclear compartment.^{170,181,182} The latter may, however, also be a result of the well-known interaction of Cxs with transcriptional regulators and/or mediators of crucial signaling pathways. A broad range of Cx-binding partners have actually been characterized, some of which are acknowledged regulators of gene expression.^{183,184} Cx43, for instance, interacts with β -catenin, a key player in Wnt signaling.¹⁸⁵ In the cell nucleus, β -catenin forms a complex with the T-cell factor that facilitates the transcription of a number of target genes that are relevant to apoptosis, including survivin,¹⁸⁶ Bcl-xL¹⁸⁷ and Bcl-2.¹⁸⁸ Cx43 expression itself is also controlled by Wnt signaling.¹⁸⁵ In fact, TNF α and TRAIL not only increase Cx43 degradation but also augment the Cx43 mRNA content in human osteosarcoma cells. The enhanced Cx43 gene transcription is thought to represent a reflexive response to apoptosis and is likely to be mediated by β -catenin.¹⁸⁹

Pannexins and Cell Death

The structural and functional similarities between Cxs and Panxs raise the possibility that Panxs might be implicated in cell responses to several pathological insults as well, for example ischemic neuronal cell death. Exposure of pyramidal neurons from cortex or hippocampus to oxygen and glucose deprivation (OGD), a condition that mimics some aspects of ischemia, triggered opening of a large conductance and the bidirectional transfer of small fluorescent tracer molecules.⁵² Long OGD exposures resulted in irreversible current activation, neuronal swelling and membrane breakdown. Based on the absence of Cxs but presence of Panx1 in these neurons, the large single channel conductance and the effect of the (non-specific) blockers carbenoxolone and the HC-inhibiting La³⁺ ions, the authors concluded that these responses are likely to be mediated by Panx1 HCs. The mechanism by which OGD leads to Panx HC opening is not clear but may involve an increase in cytoplasmic Ca²⁺, which is known to occur in hippocampal neurons after ischemic injury and to activate Panx HCs.^{56,152,153} The proposed mechanism of dephosphorylation-induced opening of Cx43 HCs in metabolically inhibited astrocytes could also account for Panx HC opening as multiple phosphorylation sites have been predicted in both the cytoplasmic loop and the C-terminal tail of the Panx protein.⁴⁵ There is, however, no clear evidence yet of functional modulation of Panx HCs by this covalent modification.

Similar to Cxs, Panx1 was found to be critical for inflammatory responses in lipopolysaccharide-stimulated macrophages pulsed with ATP⁵³ or exposed to the pore-forming toxins nigericin and maitotoxin,¹⁹⁰ through HC-dependent or HC-independent mechanisms respectively. With respect to the former, studies further demonstrated that Panx1 HC opening can be induced by ATP stimulation of P₂X₇ receptors (P₂X₇R) and thereby provides an entry pathway for bacterial inflammatory proteins to make their way to the cytosol.¹⁹¹ P₂X receptors are a family of ionotropic receptors that open non-selective cation channels when activated by

pyrimidine nucleotides. Interestingly, during prolonged agonist exposure, certain of these receptors like the P₂X₇R lead to the opening of a non-selective pore capable of passing large hydrophilic molecules, leading to cell death by apoptosis or necrosis.^{147,148,192–194} In fact, many of the characteristics of the P₂X₇R-activated pore seem to match those of HCs, including blockade by prototypical GJ channel blockers.¹⁹⁵ A number of studies have demonstrated that the large permeabilization pore may correspond to Panx1 HCs being recruited by prolonged P₂X₇R activation.^{53,190,193,196} Binding of the tyrosine kinase Src to the P₂X₇R seems to be a crucial step in this process.¹⁹⁶ In support of this finding is the observation that prolonged stimulation of the P₂X₇R by ATP or an agonist induced cell death in oocytes and human astrocytoma cells, a process that was affected by carbenoxolone. Thus, Panx1 appears to be the molecular substrate for the permeabilization pore or 'death receptor channel' recruited into the P₂X₇R-signaling complex.¹⁹³ Panx1 HCs can also be activated by ATP through the metabotropic P₂Y₁ and P₂Y₂ receptors, presumably through increases in cytoplasmic Ca²⁺.⁵⁶ This, however, is not accompanied by subsequent membrane permeabilization and is therefore thought to serve more physiological functions, such as the communication of Ca²⁺ signals between cells,⁵⁵ a communication pathway that may be further supported by Panx1 GJs connecting the cells.^{48,55,56} In addition, a recent study suggests that Panx1 may also be present in the ER, forming Ca²⁺-permeable channels and representing one of the mechanisms responsible for ER Ca²⁺ leak.⁴⁸ The latter may on its turn trigger apoptotic cell death by facilitating Ca²⁺-induced opening of the mitochondrial permeability transition pore, which has been implicated in the release of CytC from mitochondria.^{126,197}

In contrast to Panx1, a neuroprotective role was proposed for Panx2. *De novo* and transient expression of Panx2 in astrocytes was observed in the brain of adult rats following ischemia-reperfusion and in co-cultures of hippocampal neurons and astrocytes. The authors hypothesized that Panx2 mediates its neuroprotective effect through the formation of HCs allowing the release of signaling molecules devoted to influence the cellular metabolism and the redox status of the surrounding environment.¹⁹⁸

Conclusions and Perspectives

The maintenance of the homeostatic balance is controlled by the global interplay between three major communicative networks that make use of extracellular, intracellular and intercellular signaling pathways. Direct intercellular communication is mainly mediated by GJ channels composed of Cxs. The key roles of GJs in the control of cellular proliferation and differentiation have been demonstrated at numerous occasions over the last few decades.^{12,15,57} Their participation as mediators of cell death communication, particularly of apoptotic cell death, has only gained interest in recent years but has already brought up the important insight that not only GJs but also their structural precursors, the HCs and the Cx proteins as such, influence the cell death process. Strikingly, Cx-related communication has been found to both facilitate or counteract the cell death process. Such opposed functions may, to some extent, relate to differences in the model

systems used, including the mode of cell death induction. Different cell death activators may indeed recruit distinct signaling cascades and messengers that may or may not be able to pass or activate Cx channels. An additional factor that may complicate the situation is the fact that the two channel types, although composed of the same protein, are differentially regulated. Under normal conditions, GJs are open and HCs are closed and two recent papers indicate that these two channels can be differentially influenced by the same stimulus,^{28,131} for example by arachidonic acid which inhibits GJs and potentiates HC responses. The decision to stay alive or die may further depend on the condition of the cell receiving the message carried through Cx channels. Thus, if the concerned cell is concomitantly exposed to stress stimuli, then Cx-related communication may switch from a survival program to a death mode. Furthermore, the contributions of Cxs to cell death communication may be adjoined by the Panx channel family. The specifics and possible differences in the regulation of Cx *versus* Panx channels are currently largely unknown and a major challenge for future work is to elucidate the exact contribution of GJs and HCs composed of either Cxs or Panxs, to cell death processes of both the apoptotic or non-apoptotic type. An equally important challenge concerns the development of specific blockers that affect HCs or GJs only. Comparisons between Cx-expressing cells and cells treated with protocols to silence the concerned gene or obtained from knockout animals are necessary to unequivocally determine the involvement of Cx HCs.¹⁹⁵ Knockout animal models can, however, not be used to establish the involvement of Cx HCs *in vivo* because both GJs and HCs are suppressed in that case. The quest is still open but progress is, as expected from channels composed of the same building blocks, not in immediate sight. The molecular principles to distinguish GJs from HCs are still hazy. For instance, Cx mimetic peptides composed of a short stretch from one of the extracellular Cx loops are actually expected to open the HCs by a process similar to 'loop gating' but these substances turn out to inhibit the responses attributed to HCs. Clearly, further work is needed to bring progress in this complex but tantalizing field: two distinct channels formed by a single protein that in addition (and beyond its channel function) has intrinsic signaling functions is not a simple case at all! In summary, Cxs and Panxs as well as their corresponding channels represent multilevel platforms for cell death-related signaling. It is expected that, upon the introduction of appropriate experimental tools, more insight will be gained into the exact molecular mechanisms that underlie Cx- and Panx-inherent communication in (apoptotic) cell death.

Acknowledgements. This work is supported by the Fund for Scientific Research Flanders (FWO-Vlaanderen), Belgium (grant nos. G.0354.07 and G.0140.08 and G.0134.09 to LL) and the Interuniversity Attraction Poles Program (Belgian Science Policy, project P6/31) and the Research Council of the Vrije Universiteit Brussel (OZR-VUB). Special thanks go to the European FP6 projects carcinoGENOMICS and LIINTOP.

1. Elmore S. Apoptosis: a review of programmed cell death. *Toxicol Pathol* 2007; **35**: 495–516.
2. Ferrer I. Apoptosis: future targets for neuroprotective strategies. *Cerebrovasc Dis* 2006; **21**: 9–20.

3. Contreras JE, Sanchez HA, Veliz LP, Bukauskas FF, Bennett MV, Saez JC. Role of connexin-based gap junction channels and hemichannels in ischemia-induced cell death in nervous tissue. *Brain Res Brain Res Rev* 2004; **47**: 290–303.
4. Ozoren N, El-Deiry WS. Cell surface Death Receptor signaling in normal and cancer cells. *Semin Cancer Biol* 2003; **13**: 135–147.
5. Blatt NB, Glick GD. Signaling pathways and effector mechanisms pre-programmed cell death. *Bioorg Med Chem* 2001; **9**: 1371–1384.
6. Ashe PC, Berry MD. Apoptotic signaling cascades. *Prog Neuropsychopharmacol Biol Psychiatry* 2003; **27**: 199–214.
7. Earnshaw WC, Martins LM, Kaufmann SH. Mammalian caspases: structure, activation, substrates, and functions during apoptosis. *Annu Rev Biochem* 1999; **68**: 383–424.
8. Wang X. The expanding role of mitochondria in apoptosis. *Genes Dev* 2001; **15**: 2922–2933.
9. Festjens N, Vanden Berghe T, Vandenabeele P. Necrosis, a well-orchestrated form of cell demise: signalling cascades, important mediators and concomitant immune response. *Biochim Biophys Acta* 2006; **1757**: 1371–1387.
10. Vanlangenakker N, Berghe TV, Krysko DV, Festjens N, Vandenabeele P. Molecular mechanisms and pathophysiology of necrotic cell death. *Curr Mol Med* 2008; **8**: 207–220.
11. Evans WH, De Vuyst E, Leybaert L. The gap junction cellular internet: connexin hemichannels enter the signalling limelight. *Biochem J* 2006; **397**: 1–14.
12. Vinken M, Vanhaecke T, Papeleu P, Snykers S, Henkens T, Rogiers V. Connexins and their channels in cell growth and cell death. *Cell Signal* 2006; **18**: 592–600.
13. Saez JC, Berthoud VM, Branes MC, Martinez AD, Beyer EC. Plasma membrane channels formed by connexins: their regulation and functions. *Physiol Rev* 2003; **83**: 1359–1400.
14. Sohl G, Willecke K. An update on connexin genes and their nomenclature in mouse and man. *Cell Commun Adhes* 2003; **10**: 173–180.
15. Alexander DB, Goldberg GS. Transfer of biologically important molecules between cells through gap junction channels. *Curr Med Chem* 2003; **10**: 2045–2058.
16. Elfgang C, Eckert R, Lichtenberg-Frate H, Butterweck A, Traub O, Klein RA *et al*. Specific permeability and selective formation of gap junction channels in connexin-transfected HeLa cells. *J Cell Biol* 1995; **129**: 805–817.
17. Bukauskas FF, Elfgang C, Willecke K, Weingart R. Heterotypic gap junction channels (connexin26-connexin32) violate the paradigm of unitary conductance. *PLoS Arch* 1995; **429**: 870–872.
18. Kanaporis G, Mese G, Valiuniene L, White TW, Brink PR, Valiunas V. Gap junction channels exhibit connexin-specific permeability to cyclic nucleotides. *J Gen Physiol* 2008; **131**: 293–305.
19. Goldberg GS, Moreno AP, Lampe PD. Gap junctions between cells expressing connexin 43 or 32 show inverse permselectivity to adenosine and ATP. *J Biol Chem* 2002; **277**: 36725–36730.
20. Bukauskas FF, Verselis VK. Gap junction channel gating. *Biochim Biophys Acta* 2004; **1662**: 42–60.
21. Peracchia C. Chemical gating of gap junction channels; roles of calcium, pH and calmodulin. *Biochim Biophys Acta* 2004; **1662**: 61–80.
22. Lampe PD, Lau AF. The effects of connexin phosphorylation on gap junctional communication. *Int J Biochem Cell Biol* 2004; **36**: 1171–1186.
23. Bao X, Chen Y, Reuss L, Altenberg GA. Functional expression in *Xenopus* oocytes of gap-junctional hemichannels formed by a cysteine-less connexin 43. *J Biol Chem* 2004; **279**: 9689–9692.
24. Contreras JE, Saez JC, Bukauskas FF, Bennett MV. Functioning of cx43 hemichannels demonstrated by single channel properties. *Cell Commun Adhes* 2003; **10**: 245–249.
25. Thimm J, Mechler A, Lin H, Rhee S, Lal R. Calcium-dependent open/closed conformations and interfacial energy maps of reconstituted hemichannels. *J Biol Chem* 2005; **280**: 10646–10654.
26. De Vuyst E, Decrock E, Cabooter L, DUBYAK GR, Naus CC, Evans WH *et al*. Intracellular calcium changes trigger connexin 32 hemichannel opening. *EMBO J* 2006; **25**: 34–44.
27. Gomes P, Srinivas SP, Van Driessche W, Vereecke J, Himpens B. ATP release through connexin hemichannels in corneal endothelial cells. *Invest Ophthalmol Vis Sci* 2005; **46**: 1208–1218.
28. De Vuyst E, Decrock E, De Bock M, Yamasaki H, Naus CC, Evans WH *et al*. Connexin hemichannels and gap junction channels are differentially influenced by lipopolysaccharide and basic fibroblast growth factor. *Mol Biol Cell* 2007; **18**: 34–46.
29. Retamal MA, Schalper KA, Shoji KF, Bennett MV, Saez JC. Opening of connexin 43 hemichannels is increased by lowering intracellular redox potential. *Proc Natl Acad Sci USA* 2007; **104**: 8322–8327.
30. Ramachandran S, Xie LH, John SA, Subramaniam S, Lal R. A novel role for connexin hemichannel in oxidative stress and smoking-induced cell injury. *PLoS ONE* 2007; **2**: e712.
31. Retamal MA, Cortes CJ, Reuss L, Bennett MV, Saez JC. S-nitrosylation and permeation through connexin 43 hemichannels in astrocytes: induction by oxidant stress and reversal by reducing agents. *Proc Natl Acad Sci USA* 2006; **103**: 4475–4480.
32. Contreras JE, Sanchez HA, Eugenin EA, Speidel D, Theis M, Willecke K *et al*. Metabolic inhibition induces opening of unapposed connexin 43 gap junction hemichannels and reduces gap junctional communication in cortical astrocytes in culture. *Proc Natl Acad Sci USA* 2002; **99**: 495–500.

33. Retamal MA, Schalper KA, Shoji KF, Orellana JA, Bennett MV, Saez JC. Possible involvement of different connexin43 domains in plasma membrane permeabilization induced by ischemia-reperfusion. *J Membr Biol* 2007; **218**: 49–63.
34. Stong BC, Chang Q, Ahmad S, Lin X. A novel mechanism for connexin 26 mutation linked deafness: cell death caused by leaky gap junction hemichannels. *Laryngoscope* 2006; **116**: 2205–2210.
35. Gomez-Hernandez JM, de Miguel M, Larrosa B, Gonzalez D, Barrio LC. Molecular basis of calcium regulation in connexin-32 hemichannels. *Proc Natl Acad Sci USA* 2003; **100**: 16030–16035.
36. Liang GS, de Miguel M, Gomez-Hernandez JM, Glass JD, Scherer SS, Mintz M *et al*. Severe neuropathy with leaky connexin32 hemichannels. *Ann Neurol* 2005; **57**: 749–754.
37. Gerido DA, DeRosa AM, Richard G, White TW. Aberrant hemichannel properties of Cx26 mutations causing skin disease and deafness. *Am J Physiol Cell Physiol* 2007; **293**: C337–C345.
38. Li F, Sugishita K, Su Z, Ueda I, Barry WH. Activation of connexin-43 hemichannels can elevate [Ca²⁺] and [Na⁺] in rabbit ventricular myocytes during metabolic inhibition. *J Mol Cell Cardiol* 2001; **33**: 2145–2155.
39. Bruzzone S, Guida L, Zocchi E, Franco L, De Flora A. Connexin 43 hemichannels mediate Ca²⁺-regulated transmembrane NAD⁺ fluxes in intact cells. *FASEB J* 2001; **15**: 10–12.
40. Kang J, Kang N, Lovatt D, Torres A, Zhao Z, Lin J *et al*. Connexin 43 hemichannels are permeable to ATP. *J Neurosci* 2008; **28**: 4702–4711.
41. Ye ZC, Wyeth MS, Baltan-Tekkok S, Ransom BR. Functional hemichannels in astrocytes: a novel mechanism of glutamate release. *J Neurosci* 2003; **23**: 3588–3596.
42. Cherian PP, Siller-Jackson AJ, Gu S, Wang X, Bonewald LF, Sprague E *et al*. Mechanical strain opens connexin 43 hemichannels in osteocytes: a novel mechanism for the release of prostaglandin. *Mol Biol Cell* 2005; **16**: 3100–3106.
43. Rana S, Dringen R. Gap junction hemichannel-mediated release of glutathione from cultured rat astrocytes. *Neurosci Lett* 2007; **415**: 45–48.
44. Baranova A, Ivanov D, Petrushin N, Pestova A, Skoblov M, Kelmanson I *et al*. The mammalian pannexin family is homologous to the invertebrate innexin gap junction proteins. *Genomics* 2004; **83**: 706–716.
45. Shestopalov VI, Panchin Y. Pannexins and gap junction protein diversity. *Cell Mol Life Sci* 2008; **65**: 376–394.
46. Bruzzone R, Barbe MT, Jakob NJ, Monyer H. Pharmacological properties of homomeric and heteromeric pannexin hemichannels expressed in *Xenopus* oocytes. *J Neurochem* 2005; **92**: 1033–1043.
47. Bruzzone R, Hormuzdi SG, Barbe MT, Herb A, Monyer H. Pannexins, a family of gap junction proteins expressed in brain. *Proc Natl Acad Sci USA* 2003; **100**: 13644–13649.
48. Vanden Abeele F, Bidaux G, Gordienko D, Beck B, Panchin YV, Baranova AV *et al*. Functional implications of calcium permeability of the channel formed by pannexin 1. *J Cell Biol* 2006; **174**: 535–546.
49. Huang Y, Grinspan JB, Abrams CK, Scherer SS. Pannexin1 is expressed by neurons and glia but does not form functional gap junctions. *Glia* 2007; **55**: 46–56.
50. Penuela S, Bhalla R, Gong XQ, Cowan KN, Celetti SJ, Cowan BJ *et al*. Pannexin 1 and pannexin 3 are glycoproteins that exhibit many distinct characteristics from the connexin family of gap junction proteins. *J Cell Sci* 2007; **120**: 3772–3783.
51. Locovei S, Bao L, Dahl G. Pannexin 1 in erythrocytes: function without a gap. *Proc Natl Acad Sci USA* 2006; **103**: 7655–7659.
52. Thompson RJ, Zhou N, MacVicar BA. Ischemia opens neuronal gap junction hemichannels. *Science* 2006; **312**: 924–927.
53. Pelegrin P, Surprenant A. Pannexin-1 mediates large pore formation and interleukin-1 β release by the ATP-gated P2X₇ receptor. *EMBO J* 2006; **25**: 5071–5082.
54. Bao L, Locovei S, Dahl G. Pannexin membrane channels are mechanosensitive conduits for ATP. *FEBS Lett* 2004; **572**: 65–68.
55. Huang YJ, Maruyama Y, Dvoryanchikov G, Pereira E, Chaudhari N, Roper SD. The role of pannexin 1 hemichannels in ATP release and cell-cell communication in mouse taste buds. *Proc Natl Acad Sci USA* 2007; **104**: 6436–6441.
56. Locovei S, Wang J, Dahl G. Activation of pannexin 1 channels by ATP through P2Y receptors and by cytoplasmic calcium. *FEBS Lett* 2006; **580**: 239–244.
57. Bruzzone R, Dermietzel R. Structure and function of gap junctions in the developing brain. *Cell Tissue Res* 2006; **326**: 239–248.
58. Krysko DV, Leybaert L, Vandenabeele P, D'Herde K. Gap junctions and the propagation of cell survival and cell death signals. *Apoptosis* 2005; **10**: 459–469.
59. Rodriguez-Sinovas A, Cabestrero A, Lopez D, Torre I, Morente M, Abellan A *et al*. The modulatory effects of connexin 43 on cell death/survival beyond cell coupling. *Prog Biophys Mol Biol* 2007; **94**: 219–232.
60. de Pina-Benabou MH, Szostak V, Kyrozis A, Rempe D, Uziel D, Urban-Maldonado M *et al*. Blockade of gap junctions *in vivo* provides neuroprotection after perinatal global ischemia. *Stroke* 2005; **36**: 2232–2237.
61. de Rivero Vaccari JC, Corriveau RA, Belousov AB. Gap junctions are required for NMDA receptor dependent cell death in developing neurons. *J Neurophysiol* 2007; **98**: 2878–2886.
62. Frank DK, Szymkowiak B, Josifovska-Chopra O, Nakashima T, Kinnally KW. Single-cell microinjection of cytochrome c can result in gap junction-mediated apoptotic cell death of bystander cells in head and neck cancer. *Head Neck* 2005; **27**: 794–800.
63. Decrock E, De Vuyst E, Vinken M, Van Moorhem M, Vranckx K, Wang N *et al*. Connexin 43 hemichannels contribute to the propagation of apoptotic cell death in a rat C6 glioma cell model. *Cell Death Differ* 2009; **16**: 151–163.
64. Elcock FJ, Chipman JK, Roberts RA. The rodent nongenotoxic hepatocarcinogen and peroxisome proliferator nafenopin inhibits intercellular communication in rat but not guinea-pig hepatocytes, perturbing S-phase but not apoptosis. *Arch Toxicol* 1998; **72**: 439–444.
65. Kolaja KL, Engelken DT, Klaassen CD. Inhibition of gap-junctional-intercellular communication in intact rat liver by nongenotoxic hepatocarcinogens. *Toxicology* 2000; **146**: 15–22.
66. Fujimoto E, Sato H, Shirai S, Nagashima Y, Fukumoto K, Hagiwara H *et al*. Connexin32 as a tumor suppressor gene in a metastatic renal cell carcinoma cell line. *Oncogene* 2005; **24**: 3684–3690.
67. Fujimoto E, Sato H, Nagashima Y, Negishi E, Shirai S, Fukumoto K *et al*. A Src family inhibitor (PP1) potentiates tumor-suppressive effect of connexin 32 gene in renal cancer cells. *Life Sci* 2005; **76**: 2711–2720.
68. Huang R, Liu YG, Lin Y, Fan Y, Boynton A, Yang D *et al*. Enhanced apoptosis under low serum conditions in human glioblastoma cells by connexin 43 (Cx43). *Mol Carcinog* 2001; **32**: 128–138.
69. Huang RP, Hossain MZ, Huang R, Gano J, Fan Y, Boynton AL. Connexin 43 (cx43) enhances chemotherapy-induced apoptosis in human glioblastoma cells. *Int J Cancer* 2001; **92**: 130–138.
70. Kalvelyte A, Imbrasaitė A, Bukauskiene A, Verselis VK, Bukauskas FF. Connexins and apoptotic transformation. *Biochem Pharmacol* 2003; **66**: 1661–1672.
71. Muramatsu A, Iwai M, Morikawa T, Tanaka S, Mori T, Harada Y *et al*. Influence of transection with connexin 26 gene on malignant potential of human hepatoma cells. *Carcinogenesis* 2002; **23**: 351–358.
72. Sato H, Fukumoto K, Hada S, Hagiwara H, Fujimoto E, Negishi E *et al*. Enhancing effect of connexin 32 gene on vinorelbine-induced cytotoxicity in A549 lung adenocarcinoma cells. *Cancer Chemother Pharmacol* 2007; **60**: 449–457.
73. Hosoya S, Lu Z, Ozaki Y, Takeuchi M, Sato T. Cytological analysis of the mother cell death process during sporulation in *Bacillus subtilis*. *J Bacteriol* 2007; **189**: 2561–2565.
74. Seul KH, Kang KY, Lee KS, Kim SH, Beyer EC. Adenoviral delivery of human connexin37 induces endothelial cell death through apoptosis. *Biochem Biophys Res Commun* 2004; **319**: 1144–1151.
75. Tanaka M, Grossman HB. Connexin 26 gene therapy of human bladder cancer: induction of growth suppression, apoptosis, and synergy with Cisplatin. *Hum Gene Ther* 2001; **12**: 2225–2236.
76. Tanaka M, Grossman HB. Connexin 26 induces growth suppression, apoptosis and increased efficacy of doxorubicin in prostate cancer cells. *Oncol Rep* 2004; **11**: 537–541.
77. Udawatte C, Rippes H. The spread of apoptosis through gap-junctional channels in BHK cells transfected with Cx32. *Apoptosis* 2005; **10**: 1019–1029.
78. Wang M, Berthoud VM, Beyer EC. Connexin43 increases the sensitivity of prostate cancer cells to TNF α -induced apoptosis. *J Cell Sci* 2007; **120**: 320–329.
79. Cusato K, Bosco A, Rozental R, Guimaraes CA, Reese BE, Linden R *et al*. Gap junctions mediate bystander cell death in developing retina. *J Neurosci* 2003; **23**: 6413–6422.
80. Krutovskikh VA, Piccoli C, Yamasaki H. Gap junction intercellular communication propagates cell death in cancerous cells. *Oncogene* 2002; **21**: 1989–1999.
81. Lin JH, Weigel H, Cotrina ML, Liu S, Bueno E, Hansen AJ *et al*. Gap-junction-mediated propagation and amplification of cell injury. *Nat Neurosci* 1998; **1**: 494–500.
82. Rawanduzy A, Hansen A, Hansen TW, Nedergaard M. Effective reduction of infarct volume by gap junction blockade in a rodent model of stroke. *J Neurosurg* 1997; **87**: 916–920.
83. Frantseva MV, Kokorovtseva L, Naus CG, Carlen PL, MacFabe D, Perez Velazquez JL. Specific gap junctions enhance the neuronal vulnerability to brain traumatic injury. *J Neurosci* 2002; **22**: 644–653.
84. Farahani R, Pina-Benabou MH, Kyrozis A, Siddiq A, Barradas PC, Chiu FC *et al*. Alterations in metabolism and gap junction expression may determine the role of astrocytes as “good samaritans” or executioners. *Glia* 2005; **50**: 351–361.
85. Andrade-Rozental AF, Rozental R, Hopperstad MG, Wu JK, Vrionis FD, Spray DC. Gap junctions: the ‘kiss of death’ and the ‘kiss of life’. *Brain Res Brain Res Rev* 2000; **32**: 308–315.
86. Mesnil M, Yamasaki H. Bystander effect in herpes simplex virus-thymidine kinase/ganciclovir cancer gene therapy: role of gap-junctional intercellular communication. *Cancer Res* 2000; **60**: 3989–3999.
87. Yamasaki H, Omori Y, Krutovskikh V, Zhu W, Mironov N, Yamakage K *et al*. Connexins in tumour suppression and cancer therapy. *Novartis Found Symp* 1999; **219**: 241–254; discussion 254–260.
88. Neijssen J, Herberts C, Drijfhout JW, Reits E, Janssen L, Neeffjes J. Cross-presentation by intercellular peptide transfer through gap junctions. *Nature* 2005; **434**: 83–88.
89. Krysko DV, Mussels S, Leybaert L, D'Herde K. Gap junctional communication and connexin43 expression in relation to apoptotic cell death and survival of granulosa cells. *J Histochem Cytochem* 2004; **52**: 1199–1207.
90. Trosko JE, Goodman JI. Intercellular communication may facilitate apoptosis: implications for tumor promotion. *Mol Carcinog* 1994; **11**: 8–12.

91. Wilson MR, Close TW, Trosko JE. Cell population dynamics (apoptosis, mitosis, and cell-cell communication) during disruption of homeostasis. *Exp Cell Res* 2000; **254**: 257–268.
92. Theiss C, Mazur A, Meller K, Mannherz HG. Changes in gap junction organization and decreased coupling during induced apoptosis in lens epithelial and NIH-3T3 cells. *Exp Cell Res* 2007; **313**: 38–52.
93. Yin X, Gu S, Jiang JX. Regulation of lens connexin 45.6 by apoptotic protease, caspase-3. *Cell Commun Adhes* 2001; **8**: 373–376.
94. Lin D, Shanks D, Prakash O, Takemoto DJ. Protein kinase C gamma mutations in the C1B domain cause caspase-3-linked apoptosis in lens epithelial cells through gap junctions. *Exp Eye Res* 2007; **85**: 113–122.
95. Lin D, Takemoto DJ. Oxidative activation of protein kinase Cgamma through the C1 domain. Effects on gap junctions. *J Biol Chem* 2005; **280**: 13682–13693.
96. Prochazka L, Turanek J, Tesarik R, Knotigova P, Polaskova P, Andrysik Z *et al*. Apoptosis and inhibition of gap-junctional intercellular communication induced by LA-12, a novel hydrophobic platinum(IV) complex. *Arch Biochem Biophys* 2007; **462**: 54–61.
97. Ogawa T, Hayashi T, Tokunou M, Nakachi K, Trosko JE, Chang CC *et al*. Suberoylanilide hydroxamic acid enhances gap junctional intercellular communication via acetylation of histone containing connexin 43 gene locus. *Cancer Res* 2005; **65**: 9771–9778.
98. Hao JL, Suzuki K, Lu Y, Hirano S, Fukuda K, Kumagai N *et al*. Inhibition of gap junction-mediated intercellular communication by TNF-alpha in cultured human corneal fibroblasts. *Invest Ophthalmol Vis Sci* 2005; **46**: 1195–1200.
99. Xie H, Laird DW, Chang TH, Hu VW. A mitosis-specific phosphorylation of the gap junction protein connexin43 in human vascular cells: biochemical characterization and localization. *J Cell Biol* 1997; **137**: 203–210.
100. Lampe PD, Kurata WE, Wam-Cramer BJ, Lau AF. Formation of a distinct connexin43 phosphoisoform in mitotic cells is dependent upon p34cdc2 kinase. *J Cell Sci* 1998; **111**: 833–841.
101. Kanemitsu MY, Jiang W, Eckhart W. Cdc2-mediated phosphorylation of the gap junction protein, connexin43, during mitosis. *Cell Growth Differ* 1998; **9**: 13–21.
102. Albright CD, Kuo J, Jeong S. cAMP enhances Cx43 gap junction formation and function and reverses choline deficiency apoptosis. *Exp Mol Pathol* 2001; **71**: 34–39.
103. Nakase T, Fushiki S, Sohl G, Theis M, Willecke K, Naus CC. Neuroprotective role of astrocytic gap junctions in ischemic stroke. *Cell Commun Adhes* 2003; **10**: 413–417.
104. Blanc EM, Bruce-Keller AJ, Mattson MP. Astrocytic gap junctional communication decreases neuronal vulnerability to oxidative stress-induced disruption of Ca²⁺ homeostasis and cell death. *J Neurochem* 1998; **70**: 958–970.
105. Cohen-Salmon M, Ott T, Michel V, Hardelin JP, Perfettini I, Eybalin M *et al*. Targeted ablation of connexin26 in the inner ear epithelial gap junction network causes hearing impairment and cell death. *Curr Biol* 2002; **12**: 1106–1111.
106. Giardina SF, Mikami M, Goubaeva F, Yang J. Connexin 43 confers resistance to hydrogen peroxide-mediated apoptosis. *Biochem Biophys Res Commun* 2007; **362**: 747–752.
107. Hutnik CM, Pocrnich CE, Liu H, Laird DW, Shao Q. The protective effect of functional connexin43 channels on a human epithelial cell line exposed to oxidative stress. *Invest Ophthalmol Vis Sci* 2008; **49**: 800–806.
108. Jin EJ, Lee SY, Jung JC, Bang OS, Kang SS. TGF-beta3 inhibits chondrogenesis of cultured chick leg bud mesenchymal cells via downregulation of connexin 43 and integrin beta4. *J Cell Physiol* 2008; **214**: 345–353.
109. Kruger O, Plum A, Kim JS, Winterhager E, Maxeiner S, Hallas G *et al*. Defective vascular development in connexin 45-deficient mice. *Development* 2000; **127**: 4179–4193.
110. Lin JH, Yang J, Liu S, Takano T, Wang X, Gao Q *et al*. Connexin mediates gap junction-independent resistance to cellular injury. *J Neurosci* 2003; **23**: 430–441.
111. Melanson-Drapeau L, Beyko S, Dave S, Hebb AL, Franks DJ, Sellitto C *et al*. Oligodendrocyte progenitor enrichment in the connexin32 null-mutant mouse. *J Neurosci* 2003; **23**: 1759–1768.
112. Ott T, Jokwitz M, Lenhard D, Romualdi A, Dombrowski F, Itrich C *et al*. Ablation of gap junctional communication in hepatocytes of transgenic mice does not lead to disrupted cellular homeostasis or increased spontaneous tumorigenesis. *Eur J Cell Biol* 2006; **85**: 717–728.
113. Paraguassu-Braga FH, Borojevic R, Bouzas LF, Barcinski MA, Bonomo A. Bone marrow stroma inhibits proliferation and apoptosis in leukemic cells through gap junction-mediated cell communication. *Cell Death Differ* 2003; **10**: 1101–1108.
114. Walker DL, Vacha SJ, Kirby ML, Lo CW. Connexin43 deficiency causes dysregulation of coronary vasculogenesis. *Dev Biol* 2005; **284**: 479–498.
115. Yasui K, Kada K, Hojo M, Lee JK, Kamiya K, Toyama J *et al*. Cell-to-cell interaction prevents cell death in cultured neonatal rat ventricular myocytes. *Cardiovasc Res* 2000; **48**: 68–76.
116. Nakase T, Sohl G, Theis M, Willecke K, Naus CC. Increased apoptosis and inflammation after focal brain ischemia in mice lacking connexin43 in astrocytes. *Am J Pathol* 2004; **164**: 2067–2075.
117. Goubaeva F, Mikami M, Giardina S, Ding B, Abe J, Yang J. Cardiac mitochondrial connexin 43 regulates apoptosis. *Biochem Biophys Res Commun* 2007; **352**: 97–103.
118. Satomi Y, Nishino H, Shibata S. Glycyrrhetic acid and related compounds induce G1 arrest and apoptosis in human hepatocellular carcinoma HepG2. *Anticancer Res* 2005; **25**: 4043–4047.
119. Wong RC, Dottori M, Koh KL, Nguyen LT, Pera MF, Pebay A. Gap junctions modulate apoptosis and colony growth of human embryonic stem cells maintained in a serum-free system. *Biochem Biophys Res Commun* 2006; **344**: 181–188.
120. Lee NP, Leung KW, Wo JY, Tam PC, Yeung WS, Luk JM. Blockage of testicular connexins induced apoptosis in rat seminiferous epithelium. *Apoptosis* 2006; **11**: 1215–1229.
121. Evans WH, Boitano S. Connexin mimetic peptides: specific inhibitors of gap-junctional intercellular communication. *Biochem Soc Trans* 2001; **29**: 606–612.
122. Cusato K, Ripps H, Zakevicius J, Spray DC. Gap junctions remain open during cytochrome c-induced cell death: relationship of conductance to 'bystander' cell killing. *Cell Death Differ* 2006; **13**: 1707–1714.
123. Boitano S, Dirksen ER, Sanderson MJ. Intercellular propagation of calcium waves mediated by inositol trisphosphate. *Science* 1992; **258**: 292–295.
124. Chen R, Valencia I, Zhong F, McColl KS, Roderick HL, Bootman MD *et al*. Bcl-2 functionally interacts with inositol 1,4,5-trisphosphate receptors to regulate calcium release from the ER in response to inositol 1,4,5-trisphosphate. *J Cell Biol* 2004; **166**: 193–203.
125. Rong YP, Aromolaran AS, Bultynck G, Zhong F, Li X, McColl K *et al*. Targeting Bcl-2-IP3 receptor interaction to reverse Bcl-2's inhibition of apoptotic calcium signals. *Mol Cell* 2008; **31**: 255–265.
126. Orrenius S, Zhivotovskiy B, Nicotera P. Regulation of cell death: the calcium-apoptosis link. *Nat Rev Mol Cell Biol* 2003; **4**: 552–565.
127. Boehning D, Patterson RL, Snyder SH. Apoptosis and calcium: new roles for cytochrome c and inositol 1,4,5-trisphosphate. *Cell Cycle* 2004; **3**: 252–254.
128. Rong Y, Distelhorst CW. Bcl-2 protein family members: versatile regulators of calcium signaling in cell survival and apoptosis. *Annu Rev Physiol* 2008; **70**: 73–91.
129. Dupont G, Combettes L, Leybaert L. Calcium dynamics: spatio-temporal organization from the subcellular to the organ level. *Int Rev Cytol* 2007; **261**: 193–245.
130. Ebihara L. New roles for connexons. *News Physiol Sci* 2003; **18**: 100–103.
131. Retamal MA, Froger N, Palacios-Prado N, Ezan P, Saez PJ, Saez JC *et al*. Cx43 hemichannels and gap junction channels in astrocytes are regulated oppositely by proinflammatory cytokines released from activated microglia. *J Neurosci* 2007; **27**: 13781–13792.
132. Abrams CK, Bennett MV, Verselis VK, Bargiello TA. Voltage opens unopposed gap junction hemichannels formed by a connexin 32 mutant associated with X-linked Charcot-Marie-Tooth disease. *Proc Natl Acad Sci USA* 2002; **99**: 3980–3984.
133. Essenfelder GM, Bruzzone R, Lamartine J, Charollais A, Blanchet-Bardon C, Barbe MT *et al*. Connexin30 mutations responsible for hidrotic ectodermal dysplasia cause abnormal hemichannel activity. *Hum Mol Genet* 2004; **13**: 1703–1714.
134. Schock SC, Leblanc D, Hakim AM, Thompson CS. ATP release by way of connexin 36 hemichannels mediates ischemic tolerance *in vitro*. *Biochem Biophys Res Commun* 2008; **368**: 138–144.
135. Stridh MH, Tranberg M, Weber SG, Blomstrand F, Sandberg M. Stimulated efflux of amino acids and glutathione from cultured hippocampal slices by omission of extracellular calcium: likely involvement of connexin hemichannels. *J Biol Chem* 2008; **283**: 10347–10356.
136. Takeuchi H, Jin S, Wang J, Zhang G, Kawanokuchi J, Kuno R *et al*. Tumor necrosis factor-alpha induces neurotoxicity via glutamate release from hemichannels of activated microglia in an autocrine manner. *J Biol Chem* 2006; **281**: 21362–21368.
137. Hur KC, Shim JE, Johnson RG. A potential role for cx43-hemichannels in staurosporin-induced apoptosis. *Cell Commun Adhes* 2003; **10**: 271–277.
138. Leist M, Single B, Castoldi AF, Kuhnle S, Nicotera P. Intracellular adenosine triphosphate (ATP) concentration: a switch in the decision between apoptosis and necrosis. *J Exp Med* 1997; **185**: 1481–1486.
139. Saez JC, Contreras JE, Bukauskas FF, Retamal MA, Bennett MV. Gap junction hemichannels in astrocytes of the CNS. *Acta Physiol Scand* 2003; **179**: 9–22.
140. Turner MS, Haywood GA, Andreaka P, You L, Martin PE, Evans WH *et al*. Reversible connexin 43 dephosphorylation during hypoxia and reoxygenation is linked to cellular ATP levels. *Circ Res* 2004; **95**: 726–733.
141. Vergara L, Bao X, Bello-Reuss E, Reuss L. Do connexin 43 gap-junctional hemichannels activate and cause cell damage during ATP depletion of renal-tubule cells? *Acta Physiol Scand* 2003; **179**: 33–38.
142. Tran Van Nhiu G, Clair C, Bruzzone R, Mesnil M, Sansonetti P, Combettes L. Connexin-dependent inter-cellular communication increases invasion and dissemination of Shigella in epithelial cells. *Nat Cell Biol* 2003; **5**: 720–726.
143. Schrier SM, Florea BI, Mulder GJ, Nagelkerke JF, AP JJ. Apoptosis induced by extracellular ATP in the mouse neuroblastoma cell line N1E-115: studies on involvement of P2 receptors and adenosine. *Biochem Pharmacol* 2002; **63**: 1119–1126.
144. Dunwiddie TV, Masino SA. The role and regulation of adenosine in the central nervous system. *Annu Rev Neurosci* 2001; **24**: 31–55.
145. Franke H, Krugel U, Illes P. P2 receptors and neuronal injury. *Pflügers Arch* 2006; **452**: 622–644.
146. Chow SC, Kass GE, Orrenius S. Purines and their roles in apoptosis. *Neuropharmacology* 1997; **36**: 1149–1156.
147. Ferrari D, Los M, Bauer MK, Vandenabeele P, Wesselborg S, Schulze-Osthoff K. P2Z purinoreceptor ligation induces activation of caspases with distinct roles in apoptotic and necrotic alterations of cell death. *FEBS Lett* 1999; **447**: 71–75.

148. Tsukimoto M, Harada H, Ikari A, Takagi K. Involvement of chloride in apoptotic cell death induced by activation of ATP-sensitive P2X7 purinoceptor. *J Biol Chem* 2005; **280**: 2653–2658.
149. Jabbour HN, Kelly RW, Boddy SC. Autocrine/paracrine regulation of apoptosis in epithelial cells by prostaglandin E2. *Prostaglandins Leukot Essent Fatty Acids* 2002; **67**: 357–363.
150. Vajda FJ. Neuroprotection and neurodegenerative disease. *J Clin Neurosci* 2002; **9**: 4–8.
151. Arthur DB, Georgi S, Akassoglou K, Insel PA. Inhibition of apoptosis by P2Y2 receptor activation: novel pathways for neuronal survival. *J Neurosci* 2006; **26**: 3798–3804.
152. Bano D, Nicotera P. Ca²⁺ signals and neuronal death in brain ischemia. *Stroke* 2007; **38**: 674–676.
153. Zipfel GJ, Babcock DJ, Lee JM, Choi DW. Neuronal apoptosis after CNS injury: the roles of glutamate and calcium. *J Neurotrauma* 2000; **17**: 857–869.
154. Abbraccio MP, Verderio C. Pathophysiological roles of P2 receptors in glial cells. *Novartis Found Symp* 2006; **276**: 91–103; discussion 103–112, 275–181.
155. Lahne M, Gale JE. Damage-induced activation of ERK1/2 in cochlear supporting cells is a hair cell death-promoting signal that depends on extracellular ATP and calcium. *J Neurosci* 2008; **28**: 4918–4928.
156. Lin JH, Lou N, Kang N, Takano T, Hu F, Han X *et al*. A central role of connexin 43 in hypoxic preconditioning. *J Neurosci* 2008; **28**: 681–695.
157. Yawata I, Takeuchi H, Doi Y, Liang J, Mizuno T, Suzumura A. Macrophage-induced neurotoxicity is mediated by glutamate and attenuated by glutamine inhibitors and gap junction inhibitors. *Life Sci* 2008; **82**: 1111–1116.
158. Naus CC, Ozog MA, Bechberger JF, Nakase T. A neuroprotective role for gap junctions. *Cell Commun Adhes* 2001; **8**: 325–328.
159. O'Carroll SJ, Alkadi M, Nicholson LF, Green CR. Connexin 43 mimetic peptides reduce swelling, astrogliosis, and neuronal cell death after spinal cord injury. *Cell Commun Adhes* 2008; **15**: 27–42.
160. Danesh-Meyer HV, Huang R, Nicholson LF, Green CR. Connexin43 antisense oligodeoxynucleotide treatment down-regulates the inflammatory response in an *in vitro* interphase organotypic culture model of optic nerve ischaemia. *J Clin Neurosci* 2008; **15**: 1253–1263.
161. Krysko DV, D'Herde K, Vandenabeele P. Clearance of apoptotic and necrotic cells and its immunological consequences. *Apoptosis* 2006; **11**: 1709–1726.
162. Krysko DV, Vandenabeele P. From regulation of dying cell engulfment to development of anti-cancer therapy. *Cell Death Differ* 2008; **15**: 29–38.
163. Plotkin LI, Manolagas SC, Bellido T. Transduction of cell survival signals by connexin-43 hemichannels. *J Biol Chem* 2002; **277**: 8648–8657.
164. Boengler K, Dodoni G, Rodriguez-Sinovas A, Cabestrero A, Ruiz-Meana M, Gres P *et al*. Connexin 43 in cardiomyocyte mitochondria and its increase by ischemic preconditioning. *Cardiovasc Res* 2005; **67**: 234–244.
165. Rodriguez-Sinovas A, Boengler K, Cabestrero A, Gres P, Morente M, Ruiz-Meana M *et al*. Translocation of connexin 43 to the inner mitochondrial membrane of cardiomyocytes through the heat shock protein 90-dependent TOM pathway and its importance for cardioprotection. *Circ Res* 2006; **99**: 93–101.
166. Ruiz-Meana M, Rodriguez-Sinovas A, Cabestrero A, Boengler K, Heusch G, Garcia-Dorado D. Mitochondrial connexin43 as a new player in the pathophysiology of myocardial ischaemia-reperfusion injury. *Cardiovasc Res* 2008; **77**: 325–333.
167. Schulz R, Boengler K, Totzeck A, Luo Y, Garcia-Dorado D, Heusch G. Connexin 43 in ischemic pre- and postconditioning. *Heart Fail Rev* 2007; **12**: 261–266.
168. Fujimoto E, Satoh H, Negishi E, Ueno K, Nagashima Y, Hagiwara K *et al*. Negative growth control of renal cell carcinoma cell by connexin 32: possible involvement of Her-2. *Mol Carcinog* 2004; **40**: 135–142.
169. Goldberg GS, Bechberger JF, Tajima Y, Merritt M, Omori Y, Gawinowicz MA *et al*. Connexin43 suppresses MFG-E8 while inducing contact growth inhibition of glioma cells. *Cancer Res* 2000; **60**: 6018–6026.
170. Huang RP, Fan Y, Hossain MZ, Peng A, Zeng ZL, Boynton AL. Reversion of the neoplastic phenotype of human glioblastoma cells by connexin 43 (cx43). *Cancer Res* 1998; **58**: 5089–5096.
171. Princen F, Robe P, Gros D, Jarry-Guichard T, Gielen J, Merville MP *et al*. Rat gap junction connexin-30 inhibits proliferation of glioma cell lines. *Carcinogenesis* 2001; **22**: 507–513.
172. Zhang YW, Kaneda M, Morita I. The gap junction-independent tumor-suppressing effect of connexin 43. *J Biol Chem* 2003; **278**: 44852–44856.
173. Gu S, Yu XS, Yin X, Jiang JX. Stimulation of lens cell differentiation by gap junction protein connexin 45.6. *Invest Ophthalmol Vis Sci* 2003; **44**: 2103–2111.
174. Jiang JX, Gu S. Gap junction- and hemichannel-independent actions of connexins. *Biochim Biophys Acta* 2005; **1711**: 208–214.
175. Kanczuga-Koda L, Sulkowski S, Tomaszewski J, Koda M, Sulkowska M, Przystupa W *et al*. Connexins 26 and 43 correlate with Bak, but not with Bcl-2 protein in breast cancer. *Oncol Rep* 2005; **14**: 325–329.
176. Kanczuga-Koda L, Sulkowski S, Koda M, Skrzydlewska E, Sulkowska M. Connexin 26 correlates with Bcl-xL and Bax proteins expression in colorectal cancer. *World J Gastroenterol* 2005; **11**: 1544–1548.
177. Fujimoto E, Sato H, Shirai S, Nagashima Y, Virgona N, Hagiwara K *et al*. Inhibition of Src activity enhances the tumor-suppressive effect of the connexin 32 gene in Caki-1 renal cancer cells. *Oncol Rep* 2006; **15**: 1359–1365.
178. Iacobas DA, Urban-Maldonado M, Iacobas S, Scemes E, Spray DC. Array analysis of gene expression in connexin-43 null astrocytes. *Physiol Genomics* 2003; **15**: 177–190.
179. Stains JP, Civitelli R. Gap junctions regulate extracellular signal-regulated kinase signaling to affect gene transcription. *Mol Biol Cell* 2005; **16**: 64–72.
180. Stains JP, Lecanda F, Screen J, Towler DA, Civitelli R. Gap junctional communication modulates gene transcription by altering the recruitment of Sp1 and Sp3 to connexin-response elements in osteoblast promoters. *J Biol Chem* 2003; **278**: 24377–24387.
181. Dang X, Doble BW, Kardami E. The carboxy-tail of connexin-43 localizes to the nucleus and inhibits cell growth. *Mol Cell Biochem* 2003; **242**: 35–38.
182. Moorby C, Patel M. Dual functions for connexins: Cx43 regulates growth independently of gap junction formation. *Exp Cell Res* 2001; **271**: 238–248.
183. Herve JC, Bourmeyster N, Sarrouilhe D, Duffy HS. Gap junctional complexes: from partners to functions. *Prog Biophys Mol Biol* 2007; **94**: 29–65.
184. Kardami E, Dang X, Iacobas DA, Nickel BE, Jayaraman M, Srisakuldee W *et al*. The role of connexins in controlling cell growth and gene expression. *Prog Biophys Mol Biol* 2007; **94**: 245–264.
185. Ai Z, Fischer A, Spray DC, Brown AM, Fishman GI. Wnt-1 regulation of connexin43 in cardiac myocytes. *J Clin Invest* 2000; **105**: 161–171.
186. Kim PJ, Plescia J, Clevers H, Fearon ER, Altieri DC. Survivin and molecular pathogenesis of colorectal cancer. *Lancet* 2003; **362**: 205–209.
187. Xie H, Huang Z, Sadim MS, Sun Z. Stabilized beta-catenin extends thymocyte survival by up-regulating Bcl-xL. *J Immunol* 2005; **175**: 7981–7988.
188. Li Q, Dashwood WM, Zhong X, Nakagama H, Dashwood RH. Bcl-2 over-expression in PhIP-induced colon tumors: cloning of the rat Bcl-2 promoter and characterization of a pathway involving beta-catenin, c-Myc and E2F1. *Oncogene* 2007; **26**: 6194–6202.
189. Sharrow AC, Li Y, Micsenyi A, Griswold RD, Wells A, Monga SS *et al*. Modulation of osteoblast gap junction connectivity by serum, TNFalpha, and TRAIL. *Exp Cell Res* 2008; **314**: 297–308.
190. Pelegrin P, Surprenant A. Pannexin-1 couples to maitotoxin- and nigericin-induced interleukin-1beta release through a dye uptake-independent pathway. *J Biol Chem* 2007; **282**: 2386–2394.
191. Kanneganti TD, Lamkanfi M, Kim YG, Chen G, Park JH, Franchi L *et al*. Pannexin-1-mediated recognition of bacterial molecules activates the cryopyrin inflammasome independent of Toll-like receptor signaling. *Immunity* 2007; **26**: 433–443.
192. Smart ML, Gu B, Panchal RG, Wiley J, Cromer B, Williams DA *et al*. P2X7 receptor cell surface expression and cytolytic pore formation are regulated by a distal C-terminal region. *J Biol Chem* 2003; **278**: 8853–8860.
193. Locovei S, Scemes E, Qiu F, Spray DC, Dahl G. Pannexin1 is part of the pore forming unit of the P2X(7) receptor death complex. *FEBS Lett* 2007; **581**: 483–488.
194. Egan TM, Samways DS, Li Z. Biophysics of P2X receptors. *PLoS Arch* 2006; **452**: 501–512.
195. Spray DC, Ye ZC, Ransom BR. Functional connexin "hemichannels": a critical appraisal. *Glia* 2006; **54**: 758–773.
196. Iglesias R, Locovei S, Roque AP, Alberto AP, Dahl G, Spray DC *et al*. P2X7 receptor-Pannexin1 complex: Pharmacology and signaling. *Am J Physiol Cell Physiol* 2008; **295**: C752–C760.
197. Szalai G, Krishnamurthy R, Hajnoczky G. Apoptosis driven by IP(3)-linked m. *EMBO J* 1999; **18**: 6349–6361.
198. Zappala A, Li Volti G, Serapide MF, Pellitteri R, Falchi M, La Delia F *et al*. Expression of pannexin2 protein in healthy and ischemized brain of adult rats. *Neuroscience* 2007; **148**: 653–667.