



University of
Salford
MANCHESTER

Consequences of a large-scale fragmentation experiment for Neotropical bats : disentangling the relative importance of local and landscape-scale effects

Rocha, R, López-Baucells, A, Farneda, FZ, Groenenberg, M, Bobrowiec, PED, Cabeza, M, Palmeirim, JM and Meyer, CFJ

<http://dx.doi.org/10.1007/s10980-016-0425-3>

Title	Consequences of a large-scale fragmentation experiment for Neotropical bats : disentangling the relative importance of local and landscape-scale effects
Authors	Rocha, R, López-Baucells, A, Farneda, FZ, Groenenberg, M, Bobrowiec, PED, Cabeza, M, Palmeirim, JM and Meyer, CFJ
Publication title	Landscape Ecology
Publisher	Springer
Type	Article
USIR URL	This version is available at: http://usir.salford.ac.uk/id/eprint/39550/
Published Date	2017

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: library-research@salford.ac.uk.

**Consequences of a large-scale fragmentation experiment for Neotropical bats:
disentangling the relative importance of local and landscape-scale effects**

Ricardo Rocha^{1,2,3,4,*}, Adrià López-Baucells^{1,2,5,6}, Fábio Z. Farneda^{1,2,7}, Milou Groenenberg^{2,8}, Paulo E.D. Bobrowiec², Mar Cabeza³, Jorge M. Palmeirim^{1,2} and Christoph F.J. Meyer^{1,2,5}

¹*Centre for Ecology, Evolution and Environmental Changes, Faculty of Sciences, University of Lisbon, Lisbon, Portugal*

²*Biological Dynamics of Forest Fragments Project, National Institute for Amazonian Research and Smithsonian Tropical Research Institute, Manaus, Brazil*

³*Metapopulation Research Centre, Faculty of Biosciences, University of Helsinki, Helsinki, Finland*

⁴*Faculty of Life Sciences, University of Madeira, Portugal*

⁵*Ecosystems and Environment Research Centre (EERC), School of Environment and Life Sciences, University of Salford, Salford, M5 4WT, United Kingdom*

⁶*Museu de Ciències Naturals de Granollers, Catalunya, Spain*

⁷*Department of Ecology/PPGE, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil*

⁸*Department of Life Sciences, Imperial College London, London, United Kingdom*

***Correspondence:** Ricardo Rocha; **E-mail:** ricardo.nature@gmail.com; **Phone:** +351 217500577; **Fax:** +351 217500028

Type of manuscript: Original research

Date: 15-05-2015

Manuscript word count: 6206

Abstract

Context Habitat loss, fragmentation and degradation are widespread drivers of biodiversity decline. Understanding how habitat quality interacts with landscape context, and how they jointly affect species in human-modified landscapes, is of great importance for informing conservation and management.

Objectives We used a whole-ecosystem manipulation experiment in the Brazilian Amazon to investigate the relative roles of local and landscape attributes in affecting bat assemblages at an interior-edge-matrix disturbance gradient.

Methods We surveyed bats in 39 sites, comprising continuous forest, fragments, forest edges and intervening secondary regrowth. For each site, we assessed vegetation structure (local-scale variable) and, for five focal scales, quantified habitat amount and four landscape configuration metrics.

Results Smaller fragments, edges and regrowth sites had fewer species and higher levels of dominance than continuous forest. Regardless of the landscape scale analysed, species richness and evenness were mostly related to the amount of forest cover. Vegetation structure and configurational metrics were important predictors of abundance, whereby the magnitude and direction of response to configurational metrics were scale-dependent. Responses were ensemble-specific with local-scale vegetation structure being more important for frugivorous than for gleaning animalivorous bats.

Conclusions Our study indicates that scale-sensitive measures of landscape structure are needed for a more comprehensive understanding of the effects of fragmentation on tropical biota. Although forest fragments and regrowth habitats can be of conservation significance for tropical

bats our results further emphasize that primary forest is of irreplaceable value, underlining that their conservation can only be achieved by the preservation of large expanses of pristine habitat.

25

Keywords Amazon; edge effects; FRAGSTATS; landscape context; matrix; secondary forest; spatial scale; vegetation structure.

Introduction

Anthropogenic forest loss and fragmentation are key drivers of the ongoing
30 defaunation crisis (Dirzo et al 2014). This erosion of biological diversity has repeatedly been
associated with human population growth and rising per-capita consumption (Laurance et al
2014) whose future increases are expected to be greatest in the tropics where much of the
planet's biodiversity resides (Bradshaw et al 2008; Gibson et al 2011).

In human-modified landscapes, habitat loss and fragmentation typically co-occur with
35 habitat degradation (Fischer and Lindenmayer 2007). Deterioration in habitat quality is most
noteworthy near primary forest edges and in regenerating forests, where biotic and abiotic
gradients and alternative successional pathways lead to marked differences in vegetation
structure (Williamson et al 2014; Faria et al 2009). Forest edges and regenerating forests are
ubiquitous features of tropical landscapes (Chazdon 2014). For instance, ~32000 km of new
40 forest edges are created annually in the Brazilian Amazon by deforestation alone (Broadbent
et al 2008) and, in 2000, ~140 x 10³ km² of the region's land area was composed of
regenerating forests (Carreiras et al 2006). Regenerating secondary forests profoundly
influence the spatio-temporal distribution of many species (e.g. Barlow et al 2007; Banks-
Leite et al 2010). However, studies focussing on the full disturbance gradient of continuous
45 forest (CF) and fragment interiors (I), forest edges (E) and matrix (M) habitats (hereafter IEM
gradients) in fragmented landscapes are scarce, and habitat quality metrics are rarely
incorporated into landscape-scale fragmentation studies (Galitsky and Lawler 2015). This
translates into a poor understanding of how habitat quality interacts with landscape context
and how they jointly affect species persistence and abundance in human-modified landscapes.

50 Bats comprise a large fraction of tropical mammalian fauna and play key ecological
roles in pollination, seed dispersal and insect suppression (Kunz et al 2011). They are acutely
sensitive to human-induced landscape changes (García-Morales et al 2013; Meyer et al 2016)
and their local abundance and diversity qualifies them as a well-suited indicator group to
examine the effects of fragmentation on tropical biota (Jones et al 2009).

55 MacArthur and Wilson's (1967) island biogeographic theory profoundly influenced
early research on fragmented ecosystems. Studies on tropical bats were no exception to this,
with earlier work focusing mainly on the effects of patch area and isolation metrics (Cosson et
al 1999; Schulze et al 2000). As the conceptual basis of fragmentation studies matured,
landscape characteristics such as habitat amount and configuration came to be recognized as
60 important determinants of bat species persistence in modified forest landscapes and the few
studies that have explored tropical bat associations with landscape structure at multiple spatial
scales have found assemblages to respond in a scale-sensitive manner (reviewed in Meyer et
al 2016). This scale sensitivity in bat responses towards landscape structure likely reflects
interspecific differences in species ecological traits such as diet, body size and home range
65 which are linked to the scale at which individual species perceive and interact with their
environment (Pinto and Keitt 2008; Meyer et al 2016). Scale dependency is also indicative of
the influence of smaller scale drivers upon ecological processes that operate at larger spatial
scales (McGill 2010). By imposing limitations on mobility and food detection, microhabitat
characteristics such as vegetation structure strongly influence the type and number of bat
70 species co-occurring on a local scale (Marciente et al 2015). However, vegetation structure is
rarely included in multi-scale fragmentation studies although it has been suggested to
constitute a better predictor of the activity of forest-dwelling bats than landscape-level

features (Erickson and West 2003; Charbonnier et al 2016) and is likely to modulate ecological responses to fragmentation at the landscape level.

75 The Biological Dynamics of Forest Fragments Project (BDFFP) in the Brazilian Amazon is the world's largest and longest-running experimental study of forest fragmentation (Laurance et al 2011). Vertebrate assemblages at the BDFFP have been found to be sensitive to fragment size (Ferraz et al 2007; Boyle and Smith 2010), edge effects (Lenz et al 2014; Powell et al 2015), matrix composition (Antongiovanni and Metzger 2005; Bobrowiec and
80 Gribel 2010), local vegetation structure (Stratford and Stouffer 2013; Mokross et al 2014) and landscape-scale characteristics (Stouffer et al 2006; Boyle and Smith 2010). However, no study has jointly investigated how vegetation structure and landscape composition and configuration affect the occurrence and abundance patterns of its vertebrate assemblages. Here we address this gap by examining how BDFFP bat assemblages respond to an IEM
85 disturbance gradient in a landscape where fragments are embedded in a "soft" matrix composed of advanced secondary vegetation. Specifically, we address the following questions:

(1) How do bat species richness, evenness, abundance and assemblage composition change along IEM (interior, edge and matrix) and size (CF, 100, 10 and 1 ha fragments)
90 gradients? Relative to CF interiors we expected forest fragments to exhibit reduced species richness and evenness and we hypothesized that differences in response metrics (species richness, evenness, abundance and assemblage composition) between IEM habitats would decrease with fragment size. Additionally, we predicted frugivores to be more edge- and matrix-tolerant than gleaning animalivorous bats.

95 (2) What is the relative importance of local vegetation structure versus landscape-scale
primary forest cover and configuration as determinants of bat species richness, evenness and
abundance? And how does it vary across multiple spatial scales? We anticipated different
response metrics to relate differently to vegetation structure and landscape characteristics and
predicted that responses would be scale-dependent with all three assemblage attributes
100 (species richness, evenness and abundance) responding predominantly to forest cover. We
also expected gleaning animalivores to present stronger negative effects towards
configuration metrics than frugivorous bats.

Material and methods

105 Study area and site selection

Fieldwork was conducted at the BDFFP, located ~80 km north of Manaus (2°30'S,
60°W), Brazil (see Fig. S1 in the online supplementary material). The area is classified as
tropical moist forest, and is characterized by a mosaic of *terra firme* rainforest, secondary
regrowth, and primary forest fragments. Annual rainfall varies from 1900-3500 mm, with a
110 dry season between June and October (Laurance et al 2011). The forest fragments were
isolated from continuous forest by distances of 80-650 m in the early 1980s, and are
categorized into size classes of 1, 10 and 100 ha. Each fragment was re-isolated on 3-4
occasions prior to this study, most recently between 1999 and 2001 (Laurance et al 2011). The
matrix is composed of tall secondary forest dominated mainly by *Vismia* spp. and *Cecropia*
115 spp. (Mesquita et al 2001).

Bats were sampled in eight forest fragments - three of 1 ha, three of 10 ha and two of
100 ha (Colosso, Porto Alegre and Dimona camps) - and nine control sites in three areas of

CF (Cabo Frio, Florestal and Km 41 camps) (Fig. S1). Sampling was conducted in the interiors and at the edges of all eight fragments, as well as at eight sites located in the nearby secondary regrowth, 100 m away from the edge of each fragment. A similar sampling scheme was employed for CF, with nine sampling sites in the interior, three at the edge, and three matrix sites located 100 m away from the forest edge. Therefore, a total of 39 sites were sampled. Distances between interior and edge sites of CF and fragments were respectively 1118 ± 488 and 245 ± 208 m (mean \pm SD).

125

Bat surveys

Each sampling site was visited eight times over a 2-year period, between August 2011 and June 2013. Bats were captured using 14 ground-level mist nets (12 x 2.5 m, 16 mm mesh, ECOTONE, Poland) in CF and fragment interiors, and seven ground-level mist nets at the edge and matrix sites. Mist nets were deployed along existing trails which are known to be used by Neotropical bats as commuting flyways (Palmeirim and Etheridge 1985). At edge sites, these trails ran parallel to the border between primary forest and secondary regrowth. In our study area mist netting efficiency was found to be highest in the first few hours after sunset (Bernard 2002). Sampling therefore started at dusk and was performed for six hours during which nets were visited at intervals of ~20 minutes. Mist netting at the same location for consecutive days can lead to diminishing capture efficiency over time (Marques et al. 2013). Such net-shyness related bias was avoided by spacing visits to the same site three to four weeks apart. Species were identified following Gardner (2007) and Lim and Engstrom (2001), and taxonomy follows Gardner (2007). Most adult bats were marked with individually numbered ball-chain necklaces (frugivores and *Pteronotus parnellii*) or subcutaneous transponders (gleaning animalivores). We restricted analyses to phyllostomids and *P.*

140

parnellii, since all other species in Neotropical bat assemblages are known to be inadequately sampled with mist-nets (Kalko 1998).

145 Influence of fragment-size and interior-edge-matrix gradient

Species richness, evenness and abundance

Differences in species richness, evenness, and abundance between size- (CF, 100, 10 and 1 ha fragments) and IEM-gradients were assessed using generalized linear mixed-effects models (GLMMs), fitted in the R package “*lme4*” (Bates 2010). A categorical variable combining
150 information of both the size- and IEM-gradient was specified as a fixed effect, and a random term nesting “site” within “location” (the latter referring to the six research camps; Fig. S1) was incorporated. This approach accounts for potential autocorrelation between sites within the same location (Bolker et al 2009). For each size category and IEM, species richness and evenness, the latter quantified as Hurlbert's probability of interspecific encounter (PIE), were
155 computed using rarefaction. Rarefaction was performed using EcoSim software v.7 (Gotelli and Entsminger 2004) based on 1000 random rearrangements and independent sampling of individuals, rarefying to the abundance level of the site with the lowest number of captures. Total number of captures per site was used to compare differences in abundance using a Poisson GLMM, with the site's total number of mist-net hours (1 mist-net hour [mnh] equals
160 one 12-m net open for 1 h) specified as an offset. High inter-fragment variation in capture rates precluded robust inference about how fragment size affects capture rates.

Differences in abundance between size- and IEM-gradients were therefore analysed by both considering the distinct fragment size categories (100, 10 and 1 ha fragments) independently and by lumping the capture data from all fragments. Significant effects were further evaluated

165 via multiple comparison tests with Tukey contrasts (adjusted *P*-values reported) using the R
package “*multcomp*” (Hothorn et al 2013).

Assemblage composition

Differences in assemblage composition were characterized by means of a non-metric
multidimensional scaling (NMDS) ordination based on a Bray-Curtis dissimilarity matrix,
170 using the number of captures standardized by the site’s effort (bats per mnh) and scaled to a
mean of 0 and standard deviation of 1. Compositional differences between size- and IEM-
gradient habitat types were evaluated with a permutational multivariate analysis of variance
(PERMANOVA). Both analyses were conducted using the “*vegan*” package in R (Oksanen
et al 2013).

175 *Ensemble-specific responses*

According to available literature (Bernard 2001, 2002; Giannini and Kalko, 2004;
Pereira et al 2010) species were grouped into frugivores (subdivided into shrub and canopy
frugivores), gleaning animalivores, aerial insectivores and nectarivores (Table S1). The same
approach used to compare total abundance was used to explore ensemble-specific differences
180 in abundance across the size- and IEM-gradients.

Influence of local and landscape-scale variables

Vegetation structure

Vegetation structure was characterized within three 100 m² (5 x 20 m) plots
185 established 5 m from each side of the mist net transects. In each plot, nine variables were
quantified: (i) number of trees (diameter at breast height [DBH] ≥ 10 cm), (ii) number of

woody stems (DBH < 10 cm), (iii) average DBH of trees \geq 10 cm, (iv) percent canopy cover (estimated as the average of four spherical densiometer readings), (v) number of palms, (vi) number of *Vismia* spp. and *Cecropia* spp. (the fruits of both genera are consumed by several frugivorous bat species, e.g. Bernard 2002; Giannini and Kalko 2004), (vii) liana density (visually classified every 5 m in five categories varying from no lianas to very high liana density), (viii) tree height (based on visual estimates of 25 trees \geq 10 cm DBH) and (ix) vertical foliage density (calculated as the sum of the values obtained by visual estimation at seven height intervals [0-1 m, 1-2 m, 2-4 m, 4-8 m, 8-16 m, 16-24 m, 24-32 m] using 6 categorical classes [0 = no foliage, 1 = very sparse 0-20%, 2 = sparse 20-40%, 3 = medium 40-60%, 4 = dense 60-80%, 5 = very dense 80-100%]). Values for each sampling site were calculated as the average across replicated plots (Table S2).

Vegetation variables were $\log(x + 1)$ transformed, standardized to a mean of zero and a standard deviation of one, and submitted to a Principal Components Analysis (PCA). The scores of the first axis (PCA1) were then used as predictor variable summarizing vegetation structure (Fig. S2 and Table S3) in modelling bat responses to local habitat and landscape structure.

Landscape structure

To quantify compositional and configurational aspects of landscape structure we used a detailed forest vs non-forest map of the BDFFP landscape based on 2004 LandSat Thematic Mapper (TM) satellite images (30 m spatial resolution). Land cover classification was obtained through supervised classification (bands 7, 5, 4, 3, 2 and 1), with thorough field checking performed to validate map accuracy. Primary forest (hereafter simply “forest”) was clearly distinguished from second growth. Landscape metrics were chosen based on their

210 reported influence on Neotropical bats (Gorresen and Willig 2004; Meyer and Kalko 2008; Klingbeil and Willig 2009; Avila-Cabadilla et al 2012) and were calculated using Fragstats v.4.1 (McGarigal et al 2012) for buffers with radii of 250, 500, 750, 1000 and 1500 m around each of the 39 sampling points. Buffer scales were selected so as to encompass the home ranges of different-sized bat species while at the same time minimizing spatial overlap
215 between neighboring sites (Meyer et al 2008). Five metrics were selected to represent: (a) habitat amount (forest cover) and (b) habitat configuration (mean patch area, patch density, edge density and mean shape index).

Relative importance of local and landscape-scale predictors of bat responses

The relative importance of local (vegetation structure) vs landscape-scale
220 characteristics in determining species richness, evenness and abundance were investigated at the five different focal scales using GLMMs. For this, we used rarefied species richness, evenness and total number of captures at each site as response variables. Additionally, GLMMs using total number of captures per site of frugivores and gleaning animalivores, respectively, were used to explore ensemble-specific relationships. Low number of captures
225 precluded separate analyses for the other ensembles. Severe collinearity between predictor variables can undermine statistical inference in GLMMs (Dormann et al 2013). We therefore quantified collinearity by calculating each predictor's variance inflation factor (VIF) within a set of predictors that always included vegetation structure and habitat amount (forest cover). As VIFs > 10 are known to indicate "severe" collinearity (Neter et al 1990) we reduced our
230 set of predictor variables (by excluding mean patch area and mean shape index) so that those included in the final set presented a VIF < 6 in all analysed buffers.

Analyses were restricted to a subset of a priori selected models comprising plausible combinations of local (vegetation structure) and landscape predictors (forest cover, edge density and patch density). For each response variable and landscape-scale separate sets of models were defined, which considered (i) each metric independently, (ii) vegetation structure and each landscape metric independently, (iii) vegetation structure and habitat configuration metrics, (iv) forest cover and each habitat configuration metric independently and (v) configuration metrics only. Each model included a random term accounting for the nested sampling design (site within location). Model goodness-of-fit was assessed as the marginal R^2_m and conditional R^2_c (Nakagawa and Schielzeth 2013). Following Burnham and Anderson (2002), the most parsimonious models were selected using Akaike's Information Criterion corrected for small samples sizes (AIC_c). Model-averaging was used to obtain parameter estimates for the models with an AIC_c difference from the best model (Δ_i) < 2 (Burnham and Anderson 2002). Residual spatial autocorrelation in the best-fit GLMMs was inspected by means of Moran's I test. For these best-fit models, the variation independently explained by each explanatory variable was then determined by hierarchical partitioning analysis using the "hier.part" package (Walsh and Mac Nally 2013), modified to accommodate a model offset [log(effort)] for abundance data (Jeppsson et al 2010). Following Benchimol and Peres (2015), hierarchical partitioning was conducted only considering fixed effects. Unless otherwise stated, all analyses were conducted in R v3.0.2 (R Development Core Team 2013).

Results

During 18650 mnh we captured 4210 bats belonging to six families and 55 species (Table S1). Phyllostomids and mormoopids (*P. parnellii*) were the dominant groups, accounting respectively for 90.9% (3827) and 6.5% (272) of total captures.

Responses to size- and interior-edge-matrix gradients

Species richness, evenness and abundance

Species richness was significantly higher in CF interiors than in any fragment size class, with the exception of the 100 ha fragment interiors (Fig. 1). Similarly, edge and matrix sites adjoining CF were more species-rich than those adjacent to fragments and, for both interior and matrix habitats, species richness tended to increase with fragment size. Conversely, this pattern did not hold for edge habitats as the edges of the 1 ha fragments were surprisingly diverse, attaining comparable richness to those of CF (Table S4).

265

#Figure 1 approximately here#

Evenness showed a similar pattern to species richness (Fig. 1), with all habitats other than the 100 ha fragment interiors and CF edges being significantly less even than CF interiors (Table S4). Evenness was higher for edge and matrix habitats adjoining CF and 100 ha fragments, and again, the edges of 1 ha fragments had significantly higher evenness than those of the other size classes.

For both CF and fragments, total species abundance increased progressively from interior to edge and matrix habitats, with capture rates in the edge and matrix habitats being

275 significantly higher than in CF interiors. However, when the most common species (*Carollia perspicillata*) was excluded, the effect only remained significant for the comparison with CF edges (Fig. 2 and Table S5; see Fig. S3 for results by fragment size).

#Figure 2 approximately here#

280

Assemblage composition

The NMDS ordination had a stress value of 0.095, conveying a good representation of the data along two dimensions. Bat assemblage composition differed significantly across the size- and IEM-gradients (Fig. 3; $F_{11} = 2.316$, $R^2 = 0.486$, $P = 0.001$). The interiors of each
285 fragment size category formed a distinct cluster and the 100 ha fragment interiors grouped closely to the cluster formed by CF interiors, indicating high assemblage similarity. Edge and matrix sites clustered independently from CF and 100 ha fragment interiors and presented a large spread along the first ordination axis.

290 #Figure 3 approximately here#

Ensemble-specific responses

Shrub frugivores was the ensemble with the most captures (69%), followed by gleaning animalivores with 12.5%. Shrub and canopy frugivores showed similar patterns of
295 relative abundance, with higher capture rates in edge and matrix habitats compared to habitat interiors (Fig. 4; see Table S6 for results by fragment size). Capture rates for all frugivores

and shrub frugivores were significantly lower at CF interiors than in any other habitat category. For canopy frugivores there was no significant difference between CF interiors and CF edges and fragment interiors. Conversely, compared with the same IEM habitat type, gleaning animalivores in fragments had significantly lower capture rates than in CF. The capture rate of aerial insectivores (*P. parnellii*) was lower in fragment interior, edge and matrix habitats than in CF interiors, and the abundance of nectarivores peaked in fragment interiors and CF edges. However, none of these differences were significant (Table S7).

#Figure 4 approximately here#

Influence of local and landscape-scale variables

Assemblage and ensemble associations with local and landscape metrics were scale-sensitive and varied according to the response variable analysed (Fig. 5 and 6). However, for most response metrics and spatial scales, relatively high model selection uncertainty made it difficult to unequivocally pinpoint either local vegetation structure or landscape-scale attributes as best predictors (Table S8 and Table S9). GLMM residuals were not significantly spatially autocorrelated for evenness or any of the abundance models (Moran's I from -0.23 to 0.02, $P > 0.05$). However, for species richness and at the smallest spatial scale (250 m), one of the models included in the most parsimonious set ($\Delta AIC_c < 2$) presented spatially structured residuals (Moran's $I = -0.23$, $P < 0.05$) (Table S10). Consequently, for this scale and response variable the results should be interpreted with caution as autocorrelation may prompt to an elevated Type I error.

320 #Figure 5 approximately here#

Vegetation structure, as represented by PCA1, was a particularly relevant predictor of total abundance and abundance of frugivores. Other than for the smallest (250, 500 and 750 m) spatial scales, with species richness and the abundance of gleaning animalivores as response variables and for the smallest spatial scale for evenness, vegetation structure was always included in the most parsimonious models. The PCA1 explained 42.02% of the total variance and represented a gradient from simpler vegetation structural complexity, typical of secondary forest (higher density of pioneer trees [*Vismia* spp. and *Cecropia* spp.] and woody stems [DBH <10 cm]; negative values), to higher structural complexity, typical of primary forest sites (more closed canopy cover and higher density of trees [DBH >10]; positive values) (Fig. S2 and Table S2). Its relationship was positive with respect to all response metrics analysed, indicating that more complex sites in terms of vegetation structure presented higher species richness, evenness and abundance. Forest cover emerged as the most important predictor of species richness and evenness, being positively associated with both and with the abundance of gleaning animalivores. Its influence on total and frugivore abundance was negative across all scales. The effect of edge density was particularly sensitive to scale, being positively correlated with total abundance and abundance of frugivores and gleaning animalivores at the smallest scales (≤ 500 m) and negatively at larger spatial scales. Lastly, patch density showed greater consistency as predictor across scales, being negatively associated with species richness, total abundance and, except at one spatial scale, with the abundance of frugivores and gleaning animalivores.

#Figure 6 approximately here#

Discussion

345 Our analyses revealed that patterns of bat species richness, evenness and abundance varied across the BDFFP landscape, and were affected by local- and landscape-level habitat attributes in a scale-dependent and ensemble-specific manner. This despite the low structural contrast between CF, forest fragments and surrounding secondary regrowth.

350 Responses to size- and interior-edge-matrix gradients

Compared with CF, smaller (≤ 10 ha) fragments harboured fewer species and their assemblages were characterized by higher levels of dominance, results consistent with previous studies addressing the impacts of fragmentation on tropical bats (Cosson et al 1999; Meyer and Kalko 2008; Struebig et al 2008; Estrada-Villegas et al 2010) and other taxa at the BDFFP (Laurance et al 2006b; Ferraz et al 2007; Boyle and Smith 2010) as well as elsewhere in the tropics (Benchimol and Venticinque 2014; Bregman et al 2014). These differences, though remarkable given the low fragment-matrix contrast at the time of our study and the relatively short distance between forest fragments and nearby CF, seem to result from the strong effect of trait-mediated environmental filters that selectively benefit bat species with reduced body mass and a phytophagous diet (Farneda et al 2015). This is likely attributable to the elevated abundance of pioneer plants in early successional habitats, which benefit many small-bodied nectarivorous and frugivorous phyllostomids (e.g., *Glossophaga* spp., *Carollia* spp., *Sturnira* spp.), but fail to provide enough food resources to fulfil the energetic requirements of larger species and those of higher trophic levels.

365 For both CF and forest fragments, edge habitats had fewer species and higher levels of dominance. These differences were more noticeable in larger fragments (100 ha) and CF,

suggesting an area effect on the magnitude of contrast between interior and edge assemblages. Edge effects have long been identified as having major impacts on species distributions and dynamics (Ewers and Didham 2006). In the BDFFP fragments, edge effects are predominant
370 drivers of ecological change (Laurance et al 2011), affecting vegetation structure (Didham and Lawton 1999) and acting synergistically with area effects (Laurance et al 2006a).

Neotropical bats are known to respond to habitat edges (Gorresen and Willig 2004; Meyer and Kalko 2008; Klingbeil and Willig 2009; Klingbeil and Willig 2010; Bolívar-Cimé et al 2013) and the few studies that have compared phyllostomid assemblages at the interiors
375 and edges of fragments have reported declines in species richness (Faria 2006; Meyer and Kalko 2008). This pattern might result from the avoidance of these habitats by gleaning animalivorous bats, an ensemble identified as edge-sensitive in both high- (Meyer et al 2008) and low-contrast systems (Faria 2006). The underlying drivers of the higher edge-sensitivity exhibited by this ensemble remain to be tested but they might relate to changes in the
380 densities of preferred arthropod prey or to restrictions to flight maneuverability imposed by denser understory vegetation near edges.

Although the most conspicuous edge effects at the BDFFP have been detected within 100 m of forest edges (Laurance et al 2002), results from French Guiana indicate that edge-mediated changes in bat assemblage structure may be noticeable as far as 3 km from the forest
385 edge (Delaval and Charles-Dominique 2006). This suggests that even our CF interior sites (located on average more than 1 km away from the forest edge) are likely to suffer from the effects of edge penetration and consequently their bat assemblages may reflect the influence of the modified secondary forest matrix.

Bat assemblages in secondary regrowth adjoining CF and 100 ha fragments were also
390 richer and more even than assemblages adjacent to smaller fragments, suggesting a spillover
of species from the more diverse CF and 100 ha fragment interiors into the matrix. Low-
contrast matrix habitats are known to harbour diverse bat assemblages, both at the BDFFP
(Bobrowiec and Gribel 2010) and elsewhere in the Neotropics (Avila-Cabadilla et al 2009;
Avila-Cabadilla et al 2014; Mendenhall et al 2014). Our results indicate not only that matrix
395 habitats influence fragment ecology at the BDFFP (Gascon et al 1999; Laurance et al 2011)
but that the influence is bidirectional and that, similarly to birds (Powell et al 2013), bat
species dependent on old-growth stands may exploit the nearby secondary regrowth for
feeding or as flyways between food patches.

Human-induced habitat changes, including fragmentation, act as non-random filters
400 selecting those species with the best combination of traits to survive in modified ecosystems
(Smart et al 2006). In the humid Neotropics, capture rates of frugivores generally increase in
fragmented or disturbed areas, whereas gleaning animalivores tend to decline, if not
disappear, in modified habitats (Meyer et al 2016). Our results are consistent with this pattern.
The *Vismia*- and *Cecropia*-dominated secondary vegetation that surrounds the fragments in
405 our study landscape provides additional food resources that augments the abundance of
frugivores such as *C. perspicillata* (Bobrowiec and Gribel 2010). However, regrowth habitats
and forest fragments are structurally less complex than CF, and constitute less suitable habitat
conditions for most gleaning animalivores due to insufficient roosting and prey resources
(Gorresen and Willig 2004; Meyer and Kalko 2008). Nectarivorous bats have been
410 documented to remain stable or increase in forest remnants and edge habitats, owing to
elevated densities of food resources following forest clearance and subsequent succession
(García-Morales et al 2013; Meyer et al 2016; Chambers et al 2016, in press). At the BDFFP,

both nectarivorous birds (Stouffer et al 2006) and bats follow this pattern, adding to the evidence that nectar-feeders, together with frugivores, are the most resilient ensembles to
415 habitat modification.

Influence of local and landscape-scale variables

Our data suggest that both local and landscape metrics are important in explaining the effects of fragmentation on tropical bat assemblages. At the local-scale, we observed that sites
420 that are more similar to CF in terms of vegetation structure are able to support assemblages that are richer, more even and comprised of greater abundances of both frugivorous and gleaning animalivorous bats. These results agree with several other studies on aerial and terrestrial forest-dependent tropical vertebrates in modified landscapes (e.g. Benchimol and Venticinque 2014; Rocha et al 2015). However, they contrast with the findings from a study
425 of bat assemblages in a land-bridge island system in Panama (Meyer and Kalko 2008), which provided little evidence for an effect of vegetation structure on bat species richness and assemblage composition. This may reflect the wider environmental gradient of our study (which encompassed CF and fragment interiors, edges and matrix habitats) in relation to the one analysed in Panama (limited to CF and fragment interiors and CF edges). Vegetation
430 structure may therefore be a stronger determinant of assemblage diversity and composition in systems with high vegetation heterogeneity such as the ones comprising present-day agricultural and countryside ecosystems.

Across taxa, habitat loss has consistently been found to have a strong negative impact on species persistence whereas the effects of habitat fragmentation per se appear to be weaker
435 and more variable, both in terms of magnitude and direction (Fahrig 2013). This general

pattern has also been repeatedly observed in tropical bat assemblages (Gorresen and Willig 2004; Meyer and Kalko 2008; Struebig et al 2008; Klingbeil and Willig 2009; Arroyo-Rodríguez et al 2016, in press; but see Cisneros et al 2015) and held true for our study. Regardless of landscape-scale, forest cover was the best predictor of species richness, having
440 a strong positive effect, whereas the influence of configurational metrics varied in magnitude, but was consistently negative. These results mirror previous findings regarding the influence of landscape-scale forest cover in fragmented systems with an aquatic matrix (Meyer and Kalko 2008; Henry et al 2010), however, contrast with results from unflooded rainforest in Paraguay, Peru, Costa Rica and Mexico, where species richness was highest in partly
445 deforested landscapes (Gorresen and Willig 2004; Klingbeil and Willig 2009; Cisneros et al 2015; Arroyo-Rodríguez et al 2016, in press). These opposing results may reflect an interaction between regional species pools and landscape-specific environmental filters, especially the ones associated with the nature of the matrix habitats in which fragments are embedded. The matrix at the BDFFP is relatively homogeneous, being composed almost
450 entirely of secondary forests (Laurance et al 2011). The higher compositional diversity of the humanized matrix habitats studied by Gorresen and Willig (2004), Klingbeil and Willig (2009), Cisneros et al (2015) and Arroyo-Rodríguez et al (2016, in press) is probably associated with a greater variety of resources, which may consequently augment species diversity in moderately fragmented landscapes. These results highlight that the influence of
455 matrix habitats on bat assemblages in forest fragments is highly context-specific (Meyer et al 2016), and are in line with previous findings that some agricultural habitats such as shade plantations can support a higher number of bat species than secondary forests (Faria 2006).

Edge density was positively correlated with total abundance and the abundances of both frugivores and gleaning animalivores at the smallest spatial scales, whereas the direction

460 of the effect was negative at larger scales. This pattern matches recent findings from temperate areas, suggesting that at smaller scales edges may translate into increased foraging opportunities and promote connectivity between roosting and foraging areas, whereas at larger scales higher edge density implies increased habitat fragmentation and therefore negative effects on bat assemblages (Kalda et al 2015). Our modelling results regarding the response of gleaning animalivores to forest cover are also congruent with previous evidence 465 that this ensemble is more sensitive to habitat modification than frugivores (Meyer et al 2016). However, contrary to our expectation, the associations of gleaning animalivores to configurational metrics were very similar to the ones observed for frugivorous bats. This indicates that the secondary regrowth habitats surrounding the BDFFP fragments may be 470 buffering the impacts of forest fragmentation on these matrix-sensitive bats and suggests that fragment connectivity is of the utmost importance for the persistence of forest-associated species in modified landscapes.

Conclusions

475 The observed effects of fragment area on bat assemblages in the adjacent matrix highlights the importance of larger (> 10 ha) forest patches in the conservation of bat diversity, and in the regeneration and ecological recovery of anthropogenically disturbed forest habitats. Our results also emphasize that, although forest fragments and secondary forest habitats can be of conservation significance for tropical bats, old-growth forest is of 480 irreplaceable value, adding to an increasing body of evidence that tropical biodiversity is overwhelmingly dependent on the maintenance of vast tracts of primary habitat (Barlow et al 2007; Gibson et al 2011).

Fragmentation effects operate at multiple spatial scales and consequently the relative influence of local- and landscape-scale attributes on tropical biota can only be better understood through a multi-scale analysis as presented here. Considering multiple spatial scales can bridge apparently contradictory results of landscape features influencing assemblages differently at distinct spatial scales and therefore greatly benefit the successful delineation of landscape-level management actions aimed at abating the wave of habitat loss and fragmentation currently eroding the biodiversity of our planet's tropical regions.

490

Acknowledgments

We would like to thank the many volunteers and field assistants that helped during fieldwork, Tobias Jeppsson for providing a modified version of the hier.part function for the hierarchical partitioning analysis, and the BDFFP management team for logistic support. Funding was provided by a Portuguese Foundation for Science and Technology (FCT) project grant (PTDC/BIA-BIC/111184/2009). R.R. was supported by FCT (SFRH/BD/80488/2011), A.L.-B. by (FCT PD/BD/52597/2014) and CNPq (160049/2013-0), P.E.D.B. by CAPES and M.C. by Academy of Finland (grant #257686). Research was conducted under ICMBio permit (26877-2) and is publication XXX in the BDFFP technical series.

500

Figure Captions

Fig. 1 Comparison of rarefied species richness and evenness (mean \pm SE) across the interior-edge-matrix as well as fragment-size gradient. Asterisks denote significant differences relative to continuous forest interiors (*** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$). See Table S4 for full results of multiple pairwise comparisons.

Fig. 2 Mean (\pm SE) capture rate (bats/mnh) for interior, edge and matrix sites of continuous forest and forest fragments. Asterisks denote significant differences relative to continuous forest interiors (*** $P < 0.001$). See Table S5 for full results of multiple pairwise comparisons.

Fig. 3 Arrangement of the 39 sampling sites along the axes of a nonmetric multidimensional scaling (NMDS) ordination based on Bray-Curtis dissimilarity.

Fig. 4 Mean (\pm SE) capture rate (bats/mnh) for total frugivores, shrub frugivores, canopy frugivores, gleaning animalivores, aerial insectivores (*Pteronotus parnellii*) and nectarivores for the interior, edge and matrix habitats of continuous forest and forest fragments. Asterisks
505 denote significant differences relative to continuous forest interiors (*** $P < 0.001$; * $P < 0.05$). See Table S7 for full results of multiple pairwise comparisons.

Fig. 5 Summary results of model averaging of the most parsimonious generalized linear mixed models (Akaike differences < 2 from the best model) investigating the relationship
510 between local and landscape-scale attributes and various response metrics (species richness, evenness and total abundance) at five focal scales across the BDFFP, Central Amazon, Brazil. Symbol size is proportional to the variation explained by the respective predictor variable based on hierarchical partitioning and colour denotes the direction of the relationship: black = positive, white = negative. See Tables S8 and S9 for additional modelling result.

515

Fig. 6 Summary results of model averaging of the most parsimonious generalized linear mixed models (Akaike differences < 2 from the best model) investigating the relationship
between local and landscape-scale attributes and the abundance of frugivores and gleaning
animalivores at five focal scales across the BDFFP, Central Amazon, Brazil. Symbol size is
520 proportional to the variation explained by the respective predictor variable based on hierarchical partitioning and colour denotes the direction of the relationship: black = positive, white = negative. See Tables S8 and S9 for additional modelling result.

Figures




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

References

- 530 Antongiovanni M, Metzger JP (2005) Influence of matrix habitats on the occurrence of insectivorous bird species in Amazonian forest fragments. *Biol Conserv* 122:441-451
- Arroyo-Rodríguez V, Rojas C, Saldaña-Vázquez RA, Stoner KE (2016) Landscape composition is more important than landscape configuration for phyllostomid bat assemblages in a fragmented biodiversity hotspot. *Biol Conserv* (in press)
- 535 Avila-Cabadilla LD, Stoner KE, Henry M, Añorve MYA (2009) Composition, structure and diversity of phyllostomid bat assemblages in different successional stages of a tropical dry forest. *Forest Ecol Manag* 258:986-996
- Avila-Cabadilla LD, Sanchez-Azofeifa GA, Stoner KE, Alvarez-Anorve MY, Quesada M, Portillo-Quintero CA (2012) Local and landscape factors determining occurrence of
540 phyllostomid bats in tropical secondary forests. *PLoS ONE* 7:e35228
- Avila-Cabadilla LD, Stoner KE, Nassar JM, Espírito-Santo MM, Alvarez-Añorve MY, Aranguren CI, Henry M, González-Carcacia JA, Falcão LAD, Sanchez-Azofeifa GA (2014) Phyllostomid bat occurrence in successional stages of Neotropical dry forests. *PLoS ONE* 9:e84572
- 545 Banks-Leite C, Ewers RM, Metzger JP (2010) Edge effects as the principal cause of area effects on birds in fragmented secondary forest. *Oikos* 119:918-926
- Barlow J, Gardner TA, Araujo IS, Ávila-Pires TC, Bonaldo AB, Costa JE, Esposito MC, Ferreira LV, Hawes J, Hernandez MIM, Hoogmoed MS, Leite RN, Lo-Man-Hung NF, Malcolm JR, Martins MB, Mestre LAM, Miranda-Santos R, Nunes-Gutjahr AL, Overall
550 WL, Parry L, Peters SL, Ribeiro-Junior MA, da Silva MNF, da Silva Motta C, Peres CA

- (2007) Quantifying the biodiversity value of tropical primary, secondary, and plantation forests. *Proc Natl Acad Sci USA* 104:18555–18560
- Bates DM (2010) *lme4: mixed-effects modeling with R*. Springer, New York. <http://lme4.r-forge.r-project.org/book/>
- 555 Benchimol M, Peres CA (2015) Edge-mediated compositional and functional decay of tree assemblages in Amazonian forest islands after 26 years of isolation. *Journal of Ecology* 103:408-420
- Benchimol M, Venticinque EM (2014) Responses of primates to landscape change in Amazonian land-bridge islands—a multi-scale analysis. *Biotropica* 46:470-478
- 560 Bernard E (2001) Vertical stratification of bat communities in primary forests of Central Amazon, Brazil. *J Trop Ecol* 17:115-126
- Bernard E (2002) Diet, activity and reproduction of bat species (Mammalia, Chiroptera) in Central Amazonia, Brazil. *Rev Bras Zool* 19:173-188
- Bobrowiec P, Gribel R (2010) Effects of different secondary vegetation types on bat 565 community composition in Central Amazonia, Brazil. *Anim Conserv* 13:204-216
- Bolívar-Cimé B, Laborde J, MacSwiney G MC, Muñoz-Robles C, Tun-Garrido J (2013) Response of phytophagous bats to patch quality and landscape attributes in fragmented tropical semi-deciduous forest. *Acta Chirop* 15:399-409
- Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White J-SS 570 (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends Ecol Evolut* 24:127-135

- Boyle SA, Smith AT (2010) Can landscape and species characteristics predict primate presence in forest fragments in the Brazilian Amazon? *Biol Conserv* 143:1134-1143
- 575 Bradshaw CJA, Sodhi NS, Brook BW (2008) Tropical turmoil: a biodiversity tragedy in progress. *Front Ecol Environ* 7:79-87
- Bregman TP, Sekercioglu CH, Tobias JA (2014) Global patterns and predictors of bird species responses to forest fragmentation: Implications for ecosystem function and conservation. *Biol Conserv* 169:372–383
- 580 Broadbent EN, Asner GP, Keller M, Knapp DE, Oliveira PJC, Silva JN (2008) Forest fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biol Conserv* 141:1745-1757
- Burnham KP, Anderson DR (2002) *Model selection and inference: a practical information-theoretic approach*, 2nd edn. Springer, New York
- 585 Carreiras JMB, Pereira JMC, Campagnolo ML, Shimabukuro YE (2006) Assessing the extent of agriculture/pasture and secondary succession forest in the Brazilian Legal Amazon using SPOT VEGETATION data. *Remote Sens Environ* 101:283-298
- Chambers CL, Cushman SA, Medina-Fitoria A, Martínez-Fonseca J, Chávez-Velásquez M (2016) Influences of scale on bat habitat relationships in a forested landscape in Nicaragua. *Landscape Ecol* (in press)
- 590 Charbonnier Y, Gaüzère P, van Halder I, Nezan J, Barnagaud JY, Jactel H, Barbaro L (2016) Deciduous trees increase bat diversity at stand and landscape scales in mosaic pine plantations. *Landscape Ecol* 31:291-300

- Chazdon RL (2014) *Second growth: The promise of tropical forest regeneration in an age of deforestation*. University of Chicago Press, Chicago
- 595 Cisneros LM, Fagan ME, Willig MR (2015) Effects of human-modified landscapes on taxonomic, functional and phylogenetic dimensions of bat biodiversity. *Divers. Distrib* 5:523-533
- Cosson J-F, Pons J-M, Masson D (1999) Effects of forest fragmentation on frugivorous and nectarivorous bats in French Guiana. *J Trop Ecol* 15:515-534
- 600 Delaval M, Charles-Dominique P (2006) Edge effects on frugivorous and nectarivorous bat communities in a neotropical primary forest in French Guiana. *Rev Ecol-Terre Vie* 61:343-352
- Didham RK, Lawton JH (1999) Edge structure determines the magnitude of changes in microclimate and vegetation structure in tropical forest fragments. *Biotropica* 31:17-30
- 605 Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJ, Collen B (2014) Defaunation in the Anthropocene. *Science* 345:401-406
- Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Marquéz JRG, Gruber B, Lafourcade B, Leitão PJ, Münkemüller T, McClean C, Osborne PE, Reineking B, Schröder B, Skidmore AK, Zurell D, Lautenbach S (2013) Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36:027-046
- 610 Erickson JL, West SD (2003) Associations of bats with local structure and landscape features of forested stands in western Oregon and Washington. *Biol Conserv* 109:95-102
- Estrada-Villegas S, Meyer CF, Kalko EK (2010) Effects of tropical forest fragmentation on aerial insectivorous bats in a land-bridge island system. *Biol Conserv* 143:597-608

- 615 Ewers RM, Didham RK (2006) Confounding factors in the detection of species responses to
habitat fragmentation. *Biol Rev* 81:117-142
- Fahrig L (2013) Rethinking patch size and isolation effects: the habitat amount hypothesis. *J
Biogeogr* 40:1649-1663
- Faria D (2006) Phyllostomid bats of a fragmented landscape in the north-eastern Atlantic
620 forest, Brazil. *J Trop Ecol* 22:531-542
- Faria D, Mariano-Neto E, Martini AMZ, Ortiz JV, Montingelli R, Rosso S, Paciencia MLB,
Baumgarten J (2009) Forest structure in a mosaic of rainforest sites: The effect of
fragmentation and recovery after clear cut. *Forest Ecol Manag* 257:2226-2234
- Farneda FZ, Rocha R, López-Baucells A, Groenenberg M, Silva I, Palmeirim JM, Bobrowiec
625 PED, Meyer CFJ (2015) Trait-related responses to habitat fragmentation in Amazonian
bats. *J Appl Ecol* 52:1381-1391
- Ferraz G, Nichols JD, Hines JE, Stouffer PC, Bierregaard RO, Lovejoy TE (2007) A large-
scale deforestation experiment: effects of patch area and isolation on amazon birds.
Science 315:238-241
- 630 Fischer J, Lindenmayer DB (2007) Landscape modification and habitat fragmentation: a
synthesis. *Global Ecol Biogeogr* 16:265-280
- García-Morales R, Badano EI, Moreno CE (2013) Response of Neotropical bat assemblages
to human land use. *Conserv Biol* 27:1096-1106
- Gardner A (2007) *Mammals of South America, Vol. 1: Marsupials, xenarthrans, shrews, and*
635 *bats*. The University of Chicago Press, Chicago

- Gascon C, Lovejoy TE, Bierregaard Jr RO, Malcolm JR, Stouffer PC, Vasconcelos HL, Laurance WF, Zimmerman B, Tocher M, Borges S (1999) Matrix habitat and species richness in tropical forest remnants. *Biol Conserv* 91:223-229
- Galitsky C, Lawler JJ (2015) Relative influence of local and landscape factors on bird
640 communities vary by species and functional group. *Landscape Ecol* 30:287-299
- Giannini NP, Kalko EKV (2004) Trophic structure in a large assemblage of phyllostomid bats in Panama. *Oikos* 105:209-220
- Gibson L, Lee TM, Koh LP, Brook BW, Gardner TA, Barlow J, Peres CA, Bradshaw CJ, Laurance WF, Lovejoy TE (2011) Primary forests are irreplaceable for sustaining tropical
645 biodiversity. *Nature* 478:378-381
- Gorresen PM, Willig MR (2004) Landscape responses of bats to habitat fragmentation in Atlantic forest of Paraguay. *J Mammal* 85:688-697
- Gotelli NJ, Entsminger GL (2004) EcoSim: Null models software for ecology. Version 7. Acquired Intelligence Inc. & Kesey-Bear. Jericho, VT 05465.
650 <http://garyentsminger.com/ecosim/index.htm>
- Henry M, Cosson JF, Pons JM (2010) Modelling multi-scale spatial variation in species richness from abundance data in a complex neotropical bat assemblage. *Ecol Model* 221:2018-2027
- Hothorn T, Bretz F, Westfall P, Heiberger RM, Schuetzenmeister A, Scheibe S (2014)
655 multcomp: simultaneous inference in general parametric models. R package version 1.3-2.
<http://cran.r-project.org/web/packages/multcomp/>

- Jeppsson T, Lindhe A, Gärdenfors U, Forslund P (2010) The use of historical collections to estimate population trends: a case study using Swedish longhorn beetles (Coleoptera: Cerambycidae). *Biol Conserv* 143:1940-1950
- 660 Jones G, Jacobs DS, Kunz TH, Willig MR, Racey PA (2009) Carpe noctem: the importance of bats as bioindicators. *Endanger Species Res* 8:93-115
- Kalda R, Kalda O, Lõhmus K, Liira J (2015) Multi-scale ecology of woodland bat the role of species pool, landscape complexity and stand structure. *Biodivers Conserv* 24:337-353
- Kalko EKV (1998) Organisation and diversity of tropical bat communities through space and
665 time. *Zoology* 101:281-297
- Klingbeil BT, Willig MR (2009) Guild-specific responses of bats to landscape composition and configuration in fragmented Amazonian rainforest. *J Appl Ecol* 46:203-213
- Klingbeil BT, Willig MR (2010) Seasonal differences in population-, ensemble- and
community-level responses of bats to landscape structure in Amazonia. *Oikos* 119:1654-
670 1664
- Kunz TH, Braun de Torrez E, Bauer D, Lobova T, Fleming TH (2011) Ecosystem services provided by bats. *Ann N Y Acad Sci* 1223:1-38
- Laurance WF, Camargo JL, Luizão RC, Laurance SG, Pimm SL, Bruna EM, Stouffer PC, Williamson GB, Benítez-Malvido J, Vasconcelos HL (2011) The fate of Amazonian forest
675 fragments: a 32-year investigation. *Biol Conserv* 144:56-67
- Laurance WF, Lovejoy TE, Vasconcelos HL, Bruna EM, Didham RK, Stouffer PC, Gascon C, Bierregaard RO, Laurance SG, Sampaio E (2002) Ecosystem decay of Amazonian forest fragments: a 22-year investigation. *Conserv Biol* 16:605-618

- Laurance WF, Nascimento HE, Laurance SG, Andrade AC, Fearnside PM, Ribeiro JE,
680 Capretz RL (2006a) Rain forest fragmentation and the proliferation of successional trees.
Ecology 87:469-482
- Laurance WF, Nascimento HE, Laurance SG, Andrade AC, Ribeiro JE, Giraldo JP, Lovejoy
TE, Condit R, Chave J, Harms KE (2006b) Rapid decay of tree-community composition in
Amazonian forest fragments. Proc Natl Acad Sci USA 103:19010-19014
- 685 Laurance WF, Sayer J, Cassman KG (2014) Agricultural expansion and its impacts on
tropical nature. Trends Ecol Evolut 29:107-116
- Lenz BB, Jack KM, Spironello WR (2014) Edge effects in the primate community of the
biological dynamics of forest fragments project, Amazonas, Brazil. Am J Phys Anthropol
155:436-446
- 690 Lim BK, Engstrom MD (2010) Species diversity of bats (Mammalia: Chiroptera) in
Iwokrama Forest, Guyana, and the Guianan subregion: implications for conservation.
Biodivers Conserv 10:613-657
- MacArthur RH, Wilson EO (1967) The Theory of Island Biogeography. Princeton University
Press, Princeton
- 695 Marciente R, Bobrowiec PED, Magnusson WE (2015) Ground-Vegetation Clutter Affects
Phyllostomid Bat Assemblage Structure in Lowland Amazonian Forest. PLoS
ONE 10:e0129560
- Marques JT, Ramos Pereira MJ, Marques TA, Santos CD, Santana J, Beja P, Palmeirim JM
(2013) Optimizing sampling design to deal with mist-net avoidance in Amazonian birds
700 and bats. PLoS ONE 8:e74505

- McGarigal K, Cushman SA, Ene E (2012) FRAGSTATS v4: spatial pattern analysis program for categorical and continuous maps. University of Massachusetts, Amherst, MA, USA.
<http://www.umass.edu/landeco/research/fragstats/fragstats.html>
- McGill BJ (2010) Matters of scale. *Science* 328:575-576.
- 705 Mendenhall CD, Karp DS, Meyer CF, Hadly EA, Daily GC (2014) Predicting biodiversity change and averting collapse in agricultural landscapes. *Nature* 509:213-217
- Mesquita RCG, Ickes K, Ganade G, Williamson GB (2001) Alternative successional pathways in the Amazon Basin. *J Ecol* 89:528-537
- Meyer CFJ, Struebig M, Willig MR (2016) Responses of tropical bats to habitat
710 fragmentation, logging, and deforestation. In: Voigt CC and Kingston T (eds) *Bats in the Anthropocene: Conservation of bats in a changing world*. Springer, New York. pp. 63-103
- Meyer CFJ, Kalko EKV (2008) Assemblage-level responses of phyllostomid bats to tropical forest fragmentation: land-bridge islands as a model system. *J Biogeogr* 35:1711-1726
- Meyer CFJ, Freund J, Lizano WP, Kalko EKV (2008) Ecological correlates of vulnerability to
715 fragmentation in Neotropical bats. *J Appl Ecol* 45:381-391
- Mokross K, Ryder TB, Côrtes MC, Wolfe JD, Stouffer PC (2014) Decay of interspecific avian flock networks along a disturbance gradient in Amazonia. *Proc R Soc Lond Ser B Biol Sci* 281:20132599
- Nakagawa S, Schielzeth H (2013) A general and simple method for obtaining R^2 from
720 generalized linear mixed-effects models. *Methods Ecol Evol* 4:133-142
- Neter J, Wasserman W, Kutner MH (1990) *Applied linear statistical models*. Irwin, New Jersey

- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Wagner H (2015) Vegan: community ecology. R package version 2.2-1. <http://cran.r-project.org/web/packages/vegan/>
- 725
- Palmeirim JM, Etheridge K (1985) The influence of man-made trails on foraging by tropical frugivorous bats. *Biotropica* 17:82-83
- Pereira MJR, Marques JT, Palmeirim JM (2010) Vertical stratification of bat assemblages in flooded and unflooded Amazonian forests. *Curr Zool* 56:469-478
- 730
- Powell LL, Zurita G, Wolfe JD, Johnson EI, Stouffer PC (2015) Changes in habitat use at rain forest edges through succession: a case study of understory birds in the Brazilian Amazon. *Biotropica* 47:723-732
- Powell LL, Stouffer PC, Johnson EI (2013) Recovery of understory bird movement across the interface of primary and secondary Amazon rainforest. *The Auk* 130:459-468
- 735
- R Development Core Team (2013) R: a Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria
- Rocha R, Tarmo V, Cabeza M (2015) Bird assemblages in a Malagasy forest-agricultural frontier: effects of habitat structure and landscape-scale forest cover. *Trop Conserv Sci* 8:681-710
- 740
- Schulze MD, Seavy NE, Whitacre DF (2000) A comparison of the phyllostomid bat assemblages in undisturbed Neotropical forest and in forest fragments of a slash-and-burn farming mosaic in Petén, Guatemala. *Biotropica* 32:174-184

- Smart SM, Thompson K, Marrs RH, Le Duc MG, Maskell LC, Firbank LG (2006) Biotic homogenization and changes in species diversity across human-modified ecosystems. Proc R Soc Lond Ser B Biol Sci 273:2659-2665
- 745
- Stouffer PC, Bierregaard RO, Strong C, Lovejoy TE (2006) Long-term landscape change and bird abundance in Amazonian rainforest fragments. Conserv Biol 20:1212-1223
- Stratford JA, Stouffer PC (2013) Microhabitat associations of terrestrial insectivorous birds in Amazonian rainforest and second-growth forests. J Field Ornithol 84:1-12
- 750
- Struebig MJ, Kingston T, Zubaid A, Mohd-Adnan A, Rossiter SJ (2008) Conservation value of forest fragments to Palaeotropical bats. Biol Conserv 141:2112-2126
- Walsh C, Mac Nally R (2013) Hier.part: variance partition of a multivariate data set. R package version 1.0-4. <http://cran.r-project.org/web/packages/hier.part/>
- Williamson GB, Bentes TV, Longworth JB, Mesquita RCG (2014) Convergence and divergence in alternative successional pathways in Central Amazonia. Plant Ecol Divers 7:341
- 755