

CONSTRUCTING GRAPHS WHICH ARE $1/2$ -TRANSITIVE

BRIAN ALSPÁCH, DRAGAN MARUŠIČ and LEWIS NOWITZ

(Received 26 July 1991; revised 5 February 1992)

Communicated by L. Caccetta

Abstract

An infinite family of vertex- and edge-transitive, but not arc-transitive, graphs of degree 4 is constructed.

1991 *Mathematics subject classification* (*Amer. Math. Soc.*): 05 C 25.

1. Introduction

If G is a graph, then the *arcs* of G are obtained by taking one arc for each orientation of each edge of G so that there are twice as many arcs as edges. A k -arc of G is a directed walk of length k using the arcs of G except that one cannot follow an arc (u, v) by the arc (v, u) in the directed walk. A k -path P of G with end vertices u and v is a walk (undirected) of length k such that each of u and v is incident with one edge of P , and all other vertices of P are incident with two edges of P . A graph G is said to be *vertex-transitive*, *edge-transitive* and *arc-transitive* provided its automorphism group $\text{Aut}(G)$ acts transitively on the vertices, edges and arcs of G , respectively. (The terms *symmetric* and *1-transitive* also have been used instead of arc-transitive.) In general, a graph is said to be *k-arc-transitive* if $\text{Aut}(G)$ acts transitively on the k -arcs of G . Biggs [3] defines a graph G to be *k-transitive* if $\text{Aut}(G)$ acts transitively on the k -arcs of G , but does not act transitively on the $(k + 1)$ -arcs. Since we

© 1994 Australian Mathematical Society 0263-6115/94 \$A2.00 + 0.00

This research was partially supported by the Natural Sciences and Engineering Research Council of Canada under Grant A-4792

deal with graphs G for which $\text{Aut}(G)$ is vertex- and edge-transitive but not arc-transitive and we wish to avoid that cumbersome phrase, in keeping with Biggs' definition (and viewing the vertices of G as 0-arcs), we shall call such graphs *1/2-transitive*. In general, a graph G is $(2m + 1)/2$ -transitive if $\text{Aut}(G)$ acts transitively on the m -arcs and the $(m + 1)$ -paths of G , but does not act transitively on the $(m + 1)$ -arcs of G .

Of course, there exist vertex-transitive graphs which are not edge-transitive. Likewise, an edge-transitive graph is not necessarily regular, and thus not necessarily vertex-transitive. Even more, there are regular edge-transitive graphs which are not vertex-transitive [6]. The only general result linking vertex- and edge-transitivity to arc-transitivity is due to Tutte [15] who proved that a vertex- and edge-transitive graph of odd degree is necessarily arc-transitive. With regard to even degree, Bower [4] in 1970 constructed an infinite family of $1/2$ -transitive graphs. The smallest order graph in his family has 54 vertices. More recently, Holt [7] found one with 27 vertices.

The automorphism groups of Holt's graph and Bower's family of graphs act imprimitively on the vertex-set. This prompted Holton [8] to ask if the automorphism group of every $1/2$ -transitive graph is necessarily imprimitive. This question has been answered by Xu and Praeger [12] in the negative. They found several examples of $1/2$ -transitive graphs, the smallest of which has 253 vertices and is of degree 24, whose automorphism groups are primitive.

As mentioned above, Holt has produced a $1/2$ -transitive graph on 27 vertices. Using the facts that every vertex- and edge-transitive Cayley graph on an abelian group or with $2p$ vertices, p a prime, is also arc-transitive, McKay's list of all vertex-transitive graphs with 19 or fewer vertices [10], McKay and Royle's list of vertex-transitive graphs with 20 and 21 vertices [11], and Praeger and Royle's proof that there are no $1/2$ -transitive graphs with 24 vertices, we conclude that there are no $1/2$ -transitive graphs with fewer than 27 vertices. In his paper, Holt mentions that a referee informed him that Kornya found another example with 27 vertices. However, Xu has informed the authors (personal communication) that he has shown that Holt's graph is the only $1/2$ -transitive graph with 27 vertices and of degree 4.

The objective of this paper is to give an infinite family of $1/2$ -transitive graphs of degree 4. All of them are metacirculant graphs.

2. A family of graphs

Let $n \geq 2$. A permutation on a finite set is said to be (m, n) -semiregular if it has m cycles of length n in its disjoint cycle decomposition. We shall be sloppy and refer to the orbits of the group $\langle \alpha \rangle$ generated by α as the orbits of α . A graph G is an (m, n) -metacirculant if it has an (m, n) -semiregular automorphism α together with another automorphism β normalizing α and cyclically permuting the orbits of α such that as a permutation there is a cycle of length m in the disjoint cycle decomposition of β . Therefore, we may partition the vertex-set of an (m, n) -metacirculant into the orbits X_0, X_1, \dots, X_{m-1} of α , where $X_{i+1} = \beta(X_i)$ for all $i \in Z_m$. We shall refer to the orbits of α as the *blocks* of the metacirculant graph. It should be pointed out that the blocks of a metacirculant graph need not be blocks of imprimitivity of the automorphism group of the graph.

If G is a graph and A and B are two vertex-disjoint subsets of the vertex-set $V(G)$ of G , we let $\langle A \rangle$ denote the subgraph induced by G on A , and let $\langle A, B \rangle$ denote the bipartite subgraph induced by G on $A \cup B$, that is, all edges of G with one endvertex in A and the other endvertex in B . If $S_0 \subseteq Z_n \setminus \{0\}$ is the symbol of the subcirculant $\langle X_0 \rangle$ and, for all $i \in Z_m \setminus \{0\}$, $T_i \subseteq Z_m$ is the symbol of the bipartite subgraph $\langle X_0, X_i \rangle$, then there exists an $r \in Z_m^*$, where Z_m^* denotes the multiplicative group of units in Z_m , such that for all $j \in Z_m$, the symbol of $\langle X_j \rangle$ is $r^j S_0$ and the symbol of the bipartite graph $\langle X_j, X_{j+1} \rangle$, $i \in Z_m$, is $r^j T_i$. Moreover, for all $i \in Z_m$, we have $T_{m-i} = -r^{m-i} T_i$. Thus, the metacirculant graph is completely determined by the $\lfloor (m+4)/2 \rfloor$ -tuple $(r; S_0, T_1, T_2, \dots, T_{\lfloor m/2 \rfloor})$ which is called a *symbol* of G . (For a more detailed discussion of metacirculants, the reader is referred to [1, 2].)

We are now ready to define a family of metacirculants of degree 4 containing infinitely many 1/2-transitive graphs. The smallest among them has 27 vertices.

Let $n \geq 5$ be an integer and $r \in Z_n^*$ be an element of order m or $2m$ such that $m \geq 3$. Then let $M(r; m, n)$ denote the (m, n) -metacirculant graph with symbol $(r; \emptyset, \{1, n-1\}, \emptyset, \dots, \emptyset)$. For example, the graph $M(2; 3, 9)$ can be thought of as having vertices $\{x_i^j : 0 \leq i \leq 2 \text{ and } 0 \leq j \leq 8\}$ with x_0^j adjacent to $x_1^{j \pm 1}$, x_1^j adjacent to $x_2^{j \pm 2}$, and x_2^j adjacent to $x_0^{j \pm 4}$, where the superscripts are reduced modulo 9. This graph is in fact the Holt graph of [7]. This suggests there may be other 1/2-transitive $M(r; m, n)$ graphs, as indeed is the case, for other values of the parameters r, m and n .

Throughout the paper, α is the (m, n) -semiregular automorphism of $M = M(r; m, n)$ with orbits X_0, X_1, \dots, X_{m-1} , where $X_i = \{x_i^0, x_i^1, \dots, x_i^{n-1}\}$ and

$\alpha(x_i^j) = x_i^{j+1}$ for all $i \in Z_m$ and $j \in Z_n$. Similarly, β is the automorphism of M mapping according to the formula $\beta(x_i^j) = x_{i+1}^j$, for $i \in Z_m, j \in Z_n$. Finally, another automorphism τ of $M(r; m, n)$ which will prove useful is defined by $\tau(x_i^j) = x_i^{-j}$.

DEFINITION. A cycle of $M(r; m, n)$ of length at least m is said to be *coiled* if every subpath with m vertices intersects each one of X_0, X_1, \dots, X_{m-1} . It is easy to see that a coiled cycle must have length a multiple of m .

DEFINITION. The *coiled girth* of $M(r; m, n)$ is the length of a shortest coiled cycle in $M(r; m, n)$.

PROPOSITION 2.1. *The coiled girth of $M(r; m, n)$ is either m or $2m$.*

PROOF. Assume that $M(r; m, n)$ does not contain a coiled cycle of length m . Consider the closed trail

$$x_0^0 x_1^1 x_2^{1+r} \dots x_{m-1}^{1+r+r^2+\dots+r^{m-2}} x_0^{1+r+r^2+\dots+r^{m-1}} x_1^{r+r^2+\dots+r^{m-1}} x_2^{r^2+\dots+r^{m-1}} \dots x_0^0.$$

Since $M(r; m, n)$ has no coiled cycles of length m , all of the vertices of the closed trail are distinct. Thus, it is a coiled cycle of length $2m$ as required.

DEFINITION. We obtain natural edge-partitions of $M(r; m, n)$ using coiled girth cycles and α . If $M(r; m, n)$ has coiled girth m , let C be a coiled cycle of length m . If $C = x_0^{i_0} x_1^{i_1} \dots x_{m-1}^{i_{m-1}}$, let $\alpha(C)$ denote the cycle obtained under the substitution of $\alpha(x_j^{i_j})$ for $x_j^{i_j}, 0 \leq j \leq m-1$. Then $C, \alpha(C), \alpha^2(C), \dots, \alpha^{n-1}(C)$ is a 2-factor of $M(r; m, n)$. It is easy to see that the remaining edges also form a 2-factor made up of coiled cycles of length m . If C is a coiled cycle of length $2m$ such that of the two edges from X_j to X_{j+1} in C , one is of the form $x_j^{i_j} x_{j+1}^{i_j+r^j}$ and the other is of the form $x_j^{i_j} x_{j+1}^{i_j-r^j}$, then $C, \alpha(C), \alpha^2(C), \dots, \alpha^{n-1}(C)$ is a partition of the edge-set of $M(r; m, n)$ into $2m$ -cycles. We call both of the above the α -partition of $M(r; m, n)$ induced by C .

DEFINITION. When $M(r; m, n)$ has coiled girth m and the α -partition of $M(r; m, n)$ induced by every coiled m -cycle yields the same 2-factorization of $M(r; m, n)$, then we shall say that $M(r; m, n)$ is *tightly coiled*. Otherwise, we shall say that the graph is *loosely coiled*. A similar definition applies in the case that the coiled girth of $M(r; m, n)$ is $2m$. However, in the latter case, $M(r; m, n)$ is always loosely coiled as we shall see soon.

We now set about proving some lemmas which will be useful in establishing that certain $M(r; m, n)$ metacirculants are 1/2-transitive.

LEMMA 2.2. *If $M = M(r; m, n)$ has coiled girth $2m$, then M is loosely coiled.*

PROOF. As seen in the proof of Proposition 2.1,

$$x_0^0 x_1^1 x_2^{1+r} \dots x_{m-1}^{1+r+r^2+\dots+r^{m-2}} x_0^{1+r+r^2+\dots+r^{m-1}} x_1^{r+r^2+\dots+r^{m-1}} x_2^{r^2+\dots+r^{m-1}} \dots x_0^0$$

is a coiled cycle C of length $2m$. Obtain another coiled cycle of length $2m$ by taking the first edge from x_0^0 to x_1^{n-1} and then taking the same kind of edge as in C until reaching X_0 again. This means the vertex in X_j will be $x_j^{-1+r+r^2+\dots+r^{j-1}}$. When leaving X_0 the second time, use the edge to $x_1^{r+r^2+\dots+r^{m-1}}$ from which point the vertices will be the same as in C .

LEMMA 2.3. *Let $M = M(r; m, n)$. If $\sigma \in \text{Aut}(M)$ fixes two adjacent blocks of M pointwise, then σ is the identity.*

PROOF. Let σ fix X_i and X_{i+1} pointwise. Then X_{i-1} is fixed setwise by σ . The subgraph $\langle X_{i-1}, X_i \rangle$ is either a $2n$ -cycle or two n -cycles because $r \in Z_n^*$. The automorphism σ fixes alternate vertices of the $2n$ -cycle or the two n -cycles, and thus it fixes every vertex. This means that σ also fixes X_{i-1} pointwise. Continuing in this way establishes the result.

LEMMA 2.4. *Let $M = M(r; m, n)$. If $\sigma \in \text{Aut}(M)$ fixes some block of M pointwise, then σ is the identity.*

PROOF. Without loss of generality we may assume X_1 is the block of M which is fixed pointwise by σ . The neighbors of x_1^0 are x_2^r, x_2^{-r}, x_0^1 and x_0^{n-1} , and the neighbors of x_1^2 are $x_2^{r+2}, x_2^{-r+2}, x_0^1$ and x_0^3 . Since both x_1^0 and x_1^2 are fixed by σ , either x_0^1 is also fixed by σ or $\{x_2^r, x_2^{-r}, x_0^{n-1}\} \cap \{x_2^{r+2}, x_2^{-r+2}, x_0^3\}$ is non-empty. In the latter case, $x_0^{n-1} \neq x_0^3$ because $n \geq 5$. This forces either $x_2^r = x_2^{-r+2}$ or $x_2^{-r} = x_2^{r+2}$. In the first case $r = (n + 2)/2$ must hold and in the second case $r = (n - 2)/2$ must hold. Neither are possible when n is odd. If n is a multiple of 4, then $r^2 = 1$ in Z_n^* contradicting the fact that r has order at least 3. If n is even and not a multiple of 4, then r is even contradicting the fact that $r \in Z_n^*$. This implies x_0^1 is fixed by σ . Continuing in this way we obtain that X_0 is fixed pointwise. The preceding lemma yields the desired result.

LEMMA 2.5. *Let $M = M(r; m, n)$ and suppose that whenever $\sigma \in \text{Aut}(M)$ fixes two adjacent vertices of M , σ is the identity. Then either $\text{Aut}(M) = \langle \alpha, \beta, \tau \rangle$ or $|\text{Aut}(M)| = 2|\langle \alpha, \beta, \tau \rangle|$.*

PROOF. By hypothesis, the stabilizer of an edge of M is either the identity or has order 2. Thus, $|\text{Aut}(M)| = 2mn$ or $4mn$ and the result follows.

LEMMA 2.6. *Let $M = M(r; m, n)$, with m and n odd, have coiled girth m and be loosely coiled. If $\sigma \in \text{Aut}(M)$ fixes two adjacent vertices of M , then σ is the identity.*

PROOF. Suppose $x \in X_i$ and $y \in X_{i+1}$ are two adjacent vertices of M fixed by σ . Let y' be the other neighbor of x in X_{i+1} and let z and z' be the two neighbors of x in X_{i-1} . Since m is odd, neither of the two triples zxz' and xyy' are in m -cycles. On the other hand, since M is loosely coiled, each of the four triples zxy , zxy' , $z'xy$ and $z'xy'$ are in m -cycles. Therefore, σ must also fix y' in addition to fixing x and y . For the same reason, σ must fix the other neighbor of y' in X_i . Continuing in this way, we see that σ fixes all the vertices of X_i and X_{i+1} . By Lemma 2.3, the conclusion follows.

THEOREM 2.7. *Let $M = M(r; m, n)$, with m and n odd, have coiled girth m . If M is loosely coiled, $n > 7$ and $m \geq 3$, then M is 1/2-transitive.*

PROOF. If $\text{Aut}(M) = \langle \alpha, \beta, \tau \rangle$, then M is 1/2-transitive. Assume $\text{Aut}(M) \neq \langle \alpha, \beta, \tau \rangle$ and M is arc-transitive. Then there is a $\sigma \in \text{Aut}(M)$ interchanging two adjacent vertices, say $x \in X_i$ and $y \in X_{i+1}$. Let $a, a' \in X_{i-1}$ and $y' \in X_{i+1}$ be the remaining neighbors of x , and let $u, u' \in X_{i+2}$ and $x' \in X_i$ be the remaining three neighbors of y . The triples uyx' and $u'yx'$ are contained in an m -cycle but uyu' is not. Similarly, axy' and $a'xy'$ are contained in m -cycles but xyy' is not. We know σ interchanges $\{u, u', x'\}$ and $\{a, a', y'\}$ so that σ must interchange x' and y' . Hence, σ interchanges X_i and X_{i+1} which implies σ interchanges X_{i-1} and X_{i+2} , X_{i-2} and X_{i+3} , and so on. Thus, $\text{Aut}(M)$ acts imprimitively with the orbits of α as blocks. Since σ^2 fixes x and y , σ is an involution by Lemma 2.6.

Note that the automorphism group of $\langle X_i \cup X_{i+1} \rangle$ is dihedral of order $4n$. Hence, the restriction of $\sigma\alpha\sigma$ to $\langle X_i \cup X_{i+1} \rangle$ is the restriction of α^{-1} . Thus, the restriction of $(\sigma\alpha)^2$ to $\langle X_i \cup X_{i+1} \rangle$ is 1 implying that $(\sigma\alpha)^2 = 1$ by Lemma 2.1. This together with the fact that $\tau\alpha\tau = \alpha^{-1}$ implies that $\sigma\tau$ centralizes α . Clearly, the action of $\sigma\tau$ on the orbits of α is identical to the action of σ . Thus, there

is an orbit Z of α which is fixed by $\sigma\tau$. The restriction of $\sigma\tau$ to Z is then the same as that of some α^i . Let $\gamma = \sigma\tau\alpha^{-i}$. Note that γ restricted to Z is 1 and that γ interchanges the neighboring orbits, say U and W , of Z . However, the structure of the graph M implies that whenever two vertices of Z have a common neighbor in U , they do not have a common neighbor in W . Therefore, an automorphism such as γ cannot exist. This means that $\text{Aut}(M) = \langle \alpha, \beta, \tau \rangle$ and M is 1/2-transitive.

COROLLARY 2.8. *Let p be a prime and r a divisor of $p - 1$ whose order m in Z_p^* is odd and composite. Then $M = M(r; m, p)$ is 1/2-transitive. In particular, there are infinitely many 1/2-transitive graphs of degree 4.*

PROOF. The graph M has coiled girth m because $1 + r + r^2 + \dots + r^{m-1} \equiv 0 \pmod{p}$ implying that $x_0^0 x_1^1 x_2^{1+r} \dots x_{m-1}^{1+r+\dots+r^{m-2}} x_0^0$ is an m -cycle C . It suffices to show that M is loosely coiled. Let H be the subgroup of Z_p^* of order m generated by r . Let d be a nontrivial divisor of $p - 1$. Then the subgroup H_1 of H generated by r^d has order $(p - 1)/d$. The sum of the entries in any subgroup of Z_p^* is congruent to zero modulo p . In fact, the sum of the powers of r over each of the cosets of H_1 in H is also congruent to zero modulo p . Thus, we can either add or subtract all the elements of a given coset to give us other m -cycles of M which induce 2-factorizations of M different than that induced by C . Hence, M is loosely coiled and the corollary follows from Theorem 2.7.

3. Three blocks

In the previous section, the general case for m and n odd is covered when $M(r; m, n)$ has coiled girth m and is loosely coiled. We take care of the case when $M(r; m, n)$ is tightly coiled or has coiled girth $2m$ in this section, but only for $m = 3$. The following lemma has the same conclusion as Lemma 2.6, but the proof is completely different. Notice that if $r^3 \equiv -1 \pmod{n}$, then $(-r)^3 \equiv 1 \pmod{n}$. Since $M(r; 3, n) \cong M(-r; 3, n)$, we may as well assume r has order 3.

LEMMA 3.1. *Let $M = M(r; 3, n)$, $n \geq 9$ and n odd, let $r^3 \equiv 1 \pmod{n}$, and let M have coiled girth 3. If $\sigma \in \text{Aut}(M)$ fixes two adjacent vertices of M , then σ is the identity.*

PROOF. Since $r^3 \equiv 1 \pmod{n}$, $(r - 1)(r^2 + r + 1) \equiv 0 \pmod{n}$. This implies that either $r^2 + r + 1 \equiv 0 \pmod{n}$ or $r^2 + r + 1$ is a zero divisor in Z_n . Since 2 is not a zero divisor in Z_n when n is odd, $r^2 + r + 1 \not\equiv 2 \pmod{n}$ implying that $r^2 + r - 1 \not\equiv 0 \pmod{n}$. Similarly, $r^2 - r + 1 \not\equiv 0 \pmod{n}$ because r is a unit in Z_n . Therefore, since M has coiled girth 3, either $r^2 + r + 1 \equiv 0 \pmod{n}$ or $r^2 - r - 1 \equiv 0 \pmod{n}$ must hold.

First consider the case that $r^2 + r + 1 \equiv 0 \pmod{n}$. Without loss of generality, assume that σ fixes x_0^0 and x_1^1 . We now determine the vertices at increasing distances from $\{x_0^0, x_1^1\}$ and use them to prove the result in this case. The only problem that may arise is that for small n and certain values of r some of the apparently different vertices may be the same. The vertex x_2^{r+1} is the only vertex adjacent to both x_0^0 and x_1^1 so it too is fixed by σ . The other neighbors of x_0^0 are x_2^2 and $x_1^{r+r^2}$, and the other neighbors of x_1^1 are x_0^2 and $x_2^{r^2+2}$. The vertices x_0^2 , $x_1^{r+r^2}$, x_2^{r+1} , and $x_2^{r^2+r}$ are all distinct because $n \geq 9$, n is odd and our assumption about $r^2 + r + 1$.

We now determine the vertices at distance 2 from $\{x_0^0, x_1^1\}$. The remaining neighbors of x_2^2 are $x_0^{r^2-r-1}$ and $x_1^{r^2-r}$, of $x_1^{r+r^2}$ are x_0^{n-2} and x_2^{r-1} , of x_2^{r+1} are x_0^{2r+2} and x_1^{2r+1} , of x_0^2 are x_1^3 and x_2^{r+3} , and of $x_2^{r^2+2}$ are $x_0^{r^2-r+1}$ and $x_1^{r^2-r+2}$. We now show that we may assume all these vertices are distinct and that there is only one adjacency amongst them. This is based on the following results. The element $r \neq 0$; if $r = 2$, then $r^3 \equiv 1 \pmod{n}$ implies $n = 7$ which is a contradiction; if $r = n - 2$, then $n = 9$ so that M is Holt's graph, which is known to be 1/2-transitive; $r \neq 1$; $r \neq -1$; if $r = (n - 1)/2$, then $r^2 + r + 1 \equiv 0 \pmod{n}$ implies that $r^2 \equiv r \pmod{n}$ which is impossible; if $r = (n + 1)/2$, then $r^2 \equiv (n - 3)/2 = r - 2 \pmod{n}$ which implies $r^3 \equiv (n - 5)/2 \pmod{n}$ which, in turn, implies $n = 7$; and if $r = (n - 3)/2$, then $r^2 \equiv (n + 1)/2 = r + 2 \pmod{n}$ again leading to the contradiction that $n = 7$.

Following are a few examples showing how the above occur. Can $x_0^{n-2} = x_0^{2r+2}$, that is, can $n - 2 \equiv 2r + 2 \pmod{n}$? Since n is odd, we must have $2r = 2n - 4$ or equivalently $r = n - 2$. As another example consider whether or not $x_0^{2r+2} = x_0^{r^2-r+1}$ is possible. If the two are equal, then $4r \equiv -2 \pmod{n}$. This implies that $r = (n - 1)/2$ which is one of the above. Checking all other possibilities leads to one of the above or that $n < 9$.

We now may assume that the vertices at distance 2 from $\{x_0^0, x_1^1\}$ listed above are distinct. We now consider adjacencies amongst these vertices. Notice that there is an edge joining $x_0^{r^2-r+1}$ and $x_1^{r^2-r}$. Now consider the vertex $x_0^{r^2-r+1}$ as an example. Its other neighbor in X_2 is $x_2^{r^2-2r-2}$. For example, can $x_2^{r^2-2r-2} = x_2^{r+3}$? If so, then $r + 3 \equiv r^2 - 2r - 2 \equiv -3r - 3 \pmod{n}$. This implies that $4r \equiv$

$-6 \pmod n$ which, in turn, implies that $r = (n - 3)/2$. This means the vertex $x_2^{r^2-2r-2}$ cannot lie at distance 2 from $\{x_0^0, x_1^1\}$. Checking all other possibilities leads us to be able to assume the above edge between $x_0^{r^2-r+1}$ and $x_1^{r^2-r}$ is the only edge joining any two vertices at distance 2 from $\{x_0^0, x_1^1\}$.

This implies that σ cannot interchange $x_2^{r^2+2}$ and x_0^2 since there is a path of length 4 from $x_2^{r^2+2}$ to x_0^0 , but not from x_0^2 to x_0^0 . Similarly, σ cannot interchange $x_2^{r^2}$ and x_1^{n-1} . In particular, this implies that σ fixes x_1^{n-1} and x_0^2 . Now repeat the same argument with the adjacent vertices x_0^2 and x_1^1 leading to x_1^3 being fixed by σ . Because n is odd, continuing in this way leads to σ fixing every vertex of X_0 and X_1 . By Lemma 2.3, σ is the identity. This completes the case for $r^2 + r + 1 \equiv 0 \pmod n$.

The case for $r^2 - r - 1 \equiv 0 \pmod n$ is done in the same way. The only adjacency between two vertices at distance 2 from $\{x_0^0, x_1^1\}$ is between x_0^{-2r} and x_1^{-2r-1} . A new contradiction is used in this case. Several times $-r - 3 \equiv r + 3 \pmod n$ arises. This forces $r = -3$ which implies $n = 11$ because of $r^2 - r - 1 \equiv 0 \pmod n$. But then $r^3 \not\equiv 1 \pmod{11}$ which is a contradiction.

LEMMA 3.2. *Let $M = M(r; 3, n)$, $n \geq 9$ and n odd, let $r^3 \equiv 1 \pmod n$, and let M have coiled girth 6. If $\sigma \in \text{Aut}(M)$ fixes two adjacent vertices of M , then σ is the identity.*

PROOF. This proof hinges on the difference between coiled 6-cycles and non-coiled 6-cycles. If C is a non-coiled 6-cycle, then it must contain three successive vertices u, v, w such that $u \in X_{i+1}, v \in X_i$ and $w \in X_{i+1}$, and such that if the 6-cycle is $uvwyztu$, then $y \in X_i, z \in X_{i-1}$ and $t \in X_i$ does not happen. Call the 2-path uvw an *anchor* of C . Because of the action of α and β , it is easy to see that if M has a non-coiled 6-cycle, then there is a non-coiled 6-cycle which has the 2-path $x_1^{-1}x_0^0x_1^1$ as an anchor.

Without loss of generality, let $\sigma \in \text{Aut}(M)$ fix x_0^0 and x_1^1 . It is easy to see that each of the 2-paths $x_2^{r^2}x_0^0x_1^1, x_2^{r^2}x_0^0x_1^{-1}, x_2^{-r^2}x_0^0x_1^1$, and $x_2^{-r^2}x_0^0x_1^{-1}$ is contained in a coiled 6-cycle. If the 2-path $x_1^{-1}x_0^0x_1^1$ is not contained in a 6-cycle (that is, every 6-cycle is coiled), then σ must also fix x_1^{-1} . If the 2-path $x_1^{-1}x_0^0x_1^1$ is not in a 6-cycle, then neither is the 2-path $x_0^{-2}x_1^{-1}x_0^0$. Hence, σ also must fix x_0^{-2} . Continuing in this way leads to the conclusion that σ fixes every vertex of X_0 and X_1 . This implies that σ is the identity.

Now consider the case that there are non-coiled 6-cycles. We now describe the possible non-coiled 6-cycles with anchor $x_1^{-1}x_0^0x_1^1$. One possibility is $x_0^0x_1^{-1}x_0^2x_1^3x_0^4x_1^5x_0^6$. However, this possibility would force $n = 6$ which is a

contradiction. A second possibility is $x_0^0 x_1^1 x_0^2 x_1^3 x_2^1 x_1^{-1} x_0^0$ which implies either $r = 2$ or $r = -2$. The former implies $n = 7$, which is a contradiction, and the latter implies $n = 9$ and M is the Holt graph.

A third possibility involves completing the 4-path $x_2^{-1-r} x_1^{-1} x_0^0 x_1^1 x_2^{1+r}$ in X_0 . However, if x_2^{-1-r} and x_2^{1+r} have a common neighbor in X_0 , it must be x_0^0 . This implies M has coiled girth 3 which is a contradiction. Likewise, the possibility that the 4-path $x_2^{-1+r} x_1^{-1} x_0^0 x_1^1 x_2^{1-r}$ completes to a 6-cycle in X_0 leads to the same contradiction.

The two remaining possibilities come from completing the two preceding 4-paths in X_1 . One resulting possibility is the 6-cycle $x_0^0 x_1^1 x_2^{1+r} x_1^0 x_2^{-1-r} x_1^{-1} x_0^0$ which implies that $r = (n - 1)/2$. The other resulting possibility is the 6-cycle $x_0^0 x_1^1 x_2^{1-r} x_1^0 x_2^{-1+r} x_1^{-1} x_0^0$ which implies that $r = (n + 1)/2$. We see that the 2-path $x_1^1 x_0^0 x_1^{-1}$ lies in precisely two non-coiled 6-cycles in both cases. Thus, if $r \neq (n - 1)/2$ and $r \neq (n + 1)/2$, there are no non-coiled 6-cycles and the first part of the proof establishes the result. Suppose that $r = (n - 1)/2$. Then the 2-paths $x_2^{-r^2} x_0^0 x_1^1$ and $x_2^{r^2} x_0^0 x_1^{-1}$ lie in three 6-cycles and the remaining 2-paths with x_0^0 as central vertex lie in two 6-cycles. Thus, σ must also fix the vertex $x_2^{-r^2}$. The same argument applied to the 2-paths centered at $x_2^{-r^2}$ implies that σ must also fix $x_1^{-r-r^2}$. Continuing in this way, σ also fixes $x_0^{-1-r-r^2}$, $x_2^{-1-r-2r^2}$, $x_1^{-1-2r-2r^2}$, $x_0^{-2-2r-2r^2}$, and so on until eventually we reach x_1^1 at which point we have completed a cycle C in M . If $d = \gcd(n, 1 + r + r^2)$, then the vertices of X_0 lying in C are $x_0^0, x_0^d, x_0^{2d}, \dots, x_0^{n-d}$. It is still possible that σ interchanges $x_2^{r^2}$ and x_1^{-1} . If so, then σ must also interchange $x_0^{1+r+r^2}$ and $x_0^{-1-r-r^2}$. However, the latter two vertices are distinct and lie on C . This contradicts the fact that σ fixes every vertex of C and we conclude that σ also fixes $x_1^{-1}, x_2^{-1-r}, x_0^{-1-r-r^2}, x_1^{-2-r-r^2}, \dots, x_2^{r^2}$. By repeating the preceding argument with x_1^1 replacing x_0^0 and so on, we eventually achieve that σ must fix every vertex of M .

We are left with the case that $r = (n + 1)/2$. However, this case does not arise as it is an easy number theoretic exercise to show that if $((n + 1)/2)^3 \equiv 1 \pmod{n}$, then either $n = 3$ or $n = 7$. This contradicts our assumption that $n \geq 9$.

THEOREM 3.3. *Let $M = M(r; 3, n)$, $n \geq 9$ and n odd, and let $r^3 \equiv 1 \pmod{n}$. Then M is 1/2-transitive.*

PROOF. We know that either $\text{Aut}(M) = \langle \alpha, \beta, \tau \rangle$ or $|\text{Aut}(M)| = 12n$ because of Lemmas 3.1 and 3.2. In the former case, M is 1/2-transitive as required. In the latter case, if M is arc-transitive, there must be an automorphism $\sigma \in \text{Aut}(M)$ such that $\sigma \notin \langle \alpha, \beta, \tau \rangle$, $\text{Aut}(M) = \langle \alpha, \beta, \tau, \sigma \rangle$ and σ interchanges the vertices

x_0^0 and x_1^1 . Orient the edge $x_0^0x_1^1$ from x_0^0 to x_1^1 obtaining the arc (x_0^0, x_1^1) . The group $\langle \alpha, \beta, \tau \rangle$ is regular on edges, so orienting the edge $g(x_0^0)g(x_1^1)$ from $g(x_0^0)$ to $g(x_1^1)$ for each $g \in \langle \alpha, \beta, \tau \rangle$ gives an orientation of M which we denote by $M*$. The digraph $M*$ is arc-transitive under the group $\langle \alpha, \beta, \tau \rangle$ and the latter group acts regularly on the arcs of $M*$. Since σ interchanges x_0^0 and x_1^1 , σ must be orientation-reversing on $M*$. We now carefully examine the action of σ .

Since σ interchanges x_0^0 and x_1^1 and is orientation reversing, σ must also interchange x_1^{-1} and x_0^2 . Then it must interchange x_1^{-3} and x_0^4 . Continuing in this way, we see that σ must interchange x_1^{-k} and x_0^{k+1} for all $k \in \mathbb{Z}_n$. Now x_1^1 and x_1^{1+2r} have the common neighbor x_2^{1+r} in X_2 . Thus, x_0^0 and x_0^{-2r} must have a common neighbor in X_2 . If they have a common neighbor, it must be x_2^{-r} . But the neighbors of x_0^0 are $x_2^{r^2}$ and $x_2^{-r^2}$. Now $-r$ and $-r^2$ clearly cannot be the same. Likewise, if $-r$ and r^2 are the same, then $r = -1$ is forced which is impossible. Therefore no such σ exists and M is not arc-transitive as required.

References

- [1] B. Alspach and T. D. Parsons, 'A construction for vertex-transitive graphs', *Canad. J. Math.* **34** (1982), 307–318.
- [2] ———, 'On hamiltonian cycles in metacirculant graphs', *Ann. Discrete Math.* **15** (1982), 1–7.
- [3] N. Biggs, *Algebraic graph theory* (Cambridge University Press, Cambridge, 1974).
- [4] I. Z. Bouwer, 'Vertex and edge-transitive but not 1-transitive graphs', *Canad. Math. Bull.* **13** (1970), 231–237.
- [5] Y. Cheng and J. Oxley, 'On weakly symmetric graphs of order twice a prime', *J. Combin. Theory Ser. B* **42** (1987), 196–211.
- [6] J. Folkman, 'Regular line-symmetric graphs', *J. Combin. Theory* **3** (1967), 215–232.
- [7] D. F. Holt, 'A graph which is edge-transitive but not arc-transitive', *J. Graph Theory* **5** (1981), 201–204.
- [8] D. Holton, 'Research Problem 9', *Discrete Math.* **38** (1982), 125.
- [9] D. Marušič, 'Vertex transitive graphs and digraphs of order p^k ', *Ann. Discrete Math.* **27** (1985), 115–128.
- [10] B. D. McKay, 'Transitive graphs with fewer than 20 vertices', *Math. Comp.* **33** (1979), 1101–1121.
- [11] B. D. McKay and G. F. Royle, 'The transitive graphs with at most 26 vertices', *Ars Combin.* **30** (1990), 161–176.
- [12] C. E. Praeger and Xu M. Y., 'Vertex primitive graphs of order a product of two distinct primes', preprint.
- [13] C. E. Praeger and G. F. Royle, 'Constructing the vertex transitive graphs of order 24', *J. Symbolic Comput.* **8** (1989), 309–326.
- [14] J. Turner, 'Point symmetric graphs with a prime number of points', *J. Combin. Theory* **3** (1967), 136–145.
- [15] W. T. Tutte, *Connectivity in graphs* (University of Toronto Press, Toronto, 1966).

Department of Mathematics and Statistics
Simon Fraser University
Burnaby, B.C. V5A 1S6
Canada

Institut za Matematiko,
Fiziko in Mehaniko
Jadranska 19
61111 Ljubljana
Slovenija
Yugoslavia

Apartment 9Q
275 West 96th Street
New York, New York 10025