

to rest in the emulsion and suffer $\pi \rightarrow \mu \rightarrow e$ decay. The negative one, after 14 130 μ , gives rise to a two pronged star before the end of its range.

The energy of π_1 -meson has been deduced both from momentum balance and from direct scattering measurements.

The values obtained are the following:

	From Range	From momentum balance	From scattering measurements
E_{π_1}	—	30.7 ± 1.5 MeV	30.8 ± 6 MeV
E_{π_2}	17.06 ± 0.85 MeV	—	—
E_{π_3}	23.61 ± 0.84 MeV	—	—

The angles are:

$$\varphi_{\pi_2\pi_3} = 102^\circ 50' \pm 30'$$

$$\varphi_{\pi_1\pi_3} = 133^\circ 35' \pm 30'$$

$$\varphi_{\pi_1\pi_2} = 123^\circ 34' \pm 30'$$

The energy of π_2 and π_3 has been calculated from the range using the relation ⁽⁵⁾:

$$E = kR^{0.568} \left(\frac{m_\pi}{m_p} \right)^{0.432}$$

k has been determined from range measurements of μ coming from $\pi \rightarrow \mu$ decay,

The Q -value of the τ -meson is:

$$Q = 71.4 \pm 3.2 \text{ MeV.}$$

This is the second example of a star emitting two heavy particles observed in our laboratory ⁽⁶⁾. An analysis of these events is in progress.

We are indebted to Professors R. DEAGLIO, G. LOVERA and G. WATAGHIN for helpful discussions.

The event studied in this paper was found by Miss T. SAPPJA and Mr. M. GRECO, whom we wish to thank for the photograph.

⁽⁶⁾ A. DEBENEDETTI, C. M. GARELLI, L. TALLONE e M. VIGONE: *Nuovo Cimento*, **12**, 369 (1954).

ERRATA-CORRIGE

E. CORINALDESI — Construction of Potentials from Phase Shift and Binding Energies of Relativistic Equations, *Nuovo Cimento*, **11**, 468 (1954).

The factors $-2i$ in eqs. (30) and (A 10), and $i/2$ in eq. (31) should be omitted, being due to an oversight.