

CONTACT 3-MANIFOLDS WITH THE REEB FLOW SYMMETRY

JONG TAEK CHO

(Received February 13, 2013, revised October 16, 2013)

Abstract. We prove that the Ricci operator on a contact Riemannian 3-manifold M is invariant along the Reeb flow if and only if M is Sasakian or locally isometric to $SU(2)$ (or $SO(3)$), $SL(2, \mathbf{R})$ (or $O(1, 2)$), the group $E(2)$ of rigid motions of Euclidean 2-plane with a contact left invariant Riemannian metric.

1. Introduction. In a contact manifold (M, η) , we find a fundamental fact that the Reeb vector field ξ generates a contact transformation, that is, $\mathfrak{L}_\xi \eta = 0$. For an associated Riemannian metric g , if ξ generates an isometric flow, that is, M satisfies $\mathfrak{L}_\xi g = 0$, then M is said to be K -contact. We note that a K -contact manifold is already Sasakian in dimension three. In this paper, we study a 3-dimensional contact Riemannian manifold whose Ricci operator S is Reeb flow invariant, that is, $\mathfrak{L}_\xi S = 0$. Then, we have

MAIN THEOREM. *Let M be a 3-dimensional contact Riemannian manifold. Then $\mathfrak{L}_\xi S = 0$ if and only if M is Sasakian or locally isometric to $SU(2)$ (or $SO(3)$), $SL(2, \mathbf{R})$ (or $O(1, 2)$), $E(2)$ (the group of rigid motions of Euclidean 2-plane) with a left invariant contact Riemannian metric.*

All manifolds in the present paper are assumed to be connected and of class C^∞ .

2. Preliminaries. A 3-dimensional manifold M is said to be a contact manifold if it admits a global 1-form η such that $\eta \wedge (d\eta) \neq 0$ everywhere. Given a contact form η , we have a unique vector field ξ , which is called the characteristic vector field, satisfying $\eta(\xi) = 1$ and $\mathfrak{L}_\xi \eta = 0$ (or $i_\xi d\eta = 0$), where \mathfrak{L}_ξ denotes Lie differentiation for ξ and i_ξ denotes the interior product operator by ξ . It is well-known that there exists a Riemannian metric g and a $(1, 1)$ -tensor field φ such that

$$(1) \quad \eta(X) = g(X, \xi), \quad d\eta(X, Y) = g(X, \varphi Y), \quad \varphi^2 X = -X + \eta(X)\xi,$$

where X and Y are vector fields on M . From (1) it follows that

$$(2) \quad \varphi\xi = 0, \quad \eta \circ \varphi = 0, \quad g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y).$$

A manifold M equipped with structure tensors (φ, ξ, η, g) satisfying (1) and (2) is said to be a contact Riemannian manifold and is denoted by $M = (M; \eta, g)$. Given a contact Riemannian manifold M , we define a $(1, 1)$ -tensor field h by $h = \frac{1}{2}\mathfrak{L}_\xi \varphi$. Then h is self-adjoint and satisfies

$$(3) \quad \underline{\hspace{10em}} \quad h\xi = 0 \quad \text{and} \quad h\varphi = -\varphi h,$$

$$(4) \quad \nabla_X \xi = -\varphi X - \varphi h X,$$

where ∇ is Levi-Civita connection. From (3) and (4) we see that each trajectory of ξ is a geodesic and $\text{div}(\xi) = 0$. We denote by R the Riemannian curvature tensor defined by

$$R(X, Y)Z = \nabla_X(\nabla_Y Z) - \nabla_Y(\nabla_X Z) - \nabla_{[X, Y]}Z$$

for all vector fields X, Y, Z . Along a trajectory of ξ , the Jacobi operator $\ell = R(\cdot, \xi)\xi$ is a symmetric $(1, 1)$ -tensor field, that is, $g(\ell X, Y) = g(X, \ell Y)$. We have

$$(5) \quad \text{trace } \ell = \rho(\xi, \xi) = 2n - \text{trace}(h^2),$$

$$(6) \quad \nabla_\xi h = \varphi - \varphi \ell - \varphi h^2,$$

$$(7) \quad g(R(X, Y)\xi, Z) = g((\nabla_Z \varphi)X, Y) + g((\nabla_Y \varphi h)X - (\nabla_X \varphi h)Y, Z)$$

for all vector fields X, Y, Z on M , where $\rho(X, Y) = g(SX, Y)$. A contact Riemannian manifold for which ξ is Killing is called a K -contact Riemannian manifold. It is easy to see that a contact Riemannian manifold is K -contact if and only if $h = 0$. For a contact Riemannian manifold M one may define naturally an almost complex structure J on $M \times \mathbf{R}$;

$$J\left(X, f \frac{d}{dt}\right) = \left(\varphi X - f\xi, \eta(X) \frac{d}{dt}\right),$$

where X is a vector field tangent to M , t the coordinate of \mathbf{R} and f a function on $M \times \mathbf{R}$. If the almost complex structure J is integrable, M is said to be normal or Sasakian. It is known that M is normal if and only if M satisfies

$$[\varphi, \varphi] + 2d\eta \otimes \xi = 0,$$

where $[\varphi, \varphi]$ is the Nijenhuis torsion of φ . A Sasakian manifold is characterized by a condition

$$(8) \quad (\nabla_X \varphi)Y = g(X, Y)\xi - \eta(Y)X$$

for all vector fields X and Y on the manifold. For more details about contact Riemannian manifolds we refer to [1].

3. Contact 3-manifolds with the Reeb flow symmetry. In this section, we prove the Main Theorem. First we recall that a contact Riemannian 3-manifold M satisfies

$$(9) \quad (\nabla_X \varphi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX)$$

(cf. [5]). From (7) and (9) we have

$$(10) \quad R(X, Y)\xi = \eta(Y)(X + hX) - \eta(X)(Y + hY) + \varphi((\nabla_Y h)X - (\nabla_X h)Y)$$

for all vector fields X and Y . From (8) and (9), we have at once

LEMMA 1. *A 3-dimensional contact Riemannian manifold is Sasakian if and only if $h = 0$.*

Moreover, we have

PROPOSITION 2. *A Sasakian 3-manifold is η -Einstein, that is, $S = \alpha I + \beta \eta \otimes \xi$, where α and β are functions with $d\alpha(\xi) = d\beta(\xi) = 0$.*

Since $\mathfrak{L}_\xi \xi = \mathfrak{L}_\xi \eta = 0$, we have

COROLLARY 3. For a Sasakian 3-manifold, $\mathfrak{L}_\xi S = 0$.

Now, we prove the Main Theorem.

PROOF OF MAIN THEOREM. Let $M = (M^3; \eta, g)$ be a 3-dimensional contact Riemannian manifold. Then it is well-known that the curvature tensor R of a 3-dimensional Riemannian manifold is expressed by

$$(11) \quad \begin{aligned} R(Y, X)Z = & \rho(X, Z)Y - \rho(Y, Z)X + g(X, Z)SY - g(Y, Z)SX \\ & - \frac{\tau}{2}\{g(X, Z)Y - g(Y, Z)X\} \end{aligned}$$

for all vector fields X, Y, Z , where τ denotes the scalar curvature. If $h = 0$ on M , then from Lemma 1 we see that M is Sasakian. Moreover, M satisfies $\mathfrak{L}_\xi S = 0$ (Corollary 3). So, we consider on M the maximal open subset U_1 on which $h \neq 0$ and the maximal open subset U_2 on which h is identically zero. (U_2 is the union of all points p in M such that $h = 0$ in a neighborhood of p). $U_1 \cup U_2$ is open and dense in M . Suppose that M is non-Sasakian. Then U_1 is non-empty and there is a local orthonormal frame field $\{e_1 = e, e_2 = \varphi e, e_3 = \xi\}$ on U_1 such that $h(e_1) = \lambda e_1, h(e_2) = -\lambda e_2$ for some positive function λ . We denote $\Gamma_{ijk} = g(\nabla_{e_i} e_j, e_k), \rho_{ij} = \rho(e_i, e_j)$ for $i, j, k = 1, 2, 3$. Then from (4) we get

$$(12) \quad \Gamma_{132} = -\Gamma_{123} = -(1 + \lambda), \quad \Gamma_{231} = -\Gamma_{213} = 1 - \lambda$$

and

$$(13) \quad \Gamma_{131} = \Gamma_{113} = \Gamma_{232} = \Gamma_{223} = 0.$$

Also, from (6) and taking account of (5) and (11), we have

$$(14) \quad \xi \lambda = \rho_{12}$$

and

$$(15) \quad 4\lambda \Gamma_{312} = \rho_{22} - \rho_{11}.$$

LEMMA 4. In $U_1, \mathfrak{L}_\xi S = 0 \Leftrightarrow \nabla_\xi S = 0$ and $S\xi = \sigma\xi$, where σ is a function.

PROOF. Suppose that M satisfies $\mathfrak{L}_\xi S = 0$. Then, we compute

$$\begin{aligned} 0 &= \mathfrak{L}_\xi(SX) - S(\mathfrak{L}_\xi X) \\ &= [\xi, SX] - S[\xi, X]. \end{aligned}$$

From this using (4) we get an equivalent equation to $\mathfrak{L}_\xi S = 0$:

$$(16) \quad (\nabla_\xi S)X = (S\varphi - \varphi S)X + (S\varphi h - \varphi h S)X.$$

Since $\nabla_\xi S$ and $S\varphi - \varphi S$ are self-adjoint operators, we get

$$S\varphi h - \varphi h S = Sh\varphi - h\varphi S.$$

Since $h\varphi = -\varphi h$, it follows that

$$(17) \quad S\varphi h = \varphi h S.$$

Since $h\xi = 0$, from (17) we see that $hS\xi = 0$. From this and (5), we obtain $S\xi = \sigma\xi$, $\sigma = 2 - 2\lambda^2$ on U_1 . And from (16) and (17) we get

$$(18) \quad \nabla_{\xi} S = S\varphi - \varphi S.$$

So, we get $(\nabla_{\xi}\rho)(\xi, \xi) = 0$, and then $\xi\lambda = 0$, where we have used $\nabla_{\xi}\xi = 0$. Then from (14) we have

$$(19) \quad \rho_{12} = \rho_{21} = 0.$$

Applying e_1 to (17) and taking an inner product with e_2 (with respect to g), we get

$$(20) \quad \rho_{22} = \rho_{11}$$

on U_1 . Since $S\xi = \sigma\xi$, together with (19), we have $S\varphi = \varphi S$ on U_1 . Thus, from (18) we obtain $\nabla_{\xi} S = 0$ on U_1 . Conversely, we assume that $\nabla_{\xi} S = 0$ and $S\xi = \sigma\xi$ on U_1 . Then, it follows from (5) that $\sigma = 2 - 2\lambda^2$, and

$$(21) \quad \rho_{13} = \rho_{31} = 0, \quad \rho_{23} = \rho_{32} = 0.$$

And from $(\nabla_{\xi}\rho)(\xi, \xi) = 0$ and $\nabla_{\xi}\xi = 0$ we have

$$(22) \quad \xi\lambda = 0,$$

which together with (14) yields

$$(23) \quad \rho_{12} = \rho_{21} = 0.$$

Using $\nabla_{\xi} S = 0$ again, we obtain from (23)

$$(24) \quad \Gamma_{312}(\rho_{11} - \rho_{22}) = 0.$$

By (15) and (24) we find that

$$(25) \quad \rho_{11} = \rho_{22}.$$

Since $S\xi = \sigma\xi$, equations (23) and (25) give $S\varphi = \varphi S$. Moreover, we see that $S\varphi h = \varphi h S$ on U_1 . Therefore, by (16) we find that $\xi_{\xi} S = 0$. This completes the proof of Lemma 4. \square

Now we prove

LEMMA 5. λ is constant.

PROOF. Among the proof of Lemma 4, from (15) and (25) we get in addition

$$(26) \quad \Gamma_{312} = \Gamma_{321} = 0.$$

From (11) with the help of Lemma 4 we have in U_1 :

$$\begin{aligned}
 R(e_1, e_2)e_2 &= Se_1 - (1 - \lambda^2)e_1, \\
 R(e_1, e_2)e_1 &= -Se_2 + (1 - \lambda^2)e_2, \\
 R(e_2, e_3)e_2 &= R(e_1, e_3)e_1 = -(1 - \lambda^2)e_3, \\
 R(e_1, e_3)e_3 &= (1 - \lambda^2)e_1, \\
 R(e_2, e_3)e_3 &= (1 - \lambda^2)e_2, \\
 R(e_i, e_j)e_k &= 0 \text{ for } i \neq j \neq k \neq i.
 \end{aligned}
 \tag{27}$$

Using (12), (13), (26), and (27), we have

$$\begin{aligned}
 (\nabla_{e_1} R)(e_2, e_3)e_2 &= e_1(\lambda^2 - 1)e_3, \\
 (\nabla_{e_2} R)(e_3, e_1)e_2 &= (\nabla_{e_3} R)(e_1, e_2)e_2 = 0
 \end{aligned}
 \tag{28}$$

and

$$\begin{aligned}
 (\nabla_{e_2} R)(e_1, e_3)e_1 &= e_2(\lambda^2 - 1)e_3, \\
 (\nabla_{e_1} R)(e_3, e_2)e_1 &= (\nabla_{e_3} R)(e_2, e_1)e_1 = 0.
 \end{aligned}
 \tag{29}$$

By the second Bianchi identity, (28) and (29) yield that $e_1(\lambda) = 0$ and $e_2(\lambda) = 0$ respectively. Hence, together with (22), we see that λ is constant on M , where we have used the continuity argument of λ . □

On account of (27) we find that $R(e_1, e_2)\xi = 0$ in M . Here we use (10). Since λ is constant, we have

$$\Gamma_{212}e_1 - \Gamma_{121}e_2 = 0.
 \tag{30}$$

From (30) we get

$$\Gamma_{212} = \Gamma_{221} = \Gamma_{121} = \Gamma_{112} = 0.
 \tag{31}$$

Thus, together with (12), (13), (26), and (31), we have

$$[e_1, e_2] = 2e_3, \quad [e_2, e_3] = (1 - \lambda)e_1, \quad [e_3, e_1] = (1 + \lambda)e_2.
 \tag{32}$$

Actually, from (32) we compute the Ricci operator S :

$$\begin{aligned}
 Se_1 &= 0, \\
 Se_2 &= 0, \\
 Se_3 &= (2 - 2\lambda^2)e_3.
 \end{aligned}
 \tag{33}$$

Moreover, we can check $(\mathfrak{L}_\xi S)e_1 = (\mathfrak{L}_\xi S)e_2 = 0$.

After all, owing to J. Milnor's result (Section 4 or [2]), we see from (32) that M is locally isometric to one of the following Lie groups:

- (i) $SU(2)$ (or $SO(3)$) with a left invariant metric when $0 < \lambda < 1$;
- (ii) $SL(2, \mathbf{R})$ (or $O(1, 2)$) with a left invariant metric when $\lambda > 1$;
- (iii) $E(2)$ when $\lambda = 1$.

Thus we have proved our Main Theorem. □

We see from Proposition 2 that a Sasakian 3-manifold satisfies $S\xi = \sigma\xi$ and $\nabla_\xi S = 0$, where σ is a function.

COROLLARY 6. *Let M be a 3-dimensional contact Riemannian manifold. Then $S\xi = \sigma\xi$ and $\nabla_\xi S = 0$ if and only if M is Sasakian or locally isometric to $SU(2)$ (or $SO(3)$), $SL(2, \mathbb{R})$ (or $O(1, 2)$), the group $E(2)$ of rigid motions of Euclidean 2-plane with a left invariant contact Riemannian metric.*

We can not remove the condition $S\xi = \sigma\xi$ in Corollary 6. Indeed, we have a counter example. See Remark 2 in the next section.

4. 3-dimensional Lie groups. By a theorem due to K. Sekigawa [4] and the classification due to J. Milnor [2] of 3-dimensional Lie groups with a left invariant metric, Perrone [3] classified all simply connected homogeneous contact Riemannian 3-manifolds. Recall that M is called unimodular if its left invariant Haar measure is also right invariant. In terms of the Lie algebra \mathfrak{m} , M is unimodular if and only if the adjoint transformation ad_X has trace zero for every $X \in \mathfrak{m}$. Then we have

PROPOSITION 7 ([5]). *Let M be a 3-dimensional unimodular Lie group with a left invariant contact Riemannian structure, then there exists an orthonormal basis $\{e_1, e_2 = \varphi e_1, e_3 = \xi\} \in \mathfrak{m}$ such that*

$$(34) \quad [e_1, e_2] = 2e_3, [e_2, e_3] = c_2e_1, [e_3, e_1] = c_3e_2.$$

REMARK 1 ([3]). In fact, every three-dimensional unimodular Lie group, with only exception of the commutative Lie group \mathbb{R}^3 , admits a left-invariant contact metric structure.

We put

$$\Gamma_{ijk} = g(\nabla_{e_i}e_j, e_k) \quad \text{for } i, j, k = 1, 2, 3.$$

Then by using the Koszul formula we have

$$(35) \quad \begin{cases} \Gamma_{123} = \frac{1}{2}(c_3 - c_2 + 2), \\ \Gamma_{213} = \frac{1}{2}(c_3 - c_2 - 2), \\ \Gamma_{312} = \frac{1}{2}(c_3 + c_2 - 2), \\ \text{all others are zero.} \end{cases}$$

From (35) we easily see that M is K -contact (or Sasakian) if and only if $c_2 = c_3$. Then, using (35), we find by a direct calculation

$$\begin{aligned}
 R(e_1, e_2)e_2 &= \left(\frac{1}{4}(c_3 - c_2)^2 - 3 + c_3 + c_2\right) e_1 \\
 R(e_1, e_3)e_3 &= \left(-\frac{1}{4}(c_3 - c_2)^2 - \frac{1}{2}(c_3^2 - c_2^2) + 1 - c_2 + c_3\right) e_1 \\
 R(e_2, e_1)e_1 &= \left(\frac{1}{4}(c_3 - c_2)^2 - 3 + c_3 + c_2\right) e_2 \\
 R(e_2, e_3)e_3 &= \left(\frac{1}{4}(c_3 + c_2)^2 - c_2^2 + 1 + c_2 - c_3\right) e_2 \\
 R(e_3, e_1)e_1 &= \left(-\frac{1}{4}(c_3 - c_2)^2 - \frac{1}{2}(c_3^2 - c_2^2) + 1 - c_2 + c_3\right) e_3 \\
 R(e_3, e_2)e_2 &= \left(\frac{1}{4}(c_3 + c_2)^2 - c_2^2 + 1 + c_2 - c_3\right) e_3.
 \end{aligned}
 \tag{36}$$

By using (36) we get

$$\begin{aligned}
 Se_1 &= \left(-\frac{1}{2}(c_3^2 - c_2^2) - 2 + 2c_3\right) e_1 \\
 Se_2 &= \left(\frac{1}{2}(c_3^2 - c_2^2) - 2 + 2c_2\right) e_2 \\
 Se_3 &= \left(-\frac{1}{2}(c_3 - c_2)^2 + 2\right) e_3.
 \end{aligned}
 \tag{37}$$

Since $c_1 = 2 > 0$, the possible combinations of the signs of c_1, c_2 and c_3 and the associated Lie groups are indicated in the following table (see [2]):

Signature of (c_1, c_2, c_3)	Associated Lie group
$(+, +, +)$	$SU(2)$ or $SO(3)$
$(+, +, -)$	$SL(2, \mathbf{R})$ or $O(1, 2)$
$(+, +, 0)$	$E(2)$
$(+, -, -)$	$SL(2, \mathbf{R})$ or $O(1, 2)$
$(+, -, 0)$	$E(1, 1)$
$(+, 0, 0)$	Heisenberg group Nil_3

$SU(2)$: group of 2×2 unitary matrices of determinant 1; homeomorphic to the unit 3-sphere.

$SO(3)$: rotation group of Euclidean 3-space, isomorphic to $SU(2)/\{\pm I\}$.

$SL(2, \mathbf{R})$: group of 2×2 real matrices of determinant 1.

$O(1, 2)$: Lorentz group consisting of linear transformations preserving the quadratic form $t^2 - x^2 - y^2$. Its identity component is isomorphic to $SL(2, \mathbf{R})/\{\pm I\}$, or to the group of rigid motions of hyperbolic 2-space.

$E(2)$: group of rigid motions of Euclidean 2-space.

$E(1, 1)$: group of rigid motions of Minkowski 2-space.

Finally, the Heisenberg group can be described as the group of all 3×3 real matrices of the form

$$\begin{pmatrix} 1 & * & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{pmatrix}.$$

PROPOSITION 8 ([3]). *Let M be a 3-dimensional non-unimodular Lie group with left invariant contact Riemannian structure. Then there exists an orthonormal basis $\{e_1, e_2 = \varphi e_1, e_3 = \xi\} \in \mathfrak{m}$ such that*

$$(38) \quad [e_1, e_2] = \alpha e_2 + 2e_3, \quad [e_2, e_3] = 0, \quad [e_3, e_1] = \gamma e_2,$$

where $\alpha \neq 0$. Moreover, M is Sasakian if and only if $\gamma = 0$.

By using the Koszul formula we see from (38)

$$(39) \quad \begin{cases} \Gamma_{123} = \frac{\gamma + 2}{2} \\ \Gamma_{212} = -\alpha \\ \Gamma_{213} = \frac{\gamma - 2}{2} \\ \Gamma_{312} = \frac{\gamma - 2}{2} \\ \text{all others are zero.} \end{cases}$$

Then, using (39), we obtain by a direct calculation

$$(40) \quad \begin{aligned} R(e_1, e_2)e_2 &= \left(\frac{\gamma^2 + 4\gamma - 12}{4} - \alpha^2\right) e_1 \\ R(e_1, e_3)e_3 &= \left(\frac{-3\gamma^2 + 4\gamma + 4}{4}\right) e_1 \\ R(e_2, e_1)e_1 &= \left(\frac{\gamma^2 + 4\gamma - 12}{4} - \alpha^2\right) e_2 + \alpha\gamma e_3 \\ R(e_2, e_3)e_3 &= \frac{(\gamma - 2)^2}{4} e_2 \\ R(e_3, e_1)e_1 &= \alpha\gamma e_2 + \left(\frac{-3\gamma^2 + 4\gamma + 4}{4}\right) e_3 \\ R(e_3, e_2)e_2 &= \frac{(\gamma - 2)^2}{4} e_3, \end{aligned}$$

and thus

$$(41) \quad \begin{aligned} Se_1 &= \left(-\alpha^2 - 2 + 2\gamma - \frac{\gamma^2}{2}\right) e_1 \\ Se_2 &= \left(-\alpha^2 - 2 + \frac{\gamma^2}{2}\right) e_2 + \alpha\gamma e_3 \\ Se_3 &= \alpha\gamma e_2 + \left(2 - \frac{\gamma^2}{2}\right) e_3. \end{aligned}$$

THEOREM 9. *Let M be a 3-dimensional Lie group with left invariant contact Riemannian structure. Suppose that M satisfies $\nabla_{\xi}S = 0$.*

- (a) *If M is unimodular, then M is isometric to one of the following Lie groups:*
- (i) *SU(2) (or SO(3)) with Sasakian metric or (non-Sasakian) contact Riemannian metric,*
 - (ii) *SL(2, \mathbf{R}) (or $O(1, 2)$) with Sasakian metric or (non-Sasakian) contact Riemannian metric,*
 - (iii) *Heisenberg group with Sasakian metric,*
 - (iv) *$E(2)$ with contact Riemannian metric.*
- (b) *If M is non-unimodular, then the Lie algebra structure is given by (38) with $\gamma = 0$ (Sasakian) or $\gamma = 2$.*

PROOF. (a) By using (35) and (37), we obtain

$$(\nabla_{e_3}S)e_1 = \frac{1}{2}(c_2 - c_3)(c_3 + c_2 - 2)^2e_2$$

and

$$\begin{aligned}(\nabla_{e_3}S)e_2 &= \frac{1}{2}(c_2 - c_3)(c_3 + c_2 - 2)^2e_1, \\ (\nabla_{e_3}S)e_3 &= 0.\end{aligned}$$

Thus we see that $\nabla_{\xi}S = 0$ if and only if $c_3 = c_2$ or $c_3 + c_2 = 2$. Then, referring the Table we obtain (a).

(b) By using (39) and (41), we obtain

$$\begin{aligned}(\nabla_{e_3}S)e_1 &= -\frac{1}{2}\gamma(\gamma - 2)^2e_2 - \frac{1}{2}\alpha\gamma(\gamma - 2)e_3, \\ (\nabla_{e_3}S)e_2 &= -\frac{1}{2}\gamma(\gamma - 2)^2e_1, \\ (\nabla_{e_3}S)e_3 &= -\frac{1}{2}\alpha\gamma(\gamma - 2)e_1.\end{aligned}$$

Since $\alpha \neq 0$ from the above equations, we see that $\nabla_{\xi}S = 0$ if and only if $\gamma = 0$ or $\gamma = 2$. □

REMARK 2. From Theorem 9, we find that the non-unimodular Lie group whose Lie algebra structure is given by (38) with $\gamma = 2$ satisfies $\nabla_{\xi}S = 0$, but $S\xi \neq \sigma\xi$. In fact, we see from (41) that $S\xi = 2\alpha e_2$ ($\alpha \neq 0$).

Acknowledgement. The author would like to thank the referee and the editor for their careful reading of the manuscript and giving useful comments for the revised version. This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012R1A1B3003930).

REFERENCES

- [1] D. E. BLAIR, Riemannian geometry of contact and symplectic manifolds, Second edition, Progr. Math. 203, Birkhäuser Boston, Inc., Boston, MA, 2010.
- [2] J. MILNOR, Curvature of left invariant metrics on Lie groups, Advances in Math. 21 (1976), 293–329.
- [3] D. PERRONE, Homogeneous contact Riemannian three-manifolds, Illinois J. Math. 42 (1998), 243–256.
- [4] K. SEKIGAWA, On some 3-dimensional curvature homogeneous spaces, Tensor (N.S.) 31 (1977), 87–97.
- [5] S. TANNO, Variational problems on contact Riemannian manifolds, Trans. Amer. Math. Soc. 314 (1989), 349–379.

DEPARTMENT OF MATHEMATICS
CHONNAM NATIONAL UNIVERSITY
GWANGJU 500–757
KOREA

E-mail address: jtcho@chonnam.ac.kr