

Vol. 4 (1999) Paper no. 13, pages 1–13.

Journal URL http://www.math.washington.edu/~ejpecp Paper URL http://www.math.washington.edu/~EjpVol4/paper13.abs.html

CONVERGENCE OF STOPPED SUMS OF WEAKLY DEPENDENT RANDOM VARIABLES

Magda Peligrad Department of Mathematical Sciences University of Cincinnati P. O. Box 210025

Cincinnati, OH 45221-0025 peligram@math.uc.edu

Abstract In this paper we investigate stopped partial sums for weak dependent sequences. In particular, the results are used to obtain new maximal inequalities for strongly mixing sequences and related almost sure results.

 ${\bf Keywords}$ Partial sums, maximal inequalities, weak dependent sequences, stopping times, amarts.

AMS Subject Classification 60E15, 60F05.

This research was supported, in part, by grants from the NSF and cost sharing at the University of Cincinnati.

Submitted to EJP on July 23, 1996. Final version accepted on April 6, 1999.

Convergence of stopped sums of weakly dependent random variables

Magda Peligrad* University of Cincinnati

Abbreviated title: Convergence of stopped sums

Summary

In this paper we investigate stopped partial sums for weak dependent sequences. In particular, the results are used to obtain new maximal inequalities for strongly mixing sequences and related almost sure results.

1 Introduction

A random walk with weak dependent increments is a sequence $\{S_n; n \ge 0\}$ of random variables with $S_0 = 0$ and increments $\{X_k; k \ge 1\}$, which are weakly dependent in some sense. The applications of these random walks in renewal theory was studied by Berbee (1979).

Motivated by applications to sequential analysis, we shall investigate stopped random walks $\{S_{\tau_n}; n \ge 1\}$ where τ_n are stopping times.

For the case when increments are independent a variety of convergence results and applications are surveyed in Gut (1986).

In this paper we assume the increments are weak dependent. Mixing type of dependence gives general models for which we can prove convergence theorems of both weak and strong type with a large applicability to Markov chains, Gaussian processes, time series, number theory, etc. Various examples are contained in Bradley (1986) and Doukhan (1994).

The problem we investigate here also reveals the relation of mixing sequences with some generalized notions of martingales such as amarts (Austin, Edgar and Ionescu Tulcea (1974)) and semiamarts (Edgar and Sucheston (1976), Krengel and Sucheston (1978)). Generalized martingales are important in getting martingale-like maximal inequalities, tightness of stochastic processes associated to partial sums, as well as various strong and weak limit theorems. Semiamarts have proven useful in connection with the optimal stopping rules and the reversed semiamart property is useful in proving uniform integrability of partial sums. In addition, the amarts admit

^{*}Supported in part by a NSF grant and cost sharing at the University of Cincinnati

AMS 1991 subject classification: 60E15, 60F05

Key words and phrases: Partial sums, maximal inequalities, weak dependent sequences, stopping times, amarts

the Riesz decomposition in a martingale and a potential which makes it easy for the martingale tool to be applied to all examples of amarts we shall give in this paper. The properties of amarts and semiamarts are surveyed in Gut and Schmidt (1983) and Edgar and Sucheston (1992).

In this paper, the main application of amarts are maximal inequalities for partial sums of weakly dependent random variables. There is a large body of work dealing with this problem. Among the papers containing various maximal inequalities for dependent sequences I would like to mention Houdre (1995), McLeish (1975), Moricz (1976), Peligrad (1985a), (1985b), (1992), Roussas (1991), (1992), Rio (1994), Shao (1993), (1994) and Utev (1991).

Let (Ω, K, P) be a probability space. We shall introduce some measures of dependence between two sub σ -algebras of K, \mathcal{A} and \mathcal{B} .

$$\begin{aligned} \alpha(\mathcal{A},\mathcal{B}) &= \sup\{P(A \cap B) - P(A)P(B); \quad A \in \mathcal{A}, B \in \mathcal{B}\}, \\ \rho(\mathcal{A},\mathcal{B}) &= \sup\{\operatorname{corr}(f,g); \quad f \in L_2(\mathcal{A}), B \in L_2(\mathcal{B})\}, \\ \varphi(\mathcal{A},\mathcal{B}) &= \sup\left\{\frac{P(A \cap B) - P(A)P(B)}{P(A)}; \quad A \in \mathcal{A}, P(A) \neq 0, B \in \mathcal{B}\right\} \end{aligned}$$

Let now $(X_n)_{n\geq 1}$ be a sequence of random variables on (Ω, K, P) , and denote $\mathcal{F}_m^n = \sigma(X_k; m \leq k \leq n)$.

DEFINITION 1.1 We say that $(X_n)_{n\geq 1}$ is strongly mixing if

(1.1)
$$\alpha(n) \to 0 \quad where \quad \alpha(n) = \sup_{m \ge 1} \alpha(\mathcal{F}_1^m, \mathcal{F}_{m+n}^\infty),$$

 ρ -mixing if

(1.2)
$$\rho(n) \to 0 \quad where \quad \rho(n) = \sup_{m \ge 1} \rho(\mathcal{F}_1^m, \mathcal{F}_{m+n}^\infty),$$

 φ -mixing if (1.3)

$$arphi(n) o 0 \quad where \quad arphi(n) = \sup_{m \ge 1} \left(\mathcal{F}_1^m, \mathcal{F}_{n+m}^\infty
ight).$$

It is well known that $\varphi(n) \to 0$ implies $\rho(n) \to 0$ which in turn implies $\alpha(n) \to 0$ as $n \to \infty$.

Denote by $S_n = \sum_{i=1}^n X_i$ and by T the set of bounded stopping times adapted to \mathcal{F}_1^n , i.e.: $\tau \in T$ if and only if $\{\tau = n\} \in \mathcal{F}_1^n$ for all n and $P(\tau \leq M) = 1$ for some integer M depending on τ . A net $(a_{\tau})_{\tau \in T}$ is said to converge to a if and only if for every $\varepsilon > 0$ there exists $\tau_0 \in T$ such that $|a_{\tau} - a| < \varepsilon$ for all $\tau \in T$, $\tau \geq \tau_0$.

DEFINITION 1.2 We say that $(S_n)_{n\geq 1}$ is an amart if and only if the net $(ES_{\tau})_{\tau\in T}$ is convergent. We say that $(S_n)_{n\geq 1}$ is a semiamart if and only if the net $(ES_{\tau})_{\tau\in T}$ is bounded.

We shall consider in this paper sums of mixing sequences of random variables and we shall study the convergence of these stopped sums. We shall give sufficient conditions for the convergence of the stopped net which will relate the concept of mixing sequences to those of amarts and semiamarts. In particular, if the stopping time is defined as the first crossing of an interval by sums we get maximal inequalities for partial sums. Some applications of the results to the convergence of series of weakly dependent random variables will be given. Some of these results can be formulated in the context of Banach space valued random variables. The maximal inequalities can be used to obtain the invariance principles for random elements associated to partial sums of weakly dependent random variables by using Theorem (8.3) in Billingsley (1968). Other inequalities for rank orders of partial sums can be obtained by defining other stopping times (like in Newman and Wright (1982)). Also generalizations of these results to random fields can be considered. However we shall not follow in this paper all these implications and directions and we will leave them for further research.

In the following text, [x] denotes the integer part of x, and $\tilde{\alpha}(n) = \sup_{m \ge 1} \alpha(\mathcal{F}_1^m, \mathcal{F}_{n+m}^{n+m}).$

2 The results

Our first theorem gives an upper bound for a second moment of stopped random sums in terms of the ρ -mixing coefficients and second moments of the individual summands. It is well known that if the sequence is ρ -mixing then $c(\rho, n)$ defined in the following theorem is a slowly varying function of n when $n \to \infty$. If $\rho(n)$ is decreasing to zero logarithmically slowly such that $\sum_{i} \rho(2^{i}) < \infty$, then $c(\rho, n)$ is a numerical constant which does not depend on n.

THEOREM 2.1 Let $(X_i)_{i\geq 1}$ be a centered sequence of random variable such that $EX_i^2 < \infty$ for every $i \geq 1$. Denote

(2.1)
$$c(\rho, n) = \exp\left(2\sum_{j=0}^{\lfloor \log_2 n \rfloor} \rho\left(\left[2^{j/3}\right]\right)\right)$$

Let τ be a stopping time. Then there is an absolute constant K such that for every $n \ge 1$:

(2.2)
$$ES_{\tau}^{2}I(\tau \leq n) \leq Kc^{4}(\rho, n) \sum_{j=1}^{n} EX_{j}^{2}.$$

As a first corollary of Theorem 2.1 we can formulate the following maximal inequality.

COROLLARY 2.1 (Bradley and Utev (1994)) Under the conditions of Theorem 2.1 there is a constant K such that for every $n \ge 1$ and every $\lambda > 0$:

$$\lambda^2 P\left(\max_{1 \le i \le n} |S_i| \ge \lambda\right) \le K c^4(\rho, n) \sum_{j=1}^n E X_j^2,$$

where $c(\rho, n)$ is defined by (2.1).

Another consequence of Theorem 2.1 is the following result which gives a sufficient condition for the almost sure convergence of series in terms of ρ -mixing coefficients and provides a class of examples of amarts which include Markov processes satisfying an L_2 operator condition and functions of some Gaussian sequences (see Bradley (1986)). **COROLLARY 2.2** Let $(X_i)_{i\geq 1}$ be a centered sequence of random variables satisfying

(2.3)
$$\sum_{i} \rho(2^{i}) < \infty \quad and \quad \sum_{i} EX_{i}^{2} < \infty.$$

Then $(S_{\tau})_{\tau \in T}$ is convergent in L_2 and therefore $(S_n)_{n \geq 1}$ is an amart which is convergent a.s. and in L_2 .

REMARK 2.1 As a consequence of Corollaries 2.1 and 2.2 we can easily see that the condition $\sum_{n}^{n} \rho(2^{n}) < \infty$ can replace the condition of independence in many almost sure results for sums of random variables. For instance most of the laws of large numbers from the Chapter IX in Petrov (1975) hold with about the same proof, including the sufficiency part of the three series theorem, the sufficiency part of the Kolmogorov and Feller strong laws of large numbers, almost sure convergence of a kernel based recursive procedures. See, for instance Roussas (1991), (1992). We denote by $Q_X(u) = \inf\{t : P(|X| > t) \le u\}$, the quantile function of X. We shall establish next

PROPOSITION 2.1 Let $(X_i)_{i\geq 1}$ be a centered sequence of random variables, such that $EX_i^2 < \infty$ for all $i \geq 1$. Let τ be a stopping time. Then

$$ES_{\tau}^{2}I(\tau \leq n) \leq 4\sum_{k=1}^{n} EX_{k}^{2}$$

+16 $\int_{0}^{1}\sum_{i=1}^{n} I(2\tilde{\alpha}_{i} > u) \sum_{j=1}^{n} Q_{X_{j}}^{2}(u) du.$

We shall estimate now the second moment of stopped partial sums in terms of the strong mixing coefficients and moments of the individual summands. Notice that if $\sum_{k} k^{2/\delta} \alpha(k) < \infty$

then $c(\alpha, n, \delta)$ in the next theorem is a numerical constant which does not depend on n. This summability condition imposed on the strong mixing coefficients is implied by the condition $\sum_{k} \alpha(k)^{\delta/(2+\delta)} < \infty$ which is widely used to bound the variance of S_n . This improvement is possible due to a recent result by Rio (1993). Next theorem gives an estimate for the second

moment of $S_{\tau \wedge n}$ for strong mixing sequences which is up to a multiplicative constant the same as for the Var (S_n) .

THEOREM 2.2 Let $(X_i)_{i\geq 1}$ be a centered sequence of random variables such that for a certain $\delta > 0$, $E|X_i|^{2+\delta} < \infty$ for every $i \geq 1$. Let τ be a stopping time.

(2.4)
$$c(\alpha, n, \delta) = 16 \left[(4\delta^{-1} + 2) \sum_{k=1}^{n} (k+1)^{2/\delta} \tilde{\alpha}(k) \right]^{\delta/(2+\delta)}$$

Then for every $n \ge 1$ we have

Domoto

(2.5)
$$ES_{\tau}^{2}I(\tau \le n) \le 4\sum_{k=1}^{n} EX_{k}^{2} + c(\alpha, n, \delta)\sum_{k=1}^{n} \|X_{k}\|_{2+\delta}^{2}.$$

With a proof similar to that of Corollary 2.2 we obtain the following class of convergent amarts which is a subclass of strongly mixing sequences.

COROLLARY 2.3 Assume $(X_i)_{i\geq 1}$ is a centered sequence of random variables such that for a certain $\delta > 0$, $E|X_i|^{2+\delta} < \infty$ for every $i \geq 1$. Assume that $\sum_k k^{2/\delta} \tilde{\alpha}(k) < \infty$ and $\sum_k ||X_k||^2_{2+\delta} < \infty$. Then $(S_{\tau})_{\tau \in T}$ is convergent in L_2 and therefore $(S_n)_{n\geq 1}$ is an amart which is convergent a.s. and in L_2 .

As an important consequence of Proposition 2.1 and of Theorem 2.2 we obtain the following maximal inequalities for partial sums of a strongly mixing sequence. Related results were obtained by Shao (1993) and Rio (1995).

COROLLARY 2.4 Let $(X_i)_{i\geq 1}$ be a centered sequence of random variables such that $E|X_i|^2 < \infty$ for every $i \geq 1$. Then, for every $\lambda > 0$ and every $n \geq 1$ we have:

(2.6)
$$\lambda^2 P(\max_{1 \le i \le n} |S_i| > \lambda) \le 4 \sum_{i=1}^n EX_i^2 + 16 \int_0^1 \sum_{i=1}^n I(2\tilde{\alpha}_i > u) \sum_{j=1}^n Q_{X_j}^2(u) du.$$

If $E|X_i|^{2+\delta} < \infty$ for every $i \ge 1$ we have:

(2.7)
$$\lambda^2 P\left(\max_{1 \le i \le n} |S_i| > \lambda\right) \le 4 \sum_{i=1}^n E X_i^2 + c(\alpha, n, \delta) \sum_{j=1}^n \|X_j\|_{2+\delta}^2,$$

where $c(\alpha, n, \delta)$ is defined in Theorem 2.2.

For the case of φ -mixing coefficients we shall establish:

THEOREM 2.3 Assume $(X_i)_{i\geq 1}$ is a sequence of random variables such that $\varphi(1) < 1$. Then

- (i) $\sup_{n \to \infty} E|S_n| < \infty$ implies $(S_n)_{n \ge 1}$ is a semiamart;
- (ii) $(S_n)_{n\geq 1}$ is convergent in L_1 then $(S_{\tau})_{\tau\in T}$ is convergent in L_1 and therefore $(S_n)_{n\geq 1}$ is an amart which is convergent a.s.

3 Proofs

The proofs of Theorems 2.1 and 2.2 are based on the following lemma which is inspired by Garsia's version of Doob's maximal inequality (1973).

LEMMA 3.1 Let $(X_k)_{k\geq 1}$ be a sequence of random variables and let τ be a stopping time. Then for every $n \geq 1$,

(i)

$$S_{\tau}I(\tau \le n) = S_nI(\tau \le n) - \sum_{k=1}^{n-1} X_{k+1}I(\tau \le k),$$

(ii)

$$S_{\tau}^{2}I(\tau \le n) \le S_{n}^{2}I(\tau \le n) - 2\sum_{k=1}^{n-1} S_{k}I(\tau = k)(S_{n} - S_{k}),$$

and

(iii)

$$S_{\tau}^{2}I(\tau \leq n) \leq 2S_{n}^{2}I(\tau \leq n) + 2\sum_{k=1}^{n-1} X_{k+1}^{2}I(\tau \leq k) + 4\sum_{k=1}^{n-2} X_{k+1}I(\tau \leq k)(S_{n} - S_{k+1}).$$

PROOF. In order to prove (i) we just observe that

$$S_n I(\tau \le n) = \sum_{k=0}^{n-1} S_{k+1} (I(\tau \le k+1) - I(\tau \le k)) + \sum_{k=1}^{n-1} (S_{k+1} - S_k) I(\tau \le k) = \sum_{k=0}^{n-1} S_{k+1} I(\tau = k+1) + \sum_{k=1}^{n-1} X_{k+1} . I(\tau \le k)$$

Now (iii) results from (i) by trivial computations and the fact that $(\tau \leq i) \subset (\tau \leq j)$ for every $i \leq j$.

In order to prove (ii) we have to remark only that (i) also implies

$$S_n^2 I(\tau \le n) \ge S_\tau^2 I(\tau \le n) + 2S_\tau \sum_{j=1}^{n-1} X_{j+1} I(\tau \le j)$$

= $S_\tau^2 I(\tau \le n) + 2\sum_{j=1}^{n-1} \sum_{i=1}^j X_{j+1} S_i I(\tau = i).$

In order to prove Theorem 2.1 we also need the following analytical result which is a reformulation of Theorem 7 in Bradley and Utev (1994) to incorporate Remark 1 in the same paper.

LEMMA 3.2 For any two L_2 -integrable random variables U and V, denote $r(U,V) = EUV/||U||_2||V||_2$. Given two sequences of L_2 integrable random variables $(Y_i)_{i\geq 1}$ and $(Z_i)_{i\geq 1}$ define

$$r_{1}(s) = \sup \left\{ r\left(\sum_{i=a+1}^{a+t} Y_{i}, \sum_{j=a+t+s+1}^{a+t+s+\ell} Y_{j}\right), a \ge 1, t \ge 1, \ell \ge 1 \right\}, r_{2}(s) = \sup \left\{ r\left(\sum_{i=a+1}^{a+t} Z_{i}, \sum_{j=a+t+s+1}^{a+t+s+\ell} Z_{j}\right), a \ge 1, t \ge 1, \ell \ge 1 \right\}, r_{3}(s) = \sup \left\{ r\left(\left(\sum_{i=a+1}^{a+t} Y_{i}, \sum_{j=a+t+s+1}^{a+t+s+\ell} Z_{j}\right), a \ge 1, t \ge 1, \ell \ge 1 \right\},$$

and $r(u) = \max(r_1(u), r_2(u), r_3(u))$. Denote $c(u, n) = \exp\left(2\sum_{j=0}^{\lfloor \log_2 n \rfloor} r(\lfloor 2^{j/3} \rfloor)\right)$. Then there is an absolute constant K such that for every $a \ge 1, n \ge 1$:

$$\left|\sum_{i=a+1}^{a+n} EY_i(Z_{i+1} + \ldots + Z_n)\right| \le Kc^2(u,n) \left(\sum_{i=a+1}^{a+n} EY_i^2\right)^{1/2} \left(\sum_{j=a+1}^{a+n} EZ_j^2\right)^{1/2}$$

PROOF OF THEOREM 2.1

We shall apply the inequality (ii) of Lemma 3.1 to the sequence $(X_i)_{i\geq 1}$ and get:

(3.1)
$$ES_{\tau}^{2}I(\tau \leq n) \leq ES_{n}^{2} + 2\left|\sum_{k=1}^{n-1} ES_{k}I(\tau = k)(S_{n} - S_{k})\right|.$$

By Theorem 1.1 in Utev (1991), we can find a constant C_1 such that

(3.2)
$$ES_n^2 \le C_1 \cdot c(\rho, n) \sum_{k=1}^n EX_k^2.$$

We apply now Lemma 3.2 to the sequences

$$Y_k = S_k I(\tau = k)$$
 and $Z_k = X_k$

and observe that $r_1(u) = 0$ for every $u \ge 1$ and since all Z_k are all centered, it follows that $r_2(u) \le \rho(u)$ and $r_3(u) \le \rho(u)$ for every $u \ge 1$. Therefore we can find a positive constant K such that:

(3.3)
$$\left| \sum_{k=1}^{n-1} ES_k I(\tau = k) (S_n - S_k) \right| \leq Kc^2(\rho, n) \left(\sum_{k=1}^n ES_k^2 I(\tau = k) \sum_{j=1}^n EX_j^2 \right)^{1/2}$$
$$= Kc^2(\rho, n) \left(ES_\tau^2 I(\tau \le n) \sum_{j=1}^n EX_j^2 \right)^{1/2}.$$

By the trivial inequality $2ab \leq a^2d^{-1} + b^2d$, for every d > 0, applied with $a^2 = ES_{\tau}^2I(\tau \leq n)$, $b^2 = \sum_{k=1}^n EX_k^2$ and $d = 2Kc^2(\rho, n)$ we get

(3.4)
$$\left|\sum_{k=1}^{n-1} ES_k I(\tau = k) (S_n - S_k)\right| \le 4^{-1} ES_{\tau}^2 I(\tau \le n) + K^2 c^4(\rho, n) \sum_{k=1}^n EX_k^2,$$

and the result follows by combining (3.1), (3.2) and (3.4). \Box

PROOF OF COROLLARY 2.1

Let $\lambda > 0$, and denote by $\tau = \inf\{i \ge 1; |S_i| \ge \lambda\}$. According to Theorem 2.1 there is a constant K such that for every $n \ge 1$:

$$ES_{\tau}^{2}I(\tau \leq n) \leq Kc^{4}(\rho, n) \sum_{j=1}^{n} EX_{j}^{2}.$$

We have only to remark that by the definition of τ

$$I(\tau \le n) = I\left(\max_{1 \le i \le n} |S_i| \ge \lambda\right),$$

and

$$|S_{\tau}|I(\tau \le n) \ge \lambda I(\tau \le n)$$

for every $n \geq 1$. \Box

PROOF OF COROLLARY 2.2

Let τ be a finite stopping time, $\tau \geq m$. By Theorem 2.1 we can find a constant K such that:

(3.5)
$$E(S_{\tau} - S_m)^2 \le K \sum_{j \ge m} EX_j^2$$

It is easy to see that (3.5) implies that $(S_{\tau})_{\tau \in T}$ is Cauchy sequence in L_2 and therefore $(ES_{\tau})_{\tau \in T}$ is convergent. By the Definition 1.2 $(S_n)_{n \geq 1}$ is an amart. We also remark that, by (3.5), S_n is bounded in L_2 and therefore by Theorem 2 in Austin, Edgar and Ionescu Tulcea (1974), S_n is convergent a.e. By using (3.5) once again we see that S_n is also convergent in L_2 . \Box

In order to prove Proposition 2.1 we need the following lemma which is a consequence of Theorem 1.1 in Rio (1983).

LEMMA 3.3 For any square integrable random variables Y, Z

$$|cov(Y,Z)| \le 2\int_0^1 I(2\alpha > u)Q_Y(u)Q_Z(u)du,$$

where $\alpha = \alpha(\sigma(Y), \sigma(Z)).$

PROOF OF PROPOSITION 2.1

In order to establish this result, we shall use the relation (iii) of Lemma 3.1 where we expand ES_n^2 . We have

$$ES_{\tau}^{2}I(\tau \le n) \le 4\sum_{i=1}^{n} EX_{i}^{2} + 4\sum_{k=0}^{n-2}\sum_{j=k+2}^{n} (|E(X_{k+1}X_{j})| + |E(X_{k+1}I(\tau \le k)X_{j})|.$$

We have now to apply Lemma 3.3 twice and to take into account that $Q_{X_{k+1}}I(\tau \le k) \le Q_{X_{k+1}}$. We obtain

$$ES_{\tau}^{2}I(\tau \le n) \le 4\sum_{i=1}^{n} EX_{i}^{2} + 8\sum_{k=0}^{n-2}\sum_{j=k+2}^{n}\int_{0}^{1}I(2\alpha_{j-k+1} > u)(Q_{X_{k+1}}^{2}(u) + Q_{X_{j}}^{2}(u))du,$$

whence the result follows by easy computations involving the change of the order of summation.

PROOF OF THEOREM 2.2

A careful reader can easily obtain this Theorem from Proposition 2.1 and the proof of Theorem 1.2 and its consequences, in Rio (1993). For convenience we shall sketch the proof. By Cauchy-Schwartz inequality, for all $1 \le k \le n$,

$$\begin{split} \int_0^1 \sum_{j=1}^n I(2\tilde{\alpha}(j) > u) Q_{X_k}^2(u) du &\leq \\ &\leq \left[\int_0^1 \sum_{j=1}^n I(2\tilde{\alpha}(j) > u)^{\frac{(2+\delta)}{\delta}} du \right]^{\frac{\delta}{(2+\delta)}} \left(\int_0^1 Q_{X_k}^{2+\delta}(u) du \right)^{\frac{2}{2+\delta}}. \end{split}$$

We notice that $\int_0^1 Q_{X_k}^{2+\delta}(u) du = E|X_k|^{2+\delta}$. In order to estimate the first integral from the right hand side of the inequality, we change the variable μ to $\nu = \sum_{j=1}^n I(2\tilde{\alpha}(j) > u)$ and observe that $\nu = j$ iff $2\tilde{\alpha}(j+1) \leq u < 2\tilde{\alpha}(j)$. After an integration by parts, we obtain

$$\int_{1}^{0} \sum_{j=1}^{n} I(2\tilde{\alpha}(j) > u) Q_{X_{k}}^{2}(u) du \leq$$
$$\leq c(\delta) \left(\sum_{i=1}^{n} (i+1)^{\frac{2}{\delta}} \tilde{\alpha}(i) \right)^{\frac{2}{(2+\delta)}} \|X_{k}\|_{2+\delta}^{2}.$$

where $c(\delta) = (4\delta^{-1} + 2)^{\delta/(2+\delta)}$. In order to establish Theorem 2.2 we just combine this last inequality with Proposition 2.1.

PROOF OF COROLLARY 2.4

This corollary follows immediately from Proposition 2.1, Theorem 2.2 and the proof of Corollary 2.1.

PROOF OF THEOREM 2.3

Let τ be a finite stopping time such that $\tau \leq n$ a.e. for some $n \geq 1$. Then

$$P(|S_{\tau}| > 2x) \le P(|S_n| > x) + \sum_{i=1}^{n} P(|S_n - S_i| > x, |S_i| > 2x, \tau = i)$$

which implies

(3.6)
$$\left(1 - \varphi(1) - \max_{1 \le i \le n} P(|S_n - S_i| > x)\right) P(|S_\tau| > 2x) \le P(|S_n| > x).$$

If $\sup_{n} E|S_n| < \infty$, then for some constant $K \ge 0$,

$$\max_{1 \le i \le n} P(|S_n - S_i| > \lambda) \le \frac{K}{\lambda}, \quad \text{for every} \quad n \ge 1$$

and, so for λ sufficiently large we can find a constant $\eta > 0$ such that for every $x > \lambda$ we have,

$$1 - \varphi(1) - \max_{1 \le i \le n} P(|S_n - S_i| > x) > \eta.$$

As a consequence, by (3.6), for every $x \ge \lambda$ we get:

$$\eta P(|S_\tau| > 2x) \le P(|S_n| > x)$$

By integrating $P(|S_{\tau}| > 2x)$ with respect to x from 0 to ∞ , bounding the integrand by 1 on the interval $[0, \lambda]$ and by previous inequality on $[\lambda, \infty]$ we obtain

$$|E|S_{\tau}| \le 2\lambda + 2\eta^{-1}E|S_n|,$$

which is uniformly bounded for every finite stopping time. This proves that $(S_n)_{n\geq 1}$ is a semiamart.

Assume now that $(S_n)_{n\geq 1}$ is convergent in L_1 . Fix $\epsilon > 0$, $\epsilon < 1 - \varphi(1)$. We can find an integer $m \geq 1$ such that for every $k \geq m$ and every $x > \epsilon$ we have

$$(3.7) P(|S_k - S_m| > x) \le \epsilon.$$

Let τ be a finite stopping time, $m \leq \tau \leq n$ a.e. By (3.6) and (3.7) for every $x > \epsilon$, we have

(3.8)
$$(1 - \varphi(1) - \epsilon)P(|S_{\tau} - S_m| > 2x, \quad m \le \tau \le n) \le P(|S_n - S_m| > x)$$

By integrating $P(|S_{\tau} - S_m| > 2x)$ with respect to x from 0 to ∞ , bounding the integrand by 1 on the intervals $[0, \epsilon]$ and by (3.8) on $[\epsilon, \infty]$ we get

$$|E|S_{\tau} - S_m| \le 2\epsilon + 2(1 - \varphi(1) - \epsilon)^{-1}E|S_n - S_m|.$$

Therefore $(S_{\tau})_{\tau \in T}$ is Cauchy in L_1 and as a consequence $(E|S_{\tau}|)_{\tau \in T}$ is convergent. This proves that $(S_n)_{n\geq 1}$ is an amart bounded in L_1 and the result follows as in Corollary 2.2.

Acknowledgement. The author would like to thank the referee for carefully reading the paper and for many valuable comments which contributed to the improvement of the presentation of this paper.

References

- Austin, D.G., Edgar, G.A. and Ionescu Tulcea, A., (1974). Pointwise convergence in terms of expectations. Z. Wahrsch. verv Gebiete 30, 17-26.
- [2] Berbee, H.C.P. (1979). Random Walks with Stationary Increments and Renewal Theory, Mathematisch Centrum, Amsterdam.
- [3] Billingsley, P. (1968). Convergence of Probability Measures, Wiley.
- [4] Bradley, R.C. (1986). Basic properties of strong mixing conditions, Progress in Prob. and Stat., Dependence in Prob. and Stat., 11, E. Eberlein and M. Taqqu (eds.), 165-192.

- [5] Bradley, R.C., Utev, S.A. (1994). On second order properties of mixing sequences of random fields, *Prob. Theory and Math. Stat.*, B. Grigelionis et al (eds.), 99-120, VSP/TEV.
- [6] Doukhan, P. (1994). Mixing properties and examples, *Lecture Notes in Statistics* 85, Springer Verlag.
- [7] Edgar, G.A. and Sucheston, L. (1976). Amarts: A class of asymptotic martingales, J. Multiv. Anal., 6, 193-221.
- [8] Edgar, G.A., Sucheston, L. (1992). Stopping times and directed processes, *Encyclopedia of Mathematics and its Applications*, Cambridge University Press.
- [9] Garcia, A. (1973). On a convex function inequality for submartingales, Ann. Probab., 1, 171-174.
- [10] Gut, A., Schmidt, K. (1983). Amarts and set function processes, *Lecture Notes in Math* 1042, Springer-Verlag.
- [11] Gut, A. (1986). Stopped Random Walks Limit Theorems and Applications, Applied Probability, Springer-Verlag.
- [12] Houdre, C. (1995). On the almost sure convergence of series of stationary and related nonstationary variables, Ann. Probab., 23, 1204-1218.
- [13] Ibragimov, I.A., Linnik, Yu. V. (1971). Independent and stationary sequences of random variables, Walters-Noordhoff, Groningen, the Netherlands.
- [14] Krengel, V., Sucheston, L. (1978). On semiamarts, amarts and processes with finite value, Advances in Prob., 4, 197-266.
- [15] McLeish, D.L. (1975). A maximal inequality and dependent strong laws. Ann. Probab., 3, 829-39.
- [16] Moricz, F. (1976). Moment inequalities and strong laws of large numbers, Z. Wahrsch. Verw. Gebiete, 35, 299-314.
- [17] Newman, C., Wright, A. (1982). Associated random variables and martingale inequalities, Z. Wahr. verw. Gebiete, 59, 361-371.
- [18] Peligrad, M. (1985a). An invariance principle for φ -mixing sequences, Ann. Probab., 13, 1304-1313.
- [19] Peligrad, M. (1985b). Convergence rates of the strong law for stationary mixing sequences. Z. Wahrsch. verw. Gebiete, 70, 307-314.
- [20] Peligrad, M. (1992). On the central limit theorem for weakly dependent sequences with a decomposed strong mixing coefficient. *Stochastic Proc. Appl.*, 42, 181-193.
- [21] Petrov, V.V. (1975). Sums of independent random variables, Springer Verlag.
- [22] Rio, E. (1993). Covariance inequalities for strongly mixing processes, Ann. Inst. H. Poincaré, 29, 4, 589-597.

- [23] Rio, E. (1995). A maximal inequality and dependent Marcinkiewicz-Zygmund strong laws, Ann. Probab. 23, 918-937.
- [24] Roussas, G.G. (1991). Recursive estimation of the transition distribution function of a Markov process: asymptotic normality, *Statist. Probab. Lett.*, **11**, 435-447.
- [25] Roussas, G.G. (1992). Exact rates of almost sure convergence of a recursive kernel estimate of a probability density function: application to regression and hazard rate estimation. J. Nonpar. Statist., 1, 171-195.
- [26] Shao, Q. (1995). Maximal inequalities for partial sums of ρ-mixing sequences. Ann. Probab., 23, 948-965.
- [27] Shao, Q. (1993). Complete convergence for α -mixing sequences, Stat. and Prob. Letters 16, 279-287.
- [28] Utev, S.A. (1991). Sums of random variables with φ -mixing, Siberian Adv. Math. 1, 124-155.

Department of Mathematical Sciences University of Cincinnati PO Box 210025 Cincinnati, OH 45221-0025 peligrm@math.uc.edu