

Convergence rate analysis of an asynchronous space decomposition method for convex minimization

Tai, Xue Cheng; Tseng, Paul

2001

Tai, X. C., & Tseng, P. (2001). Convergence rate analysis of an asynchronous space decomposition method for convex minimization. *Mathematics of Computation*, 71(239), 1105-1135.

<https://hdl.handle.net/10356/90754>

<https://doi.org/10.1090/S0025-5718-01-01344-8>.

Mathematics of Computation © copyright 2001 American Mathematical Society. The journal's website is located at <http://www.ams.org/mcom/>.

Downloaded on 23 Aug 2022 12:56:19 SGT

CONVERGENCE RATE ANALYSIS OF AN ASYNCHRONOUS SPACE DECOMPOSITION METHOD FOR CONVEX MINIMIZATION

XUE-CHENG TAI AND PAUL TSENG

ABSTRACT. We analyze the convergence rate of an asynchronous space decomposition method for constrained convex minimization in a reflexive Banach space. This method includes as special cases parallel domain decomposition methods and multigrid methods for solving elliptic partial differential equations. In particular, the method generalizes the additive Schwarz domain decomposition methods to allow for asynchronous updates. It also generalizes the BPX multigrid method to allow for use as solvers instead of as preconditioners, possibly with asynchronous updates, and is applicable to nonlinear problems. Applications to an overlapping domain decomposition for obstacle problems are also studied. The method of this work is also closely related to relaxation methods for nonlinear network flow. Accordingly, we specialize our convergence rate results to the above methods. The asynchronous method is implementable in a multiprocessor system, allowing for communication and computation delays among the processors.

1. INTRODUCTION

With the advent of multiprocessor computing systems, there has been much work in the design and analysis of iterative methods that can take advantage of the parallelism to solve large linear and nonlinear algebraic problems. In these methods, the computation per iteration is distributed over the processors and each processor communicates the result of its computation to the other processors. In some systems, the activities of the processors are highly synchronized (possibly via a central processor), while in other systems (typically those with many processors), the processors may experience communication or computation delays. The latter lack of synchronization makes the analysis of the methods much more difficult. To aid in this analysis, Chazan and Miranker [16] proposed a model of asynchronous computation that allows for communication and computation delays among processors, and they showed that the Jacobi method for solving a diagonally dominant

Received by the editor September 27, 2000.

2000 *Mathematics Subject Classification.* Primary 65J10, 65M55, 65Y05; Secondary 65K10, 65N55.

Key words and phrases. Convex minimization, space decomposition, asynchronous computation, convergence rate, domain decomposition, multigrid, obstacle problem.

The work of the first author was supported by the Norwegian Research Council Strategic Institute Program within Inverse Problems at RF-Rogaland Research, and by Project SEP-115837/431 at Mathematics Institute, University of Bergen.

The work of the second author was supported by the National Science Foundation, Grant No. CCR-9311621.

system of linear equations converges under this model of asynchronous computation. Subsequently, there has been extensive study of asynchronous methods based on such a model (see [5, 6] and references therein). For these methods, convergence typically requires the algorithmic mapping to be either isotone or nonexpansive with respect to the L^∞ -norm or gradient-like. However, aside from the easy case where the algorithmic mapping is a contraction with respect to the L^∞ -norm, there have been few studies of the convergence rate of these methods. One such study was done in [55] for an asynchronous gradient-projection method.

In this paper, we study the convergence rate of asynchronous Jacobi and Gauss-Seidel type methods for finite- or infinite-dimensional convex minimization of the form

$$(1) \quad \min_{v_i \in K_i, i=1, \dots, m} F \left(\sum_{i=1}^m v_i \right),$$

where each K_i is a nonempty closed convex set in a real reflexive Banach space V and F is a real-valued lower semicontinuous Gâteaux-differentiable function that is strongly convex on $\sum_{i=1}^m K_i$. Our interest in these methods stems from their close connection to relaxation methods for nonlinear network flow (see [4, 5, 56] and references therein) and to domain decomposition (DD) and multigrid (MG) methods for solving elliptic partial differential equations (see [7, 8, 9, 14, 18, 19, 33, 40, 45, 52, 53, 57] and references therein). For example, the additive and the multiplicative Schwarz methods may be viewed as Jacobi and Gauss-Seidel type methods applied to linear elliptic partial differential equations reformulated as (1) [9, 57]. DD and MG methods are also useful as preconditioners and it can be shown that such preconditioning improves the condition number of the discrete approximation [7, 8, 10, 9, 14, 33, 40, 45, 57]. In addition, DD and MG methods are well suited for parallel implementation, for which both synchronous and asynchronous versions have been proposed. Of the work on asynchronous methods [21, 22, 27, 38, 37, 39, 46], we especially mention the numerical tests by Frommer et al. [22] which showed that, through improved load balancing, asynchronous methods can be advantageous in solving even simple linear equations. Although these tests did not use the coarse mesh in its implementation of the DD method, it is plausible that the asynchronous method would still be advantageous when the coarse mesh is used. However, the convergence rate analysis of the above asynchronous methods seems still missing from the literature. In the case where the equation is linear (corresponding to F being quadratic and K_1, \dots, K_m being suitable subspaces of V) or almost linear, this issue has been much studied for synchronous methods (see [7, 8, 9, 14, 18, 19, 33, 40, 45, 52, 53, 57] and references therein) but little studied for asynchronous methods. In the case where the equation is generally nonlinear (corresponding to K_1, \dots, K_m being suitable subspaces of V), there are some convergence studies for synchronous methods [15, 18, 44, 52, 53], and none for asynchronous methods. In the case where K_1, \dots, K_m are not all subspaces, there are various convergence studies for synchronous methods (see [1, 12, 23, 25, 28, 29, 30, 31, 34, 35, 36, 47, 50] and references therein) but, again, none for asynchronous methods.

The contributions of the present work are two-fold.

- We consider an asynchronous version of Jacobi and Gauss-Seidel methods for solving (1), and we show that, under a Lipschitzian assumption on the

Gâteaux derivative F' and a norm equivalence assumption on the product of K_1, \dots, K_m and their sum (see (5) and (6)), this asynchronous method attains a global linear rate of convergence with a convergence factor that can be explicitly estimated (see Theorem 1). This provides a unified convergence and convergence rate analysis for such asynchronous methods.

- We apply the above convergence result to (finite-dimensional) linearly constrained convex programs and, in particular, nonlinear network flow problems. This yields convergence rate results for some asynchronous network relaxation methods (see Section 6). Previous work studied the convergence of these methods, but no rate of convergence result was obtained. We also apply the above convergence result to certain nonlinear elliptic partial differential equations. This yields convergence rate results for some asynchronous parallel DD and MG methods for solving these equations and, in particular, the convergence factor is shown not to depend on the mesh parameters (see Section 7). When implementing multigrid methods on parallel processors, the nodal basis is often organized into different groups. The computation within each group can be sequential while the computation in different groups could be done in parallel. The asynchronous convergence rate analysis provides a convergence rate estimate when computation in different groups is not fully synchronized. Lastly, application to an overlapping DD method for obstacle problems is studied. We show that the method attains a linear rate of convergence with a convergence factor depending on the overlapping size, but not on the mesh size or the number of subdomains.

We note that alternative approaches such as Newton-type methods have also been applied to develop synchronous DD and MG methods for nonlinear partial differential equations without constraints [2, 3, 11, 26, 41, 58, 59]. However, these methods use the traditional DD and MG approach or use a special two-grid treatment. Our approach is different even for nonlinear partial differential equations without constraints.

2. PROBLEM DESCRIPTION AND SPACE DECOMPOSITION

Let V be a real reflexive Banach space with norm $\|\cdot\|$ and let V' be its dual space, i.e., the space of all real-valued linear continuous functionals on V . The value of $f \in V'$ at $v \in V$ will be denoted by $\langle f, v \rangle$, i.e., $\langle \cdot, \cdot \rangle$ is the duality pairing of V and V' . We wish to solve the minimization problem

$$(2) \quad \min_{v \in K} F(v) ,$$

where K is a nonempty closed (in the strong topology) convex set in V and $F : V \mapsto \Re$ is a lower semicontinuous convex Gâteaux-differentiable function. We assume F is strongly convex on K or, equivalently, its Gâteaux derivative $\lim_{t \rightarrow 0} (F(v + tw) - F(v))/t$, which is a well-defined linear continuous functional of w denoted by $F'(v)$ (so $F' : V \mapsto V'$), is strongly monotone on K , i.e.,

$$(3) \quad \langle F'(u) - F'(v), u - v \rangle \geq \sigma \|u - v\|^2, \quad \forall u, v \in K,$$

where $\sigma > 0$. It is known that, under the above assumptions, (2) has a unique solution \bar{u} [24, p. 23].

We assume that the constraint set K can be decomposed as the Minkowski sum

$$(4) \quad K = \sum_{i=1}^m K_i$$

for some nonempty closed convex sets K_i in V , $i = 1, \dots, m$. This means that, for any $v \in K$, we can find $v_i \in K_i$, not necessarily unique, satisfying $\sum_{i=1}^m v_i = v$ and, conversely, for any $v_i \in K_i$, $i = 1, \dots, m$, we have $\sum_{i=1}^m v_i \in K$. Following Xu [57], we call (4) a space decomposition of K , with the term “space” used loosely here. Then we may reformulate (2) as the minimization problem (1), with $(\bar{u}_1, \dots, \bar{u}_m)$ being a solution (not necessarily unique) of (1) if and only if $\bar{u}_i \in K_i$ for $i = 1, \dots, m$ and $\sum_{i=1}^m \bar{u}_i = \bar{u}$. As was noted earlier, the reformulated problem (1) is of interest because methods such as DD and MG methods may be viewed as Jacobi and Gauss-Seidel methods for its solution. The method we study will be an asynchronous version of these methods. The above reformulation was proposed in [9, 57] (for the case where F is quadratic and $K = V$) to give a unified analysis of DD and MG methods for linear elliptic partial differential equations. The general case was treated in [47, 50] (also see [48, 52] for the case of $K = V$).

For the above space decomposition, we will assume that there is a constant $C_1 > 0$ such that for any $v_i \in K_i, i = 1, \dots, m$, there exists $\bar{u}_i \in K_i$ satisfying

$$(5) \quad \bar{u} = \sum_{i=1}^m \bar{u}_i \quad \text{and} \quad \left(\sum_{i=1}^m \|\bar{u}_i - v_i\|^2 \right)^{\frac{1}{2}} \leq C_1 \left\| \bar{u} - \sum_{i=1}^m v_i \right\|$$

(see [14, p. 95], [50, 52], [57, Lemma 7.1] for similar assumptions). We will also assume F' has a weak Lipschitzian property in the sense that there is a constant $C_2 > 0$ such that

$$(6) \quad \sum_{i=1}^m \sum_{j=1}^m \langle F'(w_{ij} + u_{ij}) - F'(w_{ij}), v_i \rangle \leq C_2 \left(\sum_{j=1}^m \max_{i=1, \dots, m} \|u_{ij}\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|v_i\|^2 \right)^{\frac{1}{2}},$$

$$\forall w_{ij} \in K, u_{ij} \in K_j^\ominus, v_i \in K_i^\ominus, i, j = 1, \dots, m,$$

where we define the set difference $K_i^\ominus = \{u - v : u, v \in K_i\} \subset V$. The above assumption generalizes those in [50, 52, 53] for the case of K_i being a subspace, for which $K_i^\ominus = K_i$.

Furthermore, we will paint each of the sets K_1, \dots, K_m one of c colors, with the colors numbered from 1 up to c , such that sets painted the same color $k \in \{1, \dots, c\}$ are orthogonal in the sense that

$$(7) \quad \left\| \sum_{i \in I(k)} v_i \right\|^2 = \sum_{i \in I(k)} \|v_i\|^2, \quad \forall v_i \in K_i^\ominus, i \in I(k),$$

$$(8) \quad \left\langle F' \left(u + \sum_{i \in I(k)} v_i \right), \sum_{i \in I(k)} v_i \right\rangle \leq \sum_{i \in I(k)} \langle F'(u + v_i), v_i \rangle,$$

$$\forall u \in K, v_i \in K_i^\ominus, i \in I(k),$$

where $I(k) = \{i \in \{1, \dots, m\} : K_i \text{ is painted color } k\}$ (see [14, §4.1], [53] for similar orthogonal decompositions in the case K_i is a subspace). Thus $I(1), \dots, I(c)$ are disjoint subsets of $\{1, \dots, m\}$ whose union is $\{1, \dots, m\}$ and $I(k)$ comprises the indexes of the sets painted the color k . Although $c = m$ is always a valid choice, in

some of the applications that we will consider, it is essential that c be independent of m . In the context of a network flow problem, each set K_i may correspond to a node of the network and sets are painted different colors if their corresponding nodes are joined by an arc. In the context of a partial differential equation defined on a domain $\Omega \subset \mathbb{R}^d$, each set K_i may correspond to a subdomain of Ω and sets are painted different colors if their corresponding subdomains intersect (see Sections 6 and 7 for details).

Remark 1. It can be seen that condition (6) is implied by the following strengthened Cauchy-Schwarz inequality (also see [45, p. 155], [57] for the case of quadratic F and subspace K_i):

$$\langle F'(w_{ij} + u_{ij}) - F'(w_{ij}), v_i \rangle \leq \epsilon_{ij} \|u_{ij}\| \|v_i\|, \quad \forall w_{ij} \in K, u_{ij} \in K_j^\ominus, v_i \in K_i^\ominus,$$

with C_2 being the spectral radius of the matrix $\mathcal{E} = [\epsilon_{ij}]_{i,j=1}^m$, assumed to be symmetric.

Remark 2. For locally strongly convex problems, the constants σ, C_1, C_2 may depend on $u, v, v_i, w_{ij}, u_{ij}$. In this case, the subsequent analysis should be viewed as being local in nature, i.e., it is valid when the iterated solutions lie in a neighborhood of the true solution (see Section 7).

3. AN ASYNCHRONOUS SPACE DECOMPOSITION METHOD

Since F is lower semicontinuous and strongly convex, for each $(u_1, \dots, u_m) \in K_1 \times \dots \times K_m$ and each $i \in \{1, \dots, m\}$, there exists a unique $w_i \in K_i$ satisfying

$$(9) \quad F\left(\sum_{j \neq i} u_j + w_i\right) \leq F\left(\sum_{j \neq i} u_j + v_i\right), \quad \forall v_i \in K_i$$

(see [24, p. 23]). Let $\pi_i(u_1, \dots, u_m)$ denote this w_i . Then (π_1, \dots, π_m) may be viewed as the algorithmic mapping associated with the block Jacobi method for solving (1). Consider an asynchronous version of the block Jacobi method, parameterized by a stepsize $\gamma \in (0, 1]$, which for simplicity we assume to be fixed, that generates a sequence of iterates $(u_1(t), \dots, u_m(t)), t = 0, 1, \dots$, with $(u_1(0), \dots, u_m(0)) \in K_1 \times \dots \times K_m$ given, according to the updating formula,

$$(10) \quad u_i(t+1) = u_i(t) + \gamma s_i(t), \quad i = 1, \dots, m,$$

where we define

$$(11) \quad s_i(t) = \begin{cases} w_i(t) - u_i(t) & \text{if } t \in T^i \\ 0 & \text{otherwise,} \end{cases}$$

$$(12) \quad w_i(t) = \pi_i(u_1(\tau_1^i(t)), \dots, u_m(\tau_m^i(t))),$$

and T^i is some subset of $\{0, 1, \dots\}$ and each $\tau_j^i(t)$ is some nonnegative integer not exceeding t . Since each K_i is convex and $\gamma \in (0, 1]$, an induction argument shows that $(u_1(t), \dots, u_m(t)) \in K_1 \times \dots \times K_m$ for all $t = 0, 1, \dots$.

We will assume that the iterates are updated in a *partially asynchronous* manner [5, Chap. 7], i.e., there exists an integer $B \geq 1$ such that

$$(13) \quad \{t, t+1, \dots, t+B-1\} \cap T^i \neq \emptyset, \quad t = 0, 1, \dots, \forall i,$$

$$(14) \quad 0 \leq t - \tau_j^i(t) \leq B-1 \quad \text{and} \quad \tau_i^i(t) = t, \quad \forall t \in T^i, \forall i, j.$$

We say that a color $k \in \{1, \dots, c\}$ is *active* at time t if there exists an $i \in I(k)$ such that $t \in T^i$. Recall that $I(k)$ indexes those sets painted the color k . Denoting by c_t the total number of colors that are active at time t , we will also assume that

$$(15) \quad \gamma < \min \left\{ \frac{\sigma}{2C_2B}, \frac{1}{c_t} \right\}, \quad t = 0, 1, \dots$$

Notice that γ does not depend on m nor on C_1 . Although (15) may give a very conservative value of γ , this can be remedied by starting with a larger γ and decreasing γ whenever “sufficient progress” (defined in any reasonable way) is not made and (15) is not satisfied.

Remark 3. The above asynchronous method models a situation in which computation is distributed over m processors with the i th processor being responsible for updating u_i and communicating the updated value to the other processors. T^i is the set of “times” at which u_i is updated by processor i (by applying π_i to its current copy of (u_1, \dots, u_m)); $u_i(t)$ is the value of u_i known to processor i at time t ; and $\tau_j^i(t)$ is the time at which the value of u_j used by processor i at time t is generated by processor j , so $t - \tau_j^i(t)$ is the communication delay from processor j to processor i at time t . Thus, the processors need not wait for each other when updating $(u_i)_{i=1}^m$, and the values used in the computation may be out-of-date.

Remark 4. The assumption that $\tau_i^i(t) = t$ can perhaps be removed through a more careful analysis, though this seems to be a reasonable assumption in practice. Intuitively, (13) says that each component u_i is updated at least once every B time units, and (14) says that the information used by processor i from processor j should not be out-of-date by more than B time units. This assumption of bounded communication and computation delay is needed for a convergence rate analysis.

4. CONVERGENCE RATE OF THE ASYNCHRONOUS METHOD

In this section we prove that the iterates $(u_1(t), \dots, u_m(t))$, $t = 0, 1, \dots$, generated by the asynchronous method (10)–(15) attain linear rate of convergence, with a factor that depends on σ, C_1, C_2, c and B, γ only (see Theorem 1). While parts of our proof use ideas from the analysis of asynchronous gradient-like methods [5, §7.5], [55], a number of new proof ideas are introduced to account for different problem assumptions and different natures of the Jacobi and Gauss-Seidel algorithmic mappings. To simplify the notation in our analysis, define

$$(16) \quad u(t) = \sum_{j=1}^m u_j(t), \quad z_i(t) = \sum_{j=1}^m u_j(\tau_j^i(t)),$$

for all i and t . If $t \in T^i$, then the definition (12) of $w_i(t)$ and the fact that $\tau_i^i(t) = t$ and F is Gâteaux-differentiable imply $w_i(t)$ satisfy the optimality condition

$$(17) \quad \langle F'(z_i(t) + w_i(t) - u_i(t)), v_i - w_i(t) \rangle \geq 0, \quad \forall v_i \in K_i.$$

Our analysis will be based on estimates given in the following two key lemmas.

Lemma 1 (Descent estimate). *Let A_1 and A_2 be defined by*

$$(18) \quad A_2 = \frac{C_2^2 B^2}{\sigma}, \quad A_1 = \frac{\sigma}{4} - \gamma^2 A_2.$$

For $t = 0, 1, \dots$, we have

$$F(u(t+B)) \leq F(u(t)) - \gamma A_1 \sum_{j=1}^m \sum_{\tau=t}^{t+B-1} \|s_j(\tau)\|^2 + \gamma^3 A_2 \sum_{j=1}^m \sum_{\tau=t-B+1}^{t-1} \|s_j(\tau)\|^2.$$

Proof. Fix any time $t \in \{0, 1, \dots\}$. Recall that c_t is the total number of colors active at time t and, without loss of generality, we assume that the first c_t colors are active. Then $s_i(t) = 0$ for all $i \in I(k)$ and $k > c_t$, so by defining

$$e_k(t) = \sum_{i \in I(k)} s_i(t)$$

and using (16), (10) and the convexity of F , we have

$$\begin{aligned} F(u(t+1)) &= F\left(u(t) + \gamma \sum_{i=1}^m s_i(t)\right) \\ &= F\left(u(t) + \gamma \sum_{k=1}^{c_t} \sum_{i \in I(k)} s_i(t)\right) \\ &= F\left((1 - c_t \gamma)u(t) + \sum_{k=1}^{c_t} \gamma(u(t) + e_k(t))\right) \\ &\leq (1 - c_t \gamma)F(u(t)) + \gamma \sum_{k=1}^{c_t} F(u(t) + e_k(t)) \\ (19) \quad &= F(u(t)) + \gamma \sum_{k=1}^{c_t} \left(F(u(t) + e_k(t)) - F(u(t))\right). \end{aligned}$$

Since $u(t) \in K$ and $u(t) + e_k(t) \in K$, the strong monotonicity of F' on K given in (3) implies

$$(20) \quad F(u(t)) \geq F(u(t) + e_k(t)) - \langle F'(u(t) + e_k(t)), e_k(t) \rangle + \frac{\sigma}{2} \|e_k(t)\|^2.$$

Define

$$\phi_j^i(t) = \sum_{k=1}^j u_k(\tau_k^i(t)) + \sum_{k=j+1}^m u_k(t), \quad j = 0, 1, \dots, m.$$

Then $\phi_0^i(t) = u(t)$ and $\phi_m^i(t) = z_i(t)$ and

$$\phi_j^i(t) - \phi_{j-1}^i(t) = u_j(\tau_j^i(t)) - u_j(t) \in K_j^\ominus, \quad j = 1, \dots, m.$$

If $t \in T^i$, then setting $v_i = u_i(t)$ in (17) and noting that $s_i(t) = w_i(t) - u_i(t)$ (see (11)), we obtain that

$$\begin{aligned} 0 &\leq -\langle F'(z_i(t) + s_i(t)), s_i(t) \rangle \\ &= -\langle F'(z_i(t) + s_i(t)) - F'(u(t) + s_i(t)), s_i(t) \rangle - \langle F'(u(t) + s_i(t)), s_i(t) \rangle \\ &= -\sum_{j=1}^m \langle F'(\phi_j^i(t) + s_i(t)) - F'(\phi_{j-1}^i(t) + s_i(t)), s_i(t) \rangle \\ &\quad - \langle F'(u(t) + s_i(t)), s_i(t) \rangle. \end{aligned}$$

If $t \notin T^i$, then $s_i(t) = 0$ and the above inequality holds trivially. Combining the above inequality with (7) and (9) and (20), we obtain that

$$\begin{aligned}
 (21) \quad & \sum_{k=1}^{c_t} \left(F(u(t) + e_k(t)) - F(u(t)) \right) \\
 & \leq \sum_{k=1}^{c_t} \sum_{i \in I(k)} \langle F'(u(t)) + s_i(t), s_i(t) \rangle - \frac{\sigma}{2} \sum_{k=1}^{c_t} \sum_{i \in I(k)} \|s_i(t)\|^2 \\
 & = \sum_{i=1}^m \langle F'(u(t)) + s_i(t), s_i(t) \rangle - \frac{\sigma}{2} \sum_{i=1}^m \|s_i(t)\|^2 \\
 & \leq - \sum_{i=1}^m \sum_{j=1}^m \langle F'(\phi_j^i(t) + s_i(t)) - F'(\phi_{j-1}^i(t) + s_i(t)), s_i(t) \rangle - \frac{\sigma}{2} \sum_{i=1}^m \|s_i(t)\|^2.
 \end{aligned}$$

Substituting (21) into (19) and using (6) yields

$$\begin{aligned}
 (22) \quad & F(u(t+1)) \leq F(u(t)) \\
 & \quad + \gamma C_2 \left(\sum_{j=1}^m \max_{i=1, \dots, m} \|u_j(\tau_j^i(t)) - u_j(t)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|s_i(t)\|^2 \right)^{\frac{1}{2}} \\
 & \quad - \gamma \frac{\sigma}{2} \sum_{i=1}^m \|s_i(t)\|^2.
 \end{aligned}$$

Since $t - B + 1 \leq \tau_j^i(t) \leq t$ for all i and j , we also have from (10) and the triangle inequality that

$$(23) \quad \|u_j(\tau_j^i(t)) - u_j(t)\|^2 \leq \gamma^2 \left(\sum_{\tau=t-B+1}^{t-1} \|s_j(\tau)\| \right)^2 \leq \gamma^2 B \sum_{\tau=t-B+1}^{t-1} \|s_j(\tau)\|^2.$$

Combining (22) and (23) yields

$$\begin{aligned}
 (24) \quad & F(u(t+1)) \leq F(u(t)) \\
 & \quad + \gamma^2 C_2 \sqrt{B} \left(\sum_{j=1}^m \sum_{\tau=t-B+1}^{t-1} \|s_j(\tau)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|s_i(t)\|^2 \right)^{\frac{1}{2}} \\
 & \quad - \gamma \frac{\sigma}{2} \sum_{i=1}^m \|s_i(t)\|^2 \\
 & \leq F(u(t)) + \gamma^3 \frac{C_2^2 B}{\sigma} \sum_{j=1}^m \sum_{\tau=t-B+1}^{t-1} \|s_j(\tau)\|^2 - \gamma \frac{\sigma}{4} \sum_{i=1}^m \|s_i(t)\|^2,
 \end{aligned}$$

where the second inequality uses the identity $ab \leq (a^2 + b^2)/2$ with a and b being the two square-root terms multiplied and divided, respectively, by $B^{1/4} \sqrt{2\gamma C_2/\sigma}$. Applying the above argument successively to $t, t + 1, \dots, t + B - 1$, we obtain

$$\begin{aligned}
 & F(u(t+B)) - F(u(t)) \\
 & \leq -\gamma \left(\frac{\sigma}{4} - \frac{\gamma^2 C_2^2 B^2}{\sigma} \right) \sum_{j=1}^m \sum_{\tau=t}^{t+B-1} \|s_j(\tau)\|^2 + \gamma^3 \frac{C_2^2 B^2}{\sigma} \sum_{j=1}^m \sum_{\tau=t-B+1}^{t-1} \|s_j(\tau)\|^2.
 \end{aligned}$$

This proves the lemma. □

The next key lemma estimates the optimality gap $F(u(t + B)) - F(\bar{u})$, where \bar{u} is the unique solution of (2).

Lemma 2 (Optimality gap estimate). *Let A_3 and A_4 be defined by*

$$(25) \quad A_4 = \frac{C_2 B^2}{2} + \frac{8C_1^2 C_2^2 B}{\sigma}, \quad A_3 = \frac{3C_2}{2} + \frac{6C_1^2 C_2^2}{\sigma} + A_4.$$

For $t = 0, 1, \dots$, we have

$$\begin{aligned} F(u(t + B)) - F(\bar{u}) &\leq (1 - \gamma)(F(u(t)) - F(\bar{u})) \\ &\quad + \gamma A_3 \sum_{j=1}^m \sum_{\tau=t}^{t+B-1} \|s_j(\tau)\|^2 + \gamma^3 A_4 \sum_{j=1}^m \sum_{\tau=t-B+1}^{t-1} \|s_j(\tau)\|^2. \end{aligned}$$

Proof. Fix any $t \in \{0, 1, \dots\}$. For each $i \in \{1, \dots, m\}$, let t^i denote the greatest element of T^i less than $t + B$. Then we have from (11) and (17) that

$$(26) \quad \langle F'(z_i(t^i) + s_i(t^i)), v_i - w_i(t^i) \rangle \geq 0, \quad \forall v_i \in K_i.$$

We also have from (10) and (16) that

$$\begin{aligned} u_i(t + B) &= u_i(t^i) + \gamma s_i(t^i), \\ u(t + B) &= \sum_{i=1}^m u_i(t^i + 1) = \sum_{i=1}^m u_i(t^i) + \gamma \sum_{i=1}^m s_i(t^i). \end{aligned}$$

For notational simplicity, define

$$w(t) = \sum_{i=1}^m w_i(t^i), \quad \hat{u}(t) = \sum_{i=1}^m u_i(t^i).$$

By assumption, there exists $\bar{u}_i \in K_i$, $i = 1, \dots, m$, such that (5) holds with $v_i = w_i(t^i)$, i.e.,

$$(27) \quad \bar{u} = \sum_{i=1}^m \bar{u}_i \quad \text{and} \quad \left(\sum_{i=1}^m \|w_i(t^i) - \bar{u}_i\|^2 \right)^{\frac{1}{2}} \leq C_1 \|w(t) - \bar{u}\|.$$

Then $(\bar{u}_1, \dots, \bar{u}_m)$ is a solution of the convex program (1) and, by F being Gâteaux-differentiable, it satisfies the optimality condition

$$(28) \quad \sum_{i=1}^m \langle F'(\bar{u}), v_i - \bar{u}_i \rangle \geq 0, \quad \forall v_i \in K_i, \quad i = 1, \dots, m.$$

Defining

$$\phi_j^i(t) = \sum_{k=1}^j w_k(t^k) + \sum_{k=j+1}^m u_k(\tau_k^i(t^i)), \quad j = 0, 1, \dots, m,$$

we have that $\phi_0^i(t) = z_i(t^i)$ and $\phi_m^i(t) = w(t)$ and

$$(29) \quad \phi_j^i(t) - \phi_{j-1}^i(t) = w_j(t^j) - u_j(\tau_j^i(t^i)) \in K_j^\ominus, \quad j = 1, \dots, m.$$

Setting $v_i = \bar{u}_i$ in (26) and $v_i = w_i(t^i)$ in (28), we obtain that

$$\begin{aligned}
& \left\langle F'(w(t)) - F'(\bar{u}), w(t) - \bar{u} \right\rangle \leq \left\langle F'(w(t)), w(t) - \bar{u} \right\rangle \\
& \leq \sum_{i=1}^m \left\langle F'(w(t)) - F'(z_i(t^i) + s_i(t^i)), w_i(t^i) - \bar{u}_i \right\rangle \\
& = \sum_{i=1}^m \left\langle F'(w(t)) - F'(z_i(t^i)), w_i(t^i) - \bar{u}_i \right\rangle \\
& \quad + \sum_{i=1}^m \left\langle F'(z_i(t^i)) - F'(z_i(t^i) + s_i(t^i)), w_i(t^i) - \bar{u}_i \right\rangle \\
& = \sum_{i=1}^m \sum_{j=1}^m \left\langle F'(\phi_j^i(t)) - F'(\phi_{j-1}^i(t)), w_i(t^i) - \bar{u}_i \right\rangle \\
& \quad + \sum_{i=1}^m \left\langle F'(z_i(t^i)) - F'(z_i(t^i) + s_i(t^i)), w_i(t^i) - \bar{u}_i \right\rangle \\
& \leq C_2 \left(\sum_{j=1}^m \max_{i=1, \dots, m} \|u_j(\tau_j^i(t^i)) - w_j(t^j)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|w_i(t^i) - \bar{u}_i\|^2 \right)^{\frac{1}{2}} \\
& \quad + C_2 \left(\sum_{i=1}^m \|s_i(t^i)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|w_i(t^i) - \bar{u}_i\|^2 \right)^{\frac{1}{2}} \\
& \leq C_1 C_2 \left(\sum_{j=1}^m \left(4\gamma^2 B \sum_{\tau=t-B+1}^{t+B-2} \|s_j(\tau)\|^2 + 2\|s_j(t^j)\|^2 \right) \right)^{\frac{1}{2}} \|w(t) - \bar{u}\| \\
(30) \quad & \quad + C_1 C_2 \left(\sum_{i=1}^m \|s_i(t^i)\|^2 \right)^{\frac{1}{2}} \|w(t) - \bar{u}\|,
\end{aligned}$$

where the third inequality uses (6) and (29); the fourth inequality uses (27) and the fact that

$$\begin{aligned}
\|u_j(\tau_j^i(t^i)) - w_j(t^j)\|^2 &= \|u_j(\tau_j^i(t^i)) - u_j(t^j) - s_j(t^j)\|^2 \\
&\leq 2\|u_j(\tau_j^i(t^i)) - u_j(t^j)\|^2 + 2\|s_j(t^j)\|^2 \\
&\leq 2\gamma^2 \left(\sum_{\tau=t-B+1}^{t+B-2} \|s_j(\tau)\| \right)^2 + 2\|s_j(t^j)\|^2 \\
&\leq 4\gamma^2 B \sum_{\tau=t-B+1}^{t+B-2} \|s_j(\tau)\|^2 + 2\|s_j(t^j)\|^2
\end{aligned}$$

(see (10), (11), (13), (14)). Also, the strong monotonicity (3) of F' on K implies

$$\left\langle F'(w(t)) - F'(\bar{u}), w(t) - \bar{u} \right\rangle \geq \sigma \|w(t) - \bar{u}\|^2,$$

which together with (30) yields

$$\begin{aligned}
 \|w(t) - \bar{u}\| &\leq \frac{C_1 C_2}{\sigma} \left(\sum_{j=1}^m \left(4\gamma^2 B \sum_{\tau=t-B+1}^{t+B-2} \|s_j(\tau)\|^2 + 2\|s_j(t^j)\|^2 \right) \right)^{\frac{1}{2}} \\
 (31) \quad &+ \frac{C_1 C_2}{\sigma} \left(\sum_{i=1}^m \|s_i(t^i)\|^2 \right)^{\frac{1}{2}}.
 \end{aligned}$$

Next, since $F'(w(t))$ is a subgradient of F at $w(t)$ [20, p. 23], we have

$$F(w(t)) - F(\bar{u}) \leq \langle F'(w(t)), w(t) - \bar{u} \rangle,$$

so putting $v_i = \bar{u}_i$ in (26) and adding it to the above inequality yields

$$\begin{aligned}
 (32) \quad &F(w(t)) - F(\bar{u}) \\
 &\leq \sum_{i=1}^m \langle F'(w(t)) - F'(z_i(t^i) + s_i(t^i)), w_i(t^i) - \bar{u}_i \rangle \\
 &\leq \frac{C_1^2 C_2^2}{\sigma} \left(\left(\sum_{j=1}^m \left(4\gamma^2 B \sum_{\tau=t-B+1}^{t+B-2} \|s_j(\tau)\|^2 + 2\|s_j(t^j)\|^2 \right) \right)^{\frac{1}{2}} \right. \\
 &\qquad \qquad \qquad \left. + \left(\sum_{i=1}^m \|s_i(t^i)\|^2 \right)^{\frac{1}{2}} \right)^2 \\
 &\leq \frac{2C_1^2 C_2^2}{\sigma} \left(4\gamma^2 B \sum_{j=1}^m \sum_{\tau=t-B+1}^{t+B-2} \|s_j(\tau)\|^2 + 3 \sum_{i=1}^m \|s_i(t^i)\|^2 \right),
 \end{aligned}$$

where the second inequality uses (30) and (31) and the last inequality follows from the identity $(a + b)^2 \leq 2(a^2 + b^2)$.

Next we estimate $F(\hat{u}(t)) - F(u(t))$. Let $\bar{t} = \max_{i=1, \dots, m} t^i$ and, for each $i \in \{1, \dots, m\}$ and $\tau \in \{t, \dots, \bar{t}\}$, define

$$(33) \quad \tilde{u}_i(\tau) = u_i(\min\{\tau, t^i\}), \quad \tilde{u}(\tau) = \sum_{i=1}^m \tilde{u}_i(\tau).$$

Then, for each $i \in \{1, \dots, m\}$ and $\tau \in \{t, \dots, \bar{t} - 1\}$, either $\tilde{u}_i(\tau + 1) = \tilde{u}_i(\tau)$ so that

$$\langle F'(z_i(\tau) + s_i(\tau)), \tilde{u}_i(\tau) - \tilde{u}_i(\tau + 1) \rangle = 0$$

or $\tilde{u}_i(\tau + 1) \neq \tilde{u}_i(\tau)$ so that $\tau \in T^i$ and $\tau < t^i$, implying by (11) and (17) that

$$\langle F'(z_i(\tau) + s_i(\tau)), u_i(\tau) - w_i(\tau) \rangle \geq 0$$

and hence, by (33), that

$$\begin{aligned}
 \langle F'(z_i(\tau) + s_i(\tau)), \tilde{u}_i(\tau) - \tilde{u}_i(\tau + 1) \rangle &= \langle F'(z_i(\tau) + s_i(\tau)), u_i(\tau) - u_i(\tau + 1) \rangle \\
 &= \gamma \langle F'(z_i(\tau) + s_i(\tau)), u_i(\tau) - w_i(\tau) \rangle \geq 0.
 \end{aligned}$$

Using this and defining

$$\phi_j^i(\tau) = \sum_{k=1}^j \tilde{u}_k(\tau + 1) + \sum_{k=j+1}^m u_k(\tau_k^i(\tau)), \quad j = 0, 1, \dots, m,$$

we obtain that

(34)

$$\begin{aligned}
& F(\tilde{u}(\tau+1)) - F(\tilde{u}(\tau)) \\
& \leq -\langle F'(\tilde{u}(\tau+1)), \tilde{u}(\tau) - \tilde{u}(\tau+1) \rangle \\
& \leq \sum_{i=1}^m \langle F'(z_i(\tau) + s_i(\tau)) - F'(\tilde{u}(\tau+1)), \tilde{u}_i(\tau) - \tilde{u}_i(\tau+1) \rangle \\
& = \sum_{i=1}^m \langle F'(z_i(\tau)) - F'(\tilde{u}(\tau+1)), \tilde{u}_i(\tau) - \tilde{u}_i(\tau+1) \rangle \\
& \quad + \sum_{i=1}^m \langle F'(z_i(\tau) + s_i(\tau)) - F'(z_i(\tau)), \tilde{u}_i(\tau) - \tilde{u}_i(\tau+1) \rangle \\
& = \sum_{i=1}^m \sum_{j=1}^m \langle F'(\phi_{j-1}^i(\tau)) - F'(\phi_j^i(\tau)), \tilde{u}_i(\tau) - \tilde{u}_i(\tau+1) \rangle \\
& \quad + \sum_{i=1}^m \langle F'(z_i(\tau) + s_i(\tau)) - F'(z_i(\tau)), \tilde{u}_i(\tau) - \tilde{u}_i(\tau+1) \rangle \\
& \leq C_2 \left(\sum_{j=1}^m \max_{i=1, \dots, m} \|\phi_{j-1}^i(\tau) - \phi_j^i(\tau)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|\tilde{u}_i(\tau) - \tilde{u}_i(\tau+1)\|^2 \right)^{\frac{1}{2}} \\
& \quad + C_2 \left(\max_{i=1, \dots, m} \|s_i(\tau)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|\tilde{u}_i(\tau) - \tilde{u}_i(\tau+1)\|^2 \right)^{\frac{1}{2}} \\
& \leq \gamma C_2 \left(\sum_{j=1}^m \left(\max_{i=1, \dots, m} \|\tilde{u}_j(\tau+1) - u_j(\tau_j^i(\tau))\|^2 \right) \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|s_i(\tau)\|^2 \right)^{\frac{1}{2}} \\
& \quad + \gamma C_2 \sum_{i=1}^m \|s_i(\tau)\|^2 \\
& \leq \gamma C_2 \left(\gamma^2 B \sum_{j=1}^m \sum_{\nu=\tau-B+1}^{\tau+1} \|s_j(\nu)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|s_i(\tau)\|^2 \right)^{\frac{1}{2}} + \gamma C_2 \sum_{i=1}^m \|s_i(\tau)\|^2 \\
& \leq \gamma^3 \frac{C_2 B}{2} \sum_{j=1}^m \sum_{\nu=\tau-B+1}^{\tau+1} \|s_j(\nu)\|^2 + \gamma \frac{3C_2}{2} \sum_{i=1}^m \|s_i(\tau)\|^2,
\end{aligned}$$

where the first inequality uses the subgradient property of $F'(\tilde{u}(\tau+1))$ [20, p. 23]; the third inequality uses (6); the fourth and fifth inequalities use (33) and (10) and an inequality analogous to (23); the last inequality uses the identity $ab \leq (a^2 + b^2)/2$ with a and b being the two square-root terms. Summing the above inequality over $\tau = t, t+1, \dots, \bar{t}-1$ and observing that $\tilde{u}(\bar{t}) = \hat{u}(t)$ and $\tilde{u}(t) = u(t)$, we then have

$$\begin{aligned}
(35) \quad F(\hat{u}(t)) - F(u(t)) & \leq \gamma^3 \frac{C_2 B}{2} \sum_{j=1}^m \sum_{\tau=t}^{\bar{t}-1} \sum_{\nu=\tau-B+1}^{\tau+1} \|s_j(\nu)\|^2 + \gamma \frac{3C_2}{2} \sum_{i=1}^m \sum_{\tau=t}^{\bar{t}-1} \|s_i(\tau)\|^2 \\
& \leq \gamma^3 \frac{C_2 B^2}{2} \sum_{j=1}^m \sum_{\tau=t-B+1}^{t+B-1} \|s_j(\tau)\|^2 + \gamma \frac{3C_2}{2} \sum_{i=1}^m \sum_{\tau=t}^{t+B-1} \|s_i(\tau)\|^2.
\end{aligned}$$

Finally, using the convexity of F and $\gamma \in [0, 1]$, we see from (11) and (32) and (35) that

$$\begin{aligned}
 & F(u(t+B)) - F(\bar{u}) \\
 &= F\left(\sum_{i=1}^m u_i(t+B)\right) - F(\bar{u}) \\
 &= F\left(\sum_{i=1}^m (u_i(t^i) + \gamma(w_i(t^i) - u_i(t^i)))\right) - F(\bar{u}) \\
 &= F((1-\gamma)\hat{u}(t) + \gamma w(t)) - F(\bar{u}) \\
 &\leq (1-\gamma)F(\hat{u}(t)) + \gamma F(w(t)) - F(\bar{u}) \\
 &= (1-\gamma)(F(\hat{u}(t)) - F(\bar{u})) + \gamma(F(w(t)) - F(\bar{u})) \\
 &\leq (1-\gamma)(F(u(t)) - F(\bar{u})) \\
 &\quad + \gamma^3 \frac{C_2 B^2}{2} \sum_{j=1}^m \sum_{\tau=t-B+1}^{t+B-1} \|s_j(\tau)\|^2 + \gamma \frac{3C_2}{2} \sum_{i=1}^m \sum_{\tau=t}^{t+B-1} \|s_i(\tau)\|^2 \\
 &\quad + \gamma^3 \frac{8C_1^2 C_2^2 B}{\sigma} \sum_{j=1}^m \sum_{\tau=t-B+1}^{t+B-2} \|s_j(\tau)\|^2 + \gamma \frac{6C_1^2 C_2^2}{\sigma} \sum_{i=1}^m \|s_i(t^i)\|^2.
 \end{aligned}$$

Using $\gamma \leq 1$ then proves the lemma. □

We will now use Lemmas 1 and 2 to prove our convergence rate result. To simplify the notations, define

$$a_k = F(u(kB)) - F(\bar{u}), \quad b_k = \sum_{j=1}^m \sum_{\tau=kB-B}^{kB-1} \|s_j(\tau)\|^2, \quad k = 1, 2, \dots$$

By Lemmas 1 and 2, we have

$$(36) \quad a_k \leq a_{k-1} - \gamma A_1 b_k + \gamma^3 A_2 b_{k-1},$$

$$(37) \quad a_k \leq (1-\gamma)a_{k-1} + \gamma A_3 b_k + \gamma^3 A_4 b_{k-1},$$

where A_1, A_2, A_3, A_4 are given by (18) and (25). By (15), we have $A_1 > 0$. Choose γ sufficiently small so that

$$(38) \quad \varrho = \max \left\{ \left(1 + \frac{A_1}{A_3}\right)^{-1} \left(1 + (1-\gamma)\frac{A_1}{A_3} + \gamma^{3/2} \left(A_2 + \frac{A_1 A_4}{A_3}\right)\right), A_1^{-1}(\gamma^{1/2} + \gamma^2 A_2) \right\} < 1.$$

Also, define $a = \max\{a_1, \gamma^{3/2} b_1\} / \varrho$. We claim that

$$(39) \quad \max\{a_n, \gamma^{3/2} b_n\} \leq a \varrho^n$$

for $n = 1, 2, \dots$. We prove this by induction on n . Clearly (39) holds for $n = 1$ by our definition of a . Suppose (39) holds for $n = k - 1$, where $k > 1$. Multiplying (37) by A_1/A_3 and adding it to (36) gives

$$\left(1 + \frac{A_1}{A_3}\right) a_k \leq \left(1 + (1-\gamma)\frac{A_1}{A_3}\right) a_{k-1} + \gamma^{3/2} \left(A_2 + \frac{A_1 A_4}{A_3}\right) (\gamma^{3/2} b_{k-1}),$$

which together with the inductive hypothesis $\max\{a_{k-1}, \gamma^{3/2}b_{k-1}\} \leq a\varrho^{k-1}$ and (38) yields

$$a_k \leq \left(1 + \frac{A_1}{A_3}\right)^{-1} \left(1 + (1 - \gamma)\frac{A_1}{A_3} + \gamma^{3/2}\left(A_2 + \frac{A_1A_4}{A_3}\right)\right) a\varrho^{k-1} \leq a\varrho^k.$$

Similarly, (36) and $a_k \geq 0$ give

$$\gamma^{3/2}A_1b_k \leq \gamma^{1/2}a_{k-1} + \gamma^2A_2(\gamma^{3/2}b_{k-1}),$$

which together with $\max\{a_{k-1}, \gamma^{3/2}b_{k-1}\} \leq a\varrho^{k-1}$ and (38) yields

$$\gamma^{3/2}b_k \leq A_1^{-1}(\gamma^{1/2} + \gamma^2A_2)a\varrho^{k-1} \leq a\varrho^k.$$

This shows that (39) holds for $n = k$, completing our induction proof.

Thus, we have shown linear rate of convergence (in the root sense) for both a_n and b_n , with a factor of ϱ . The latter implies $u_i(t)$, $t = 0, 1, \dots$, is a Cauchy sequence for each i and hence it converges strongly. This is summarized in the theorem below.

Theorem 1. *Consider the minimization problem (2) and the space decomposition (4) of Section 2 (see (3), (5)–(9)). Let $(u_1(t), \dots, u_m(t))$, $t = 0, 1, \dots$, be generated by the asynchronous space decomposition method of Section 3 (see (10)–(12) and (13), (14)) with stepsize γ satisfying (15), (38). Then, there exist $a > 0$ and $\varrho \in (0, 1)$, depending on σ, C_1, C_2 and B, γ only, such that*

$$F(u(nB)) - F(\bar{u}) \leq a\varrho^n, \quad n = 1, 2, \dots,$$

where $u(t)$ is given by (16) and \bar{u} denotes the unique solution of (2). Moreover, $u(t)$ converges strongly to \bar{u} and, for each $i \in \{1, \dots, m\}$, $u_i(t)$ converges strongly as $t \rightarrow \infty$.

5. CONVERGENCE RATE

OF THE SYNCHRONOUS SEQUENTIAL AND PARALLEL ALGORITHMS

It is readily seen that the following Jacobi version of the method is a special case of the asynchronous space decomposition method (10)–(12) with $T^i = \{0, 1, \dots\}$ and $\tau_j^i(t) = t$ for all i, j, t (so $B = 1$ and $c_t = c$). Thus, Theorem 1 can be applied to establish its linear convergence and obtain an estimate of the factor ϱ under the assumptions of Section 2. Moreover, by observing that in this case the left-hand side of (23) is zero so that Lemma 1 holds with $A_2 = 0$, the stepsize restriction (15) can be relaxed to $\gamma \leq 1/c_t$.

Algorithm 1.

Step 1. *Choose initial values $u_i(0) \in K_i$, $i = 1, \dots, m$, and stepsize $\gamma = 1/c$, where c is defined as in Section 2.*

Step 2. *For each $t = 0, 1, \dots$, find $w_i(t) \in K_i$ in parallel for $i = 1, \dots, m$ that satisfies*

$$F\left(\sum_{j \neq i} u_j(t) + w_i(t)\right) \leq F\left(\sum_{j \neq i} u_j(t) + v_i\right), \quad \forall v_i \in K_i.$$

Step 3. *Set*

$$u_i(t + 1) = u_i(t) + \gamma(w_i(t) - u_i(t)),$$

and go to the next iteration.

The following Gauss-Seidel version of the method is also a special case of the asynchronous space decomposition method (10)–(12) with $\gamma = 1$, $T^i = \{i - 1 + km\}_{k=0,1,\dots}$ and $\tau_j^i(t) = t$ for all i, j, t (so $B = m$ and $c_t = 1$). Here Theorem 1 cannot be directly applied due to $\gamma = 1$ possibly violating (15). However, by observing that in this case the left-hand side of (23) is again zero so that Lemma 1 holds with $A_2 = 0$, the proof of the theorem can be easily modified to establish linear convergence of this method under the assumptions of Section 2, with factor ρ depending on m, σ, C_1, C_2 only. Moreover, by grouping sets of the same color into one set, we can ensure that $m = c$, where c is defined as in Section 2.

Algorithm 2.

Step 1. Choose initial values $u_i(0) \in K_i, i = 1, \dots, m$.

Step 2. For each $t = 0, 1, \dots$, find $u_i(t + 1) \in K_i$ sequentially for $i = 1, \dots, m$ that satisfies

$$F \left(\sum_{j < i} u_j(t + 1) + u_i(t + 1) + \sum_{j > i} u_j(t) \right) \leq F \left(\sum_{j < i} u_j(t + 1) + v_i + \sum_{j > i} u_j(t) \right), \quad \forall v_i \in K_i .$$

Step 3. Go to the next iteration.

The above two methods for solving (2) were studied in [47] (also see [48, 49, 50]), where convergence of the methods was proved under weaker assumptions. However, no rate of convergence result was given. In [52], a linear rate of convergence for the above two methods was proved for the unconstrained case of $K = V$. In the finite-dimensional case of $K = V = \mathbb{R}^n$, the literature concerning the linear rate of convergence is very rich. However, the study for linear convergence rate for general convex sets K_i is very sparse. The linear rate of convergence for the Gauss-Seidel method for general convex sets K_i can also be inferred from the results in [34, 35] and references therein, but our estimate of the convergence factor is new.

In [52], the minimization subproblem at each iteration is solved inexactly. We can do likewise in the constrained case. In particular, the proof of Theorem 1 (see (21) and (26)) suggests that the exact minimization condition (17) can be relaxed to the inexact minimization condition

$$\langle F'(z_i(t) + w_i(t) - u_i(t)), v_i - w_i(t) \rangle \geq -\frac{\sigma_0}{2} \|w_i(t) - u_i(t)\|^2, \quad \forall v_i \in K_i,$$

with $0 < \sigma_0 < \sigma$. However, σ would need to be known explicitly and both γ and ρ would depend on σ_0 .

6. APPLICATIONS TO CONVEX PROGRAMMING

In this section we consider the Euclidean space $V = V' = \mathbb{R}^n$, which is the space of n -dimensional real column vectors with duality pairing $\langle f, x \rangle = f^T x$ and norm $\|x\| = \sqrt{x^T x}$, where x^T denotes transpose of x . We will discuss choices of the space decomposition (4) and the corresponding estimates for C_1, C_2, c in (5), (6), (7). In the case of nonlinear network flow, we will also relate our asynchronous method to those studied in [5, §7.2.3], [56].

6.1. Primal applications. Consider the problem (2), where $F : \mathfrak{R}^n \mapsto \mathfrak{R}$ is a differentiable convex function and K is a nonempty polyhedral set in \mathfrak{R}^n . Then F is continuous [42, p. 82] and continuously differentiable [42, p. 246]. We assume that the gradient $F' = (\frac{\partial F}{\partial x_j})_{j=1}^n$ is strongly monotone and Lipschitz continuous on K , and we choose a space decomposition (4) such that each K_i is a polyhedral set.

Since each K_i is a polyhedral set, a result of Hoffman on the Lipschitzian behavior of solutions of a linear system with respect to the right-hand side (see [13]) implies that, for any $v_i \in K_i$, $i = 1, \dots, m$, there exists $\bar{u}_i \in K_i$ satisfying (5), where the constant C_1 depends on m and certain condition numbers for K_i , $i = 1, \dots, m$. In cases where each K_i has a simple structure, such as the Cartesian product of closed intervals, C_1 may be estimated explicitly. For a coloring of the sets, if K_i and K_j are not orthogonal, i.e., $(v_i)^T v_j \neq 0$ for some $v_i \in K_i, v_j \in K_j$, then we paint them different colors. Let \hat{c} be the maximum number of sets K_j that are not orthogonal to an arbitrary set K_i . Then an analysis similar to that used in subsection 7.1.3 shows that (6) holds with $C_2 = L\hat{c}$, where L is the Lipschitz constant for F' .

6.2. Dual applications. Consider the linearly constrained convex program

$$(40) \quad \text{minimize } G(x) \quad \text{subject to } Ax \geq b,$$

where $G : \mathfrak{R}^n \mapsto \mathfrak{R}$ is a strictly convex differentiable function, $b \in \mathfrak{R}^m$, and $A \in \mathfrak{R}^{m \times n}$ has nonzero rows. We assume there exists $\tilde{x} \in \mathfrak{R}^n$ satisfying $A\tilde{x} = b$. By attaching Lagrange multipliers $\lambda \in \mathfrak{R}^m$ to the inequalities $Ax \geq b$ in (40), we obtain the Lagrangian dual problem

$$(41) \quad \min_{\lambda \in \mathfrak{R}_+^m} G^*(A^T \lambda) - b^T \lambda,$$

where \mathfrak{R}_+^m denotes the nonnegative orthant in \mathfrak{R}^m , and G^* is the convex conjugate (also called Legendre-Fenchel transform) of G defined by

$$G^*(u) = \sup_{x \in \mathfrak{R}^n} \{u^T x - G(x)\}$$

(see [24, 42]). The convex programs (40) and (41) are dual in the sense that one has a solution if and only if the other does, and these solutions satisfies $G'(x) = A^T \lambda$ [42, Cor. 28.3.1 and 28.4.1]. Using $b = A\tilde{x}$, we can rewrite the dual problem (41) in the form of (2) with

$$F(u) = G^*(u) - \tilde{x}^T u, \quad K = \{u \in \mathfrak{R}^n : u = A^T \lambda \text{ for some } \lambda \in \mathfrak{R}_+^m\}.$$

We assume that $(G^*)'$ is strongly monotone and Lipschitz continuous on \mathfrak{R}^n , so that F satisfies (3) for some $\sigma > 0$. If G is twice differentiable, this assumption essentially amounts to G'' having bounded eigenvalues and the Hessian $(G'')^{-1}$ having bounded entries on \mathfrak{R}^n . Let \bar{u} denote the unique solution of (2) and let A_i denote the i th row of A .

We can decompose K in the form (4) with

$$K_i = \{u_i \in \mathfrak{R}^n : u_i = A_i^T \lambda_i \text{ for some } \lambda_i \in \mathfrak{R}_+\}.$$

First we show that, for any $v_i \in K_i$, $i = 1, \dots, m$, there exists $\bar{u}_i \in K_i$ satisfying (5), where

$$(42) \quad C_1 = \max_{\emptyset \neq I \subset \{1, \dots, m\}} \|D_I^{-1} B_I (B_I^T D_I^{-2} B_I)^{-1}\|,$$

with D_I being the diagonal matrix with diagonal entries $\|A_i^T\|$, $i \in I$, and B_I being a submatrix of $A_I = [A_i]_{i \in I}$ comprising linearly independent columns of

A_I spanning the column space of A_I . To see this, notice that $\bar{u} = A^T \bar{\lambda}$ for some $\bar{\lambda} \in \mathfrak{R}_+^m$ and $v_i = A_i^T \mu_i$ for some $\mu_i \in \mathfrak{R}_+$. Moreover, $u_i \in K_i, i = 1, \dots, m$, satisfies $\sum_{i=1}^m u_i = \bar{u}$ if and only if $u_i = A_i^T \lambda_i$ and $A^T \lambda = A^T \bar{\lambda}$ for some $\lambda = (\lambda_i)_{i=1}^m \in \mathfrak{R}_+^m$. Thus, minimizing $\sum_{i=1}^m \|u_i - v_i\|^2$ subject to $u_i \in K_i$ and $\sum_{i=1}^m u_i = \bar{u}$ is equivalent to minimizing

$$\sum_{i=1}^m \|A_i^T \lambda_i - A_i^T \mu_i\|^2 = \sum_{i=1}^m \|A_i^T\|^2 |\lambda_i - \mu_i|^2 = \|D(\lambda - \mu)\|^2$$

subject to $A^T \lambda = A^T \bar{\lambda}, \lambda \geq 0$, where $\mu = (\mu_i)_{i=1}^m$. By making the variable substitution $\xi = \lambda - \mu$ and letting $r = A^T(\bar{\lambda} - \mu)$, this in turn is equivalent to

$$(43) \quad \text{minimize } \|D\xi\| \quad \text{subject to } A^T \xi = r, \xi \geq -\mu.$$

Since $\mu \geq 0$, the optimal value of (43) equals zero if $r = 0$. Suppose instead that $r \neq 0$. Since $\mu \geq 0$, the optimal value of (43) is below that of

$$\text{minimize } \|D\xi\| \quad \text{subject to } A^T \xi = r, \xi \geq 0.$$

The latter has a unique solution, which we denote by $\bar{\xi} = (\bar{\xi}_i)_{i=1}^m$. Since $r \neq 0$, then $\bar{\xi} \neq 0$. Let $I = \{i \in \{1, \dots, m\} : \bar{\xi}_i \neq 0\}$. Then, $\bar{\xi}_I = (\bar{\xi}_i)_{i \in I}$ solves the reduced problem

$$\text{minimize } \|D_I \bar{\xi}_I\| \quad \text{subject to } B_I^T \bar{\xi}_I = r_I,$$

where r_I is the subvector of r corresponding to the columns of B_I . This yields $\bar{\xi}_I = D_I^{-2} B_I (B_I^T D_I^{-2} B_I)^{-1} r_I$, and hence

$$\begin{aligned} \text{optimal value of (43)} &\leq \|D_I \bar{\xi}_I\| \\ &= \|D_I^{-1} B_I (B_I^T D_I^{-2} B_I)^{-1} r_I\| \\ &\leq C_1 \|r_I\| \\ &\leq C_1 \|r\| \\ &= C_1 \|A^T(\bar{\lambda} - \mu)\| \\ &= C_1 \left\| \bar{u} - \sum_{i=1}^m v_i \right\|. \end{aligned}$$

Since K_i and K_j lie in orthogonal subspaces if $A_i A_j^T = 0$, we can color K_1, \dots, K_m as discussed in subsection 6.1 and show that (6) holds with $C_2 = L\hat{c}$, where L is the Lipschitz constant for $(G^*)'$ and \hat{c} is the maximum number of rows A_j that are not orthogonal to an arbitrary row A_i .

If we replace the inequality $Ax \geq b$ in (40) by an equation $Ax = b$, the constraint $\lambda \in \mathfrak{R}_+^m$ in (41) would be replaced accordingly by $\lambda \in \mathfrak{R}^m$ and it suffices to fix $I = \{1, \dots, m\}$ in the estimate (42). This estimate further simplifies if A has full row rank, in which case B_I is square and invertible. If A does not have full row rank, we could remove the redundant rows, but our experience with network flow problems suggests that this removal can slow the convergence of Gauss-Seidel methods on the problem [56].

In the case of a nonlinear network flow problem [43], where A is the node-arc incidence matrix for a connected diagraph with m nodes and n arcs (i.e., every column of A has one 1 and one -1 in two of its rows, and a 0 in the remaining rows), we can estimate C_1 explicitly in terms of m and n as follows: For any $v_i \in K_i, i = 1, \dots, m$, we have $\bar{u} = A^T \bar{\lambda} = (\bar{\lambda}_k - \bar{\lambda}_l)_{j=1; j \sim (k,l)}^n$ for some $\bar{\lambda} = (\bar{\lambda}_i)_{i=1}^m \in \mathfrak{R}_+^m$

and $v_i = A_i^T \mu_i$ for some $\mu_i \in \mathfrak{R}_+$, where $k \sim (i, j)$ means that column k has a 1 in row i and a -1 in row j or, equivalently, arc k is directed from node i to node j . Choose any spanning tree for the diagraph and choose a node \bar{i} such that $\bar{\lambda}_{\bar{i}} = \min_i \bar{\lambda}_i$. Let $\lambda_i = \mu_{\bar{i}} + (\bar{\lambda}_i - \bar{\lambda}_{\bar{i}})$ and $u_i = A_i^T \lambda_i$ for all nodes i in the network. Since $\mu_{\bar{i}} \geq 0$ and $\bar{\lambda}_i \geq \bar{\lambda}_{\bar{i}}$, we have $\lambda_i \geq 0$ for all nodes i . Since each node i can be reached from \bar{i} via a simple path P_i in the spanning tree, we also have

$$\begin{aligned} |\lambda_i - \mu_i| &= \left| - \sum_{(k,l) \in P_i^+} (\lambda_k - \mu_k - \lambda_l + \mu_l) + \sum_{(k,l) \in P_i^-} (\lambda_k - \mu_k - \lambda_l + \mu_l) \right| \\ &= \left| - \sum_{(k,l) \in P_i^+} (\bar{\lambda}_k - \bar{\lambda}_l - \mu_k + \mu_l) + \sum_{(k,l) \in P_i^-} (\bar{\lambda}_k - \bar{\lambda}_l - \mu_k + \mu_l) \right| \\ &\leq \sum_{(k,l) \in P_i} |\bar{\lambda}_k - \bar{\lambda}_l - \mu_k + \mu_l| \\ &\leq \sqrt{h_i} \left(\sum_{(k,l) \in P_i} |\bar{\lambda}_k - \bar{\lambda}_l - \mu_k + \mu_l|^2 \right)^{\frac{1}{2}} \\ &\leq \sqrt{h_i} \left(\sum_{\substack{j=1 \\ j \sim (k,l)}}^n |\bar{\lambda}_k - \bar{\lambda}_l - \mu_k + \mu_l|^2 \right)^{\frac{1}{2}} \\ &= \sqrt{h_i} \left\| \bar{u} - \sum_{p=1}^m v_p \right\|, \end{aligned}$$

where P_i^+ and P_i^- denote the set of forward arcs and backward arcs in P_i and h_i denotes the number of arcs in P_i . Thus,

$$\begin{aligned} \sum_{i=1}^m \|u_i - v_i\|^2 &= \sum_{i=1}^m \|A_i^T(\lambda_i - \mu_i)\|^2 \\ &= \sum_{i=1}^m \|A_i^T\|^2 |\lambda_i - \mu_i|^2 \leq \sum_{i=1}^m d_i h_i \left\| \bar{u} - \sum_{p=1}^m v_p \right\|^2, \end{aligned}$$

where d_i is the number of arcs incident to node i . This shows that (5) holds with $C_1 = \sqrt{\sum_{i=1}^m d_i h_i}$. Notice that $\sum_{i=1}^m d_i = 2n$ and h_i is at most the diameter of the spanning tree. Since the choice of the spanning tree is arbitrary, we can choose it to minimize C_1 . Also, $A_i A_j^T = 0$ if and only if nodes i and j are not joined by an arc, so $\hat{c} = \max\{d_1, \dots, d_m\}$ and the coloring of K_1, \dots, K_m is equivalent to graph coloring on the diagraph.

In the above case of a nonlinear network flow problem, if G is also separable in the sense that $G(x) = \sum_{j=1}^n G_j(x_j)$ for all $x = (x_j)_{j=1}^n$ and $G_j : \mathfrak{R} \mapsto \mathfrak{R}$, then $\pi_i(u_1, \dots, u_m)$ given by (9) depends on only those u_k for which node k is a neighbor of node i and the asynchronous method (10)–(12) reduces to the asynchronous network relaxation method studied in [5, §7.2.3] and [56]. It is known that iterates generated by this method converge for any stepsize $\gamma \in (0, 1)$, assuming G^* is convex differentiable and (41) has a solution (G need not be defined everywhere on \mathfrak{R}^n and $(G^*)'$ need not be strongly monotone or Lipschitz continuous). However, no rate of convergence result was known. By applying Theorem 1, we obtain that this method has a linear rate of convergence, assuming $(G^*)'$ is strongly monotone and Lipschitz continuous and the stepsize is sufficiently small.

7. APPLICATIONS TO PARTIAL DIFFERENTIAL EQUATIONS
 WITHOUT CONSTRAINTS

In this section we consider the Sobolev space $V = H_0^1(\Omega) = \{v \in H^1(\Omega) : v = 0 \text{ on } \partial\Omega\}$ with duality pairing $\langle u, v \rangle = \int_{\Omega} (\sum_{i=1}^d \partial_i u \partial_i v + uv) dx$ and norm $\|v\| = \|v\|_{H^1(\Omega)} = \langle v, v \rangle^{\frac{1}{2}}$, where Ω is an open, bounded, and connected subset of \mathfrak{R}^d with Lipschitz continuous boundary $\partial\Omega$, $H^1(\Omega) = \{v \in L^2(\Omega) : \partial_i v \in L^2(\Omega), i = 1, \dots, d\}$, and $\partial_i v$ is the locally Lebesgue integrable real function defined on Ω satisfying $\int_{\Omega} \partial_i v \phi dx = - \int_{\Omega} v \frac{\partial \phi}{\partial x_i} dx$ for all $\phi \in C_0^\infty(\Omega) = \{\phi \in C^\infty(\Omega) : \phi \text{ has compact support}\}$ [17, pp. 10-13]. We will consider two nonlinear elliptic partial differential equations formulated as the minimization problem (2) and, for each, we will consider the space decomposition (4) corresponding to, respectively, DD and MG methods, and we will develop corresponding estimates for C_1 in (5), for C_2 in (6) and for c in (7)–(9). Throughout, we denote $|x| = (\sum_{i=1}^d x_i^2)^{\frac{1}{2}}$ for any $x = (x_i)_{i=1}^d \in \mathfrak{R}^d$.

The first partial differential equation corresponds to the minimization problem (2) with

$$(44) \quad K = H_0^1(\Omega), \quad \langle F'(u), v \rangle = \int_{\Omega} \left(\sum_{i=1}^d a_i(x, u, \nabla u) \partial_i v + a_0(x, u, \nabla u) v - f v \right) dx,$$

where $f \in L^2(\Omega)$ and $\nabla u = (\partial_i u)_{i=1}^d$ is the gradient of u [18, p. 302]. It is assumed that each nonlinear coefficient $a_i(x, p)$ is a real-valued function of $x = (x_j)_{j=1}^d$ and $p = (p_k)_{k=0}^d$ and is sufficiently smooth in the sense that

$$(45) \quad a_i \in C^1(\Omega \times \mathfrak{R}^{d+1}),$$

$$(46) \quad \max_{\substack{j=1,2,\dots,d \\ k=0,1,\dots,d}} \left\{ |a_i(x, p)|, \left| \frac{\partial a_i}{\partial x_j}(x, p) \right|, \left| \frac{\partial a_i}{\partial p_k}(x, p) \right| \right\} \leq L$$

for all $(x, p) \in \Omega \times \mathfrak{R}^{d+1}$ and $i = 0, 1, \dots, d$, with L a constant. In addition, the matrix $\left[\frac{\partial a_i}{\partial p_k}(x, p) \right]_{i,k=0}^d$ is assumed to be uniformly positive definite, i.e.,

$$(47) \quad \sum_{i=0}^d \sum_{k=0}^d \frac{\partial a_i}{\partial p_k}(x, p) \xi_i \xi_k \geq \sigma \sum_{i=0}^d \xi_i^2, \quad \forall \xi_i \in \mathfrak{R}, \quad i = 0, 1, \dots, d,$$

for all $(x, p) \in \Omega \times \mathfrak{R}^{d+1}$, with $\sigma > 0$ a constant. Under these assumptions, the problem (2), which has the equation formulation

$$(48) \quad \langle F'(u), v \rangle = 0, \quad \forall v \in H_0^1(\Omega),$$

is well posed and has a unique solution $u \in H_0^1(\Omega)$ (see [18, p. 302] and [32]). Moreover, straightforward calculation shows that

$$(49) \quad \begin{aligned} \langle F'(u) - F'(v), u - v \rangle &\geq \sigma \|u - v\|^2, \\ \langle F'(u) - F'(v), w \rangle &\leq L(d+1) \|u - v\| \|w\|, \end{aligned}$$

for all $u, v, w \in H^1(\Omega)$, so F' is strongly monotone and Lipschitz continuous.

The second partial differential equation corresponds to the minimization problem (2) with

$$(50) \quad K = H_0^1(\Omega), \quad F(v) = \int_{\Omega} \left(\frac{1}{2} |\nabla v|^2 + \frac{1}{4} v^4 - f v \right) dx,$$

where $f \in L^2(\Omega)$ and $d \in \{2, 3\}$. The corresponding equation is the simplified Ginzburg-Landau equation for superconductivity:

$$(51) \quad \begin{aligned} -\Delta u + u^3 &= f && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned}$$

where u is the wave function which is valid in the absence of internal magnetic field [54], and $\Delta u = \sum_{i=1}^d \partial_i(\partial_i u)$ denotes the Laplacian of u . Notice that F' has the form (44), with $a_0(x, p) = p_0^3$ and $a_i(x, p) = p_i$, $i = 1, \dots, d$, which does not satisfy (47). Nevertheless, straightforward calculation shows

$$\begin{aligned} \langle F'(u) - F'(v), u - v \rangle &= \int_{\Omega} |\nabla u - \nabla v|^2 + (u^3 - v^3)(u - v) dx \\ &\geq \int_{\Omega} |\nabla u - \nabla v|^2 dx = |u - v|_{1,\Omega}^2, \end{aligned}$$

for all $u, v \in H^1(\Omega)$. Since the semi-norm $|\cdot|_{1,\Omega}$ is equivalent to the norm $\|\cdot\|$ on $H_0^1(\Omega)$ [17, p. 12], this shows F' is strongly monotone on $H_0^1(\Omega)$.

In subsections 7.1 and 7.2 below, we will study asynchronous DD and MG methods for solving the above two equations (48) and (51). We will analyze the convergence rate of the methods by estimating the constants C_1 , C_2 and c for the corresponding space decomposition of the finite element approximation subspace and then applying Theorem 1. In particular, we will show that the above two equations can be solved in parallel with a convergence factor that is independent of the finite element mesh size h , i.e., the number of iterations to reach a desired solution accuracy is independent of h .

7.1. Domain decomposition methods.

7.1.1. *Decomposition of the domain* Ω . In DD methods, the domain Ω is decomposed into the disjoint union of subdomains Ω_i , $i = 1, \dots, m$, and their boundary, i.e., $\Omega \cup \partial\Omega = \bigcup_{i=1}^m (\Omega_i \cup \partial\Omega_i)$ and $\Omega_i \cap \Omega_j = \emptyset$ for $i \neq j$. This is illustrated in Figure 1 where a rectangular-shaped domain in \mathbb{R}^2 is decomposed into the disjoint union of $m = 25$ rectangular-shaped subdomains and their boundary. The subdomains, which are assumed to form a regular quasi-uniform division (see p. 124 and Eq. (3.2.28) of [17] for definitions) with a specified maximum diameter of H , are the finite elements of the coarse mesh. To form the fine mesh for the finite element approximations, we further divide each Ω_i into finite elements of size (i.e., maximum diameter) h such that all the fine-mesh elements together form a regular finite element division of Ω . We denote this fine division by \mathcal{T}_h . For each Ω_i , we consider an enlarged subdomain $\Omega_i^\delta = \{e \in \mathcal{T}_h : \text{dist}(e, \Omega_i) \leq \delta\}$, where $\text{dist}(e, \Omega_i) = \min_{x \in e, y \in \Omega_i} |x - y|$. The union of Ω_i^δ , $i = 1, \dots, m$, covers Ω with overlap proportional to δ . Let $K_0 \subset H_0^1(\Omega)$ and $K \subset H_0^1(\Omega)$ denote the continuous, piecewise r th-order polynomial ($r \geq 1$) finite element subspaces, with zero trace on $\partial\Omega$, over the H -level and h -level subdivisions of Ω , respectively. For $i = 1, \dots, m$, let K_i denote the continuous, piecewise r th-order polynomial finite element subspace with zero trace on the boundary $\partial\Omega_i^\delta$ and extended to have zero value outside $\Omega_i^\delta \cup \partial\Omega_i^\delta$. Then $K_i^\ominus = K_i$ for $i = 0, 1, \dots, m$, and it can be shown that

$$K = \sum_{i=0}^m K_i.$$




FIGURE 1. Decomposition of a rectangular-shaped domain in \mathbb{R}^2 .

Thus the space decomposition (4), with summation index from 0 to m , holds. We assume that the overlapping subdomains are chosen such that each subdomain Ω_i^δ and its corresponding finite element subspace K_i can be painted one of n_c colors (numbered from 1 to n_c), with subdomains painted the same color being pairwise nonintersecting. The coarse mesh and its corresponding subspace K_0 are painted the color 0. Moreover, n_c should be independent of h . For general domain Ω , finding overlapping subdomains with such property is nontrivial. If Ω is the Cartesian product of intervals, we can easily find overlapping subdomains with $n_c = 2$ if $d = 1$, and $n_c \leq 4$ if $d = 2$, and $n_c \leq 6$ if $d = 3$. For the example of Figure 1, $d = 2$ and $n_c = 4$. Then the total number of colors needed for (7) and (9) to hold is $c = n_c + 1$.

7.1.2. *Estimating C_1 for equations (48) and (51).* Let $\{\theta_i\}_{i=1}^m$ be a smooth partition of unity with respect to $\{\Omega_i\}_{i=1}^m$, i.e., $\theta_i \in C^\infty(\Omega)$ with $\theta_i \geq 0$, $\theta_i = 0$ outside of Ω_i , and $\sum_{i=1}^m \theta_i = 1$. Let I_h be the finite element interpolation mapping onto K which uses the function values at the h -level nodes. For any $v \in K$, let v_0 be the projection in the L^2 -norm of v onto K_0 , i.e., $v_0 \in K_0$ and $\int_\Omega (v_0 - v)\phi \, dx = 0$ for all $\phi \in K_0$, and let $v_i = I_h(\theta_i(v - v_0))$. Then, it can be seen that $v_i \in K_i$ for $i = 0, 1, \dots, m$ and satisfy $v = \sum_{i=0}^m v_i$ [45, pp. 163-165], [57, p. 607]. Moreover, by further choosing θ_i so that $|\nabla\theta_i|$ has a certain boundedness property, it was recently shown in [53, Lem. 4.1] that, for any $s \geq 1$,

$$\left(\sum_{i=0}^m \|v_i\|^s \right)^{\frac{1}{s}} \leq C c^{\frac{1}{s}} \left(1 + \left(\frac{H}{\delta} \right)^{\frac{1}{2}} \right) \|v\|,$$

where C is a constant independent of the mesh parameters and m . Taking $s = 2$ and using the subspace nature of K_i , we obtain that, for any $v_i \in K_i$, $i = 0, 1, \dots, m$, there exists $\bar{u}_i \in K_i$ satisfying (5) (with summation index from 0 to m), where

$$C_1 = C\sqrt{c} \left(1 + \left(\frac{H}{\delta} \right)^{\frac{1}{2}} \right).$$

(also see [14, Thm. 16] and a work of Dryja and Widlund cited therein for related results). By choosing the overlapping size δ proportional to the coarse-mesh size H , the constant C_1 will be independent of the mesh parameters and the number of subdomains m .

7.1.3. Estimating C_2 for equations (48) and (51). Consider F given by (50), associated with the equation (51). By the mean value theorem, for any $u \in \mathfrak{R}$, $v \in \mathfrak{R}$, we have $|u^3 - v^3| = 3|\theta u + (1 - \theta)v|^2|u - v| \leq 3(|u| + |v|)^2|u - v| \leq 6(|u|^2 + |v|^2)|u - v|$ for some $\theta \in [0, 1]$. Thus, using the continuous embedding of $H^1(\Omega)$ in $L^p(\Omega)$ for $p < 2d/(d - 2)$ and $d = 2, 3$ (see [17, p. 114], [24, p. 21]), we have for any $u, v \in H^1(\Omega)$ and any subdomain Ω' of Ω that $u, v \in L^4(\Omega)$ and

$$\begin{aligned} \left| \int_{\Omega'} (u^3 - v^3)w \, dx \right| &\leq 6 \int_{\Omega'} |u|^2|u - v||w| + |v|^2|u - v||w| \, dx \\ &\leq 6 \left(\left(\int_{\Omega'} |u|^4 dx \right)^{\frac{1}{2}} + \left(\int_{\Omega'} |v|^4 dx \right)^{\frac{1}{2}} \right) \left(\int_{\Omega'} |u - v|^2|w|^2 dx \right)^{\frac{1}{2}} \\ &\leq 6 \left(\|u\|_{L^4(\Omega')}^2 + \|v\|_{L^4(\Omega')}^2 \right) \|u - v\|_{L^4(\Omega')} \|w\|_{L^4(\Omega')} \\ &\leq C \left(\|u\|_{H^1(\Omega')}^2 + \|v\|_{H^1(\Omega')}^2 \right) \|u - v\|_{H^1(\Omega')} \|w\|_{H^1(\Omega')}, \end{aligned}$$

where C depends only on the embedding constant. Also, define $\Omega_0^\delta = \Omega$ for convenience, so that every $v \in K_i$ vanishes outside of Ω_i^δ ($i = 0, 1, \dots, m$). Then, for F given by (50), we have from the above inequality that, for $i, j = 0, 1, \dots, m$,

$$\begin{aligned} a_{ij} &= \langle F'(w_{ij} + u_{ij}) - F'(w_{ij}), v_i \rangle \\ &= \int_{\Omega_i^\delta \cap \Omega_j^\delta} (\nabla u_{ij})^T \nabla v_i + u_{ij}v_i + ((w_{ij} + u_{ij})^3 - w_{ij}^3)v_i \, dx \\ &\leq \left(1 + C \|w_{ij} + u_{ij}\|_{H^1(\Omega_i^\delta \cap \Omega_j^\delta)}^2 + C \|w_{ij}\|_{H^1(\Omega_i^\delta \cap \Omega_j^\delta)}^2 \right) \\ (52) \quad &\cdot \|u_{ij}\|_{H^1(\Omega_i^\delta \cap \Omega_j^\delta)} \|v_i\|_{H^1(\Omega_i^\delta \cap \Omega_j^\delta)}, \end{aligned}$$

for any $w_{ij} \in K$, $u_{ij} \in K_j$, $v_i \in K_i$, with $a_{ij} = 0$ whenever $\Omega_i^\delta \cap \Omega_j^\delta = \emptyset$. Assume there exists a constant $\alpha > 0$ such that $\|w_{ij} + u_{ij}\|_{H^1(\Omega_i^\delta \cap \Omega_j^\delta)}^2 + \|w_{ij}\|_{H^1(\Omega_i^\delta \cap \Omega_j^\delta)}^2 \leq \alpha$ for $i, j = 0, 1, \dots, m$. Also, for $i, j = 1, \dots, m$, let $\epsilon_{ij} = 0$ if $\Omega_i^\delta \cap \Omega_j^\delta = \emptyset$ and otherwise let $\epsilon_{ij} = 1$. Let \hat{c} be the smallest integer such that every subdomain intersects at most \hat{c} other subdomains. It is not difficult to show that the symmetric matrix $\mathcal{E} = [\epsilon_{ij}]_{i,j=1}^m$ has the following estimate of its spectral radius (see [53, Corollary 4.1] for a proof):

$$\rho(\mathcal{E}) \leq \max_{i=1, \dots, m} \sum_{j=1}^m \epsilon_{ij} \leq \hat{c}.$$

This together with the estimate (52) yields

$$\begin{aligned}
 \sum_{i=1}^m \sum_{j=1}^m a_{ij} &\leq (1 + C\alpha) \sum_{i=1}^m \sum_{j=1}^m \epsilon_{ij} \|u_{ij}\| \|v_i\| \\
 &\leq (1 + C\alpha) \sum_{i=1}^m \sum_{j=1}^m \epsilon_{ij} \max_{i=1, \dots, m} \|u_{ij}\| \|v_i\| \\
 (53) \qquad &= (1 + C\alpha) \hat{c} \left(\sum_{j=1}^m \max_{i=1, \dots, m} \|u_{ij}\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|v_i\|^2 \right)^{\frac{1}{2}}.
 \end{aligned}$$

Next, by using the fact $\Omega_j^\delta, j \in I(k)$, are disjoint subsets of Ω for $k = 1, \dots, c$, the estimate (52) yields

$$\begin{aligned}
 \sum_{j=1}^m a_{0j} &\leq (1 + C\alpha) \sum_{j=1}^m \|u_{0j}\| \|v_0\|_{H^1(\Omega_j^\delta)} \\
 &\leq (1 + C\alpha) \left(\sum_{j=1}^m \|u_{0j}\|^2 \right)^{\frac{1}{2}} \left(\sum_{j=1}^m \|v_0\|_{H^1(\Omega_j^\delta)}^2 \right)^{\frac{1}{2}} \\
 &\leq (1 + C\alpha) \sqrt{c} \left(\sum_{j=1}^m \|u_{0j}\|^2 \right)^{\frac{1}{2}} \|v_0\|, \quad \forall u_{0j} \in K_j, \forall v_0 \in K_0.
 \end{aligned}$$

Similar to the above argument, the estimate (52) gives

$$\begin{aligned}
 \sum_{i=1}^m a_{i0} &\leq (1 + C\alpha) \sum_{i=1}^m \|u_{i0}\|_{H^1(\Omega_i^\delta)} \|v_i\| \\
 &\leq (1 + C\alpha) \left(\sum_{i=1}^m \|u_{i0}\|_{H^1(\Omega_i^\delta)}^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|v_i\|^2 \right)^{\frac{1}{2}}, \quad \forall u_{i0} \in K_0, \forall v_i \in K_i.
 \end{aligned}$$

We combine these estimates to obtain

$$\begin{aligned}
 \sum_{i=0}^m \sum_{j=0}^m a_{ij} &= a_{00} + \sum_{j=1}^m a_{0j} + \sum_{i=1}^m \sum_{j=1}^m a_{ij} + \sum_{i=1}^m a_{i0} \\
 &\leq (1 + C\alpha) \|u_{00}\| \|v_0\| + (1 + C\alpha) \sqrt{c} \left(\sum_{j=1}^m \|u_{0j}\|^2 \right)^{\frac{1}{2}} \|v_0\| \\
 &\quad + (1 + C\alpha) \hat{c} \left(\sum_{j=1}^m \max_{i=1, \dots, m} \|u_{ij}\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|v_i\|^2 \right)^{\frac{1}{2}} \\
 &\quad + (1 + C\alpha) \left(\sum_{i=1}^m \|u_{i0}\|_{H^1(\Omega_i^\delta)}^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|v_i\|^2 \right)^{\frac{1}{2}} \\
 &\leq \tilde{C}_2 \left(\sum_{j=0}^m \max_{i=0, 1, \dots, m} \|u_{ij}\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=0}^m \|v_i\|^2 \right)^{\frac{1}{2}} \\
 (54) \qquad &\quad + (1 + C\alpha) \left(\sum_{i=1}^m \|u_{i0}\|_{H^1(\Omega_i^\delta)}^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|v_i\|^2 \right)^{\frac{1}{2}},
 \end{aligned}$$

with \tilde{C}_2 a constant depending on $C\alpha, c, \hat{c}$ only. Compared with (6) (with $i, j = 0, 1, \dots, m$), we see that (54) has an extra term on the right-hand side. In the

appendix, we will show that this extra term does not affect the convergence rate result of Section 4. In particular, we will show that Lemmas 1 and 2 hold with $C_2 = \tilde{C}_2 + (1 + C\alpha)\sqrt{c}$, so that Theorem 1 is still valid.

For F specified by (44) and associated with the equation (48), it can be similarly proved using (49) that (54) holds, possibly with different constants C and α .

Upon applying the asynchronous method (10)–(12) with the above choice of space decomposition and under the assumptions (13)–(14), we obtain a parallel DD method for (48) and (51) whose convergence factor, according to Theorem 1 and the above estimates of C_1 and C_2 and assuming the overlapping size δ is proportional to the coarse mesh size H , is independent of the mesh parameters and the number of the subdomains.

7.2. Multigrid methods.

7.2.1. Construction of the multigrid subspaces. In MG methods, Ω is divided into a finite element triangulation \mathcal{T} by a successive refinement process. More precisely, we have $\mathcal{T} = \mathcal{T}_J$ for some $J > 1$, where \mathcal{T}_k , $k = 1, \dots, J$, is a nested sequence of regular quasi-uniform triangulation, i.e., \mathcal{T}_k is a collection of simplex $\mathcal{T}_k = \{\tau_i^k\}$ of size (i.e., maximum diameter) h_k such that $\Omega = \bigcup_i \tau_i^k$ and for which the quasi-uniformity constants are independent of k [17, Eq. (3.2.28)] and with each simplex in \mathcal{T}_{k-1} being the union of simplex in \mathcal{T}_k . We further assume that there is a constant $r < 1$, independent of k , such that h_k is proportional to r^{2k} .

For example, in the two-dimensional case of $d = 2$, if we construct \mathcal{T}_k by connecting the midpoints of the edges of the triangles of \mathcal{T}_{k-1} , with \mathcal{T}_1 being the given coarsest initial triangulation, the resulting sequence of triangulation is quasi-uniform and $r = 1/\sqrt{2}$ (see Figure 2). Corresponding to each triangulation \mathcal{T}_k , we define the finite element subspace:

$$\mathcal{M}_k = \{v \in H_0^1(\Omega) : v|_{\tau} \in \mathcal{P}_1(\tau), \quad \forall \tau \in \mathcal{T}_k\},$$




FIGURE 2. The multigrid mesh and basis functions.

where $\mathcal{P}_1(\tau)$ denotes the space of real-valued linear functions of d real variables defined on τ . We associate with \mathcal{M}_k a nodal basis, denoted by $\{\phi_i^k\}_{i=1}^{n_k}$, that satisfies $\phi_i^k \in \mathcal{M}_k$ and

$$\phi_i^k(x_j^k) = \delta_{ij}, \quad \text{the Kronecker function,}$$

where $\{x_i^k\}_{i=1}^{n_k}$ is the set of all interior nodes of the triangulation \mathcal{T}_k . For each such nodal basis function, we define the one-dimensional subspace

$$K_i^k = \text{span}(\phi_i^k).$$

Then, $(K_i^k)^\ominus = K_i^k$ and we have the space decomposition

$$K = \sum_{k=1}^J \sum_{i=1}^{n_k} K_i^k \quad \text{with} \quad K = \mathcal{M}_J.$$

On each level k , we color the nodes of \mathcal{T}_k so that neighboring nodes are always of a different color. The number of colors needed for a regular mesh is a constant independent of the mesh parameters, which we denote by n_c . Then the total number of colors needed for (7) and (9) (with summation indices adjusted accordingly) to hold is $c = n_c J$.

7.2.2. *Estimating C_1 for equations (48) and (51).* Let Q_k be the projection in the L^2 -norm onto the subspace \mathcal{M}_k , which is well defined on $H_0^1(\Omega) \subset L^2(\Omega)$. For any $v \in K$, let $v^k = (Q_k - Q_{k-1})v$, $k = 1, \dots, J$. Then, by Prop. 8.6 in [57, p. 611], we have

$$\sum_{k=1}^J \|v^k\|^2 \leq C_0 \|v\|^2,$$

where C_0 is a constant independent of the mesh parameters and J . By further decomposing each v^k as

$$v^k = \sum_{i=1}^{n_k} v_i^k \quad \text{with} \quad v_i^k = v^k(x_i^k) \phi_i^k,$$

the above estimate can be refined to show that

$$v = \sum_{k=1}^J \sum_{i=1}^{n_k} v_i^k \quad \text{and} \quad \sum_{k=1}^J \sum_{i=1}^{n_k} \|v_i^k\|^2 \leq C \|v\|^2,$$

where C is a constant independent of the mesh parameters and the number of levels J [53, §4.2]. Thus, for any $v_i^k \in K_i^k$, $i = 1, \dots, n_k$, $k = 1, \dots, J$, there exists $\tilde{u}_i^k \in K_i^k$ satisfying (5) (with summation indices adjusted accordingly), where $C_1 = \sqrt{C}$.

7.2.3. *Estimating C_2 for equations (48) and (51).* Let Λ_i^k denote the support set of the basis function ϕ_i^k , for all i and k . Also, recall the constant $r < 1$ defined earlier. Then, for any $k < l$ and $1 \leq i \leq n_k, 1 \leq j \leq n_l$, the following estimate

$$\|u\|_{H^1(\Lambda_i^k \cap \Lambda_j^l)} \leq C_0 r^{d(l-k)} \|u\|, \quad \forall u \in K_i^k,$$

can be shown, where C_0 is a constant independent of the mesh parameters and J [53, Eq. (49)]. Then, for F given by (50), we obtain as in (52) that

(55)

$$\begin{aligned} & \langle F'(w+u) - F'(w), v \rangle \\ & \leq \left(1 + C \|w+u\|_{H^1(\Lambda_i^k \cap \Lambda_j^l)}^2 + C \|w\|_{H^1(\Lambda_i^k \cap \Lambda_j^l)}^2 \right) \|u\|_{H^1(\Lambda_i^k \cap \Lambda_j^l)} \|v\|_{H^1(\Lambda_i^k \cap \Lambda_j^l)} \\ & \leq \left(1 + C \|w+u\|_{H^1(\Lambda_i^k \cap \Lambda_j^l)}^2 + C \|w\|_{H^1(\Lambda_i^k \cap \Lambda_j^l)}^2 \right) C_0 r^{d(l-k)} \|u\| \|v\|, \\ & \quad \forall w \in K, u \in K_i^k, v \in K_j^l, \end{aligned}$$

where C is the embedding constant. For any i, j, k, l , defining

$$\varepsilon_{i,j}^{k,l} = \begin{cases} C_0 \gamma^{d|l-k|}, & \text{if } \text{supp}(\phi_i^k) \cap \text{supp}(\phi_j^l) \neq \emptyset; \\ 0, & \text{otherwise.} \end{cases}$$

Assuming there exists a constant $\alpha > 0$ such that $\|w_{i,j}^{k,l} + u_{i,j}^{k,l}\|^2 + \|w_{i,j}^{k,l}\|^2 \leq \alpha$ for all i, j, k, l , the estimate (55) then yields

$$\begin{aligned} & \sum_{k=1}^J \sum_{i=1}^{n_k} \sum_{l=1}^J \sum_{j=1}^{n_l} \langle F'(w_{i,j}^{k,l} + u_{i,j}^{k,l}) - F'(w_{i,j}^{k,l}), v_i^k \rangle \\ & \leq C_0(1 + C\alpha) \sum_{i,k} \sum_{j,l} \varepsilon_{i,j}^{k,l} \|u_{i,j}^{k,l}\| \|v_i^k\| \\ & \leq C_0(1 + C\alpha) \sum_{k=1}^J \sum_{i=1}^{n_k} \sum_{l=1}^J \sum_{j=1}^{n_l} \varepsilon_{i,j}^{k,l} \max_{i,k} \|u_{i,j}^{k,l}\| \cdot \|v_i^k\|, \\ & \quad \forall u_{i,j}^{k,l} \in K_j^l, \forall v_i^k \in K_i^k. \end{aligned}$$

With proper ordering of the indices, the matrix $\mathcal{E} = [\varepsilon_{i,j}^{k,l}]$ is symmetric and its spectral radius $\rho(\mathcal{E})$ has been shown to be less than a constant independent of the mesh parameters and the number of levels [45, pp. 182–184]. Therefore,

$$\begin{aligned} & \sum_{k=1}^J \sum_{i=1}^{n_k} \sum_{l=1}^J \sum_{j=1}^{n_l} \langle F'(w_{i,j}^{k,l} + u_{i,j}^{k,l}) - F'(w_{i,j}^{k,l}), v_i^k \rangle \\ & \leq C_0(1 + C\alpha) \rho(\mathcal{E}) \left(\sum_{l=1}^J \sum_{j=1}^{n_l} \max_{i,k} \|u_{i,j}^{k,l}\|^2 \right)^{\frac{1}{2}} \left(\sum_{k=1}^J \sum_{i=1}^{n_k} \|v_i^k\|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

which shows that (6) holds, with the constant $C_2 = C_0(1 + C\alpha)\rho(\mathcal{E})$ independent of the mesh parameters and the number of levels for the MG approximation.

For F specified by (44), it can be similarly proved that (6) holds with C_2 some constant independent of the mesh parameters and the number of levels.

Upon applying the asynchronous method (10)–(12) with the above choice of space decomposition and under the assumptions (13)–(14), we obtain a parallel MG method for (48) and (51) whose convergence factor, according to the above estimates of C_1 and C_2 and Theorem 1, is independent of the mesh parameters. This method generalizes the BPX multigrid method proposed in [10], which was used as a preconditioner for linear elliptic problems. Here, the parallel MG method is used as a solver and is applicable not only to linear, but also to nonlinear elliptic problems. And it further allows for asynchronous updates.

8. APPLICATIONS TO OBSTACLE PROBLEMS

In this section, we will apply our asynchronous algorithm to the obstacle problem

$$-\Delta u \geq f \text{ in } \Omega, \quad u \geq \psi \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

where $f \in L^2(\Omega)$ and $\psi \in H^2(\Omega)$ satisfies $\psi \leq 0$ on $\partial\Omega$. This problem is equivalent to (2) with

$$K = \{v \in H_0^1(\Omega) : v \geq \psi \text{ a.e. in } \Omega\}, \quad F(v) = \int_{\Omega} \left(\frac{1}{2} |\nabla v|^2 - fv \right) dx.$$

We will use the overlapping domain decomposition without the coarse mesh. If a coarse mesh is added, it is not known how the coarse mesh obstacle can be chosen to obtain an algorithm whose convergence factor is independent of mesh parameters.

Let Ω_i^δ be defined as in subsection 7.1, $i = 1, \dots, m$. Let θ_i be the partition of unity with respect to Ω_i^δ as described in subsection 7.1.2. Accordingly, let $\psi_i = I_h(\theta_i \psi)$ for all i and let $\psi = \sum_{i=1}^m \psi_i$, where I_h is the interpolation operator using the h -level nodal values. Thus, the obstacle function ψ is replaced by its finite-element interpolation. Defining

$$K_i = \{v \in H_0^1(\Omega_i^\delta) : v \geq \psi_i, v|_e \in P_1(e), \forall e \in \mathcal{T}_h\},$$

and also assuming that K has been replaced by its finite-element analog, it is easy to see that (4) holds. Suppose we apply the asynchronous algorithm with the initial values $u_i(0) \in K_i$ chosen to satisfy $\sum_{i=1}^m u_i(0) \leq \bar{u}$. Then, it can be shown (see, e.g., [1] and [51, §4]) that

$$\sum_{i=1}^m w_i(t^i) \leq \bar{u}, \quad \forall t \geq 1,$$

where $w_i(\cdot)$ is defined by (12) and t^i denotes the greatest element of T^i less than $t + B$. Thus (27) remains valid if we assume there exist $\bar{u}_i \in K_i$ satisfying (5) only for those $v_i \in K_i$ satisfying $\sum_{i=1}^m v_i \leq \bar{u}$. Since (27) is the only point in the proofs of Lemmas 1 and 2 where (5) is used, these lemmas and Theorem 1 would remain valid. Under the condition that $\sum_{i=1}^m v_i \leq \bar{u}$, the constant C_1 in (5) can be estimated by choosing the partition of unity θ_i to satisfy $|\nabla \theta_i| \leq C/\delta$ and setting

$$\bar{u}_i = v_i + \theta_i \left(\bar{u} - \sum_{j=1}^m v_j \right).$$

Then, it is straightforward to show that $\bar{u}_i \in K_i$ and

$$\sum_{i=1}^m \bar{u}_i = \bar{u}, \quad \sum_{i=1}^m \|\bar{u}_i - v_i\|^2 \leq C(1 + \delta^{-2}) \left\| \bar{u} - \sum_{j=1}^m v_j \right\|^2,$$

with the constant C being independent of u, v_i , the mesh size h , the overlapping size δ , and the number of subdomains m (cf. subsection 7.1.2). The above estimate shows that (5) holds with $C_1 = \sqrt{C} \sqrt{1 + 1/\delta^2}$. Also, by dropping the coarse mesh and taking into account the difference between the above F and the F given by (50), we see from the proof of (53) that (6) holds with $C_2 = (1 + C)\hat{c}$, with C an embedding constant. Assuming that B is bounded by a given constant and γ is

bounded by a constant less than 1, we then obtain from the definitions in Lemmas 1 and 2 that

$$\frac{1}{D} \leq A_1 \leq D, \quad A_2 \leq D, \quad A_3 \leq D(1 + \delta^{-2}), \quad \frac{A_4}{A_3} \leq D,$$

for some $D > 0$ independent of h, δ and m . Then, the convergence factor given by (38) can be estimated by

$$\begin{aligned} \varrho &= \max \left\{ 1 - \frac{\gamma}{1 + A_1^{-1}A_3} + \gamma^{3/2} \frac{(A_2 + A_1A_4/A_3)}{1 + A_1A_3^{-1}}, \gamma^{1/2}(A_1^{-1} + \gamma^{3/2}A_2A_1^{-1}) \right\} \\ &\leq \max \left\{ 1 - \frac{\gamma}{1 + D^2(1 + \delta^{-2})} + \gamma^{3/2}(D + D^2), \gamma^{1/2}(D + D^2) \right\} \\ &\leq \max \left\{ 1 - \frac{\gamma}{D_1(1 + \delta^{-2})} + \gamma^{3/2}D_2, \gamma^{1/2}D_2 \right\}, \end{aligned}$$

where we let $D_1 = 1 + D^2$ and $D_2 = D + D^2$. Thus, for

$$\gamma < \min\{(D_1D_2(1 + \delta^{-2}))^{-2}, D_2^{-2}\},$$

we have $\varrho < 1$, independent of h and m . Finally, we note that the convergence factor for some synchronous overlapping domain decomposition without the coarse mesh has been studied in [1, 60]. The schemes obtained from our algorithms are different from those of [1, 60] in the treatment of the subproblem obstacles. The algorithms of [1, 60] use the global obstacle for the subdomain problems. In our algorithms, the subdomain obstacles can be updated dynamically during the iterations.

9. APPENDIX

In this appendix, we show that (54) can be used in place of (6) to prove Lemmas 1 and 2 for the DD method of subsection 7.1. Here, the indices i and j are understood to always range over $0, 1, \dots, m$, instead of $1, \dots, m$.

First, we note that condition (6) is used only to show (22), (30) and (34) in the proofs. For (22), if we use condition (54) instead of (6), then (22) would have \tilde{C}_2 in place of C_2 and would have the following extra term on its right-hand side:

$$E = (1 + C\alpha)\gamma \left(\sum_{i=1}^m \|u_0(\tau_0^i(t)) - u_0(t)\|_{H^1(\Omega_i^\delta)}^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|s_i(t)\|^2 \right)^{\frac{1}{2}}.$$

Correspondingly, (24) would have \tilde{C}_2 in place of C_2 and would have the above extra term on its right-hand side. Using (23) and the fact that $\Omega_i^\delta, i \in I(k)$, are disjoint subsets of Ω for $k = 1, \dots, c$, we see that

$$\begin{aligned} E &\leq (1 + C\alpha)\gamma^2\sqrt{B} \left(\sum_{i=1}^m \sum_{\tau=t-B+1}^{t-1} \|s_0(\tau)\|_{H^1(\Omega_i^\delta)}^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|s_i(t)\|^2 \right)^{\frac{1}{2}} \\ &\leq (1 + C\alpha)\gamma^2\sqrt{Bc} \left(\sum_{\tau=t-B+1}^{t-1} \|s_0(\tau)\|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^m \|s_i(t)\|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

which implies that (24) holds with $C_2 = \tilde{C}_2 + (1 + C\alpha)\sqrt{c}$. The remainder of the proof of Lemma 1 then proceeds as before.

For (30) and (34), a similar argument can be applied to show that Lemma 2 holds with the above choice of C_2 .

REFERENCES

1. L. Badea and J. Wang, *An additive Schwarz method for variational inequalities*, Math. Comp. **69** (2000), 1341–1354. MR **2001a**:65075
2. R. Bank, *Analysis of a multilevel iterative method for nonlinear finite element equations*, Math. Comp. **39** (1982), 453–465. MR **83j**:65105
3. R. Bank and T. F. Chan, PLTMGC: a multigrid continuation program for parametrized nonlinear systems, SIAM T. Sci. Statist. Comput., **7** (1986), 540–559. MR **87d**:65125
4. D. P. Bertsekas, D. Castañon, J. Eckstein, and S. A. Zenios, *Parallel computing in network optimization*, Handbooks in Operations Research and Management Science 7: Network Models (Amsterdam) (M. O. Ball, T. L. Magnanti, C. L. Monma, and G. L. Nemhauser, eds.), Elsevier Scientific, 1995, pp. 331–399. CMP 97:04
5. D. P. Bertsekas and J. N. Tsitsiklis, *Parallel and distributed computation: Numerical methods*, Prentice-Hall, Englewood Cliffs, 1989.
6. ———, *Some aspects of parallel and distributed iterative algorithms—a survey*, Automatica **27** (1991), 3–21. MR **91m**:65137
7. J. Bramble and X. Zhang, *The analysis of multigrid methods*, Handbook of numerical analysis, Vol. VII, 173–415, North-Holland, Amsterdam, 2000. CMP 2001:08
8. J. H. Bramble, J. E. Pasciak, and A. H. Schatz, *The construction of preconditioners for elliptic problems by substructuring. IV*, Math. Comp. **53** (1989), 1–24. MR **89m**:65098
9. J. H. Bramble, J. E. Pasciak, J. Wang, and J. Xu, *Convergence estimates for product iterative methods with applications to domain decomposition*, Math. Comp. **57** (1991), 1–21. MR **92d**:65094
10. J. H. Bramble, J. E. Pasciak, and J. Xu, *Parallel multilevel preconditioners*, Math. Comp. **55** (1990), 1–22. MR **90k**:65170
11. A. Brandt, *Multilevel adaptive solutions to boundary value problems*, Math. Comp. **31** (1977), 333–390. MR **55**:4714
12. A. Brandt and C. W. Cryer, *Multigrid algorithms for the solution of linear complementary problems arising from free boundary problems*, SIAM J. Sci. Comput. **4** (1983), 655–684. MR **85h**:65261
13. J. V. Burke and P. Tseng, *A unified analysis of Hoffman’s bound via Fenchel duality*, SIAM J. Optim. **6** (1996), 265–282. MR **97c**:90120
14. T. F. Chan and T. P. Mathew, *Domain decomposition algorithms*, Acta Numerica (1994), 61–143. MR **95f**:65214
15. T. F. Chan and I. Sharapov, *Subspace correction multilevel methods for elliptic eigenvalue problems*, Numer. Linear Algebra Appl., **1** (1993), 1–7.
16. D. Chazan and W. L. Miranker, *Chaotic relaxation*, Lin. Algeb. Appl. **2** (1969), 199–222. MR **40**:5114
17. P. G. Ciarlet, *The finite element method for elliptic problems*, North-Holland, Amsterdam, 1978. MR **58**:25001
18. M. Dryja and W. Hackbusch, *On the nonlinear domain decomposition method*, BIT **37** (1997), 296–311. MR **97m**:65100
19. M. Dryja and O. B. Widlund, *Towards a unified theory of domain decomposition algorithms for elliptic problems*, Third international symposium on domain decomposition methods for partial differential equations (Philadelphia, PA) (T. Chan, R. Glowinski, J. Periaux, and O. B. Widlund, eds.), SIAM, 1990. MR **91m**:65294
20. I. Ekeland and R. Temam, *Convex analysis and variational problems*, North-Holland, Amsterdam, 1976. MR **57**:3931b
21. B. H. Elton, *An asynchronous domain decomposition method for the two-dimensional Poisson equation*, Parallel processing for scientific computing, Vol. I, II (Norfolk, VA, 1993), SIAM, Philadelphia, PA, 1993, pp. 691–695. CMP 93:10
22. A. Frommer, H. Schwandt, and D. B. Szyld, *Asynchronous weighted additive Schwarz methods*, Electronic Trans. Numer. Anal. **5** (1997), 48–61. MR **98c**:65046
23. E. Gelman and J. Mandel, *On multilevel iterative methods for optimization problems*, Math. Programming **48** (1990), 1–17. MR **91b**:90180
24. R. Glowinski and P. Le Tallec, *Augmented Lagrangian and operator-splitting methods in nonlinear mechanics*, SIAM, Philadelphia, 1989. MR **91f**:73038
25. W. Hackbusch and H. D. Mittelmann, *On multigrid methods for variational inequalities*, Numer. Math. **42** (1983), 65–76. MR **84k**:49010

26. W. Hackbusch and A. Reusken, *Analysis of a damped nonlinear multilevel method*, Numer. Math. **55** (1989), 225–246. MR **90f**:65194
27. L. Hart and S. McCormick, *Asynchronous multilevel adaptive methods for solving partial differential equations on multiprocessors: basic ideas*, Parallel Comput. **12** (1989), no. 2, 131–144. MR **90m**:65250
28. R. H. Hoppe, *Multigrid solutions to the elastic plastic torsion problem in multiply connected domains*, Int. J. Numer. Methods Engin. **26** (1988), 631–646. MR **89a**:73001
29. R. H. W. Hoppe, *Two-sided approximations for unilateral variational inequalities by multigrid methods*, Optimization **16** (1987), 867–881. MR **88m**:49004
30. R. H. W. Hoppe and R. Kornhuber, *Adaptive multilevel methods for obstacle problems*, SIAM J. Numer. Anal. **31** (1994), 301–323. MR **95c**:65181
31. R. Kornhuber, *Monotone multigrid iterations for elliptic variational inequalities, II*, Numer. Math. **72** (1996), 481–499. MR **96k**:65081
32. O. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and quasilinear elliptic equations*, Academic Press, 1968. MR **39**:5941
33. P. Le Tallec, *Domain decomposition methods in computational mechanics*, Comput. Mechan. Adv. **1** (1994), 121–220. MR **95b**:65147
34. Z.-Q. Luo and P. Tseng, *On the linear convergence of descent methods for convex essentially smooth minimization*, SIAM J. Control Optim. **30** (1992), 408–425. MR **93a**:90064
35. ———, *Error bounds and convergence analysis of feasible descent methods: A general approach*, Ann. Oper. Res. **46** (1993), 157–178. MR **95a**:90138
36. J. Mandel, *A multilevel iterative method for symmetric, positive definite linear complementary problems*, Appl. Math. Optim. **11** (1984), 77–95. MR **85b**:90082
37. S. McCormick and D. Quinlan, *Asynchronous multilevel adaptive methods for solving partial differential equations on multiprocessors: performance results*, Parallel Comput. **12** (1989), no. 2, 145–156. MR **90m**:65251
38. S. F. McCormick, *Multilevel adaptive methods for partial differential equations*, Frontiers in Applied Mathematics, vol. 6, SIAM, Philadelphia, 1989. MR **91h**:65200
39. J.-C. Miellou, L. Giraud, A. Laouar, and P. Spiteri, *Subdomain decomposition methods with overlapping and asynchronous iterations*, Progress in partial differential equations: the Metz surveys, Longman Sci. Tech., Harlow, 1991, pp. 166–183. MR **94h**:65125
40. A. Quarteroni and A. Valli, *Domain decomposition methods for partial differential equations*, Oxford Science Publications, Oxford, 1999.
41. R. Rannacher, *On the convergence of the Newton-Raphson method for strongly nonlinear finite element equations*, Nonlinear computational mechanics (P. Wriggers and W. Wagner, eds.), Springer-Verlag, 1991.
42. R. T. Rockafellar, *Convex analysis*, Princeton University Press, Princeton, 1970. MR **43**:445
43. ———, *Network flows and monotropic optimization*, Wiley-Interscience, New York, 1984. MR **89d**:90156
44. I. Sharapov, *Multilevel subspace correction for large scale optimization problems*, East-West J. Numer. Math., **9** (2000), 233–252.
45. B. F. Smith, P. E. Bjørstad, and W. D. Gropp, *Domain decomposition: Parallel multilevel algorithms for elliptic partial differential equations*, Cambridge University Press, Cambridge, 1996. MR **98g**:65003
46. A. K. Stagg and G. F. Carey, *Asynchronous nonlinear iteration and domain decomposition*, Parallel processing for scientific computing (Houston, TX, 1991), SIAM, Philadelphia, PA, 1992, pp. 281–286. CMP 93:05
47. X.-C. Tai, *Parallel function decomposition and space decomposition methods with applications to optimisation, splitting and domain decomposition*, Preprint No. 231–1992, Institut für Mathematik, Technische Universität Graz (1992), See also <http://www.mi.uib.no/~tai>.
48. ———, *Parallel function and space decomposition methods*, Finite element methods, fifty years of the Courant element, Lecture notes in pure and applied mathematics (P. Neittaanmäki, ed.), vol. 164, Marcel Dekker inc., 1994, pp. 421–432. See also <http://www.mi.uib.no/~tai>. MR **95d**:65008
49. ———, *Domain decomposition for linear and nonlinear elliptic problems via function or space decomposition*, Domain decomposition methods in scientific and engineering computing (Proc. of the 7th international conference on domain decomposition, Penn. State University,

- 1993) (D. Keyes and J. Xu, eds.), Contemp. Math. 180, American Mathematical Society, 1994, pp. 355–360. MR **95k**:65121
50. ———, *Parallel function decomposition and space decomposition methods: Part II. Space decomposition*, Beijing Mathematics **1**, **part 2** (1995), 135–152, See also <http://www.mi.uib.no/~tai>.
 51. ———, *Convergence rate analysis of domain decomposition methods for obstacle problems*, East-West J. Numer. Math., **9** (2000), 233–253.
 52. X.-C. Tai and M. Espedal, *Rate of convergence of some space decomposition method for linear and nonlinear elliptic problems*, SIAM J. Numer. Anal. **35** (1998), 1558–1570. MR **99k**:65101
 53. X.-C. Tai and J.-C. Xu, *Global and uniform convergence of subspace correction methods for some convex optimization problems*, Math. Comp., posted on May 11, 2001, PII S0025-5718(01)01311-4 (to appear in print).
 54. M. Tinkham, *Introduction to superconductivity*, McGraw Hill, New York, 1975.
 55. P. Tseng, *On the rate of convergence of a partially asynchronous gradient projection algorithm*, SIAM J. Optim. **1** (1991), 603–619. MR **92k**:90096
 56. P. Tseng, D. P. Bertsekas, and J. N. Tsitsiklis, *Partially asynchronous, parallel algorithms for network flow and other problems*, SIAM J. Control Optim. **28** (1990), 678–710. MR **91e**:65075
 57. J.-C. Xu, *Iterative methods by space decomposition and subspace correction*, SIAM Rev. **34** (1992), 581–613. MR **93k**:65029
 58. ———, *A novel two-grid method for semilinear elliptic equations*, SIAM J. Sci. Comp. **15** (1994), 231–237. MR **94m**:65178
 59. ———, *Two-grid discretization techniques for linear and nonlinear PDEs*, SIAM J. Numer. Anal. **27** (1996), 1759–1777. MR **97i**:65169
 60. J. Zeng and S. Zhou, *On monotone and geometric convergence of Schwarz methods for two-sided obstacle problems*, SIAM J. Numer. Anal. **35** (1998), 600–616. MR **99d**:65199

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, JOHANNES BRUNSGATE 12, 5007, BERGEN, NORWAY

E-mail address: Xue-Cheng.Tai@mi.uib.no

DEPARTMENT OF MATHEMATICS, BOX 354350, UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON 98195

E-mail address: tseng@math.washington.edu