

**CORES OF CONVEX GAMES**

**Lloyd S. Shapley**

**April 1971**

**P-4620**

CORES OF CONVEX GAMES<sup>†</sup>

Lloyd S. Shapley\*

The Rand Corporation, Santa Monica, California

1. INTRODUCTION

The core of an  $n$ -person game is the set of feasible outcomes that cannot be improved upon by any coalition of players. A convex game is one that is based on a convex set function (see below); intuitively this means that the incentives for joining a coalition increase as the coalition grows, so that one would expect a "snowballing" or "band-wagon" effect when the game is played cooperatively.

In this paper we show that the core of a convex game is not empty--in fact, it is quite large--and that it has an especially regular structure. We further show that

---

<sup>†</sup> Presented at the Fifth Informal Conference on Game Theory, held at Princeton University in April 1965. This paper is based on a Rand Corporation research memorandum [S4], written under the sponsorship of Air Force Project Rand. The author wishes to acknowledge the stimulus of a query from Jack Edmonds concerning the properties of convex set functions.

\*Any views expressed in this paper are those of the author. They should not be interpreted as reflecting the views of The Rand Corporation or the official opinion or policy of any of its governmental or private research sponsors. Papers are reproduced by The Rand Corporation as a courtesy to members of its staff.

certain other cooperative solution concepts are related in a simple way to the core. Specifically (1) the value of a convex game is the center of gravity of the extreme points of the core, and (2) the von Neumann-Morgenstern stable set solution of a convex game is unique and coincides with the core. In a subsequent paper [M-P-S2] rather similar results will be presented for two other cooperative solutions: the kernel and the bargaining set.

### 1.1. Notation

We shall systematically use the letters  $n, s, t, \dots$  to denote the number of elements in the finite sets  $N, S, T, \dots$ . The letter " $\emptyset$ " will denote the empty set, and " $\subset$ ", " $\supset$ " will denote strict inclusion. "Payoff vectors" are elements of the  $n$ -dimensional linear space  $E^N$  with coordinates indexed by the elements of  $N$ . If  $a \in E^N$  and  $S \subseteq N$  we shall often write  $a(S)$  for  $\sum_S a_i$ , treating  $a$  as an additive function on the subsets of  $N$ . The hyperplane in  $E^N$  defined by the equation  $a(S) = v(S)$ ,  $\emptyset \subset S \subseteq N$ , will be denoted by  $H_S$ .

## 2. CONVEX GAMES

In this paper, a game is a function  $v$  from a Boolean ring  $\mathcal{N}$  to the reals, satisfying

$$(1) \quad v(0) = 0.$$

It is superadditive if

$$(2) \quad v(S) + v(T) \leq v(S \cup T), \quad \text{all } S, T \in \mathcal{N} \text{ with } S \cap T = 0.$$

It is convex\* if

$$(3) \quad v(S) + v(T) \leq v(S \cup T) + v(S \cap T), \quad \text{all } S, T \in \mathcal{N}.$$

It is strictly convex if inequality holds in (3) whenever neither  $S \subseteq T$  nor  $T \subseteq S$ , i.e., whenever  $S, T, S \cup T$ , and

---

\*In potential theory, functions satisfying the reverse inequality to (3) (i.e., negatives of convex functions) are called strongly subadditive or, when certain other properties are adduced, capacities [Ch, M]. In lattice theory and combinatorial mathematics, such functions, often restricted to be integer valued, as in the rank function of a matroid, are called submodular (or lower semi-modular), and convex functions supermodular (upper semi-modular) [Wh, E-R, C-R, E].

$S \cap T$  are all different. Note that (1) and (3) together imply (2).

To appreciate the term "convex", define for each  $R \in \mathcal{N}$  a differencing operator  $\Delta_R$  by:

$$[\Delta_R v](S) = v(S \cup R) - v(S - R), \quad \text{all } S \in \mathcal{N},$$

and let  $\Delta_{QR} v$  denote  $\Delta_Q(\Delta_R v)$ . Then (3) is equivalent to the assertion that these "second differences" are everywhere nonnegative, i.e., that

$$(4) \quad [\Delta_{QR} v](S) \geq 0, \quad \text{all } Q, R, S \in \mathcal{N}.$$

This is analogous to the nonnegative second derivatives associated with convex functions in real analysis.

Two games are termed equivalent if their difference is an additive game, i.e., obeys the equality in (2) (or (3) or (4)). It is easily seen that any game equivalent to a convex game is convex; that any positive scalar multiple of a convex game is convex; and that the sum of two or more convex games is convex. It follows that the ensemble of convex games (for fixed  $\mathcal{N}$ ) forms a convex cone in a suitable linear space (say  $E^{\mathcal{N}-\{0\}}$ ), and that this cone contains the subspace of additive games.

## 2.1. Interpretation

Throughout this paper  $\mathcal{N}$  will be the Boolean ring of subsets of a finite set  $N$ --thus,  $\mathcal{N} = 2^N$ .<sup>\*</sup> In the standard application in game theory, the elements of  $N$  are "players"; the elements of  $\mathcal{N}$  are "coalitions"; and  $v(S)$ , called the "characteristic function", gives for each coalition the best payoff it can achieve without help from other players.

Superadditivity (2) arises naturally in this interpretation, but convexity (3) is another matter. For example, in a voting situation  $S$  and  $T$ , but not  $S \cap T$ , might be winning coalitions, causing (3) to fail. To see what convexity does entail, regard the function  $m$ :

$$(5) \quad m(S, T) = v(S \cup T) - v(S) - v(T)$$

as defining the "incentive to merge" between disjoint coalitions  $S$  and  $T$ . Then it is a simple exercise to verify that (3) is equivalent to the assertion that  $m(S, T)$  is nondecreasing in each variable--whence the "snowballing" or "bandwagon" effect, mentioned in the introduction.

---

<sup>\*</sup>For the infinite case see [R].

Another condition that is equivalent to (3) (provided  $\mathcal{N}$  is finite) is to require that\*

$$(6) \quad v(S \cup \{i\}) - v(S) \leq v(T \cup \{i\}) - v(T),$$

for all individuals  $i \in N$  and all  $S \subseteq T \subseteq N - \{i\}$ . This expresses a sort of increasing marginal utility for coalition membership, and is analogous to the "increasing returns to scale" associated with convex production functions in economics.

## 2.2 Convex Measure Games

Let  $v$  be given by

$$(7) \quad v(S) = f(\mu(S)), \quad \text{all } S \in \mathcal{N},$$

where  $\mu$  is a nonnegative additive function on  $\mathcal{N}$ , i.e., a measure, and  $f$  is a real function with  $f(0) = 0$ .\*\* Then it is easy to verify that  $v$  is convex if  $f$  is convex, and

---

\*Cf. [Wh].

\*\*Actually, any game can be put into this form; we need merely choose  $\mu$  so that all the numbers  $\mu(S)$  are different and define  $f$  accordingly. Thus, the general concept of "measure game", with  $f$  and  $\mu$  not further restricted, is of no interest in the finite case. (Cf. [A-S]).

In economic applications,  $\mu$  may represent the initial distribution of some resource, while  $f$  may represent a production function.

strictly convex if  $f$  is strictly convex. A game that is a convex function of a measure is called a convex measure game.\*

Not all convex games are convex measure games. A simple counterexample is the two-person convex game with  $v(\{1\}) = -v(\{2\}) = 1$ ,  $v(\{1,2\}) = v(0) = 0$ . But this game, being additive, is equivalent to the convex measure game  $v(S) = 0$ , so we might ask whether every convex game is at least equivalent to a convex measure game. Again the answer is no. Indeed, if this were true, then every convex game would be the sum of three convex measure games, since every additive game is the sum of two.\*\* But convex games exist that are not the sum of any number of convex measure games; indeed, the first twelve games of Table 1 have this property.

The situation can be viewed geometrically. As already remarked, the convex games form a cone in  $E^n$  which is invariant under equivalence. The subcone of convex measure

---

\*Curiously, if  $f$  is a function of several variables and  $\mu$  a vector of measures, then convexity of  $f$  does not imply convexity of  $v$  in (7), or even superadditivity. Indeed, in the case of a homogeneous function  $f(\lambda x) \equiv \lambda f(x)$ , it is concavity of  $f$ , not convexity, that implies superadditivity of  $v$ .

\*\*Simply express the game as the difference of two measures,  $\mu_1 - \mu_2$ , and let  $f_1(x) \equiv -f_2(x) \equiv x$ .



games, however, is neither convex nor invariant under equivalence (unless  $n$  is trivially small). The convex hull of that subcone, which is what we obtain when we take all sums of convex measure games, is invariant under equivalence but still does not include all convex games if  $n > 3$ .

The thirty-seven extremals of the cone of four-person convex games were determined in 1965 by S. A. Cook\* (Table 1). The games in the first seven symmetry classes listed are indecomposable (see below); the remainder are obtained from smaller games by adding dummy players. For larger  $n$ , little is known about the set of all extremals.\*\*

### 2.3. Decomposable Games\*\*\*

Let  $\mathcal{P} = \{N_1, \dots, N_p\}$  be a partition of  $N$  into  $p \geq 2$  nonempty subsets. The game  $v$  on  $\mathcal{N}$  is said to be decomposable (with respect to  $\mathcal{P}$ ) if  $v$  is additive across the partition, i.e., for each  $S \in \mathcal{N}$ ,

$$(8) \quad v(S) = v(S \cap N_1) + v(S \cap N_2) + \dots + v(S \cap N_p).$$

---

\*Private communication.

\*\*For the infinite case, see [R].

\*\*\*This section is based on [M-P-S1].

Table 1

EXTREMAL CONVEX GAMES ON FOUR PLAYERS

$\emptyset$	1	2	3	4	12	13	14	23	24	34	123	124	134	234	1234
0	0	0	0	0	0	1	1	1	1	1	2	2	2	2	4
0	0	0	0	0	1	0	1	1	1	1	2	2	2	2	4
0	0	0	0	0	1	1	0	1	1	1	2	2	2	2	4
0	0	0	0	0	1	1	1	0	1	1	2	2	2	2	4
0	0	0	0	0	1	1	1	1	0	1	2	2	2	2	4
0	0	0	0	0	1	1	1	1	1	0	2	2	2	2	4
0	0	0	0	0	1	0	0	0	0	0	1	1	1	1	2
0	0	0	0	0	0	1	0	0	0	0	1	1	1	1	2
0	0	0	0	0	0	0	1	0	0	0	1	1	1	1	2
0	0	0	0	0	0	0	0	1	0	0	1	1	1	1	2
0	0	0	0	0	0	0	0	0	1	0	1	1	1	1	2
0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	2
0	0	0	0	0	0	0	1	0	1	1	1	2	2	2	3
0	0	0	0	0	0	1	0	1	0	1	2	1	2	2	3
0	0	0	0	0	1	0	0	1	1	0	2	2	1	2	3
0	0	0	0	0	1	1	1	0	0	0	2	2	2	1	3
0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	2
0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	2
0	0	0	0	0	0	0	0	0	0	0	1	1	0	1	2
0	0	0	0	0	0	0	0	0	0	0	1	1	1	0	2
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	2
0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	2
0	0	0	0	0	1	1	1	1	1	1	2	2	2	2	3
0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1
0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1
0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1
0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
0	0	0	0	0	1	1	0	1	0	0	2	1	1	1	2
0	0	0	0	0	1	0	1	0	1	0	1	2	1	1	2
0	0	0	0	0	0	1	1	0	0	1	1	1	2	1	2
0	0	0	0	0	0	0	0	1	1	1	1	1	1	2	2
0	0	0	0	0	1	0	0	0	0	0	1	1	0	0	1
0	0	0	0	0	0	1	0	0	0	0	1	0	1	0	1
0	0	0	0	0	0	0	1	0	0	0	0	1	1	0	1
0	0	0	0	0	0	0	0	1	0	0	1	0	0	1	1
0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	1
0	0	0	0	0	0	0	0	0	0	1	0	0	1	1	1

Note that  $v$  is completely determined by its values on the subsets of the  $N_i$ . The  $p$  smaller games, obtained by restricting  $v$  to the subsets of each  $N_i$  in turn, are the components of the decomposition. A decomposable game has a unique finest decomposition, that is, a decomposition in which none of the components are themselves decomposable.\*

THEOREM 1. (a) A decomposable game is convex if and only if each component is convex.

(b) A convex game  $v$  is decomposable if and only if

$$(9) \quad v(N) = v(N_1) + v(N_2) + \dots + v(N_p)$$

holds for some partition  $\{N_1, \dots, N_p\}$  of  $N$  into  $p \geq 2$  nonempty subsets.

Proof.\*\* The first statement is immediate. For the second, suppose (9) holds and let  $S \in \mathcal{N}$ . By convexity,

---

\*The notion of decomposition was introduced in [VN-M]; see also [G1, G2, P1, P2].

\*\*Several proofs can be given; this one is due to Y. Kannai.

$$v(S) + v(N_1) \leq v(S \cap N_1) + v(S \cup N_1)$$

$$v(S \cup N_1) + v(N_2) \leq v(S \cap N_2) + v(S \cup N_1 \cup N_2)$$

...

...

$$v(S \cup N_1 \cup \dots \cup N_{p-1}) + v(N_p) \leq v(S \cap N_p) + v(S \cup N_1 \cup \dots \cup N_p).$$

Adding these inequalities to each other and to (9) yields

$$(10) \quad v(S) \leq v(S \cap N_1) + v(S \cap N_2) + \dots + v(S \cap N_p).$$

But equality must hold here, by superadditivity (2). This completes the proof.

A coalition is said to be "inessential" if its characteristic-function value is no greater than required for superadditivity (2). Theorem 1(b) expresses the fact that in a convex game, decomposability is equivalent to inessentiality of the all-player coalition.

COROLLARY. A strictly convex game is indecomposable.

### 3. CORE GEOMETRY

A payoff vector  $a \in E^N$  is said to be feasible (for  $v$ ) if  $a(N) \leq v(N)$ .\* The core of  $v$  is defined as the set  $C$  of all feasible  $a \in E^N$  such that

$$(11) \quad a(S) \geq v(S), \quad \text{all } S \subseteq N.$$

The core is obviously a subset of the hyperplane  $H_N$  (see §1.1), and since the inequalities  $a_i \geq v(\{i\})$  are included in (11), the core is bounded. Thus,  $C$  is a compact convex polyhedron, possibly empty, of dimension at most  $n - 1$ .

By a "full-dimensional" core we shall mean one of dimension exactly  $n - 1$ .

In order to discuss the facial structure of  $C$ , we define  $C_S = C \cap H_S$  for  $0 \subset S \subseteq N$ ; note that  $C_N = C$ . It is convenient also to define  $C_0 = C$ . If none of the  $C_S$  are empty, we say that the configuration  $\{C_S\}$  is complete; if they all have the highest possible dimension we say that  $\{C_S\}$  is strictly complete. In the latter case, the core

---

\*In application, there may be also a lower bound  $\beta$ , such that only payoff vectors with  $\beta \leq a(N) \leq v(N)$  are truly "feasible". Only the upper bound, however, will be of any significance in the present work.




Fig. 1—Core configuration of a 3-person convex game

is full-dimensional and has exactly  $2^n - 2$  polyhedral faces of dimension  $n - 2$ .

Figure 1 shows a core configuration for  $n = 3$  that is complete but not strictly complete, since  $C_{\{2\}}$  is only a point. Figure 2 depicts a strictly complete core\* for  $n = 4$ , there being fourteen 2-dimensional faces. (The extra lines and points, included for perspective, indicate the intersections of the  $H_S$  for  $s = 3$  with the simplex  $A$  in  $H_N$  defined by the  $H_S$  for  $s = 1$ .) We see that parallel faces, on opposite sides of the core, correspond to complementary sets of players.\*\*

Note that completeness necessarily entails a "large" core, which touches all  $n$  faces of the simplex  $A$ .

### 3.1. Regular Core Configurations

Although the hyperplanes  $H_S$  have the same slopes in every game  $v$ , their positions in  $E^N$  relative to one another


---

\*Note that if we are given only the point set  $C$ , we cannot in general identify all of the sets  $C_S$ , or even tell whether any are empty. In fact, it is not difficult to show that  $C$  uniquely determines  $v$ , and hence  $\{C_S\}$ , only in the strictly complete case. Thus, it makes sense to speak of a "strictly complete core," but not a "complete core."


\*\*In [Sch] a game with a complete core configuration is called an exact game.

Key to vertices and faces:

Front:



Back:



Characteristic function:

$$v(S) = \left( \sum_{i \in S} i \right)^2$$




Fig. 2—Core of a four-person convex game



are not predetermined, and even in the case of a strictly complete core there are many ways (if  $n > 3$ ) in which the faces  $C_S$  can fit together. But one "natural" arrangement stands out from the rest; it is the one that occurs, for example, when all faces  $C_S$  are equidistant from a common center--i.e., tangent to the inscribed  $(n-2)$ -sphere.\* In this arrangement, it can be shown that two faces touch if and only if their corresponding coalitions are comparable, i.e.,

$$(12) \quad C_S \cap C_T \neq \emptyset \quad \text{if and only if} \quad S \subseteq T \quad \text{or} \quad T \subseteq S.$$

We shall call this property strict regularity. It obviously implies completeness (take  $S = T$ ); we shall see presently that it implies strict completeness. The configuration of Fig. 2 is strictly regular, but not that of Fig. 1, since  $C_{12}$  and  $C_{23}$  meet.

We shall work primarily with a weaker form of (12). A core configuration  $\{C_S\}$  will be called regular if  $C_N \neq \emptyset$  and

---

\*Such a game is  $v(S) \equiv s - \sqrt{s(n-s)/(n-1)}$ .

$$(13) \quad C_S \cap C_T \subseteq C_{S \cup T} \cap C_{S \cap T}, \quad \text{all } S, T \subseteq N.$$

We see at once that strict regularity implies regularity, as the terminology would suggest. Condition (13) is equivalent to the statement that for each  $a \in E^N$ , the family of sets

$$(14) \quad g_a = \{S : a \in C_S\}$$

is closed under union and intersection.

The formal similarity between (13) and (3) is not accidental; in due course we shall show that convexity of a game is equivalent to regularity of its core configuration. First we shall develop some geometrical consequences of (13).

### 3.2. Faces

THEOREM 2. In a regular core configuration

$\{C_S\}$  we have

$$(15) \quad C_{S_1} \cap C_{S_2} \cap \dots \cap C_{S_m} \neq \emptyset$$

for any increasing sequence  $S_1 \subset S_2 \subset \dots \subset S_m$ .

In particular (take  $m = 1$ ), regularity implies completeness.

The proof proceeds with the aid of two lemmas concerning regular cores. Let us write  $S \subset \subset T$  to mean that  $S \subset T$  and  $t - s \geq 2$ .

LEMMA 1. If  $S \subset \subset T$  and  $a \in C_S \cap C_T$ , then there exists  $Q$  and  $b$  such that  $S \subset Q \subset T$  and  $b \in C_S \cap C_Q \cap C_T$ . Moreover, we can require that  $b_i = a_i$ , all  $i \in S$ , and that  $j \in Q$ ,  $k \notin Q$  for any two preassigned elements  $j, k$  of  $T - S$ .

Proof. Fix  $j, k \in T - S$ . Define  $b \in C$  by  $b_j = a_j - \rho$ ,  $b_k = a_k + \rho$ , and  $b_i = a_i$  for  $i \neq j, k$ , giving  $\rho$  the largest possible value compatible with  $b \in C$ ; this can be done because  $C$  is compact and contains  $a$ . Clearly,  $b(S) = a(S)$ ,  $b(T) = a(T)$ ; hence  $b \in C_S \cap C_T$ . Since a larger  $\rho$  would have taken  $b$  out of the core entirely, there must be a set  $R$ , containing  $j$  and not containing  $k$ , such that  $b \in C_R$ . Let  $Q = (S \cup R) \cap T$ . Since  $\mathcal{S}_b$  (see (14)) is closed under " $\cup$ " and " $\cap$ ", we have  $b \in C_Q$ . But  $Q$  has all the required properties:  $S \subset Q \subset T$ ,  $j \in Q$ , and  $k \notin Q$ ; so the lemma is proved.

LEMMA 2. Let  $S \subset \subset N$  and  $a \in C_S$ . Then,  
for any  $j \in N - S$  there exists  $b \in C_S \cap C_{S \cup \{j\}}$   
such that  $b_i = a_i$ , all  $i \in S$ .

Proof. Set  $T = N$  and use Lemma 1 to find  $Q, b$  such  
that  $S \cup \{j\} \subseteq Q \subset N$  and  $b \in C_S \cap C_Q$ , the latter agreeing  
with  $a$  on  $S$ . If  $Q = S \cup \{j\}$  we are through; if not, set  
 $T = Q$  and repeat the argument.

Proof of Theorem 2. We may assume that the sequence  
 $\{S_k\}$  is of maximum length, i.e., that  $m = n + 1$ . Then we  
have  $C_{S_1} = C_0$ , which is nonempty by definition. Take  
 $a_{(1)} \in C_{S_1}$ , and for  $k = 1, \dots, n - 1$ , apply Lemma 2 to  
find a point  $a_{(k+1)} \in C_{S_k} \cap C_{S_{k+1}}$ , agreeing with  $a_{(k)}$  on  
 $S_k$ . The point  $a_{(n)}$  will be in all of the  $C_{S_k}$ ,  $1 \leq k \leq n$ ,  
and it will also be in  $C_{S_{n+1}} = C_N = C$ . Hence (15) follows.

### 3.3. Vertices

Let  $\omega$  represent a simple ordering of the players.  
Specifically, let  $\omega$  be one of the  $n!$  functions that map  
 $N$  onto  $\{1, 2, \dots, n\}$ . Define

$$(16) \quad S_{\omega, k} = \{i \in N : \omega(i) \leq k\}, \quad k = 0, 1, \dots, n;$$

these are the "initial segments" of the ordering. Thus,  $S_{\omega,0} = 0$  and  $S_{\omega,n} = N$ . Consider the equations  $a(S_{\omega,k}) = v(S_{\omega,k})$ ,  $k = 1, 2, \dots, n$ . Linear independence is assured, and we can proceed by direct elimination to obtain the coordinates of the intersection of the hyperplanes  $H_{S_{\omega,k}}$ , namely

$$(17) \quad a_i^{\omega} = v(S_{\omega,\omega(i)}) - v(S_{\omega,\omega(i)-1}), \quad \text{all } i \in N.$$

In this way, each ordering  $\omega$  defines a payoff vector  $a^{\omega}$ ; there is, of course, no assurance that the  $a^{\omega}$  are all distinct.

THEOREM 3.\* The vertices of the core in a regular configuration are precisely the points  $a^{\omega}$ .

Proof. We have, for any  $\omega$ ,

$$C_{S_{\omega,1}} \cap \dots \cap C_{S_{\omega,n}} \subseteq H_{S_{\omega,1}} \cap \dots \cap H_{S_{\omega,n}}.$$

---

\*Theorem 22 of [E] gives a similar result for "polymatroids", which are similar in many respects to cores of convex games. Thus, Theorem 8 of [E] may be compared with (11), Theorem 42 with (13), etc.

The left-hand side is not empty, by Theorem 2; it therefore consists of the single point  $a^w$ . Hence  $a^w$  is in the core. If it were not a vertex (i.e., an extreme point), then for some nonzero vector  $d$  we would have both  $a^w \pm d \in C$ . But this is impossible, since at least one of the hyperplanes  $H_{S_{w,k}}$  that meet at and determine  $a^w$  must pass between the two points  $a^w \pm d$ , excluding one of them from the core. Hence  $a^w$  is a vertex of  $C$ , for each  $w$ .

It remains to show that  $C$  has no other vertices. Let  $a$  be any vertex of  $C$  and let  $0 = S_1 \subset S_2 \subset \dots \subset S_m = N$  be a "longest" increasing sequence of members of  $\mathcal{S}_a$  (see (14)), i.e., one that maximizes  $m$ . If there are no gaps in this sequence, i.e., if  $m = n + 1$ , then  $a$  is of the form  $a^w$ , and we are through. Suppose therefore that a gap exists, so that  $S_k \subset \subset S_{k+1}$  for some  $k$ , and let  $i, j \in S_{k+1} - S_k$ ,  $i \nmid j$ . Since  $a$  is a vertex, it is not only a solution of all the equations  $a(S) = v(S)$ ,  $S \in \mathcal{S}_a$ , it is their unique solution. But in order to determine the coordinates  $a_i$  and  $a_j$ , and not just their sum  $a_i + a_j$ , there must be an equation that separates them, i.e., an  $R \in \mathcal{S}_a$  that contains precisely one of  $i, j$ . Since  $\mathcal{S}_a$  is closed under " $\cup$ " and " $\cap$ ", we have  $Q \in \mathcal{S}_a$ , where  $Q = (S_k \cup R) \cap S_{k+1}$ . But  $S_k \subset Q \subset S_{k+1}$ , contradicting the maximality of  $m$ . This completes the proof.

### 3.4. Strictly Regular Cores

This section is mainly descriptive, and formal proofs are omitted. Strict regularity was defined by (12) above, but the one-sided condition

$$(18) \quad C_S \cap C_T \neq \emptyset \text{ implies } S \subseteq T \text{ or } T \subseteq S,$$

together with  $C \neq \emptyset$ , would have sufficed. To see this, note that (18) with  $C \neq \emptyset$  implies regularity, which in turn implies the converse of (18) by way of Theorem 2 with  $m = 2$ .

Strict regularity implies strict completeness, and hence is a property of the point set  $C$ . A strictly regular core has  $n!$  distinct extreme points  $a^u$ . The core in a regular configuration that is not strictly regular, on the other hand, always has fewer than  $n!$  extreme points, although it may still be strictly complete, i.e., have the maximum number of  $(n-2)$ -dimensional faces.

The facial arrangement of a strictly regular core is completely determined in all dimensions. For example, an  $(n-2)$ -dimensional face  $C_S$  is bounded by exactly  $2^S + 2^{n-S} - 4$   $(n-3)$ -dimensional faces of the form  $C_S \cap C_T$  with  $\emptyset \subset T \subset S$  or  $S \subset T \subset N$ . The automorphisms of this highly symmetric

combinatorial structure are generated by the permutations of  $N$ , together with complementation, the latter automorphism mapping each vertex  $a^{\omega}$  into its antipode  $a^{\omega'}$ , where  $\omega'$  denotes the reversal of  $\omega$ . This is illustrated for  $n = 4$  in Fig. 2.

We can now distinguish several classes of games, according to the kinds of cores they possess:

Table 2

nonempty cores	full-dimensional cores
complete core configurations	strictly complete cores
regular core configurations	strictly regular cores

Each class includes those listed below it and to the right. For  $n > 3$ , the inclusions are all strict. If a game is regarded as a point in  $E^{n-\{0\}}$  (see §2.2), then each of the six classes describes a convex cone; and those on the left [right] are the closures [interiors] of those on the right [left]. In particular, every regular core is a limit of strictly regular cores--indeed it can be approximated by both increasing and decreasing nested sequences of strictly regular cores.



#### 4. SOLUTIONS OF CONVEX GAMES

Many different "solution concepts" have been proposed for  $n$ -person games in characteristic-function form. We shall be concerned here with three of them: the core, already defined; the (Shapley) value (§4.2); and the (von Neumann-Morgenstern) solutions, which we call stable sets (§4.3). We shall see that the three concepts are intimately related when the game is convex.

##### 4.1. The Core

The core of a game, already defined at (11), may be interpreted as the set of "socially stable" outcomes, in that no coalition can improve upon any of them. In a game with an empty core, at least one set of players must fail to realize its full potential, no matter how the winnings are divided.

First we require

THEOREM 4. The core of a convex game is not empty.

Proof. It suffices to show that  $a^{(w)}$  is in the core, for some  $w$  (see §3.3). Let  $T \subset N$  and let  $j$  be the " $w$ -earliest" element of  $N - T$ , so that all the  $w$ -predecessors of  $j$  are in  $T$ , but not  $j$  itself. Then we have at once

$$T \cup S_{\omega, \omega(j)} = T \cup \{j\},$$

$$T \cap S_{\omega, \omega(j)} = S_{\omega, \omega(j)-1},$$

using the notation of (16). Hence, by convexity of  $v$ ,

$$v(T) + v(S_{\omega, \omega(j)}) \leq v(T \cup \{j\}) + v(S_{\omega, \omega(j)-1})$$

or, by (17),

$$a_j^{\omega} \leq v(T \cup \{j\}) - v(T).$$

Hence,

$$a^{\omega}(T) - v(T) \geq a^{\omega}(T \cup \{j\}) - v(T \cup \{j\}).$$

By induction,

$$a^{\omega}(T) - v(T) \geq a^{\omega}(N) - v(N) = 0.$$

Since  $T$  was arbitrary,  $a^{\omega}$  is in the core.

THEOREM 5. A game is convex if and only if its core configuration is regular.

Proof. Suppose we have regularity (13). Let  $S, T \subseteq N$ . Since  $S \cup T \supseteq S \cap T$ , we can find  $a \in C_{S \cup T} \cap C_{S \cap T}$ , by Theorem 2. Then we have

$$v(S \cup T) + v(S \cap T) = a(S \cup T) + a(S \cap T) = a(S) + a(T) \geq v(S) + v(T).$$

Hence  $v$  is convex.

Conversely, suppose that  $v$  is convex. Then  $C \neq \emptyset$  by Theorem 4 and it remains only to establish (13). Suppose  $a$  is in  $C_S \cap C_T$ . Then we have

$$v(S \cup T) + v(S \cap T) \geq v(S) + v(T) = a(S) + a(T) = a(S \cup T) + a(S \cap T).$$

But we also have  $a(S \cup T) \geq v(S \cup T)$  and  $a(S \cap T) \geq v(S \cap T)$ , because  $a$  is in the core. Hence equality prevails everywhere, and  $a \in C_{S \cup T} \cap C_{S \cap T}$ , as required.

COROLLARY. A game is strictly convex if and only if its core is strictly regular.

We omit the simple proof.

THEOREM 6.\* (a) The core of an indecomposable convex game is full-dimensional. (b) The core of a decomposable convex game is the cartesian product of the cores of the components of any decomposition. Its dimension is  $n - p$ , where  $p$  is the number of components in the finest decomposition.

Proof. (a) If  $C$  is less than full-dimensional, then  $C = C_S$  for some  $0 \subset S \subset N$ . We have  $C_{N-S} \neq 0$ , by completeness. Let  $a \in C_{N-S} \subseteq C_S$ . Then

$$v(S) + v(N-S) = a(S) + a(N-S) = a(N) = v(N),$$

and decomposability follows from (b) of Theorem 1.

(b) Let  $v$  be additive across the partition  $\{N_i\}$ , and first let  $a$  be in the cartesian product of the component cores. Then for all  $i$  and  $S$  we have

$$a(N_i \cap S) \geq v(N_i \cap S) \quad \text{and} \quad a(N_i) = v(N_i).$$

---

\*Based on [M-P-S1]. Note that the "cartesian product" property does not depend on the convexity of the game.

Applying (8) we obtain  $a(S) \geq v(S)$  and  $a(N) = v(N)$ , showing that  $a$  is in the core. Next let  $a$  be a payoff vector that is not in the cartesian product of the component cores. Then for some  $i$ , either

$$a(S) < v(S) \quad \text{for some} \quad S \subseteq N_i, \quad \text{or} \quad a(N_i) > v(N_i).$$

In the first case,  $a \notin C$ . In the second case, we have

$$a(N) - a(N - N_i) = a(N_i) > v(N_i) = v(N) - v(N - N_i),$$

using (9). Hence either  $a(N) > v(N)$  or  $a(N - N_i) < v(N - N_i)$ , so again  $a \notin C$ .

Finally, to determine the dimension of  $C$  let  $\{N_1, \dots, N_p\}$  yield the finest decomposition. Then the components are all indecomposable and have full-dimensional cores, by part (a), so that their cartesian product has dimension

$$\sum_{i=1}^p (n_i - 1) = n - p.$$

This completes the proof.

#### 4.2. The Value

The value of a game  $v$  is the payoff vector  $\varphi \in E^N$  defined by

$$(19) \quad \varphi_i = \sum_{\substack{S \subseteq N \\ S \ni i}} \frac{(s-1)!(n-s)!}{n!} [v(S) - v(S-\{i\})], \quad \text{all } i \in N.$$

We have  $\varphi(N) = v(N)$  and, for superadditive games,  $\varphi_i \geq v(\{i\})$  for all  $i \in N$ .<sup>\*</sup> The quantity  $\varphi_i$  may be interpreted as the "equity value" associated with the position of the  $i$ -th player in the game.

THEOREM 7. The value of a convex game is an element of the core.

Proof. It is well known (see [S3]) that

$$(20) \quad \varphi = \frac{1}{n!} \sum_{\omega \in \Omega} a^{\omega},$$

where  $\Omega$  is the set of all orderings of  $N$ . Hence, by Theorems 3 and 5, the value belongs to the core--indeed, it is in the relative interior since it is a center of gravity of the vertices.

---

<sup>\*</sup>See [S3]. It is also shown in [S3] that the value of a decomposable game is the cartesian product of the values of its components.

For completeness, we sketch a proof of (20). Fix  $i$ . Given  $S \ni i$ , we ask: for how many orderings  $\omega$  do we have  $S_{\omega, \omega(i)} = S$ ? It is clearly both necessary and sufficient that  $\omega$  be such that  $\omega(j) < \omega(i)$  for all  $j \in S - \{i\}$  and  $\omega(j) > \omega(i)$  for all  $j \in N - S$ . Thus the number of orderings is  $(s - 1)!(n - s)!$ . The result now follows easily from (19) and (17).

#### 4.3. Stable Sets

A payoff vector  $b$  is said to be dominated by a payoff vector  $a$  if there is a nonempty  $S \in \mathcal{N}$  such that

$$(21) \quad a(S) \leq v(S), \quad \text{and} \quad a_i > b_i \quad \text{all } i \in S.$$

A set  $V$  of feasible\* payoff vectors is said to be stable if every feasible\* payoff vector is either a member of  $V$  or dominated by a member of  $V$ , but not both. A stable set may be interpreted as a "standard of behavior," i.e., a

---

\*The classical definition [VN-M] replaces "feasible" in these two places by "feasible and individually rational", where  $a \in E^N$  is individually rational if and only if  $a_i \geq v(\{i\})$ , all  $i \in N$ . This distinction (discussed at length in [S1]--see also [L-R] p. 215ff) is not important here, since it can be shown that the core is stable in the one sense if and only if it is stable in the other. (See e.g., Theorems 12 and 13 of [G2].)

set of "conventional" outcomes which are always given a chance to dominate any proposal that might be put forward during pre-play negotiations.

It is easily verified that every stable set contains the core. Since no stable set can properly include another, it follows that if the core is stable then it is the only stable set.\*

THEOREM 8. The core of a convex game is stable (i.e., is the unique Neumann-Morgenstern solution).

Proof. Let  $\{C_S\}$  be regular, and take any feasible  $b$  not in  $C$ . We shall show that  $b$  is dominated by an element of  $C$ . Define  $g(0) = 0$  and

$$g(S) = \frac{v(S) - b(S)}{s}, \quad \text{all } 0 \subset S \subseteq N,$$

and let  $g$  attain its maximum,  $g^*$ , at  $S = S^*$ . Since  $b \notin C$  but is feasible we see that  $g^* > 0$  and hence  $S^* \neq 0$ . By completeness,  $C_{S^*} \neq 0$ . Let  $c \in C_{S^*}$ , and define  $a \in E^N$  by:

---

\*The early conjectures [VN-M, S2] that stable sets exist for all superadditive games and that the core is the intersection of all stable sets have been disproved; see [L1, L2].



$$a_i = \begin{cases} b_i + g^*, & i \in S^*, \\ c_i, & i \in N - S^*. \end{cases}$$

Since  $a(S^*) = b(S^*) + s^*g^* = v(S^*)$ , condition (21) is satisfied, and we see that  $a$  dominates  $b$ . Moreover, since  $a(N) = v(S^*) + c(N-S^*) = c(N) = v(N)$ , we see that  $a$  is feasible. It remains to show that  $a$  satisfies the core inequalities (11).

Let  $T \subseteq N$  be arbitrary, and break up  $T$  into  $Q = T \cap S^*$  and  $R = T - S^*$ . Then we have, in easy steps,

$$\begin{aligned} a(T) &= a(Q) + a(R) \\ &= b(Q) + qg^* + c(R) \\ &\geq b(Q) + qg(Q) + c(T \cup S^*) - c(S^*) \\ &\geq v(Q) + v(T \cup S^*) - v(S^*) \\ &\geq v(T). \end{aligned}$$

Hence  $a$  is in the core.

#### 4.4. Remarks

1. An example of a stable core that does not come from a regular, or even complete, configuration is shown in Fig. 3. The core is a perfect cube, with faces  $\{C_s : s = 2\}$  and vertices at  $(1, 1, 1, 0)$ ,  $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{3}{2})$ , etc. The sets  $C_s$ ,  $s = 3$ , are empty.

2. We have obtained a complete theory of core stability in symmetric games, i.e., games of the form

$$v(S) \equiv f(s);$$

this will be presented elsewhere.

3. Theorem 8 implies that a "positive fraction" of all games are solvable--more precisely, that the set of solvable games on  $\mathcal{N}$  includes a full-dimensional (i.e.,  $(2^n-1)$ -dimensional) cone in  $E^{\mathcal{N}-\{0\}}$ . This was first pointed out by Gillies,\* using the cone generated by the games in 0,1-normalization that satisfy  $v(S) < 1/n$  for  $2 \leq s < n$ . This cone includes some but not all convex games, as well as some that are not convex.

---

\*[G1, G2]; see also [B].




Fig.3—Example of a stable, nonregular core

REFERENCES

- [A-S] Aumann, R. J., and L. S. Shapley, Values of Non-Atomic Games I, II, III, IV, RM-5468, RM-5842, RM-6216, RM-6260, The Rand Corporation, Santa Monica, California, 1968, 1969, 1970.
- [B] Bondareva, O. N., "Some Applications of the Methods of Linear Programming to the Theory of Cooperative Games," (Russian), Problemy Kibernetiki 10 (1963); 119-139; esp. §4.
- [Ch] Choquet, Gustave, "Theory of Capacities," Annals de l'Institut Fourier 5 (1955); 131-295.
- [C-R] Crapo, H. H., and G.-C. Rota, Combinatorial Geometries, Massachusetts Institute of Technology, Cambridge, Massachusetts, 1968.
- [E] Edmonds, Jack, "Submodular Functions, Matroids, and Certain Polyhedra," Combinatorial Structures and Their Applications (proceedings of a conference at the University of Calgary, June 1969), Gordon and Breach, New York, 1970; 69-87.
- [E-R] Edmonds, J. and G.-C. Rota, "Submodular Set Functions" (abstract, Waterloo Combinatorics Conference), University of Waterloo, Waterloo, Ontario, 1966.
- [G1] Gillies, D. B., Some Theorems on n-Person Games (dissertation), Department of Mathematics, Princeton University, 1953.
- [G2] \_\_\_\_\_ "Solutions to General Non-Zero-Sum Games," Annals of Mathematics Study 40 (1959); 47-85; esp. 77-81.
- [L1] Lucas, W. F., "A Counterexample in Game Theory," Management Science 13 (1967); 766-767.
- [L2] \_\_\_\_\_ "A Game with no Solution," Bull. Am. Math. Soc. 74 (1968); 237-239.

- [L-R] Luce, R. D., and H. Raiffa, Games and Decisions, Wiley and Sons, New York (1957).
- [M] Meyer, A., Probabilities and Potentials, Blaisdell, Waltham, Massachusetts, 1966; esp. 40ff.
- [M-P-S1] Maschler, M., B. Peleg, and L. S. Shapley, The Kernel and Bargaining Set for Convex Games, RM-5372, The Rand Corporation, Santa Monica, California, 1967; esp. 10f.
- [M-P-S2] \_\_\_\_\_ "The Kernel and Bargaining Set for Convex Games," (to appear).
- [P1] Peleg, B., Composition of General Sum Games, RM-74, Econometric Research Program, Princeton University, 1965.
- [P2] \_\_\_\_\_ Composition of Kernels of Characteristic Function Games, RM-15, Department of Mathematics, The Hebrew University of Jerusalem, 1965.
- [R] Rosenmüller, Joachim, Some Properties of Convex Set Functions, (duplicated), Mathematics Institut der Universität Erlanger-Nürnberg, 852 Erlangen, Germany, 1970.
- [Sch] Schmeidler, David, Cores of Exact Games, I, CP-329, Center for Research in Management Science, University of California, Berkeley, 1971.
- [S1] Shapley, L. S., Notes on the n-Person Game III: Some Variants of the Von Neumann-Morgenstern Definition of Solution, RM-670, The Rand Corporation, Santa Monica, California, 1951.
- [S2] \_\_\_\_\_ "Open Questions," Theory of n-Person Games (report of an informal conference), Department of Mathematics, Princeton University, 1953; 15.
- [S3] \_\_\_\_\_ "A Value for n-Person Games," Annals of Mathematics Study 28 (1953); 307-317.

- [S4]        Notes on n-Person Games VII: Cores of  
Convex Games, RM-4571, The Rand Corporation,  
Santa Monica, California, 1965.
- [VN-M]    Von Neumann, J., and O. Morgenstern, Theory of Games  
and Economic Behavior, Princeton University Press,  
1944.
- [Wh]       Whitney, Hassler, "The Abstract Properties of Linear  
Dependence," Am. J. Math. 57 (1935); 509-533;  
esp. 511.