

Erratum to: Cork oak (*Quercus suber* L.) seedlings acclimate to elevated CO₂ and water stress: photosynthesis, growth, wood anatomy and hydraulic conductivity

M. Vaz · H. Cochard · L. Gazarini ·
J. Graça · M. M. Chaves · J. S. Pereira

Published online: 9 March 2012
© Springer-Verlag 2012

Erratum to: Trees
DOI 10.1007/s00468-012-0691-x

In the original version of article, Fig. 4 was published incorrectly. The correct Fig. 4 is produced in the following page.

The online version of the original article can be found under
doi:[10.1007/s00468-012-0691-x](https://doi.org/10.1007/s00468-012-0691-x).

M. Vaz (✉) · L. Gazarini
Departamento de Biologia, Instituto de Ciências Agrárias
e Ambientais Mediterrânicas (ICAAM), Universidade de Évora,
Apartado 94, 7002-554 Évora, Portugal
e-mail: mvaz@uevora.pt

H. Cochard
INRA, UMR 547 PIAF, 63100 Clermont-Ferrand, France

H. Cochard
Clermont Université, Université Blaise Pascal, UMR 547 PIAF,
BP 10448, 63000 Clermont-Ferrand, France

J. Graça · M. M. Chaves · J. S. Pereira
Instituto Superior de Agronomia, Universidade Técnica
de Lisboa, Tapada da Ajuda, 1349-017 Lisbon, Portugal

M. M. Chaves
Instituto Tecnologia Química e Biológica,
Universidade Nova de Lisboa (ITQB-UNL),
Av. da República, 2780-157 Oeiras, Portugal




Fig. 4 Predawn leaf water potential (Ψ_{IPD} , MPa); midday leaf water potential (Ψ_{IMD} , MPa); Photosynthetic carbon assimilation (A , $\mu\text{mol m}^{-2} \text{s}^{-1}$); stomatal conductance (g_s , $\text{mol m}^{-2} \text{s}^{-1}$), of plants grown at 350 and 700 ppm of CO_2 after a period of 6, 9 and 15 months in the greenhouses. Values are means of 10 plants \pm SE, data sharing the same letter are not significantly different ($p < 0.05$). WW well watered, WS water stress