

CORRECTION

# Correction: Conditioned Medium from Hypoxic Bone Marrow-Derived Mesenchymal Stem Cells Enhances Wound Healing in Mice

Lei Chen, Yingbin Xu, Jingling Zhao, Zhaoqiang Zhang, Ronghua Yang, Julin Xie, Xusheng Liu, Shaohai Qi

The authors would like to correct [Fig 6](#), as errors were introduced in the preparation of this figure for publication. In [Fig 6A](#), the hypoCM panel appears as a duplicate of the Vehicle Medium panel. The authors have provided a corrected version of [Fig 6](#) here.




 OPEN ACCESS

**Citation:** Chen L, Xu Y, Zhao J, Zhang Z, Yang R, Xie J, et al. (2015) Correction: Conditioned Medium from Hypoxic Bone Marrow-Derived Mesenchymal Stem Cells Enhances Wound Healing in Mice. PLoS ONE 10(12): e0145565. doi:10.1371/journal.pone.0145565

**Published:** December 18, 2015

**Copyright:** © 2015 Chen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Fig 6. IHC evaluation of wounded mouse skin.** Wound sections were evaluated on day 11 by staining with anti-Ki67 and anti-F4/80 antibodies. The numbers of Ki67+ proliferating cells (A, C) and recruited F4/80+ macrophages (B, D) in each of 4 randomly chosen high-power fields in the dermis were counted. Scale bar, 100  $\mu$ m (400 $\times$ ). Data are expressed as the mean $\pm$ the SEM; \* $p$ <0.05 compared with the vehicle control or the norCM group.

doi:10.1371/journal.pone.0145565.g001

The authors confirm that these changes do not alter their findings. The authors have provided the underlying images as Supporting Information.

### Supporting Information

**S1 Fig. Ki67 Vehicle Medium.**  
(TIF)

**S2 Fig. Ki67 norCM.**  
(TIF)

**S3 Fig. Ki67 hypoCM.**  
(TIF)

**S4 Fig. F4/80 Vehicle Medium.**  
(TIF)

**S5 Fig. F4/80 norCM.**  
(TIF)

**S6 Fig. F4/80 hypoCM.**  
(TIF)

## Reference

1. Chen L, Xu Y, Zhao J, Zhang Z, Yang R, Xie J, et al. (2014) Conditioned Medium from Hypoxic Bone Marrow-Derived Mesenchymal Stem Cells Enhances Wound Healing in Mice. PLoS ONE 9(4): e96161. doi: [10.1371/journal.pone.0096161](https://doi.org/10.1371/journal.pone.0096161) PMID: [24781370](https://pubmed.ncbi.nlm.nih.gov/24781370/)