

# SCIENTIFIC REPORTS


**OPEN**

## **Corrigendum: Gene expression differences in *Longissimus* muscle of Nelore steers genetically divergent for residual feed intake**


Polyana C. Tizioto, Luiz L. Coutinho, Priscila S. N. Oliveira, Aline S. M. Cesar, Wellison J. S. Diniz, Andressa O. Lima, Marina I. Rocha, Jared E. Decker, Robert D. Schnabel, Gerson B. Mourão, Rymer R. Tullio, Adhemar Zerlotini, Jeremy F. Taylor & Luciana C. A. Regitano

*Scientific Reports* 6:39493; doi: 10.1038/srep39493; published online 22 December 2016; updated on 05 April 2017

This Article contains an error in the order of the Figures, where Figures 2 and 3 were inverted. The correct Figures 2 and 3 appear below:



**Figure 2. Network integration of the EGR1 gene and other differentially expressed genes.** Genes presented as black circles were differentially expressed (DE) between the efficient (LRFI) and inefficient (HRFI) groups. Genes presented in grey interact with the DE genes. Arrows presented in pink, green, blue and red represent co-expression relationships, genetic interactions, co-localizations and physical interactions, respectively.



**Figure 3. Venn diagram showing overlaps between differentially expressed genes found for residual feed intake (RFI), average daily gain (ADG) and dry matter intake (DMI).**



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017