



Correction to: Nest predation and the influence of habitat structure on nest predation of Wood Warbler *Phylloscopus sibilatrix*, a ground-nesting forest passerine

Paul E. Bellamy¹ · Malcolm D. Burgess¹ · John W. Mallord¹ · Andrew Cristinacce¹ · Christopher J. Orsman¹ · Tony Davis² · Philip V. Grice³ · Elisabeth C. Charman^{1,4}

Published online: 5 May 2018
© Dt. Ornithologen-Gesellschaft e.V. 2018

Correction to: Journal of Ornithology (2018) 159:493–506
<https://doi.org/10.1007/s10336-017-1527-7>

In the original publication of the article, there is a misalignment of the last two columns in Table 2. The correct Table 2 is provided below.

The original article can be found online at <https://doi.org/10.1007/s10336-017-1527-7>.

✉ Paul E. Bellamy
paul.bellamy@rspb.org.uk

¹ RSPB Centre for Conservation Science, The Lodge, Sandy, Bedfordshire SG19 2DL, UK

² 17 Orchards Way, West End, Southampton SO30 3FB, UK

³ Natural England, Mail Hub, County Hall, Spetchley Road, Worcester WR5 2NP, UK

⁴ Present Address: RSPB, 1 Sirius House, Amethyst Road, Newcastle Business Park, Newcastle-upon-Tyne NE4 7YL, UK

Table 2 The identified predators from 95 predation events out of 144 Wood Warbler nests by study site monitored by nest camera

	Dartmoor	New forest	Mid-Wales	Total
Birds				
Jay <i>Garrulus glandarius</i>	7	5	18	30
Buzzard <i>Buteo buteo</i>	8	1	4	13
Sparrowhawk <i>Accipiter nisus</i>	4	1	2	7
Tawny owl <i>Strix aluco</i>	0	2	0	2
Great spotted woodpecker <i>Dendrocopos major</i>	0	0	1 (Plus one partial ^a)	1
Mammals				
Badger <i>Meles meles</i>	2	4	1	7
Fox <i>Vulpes vulpes</i>	1	2	1	4
Weasel <i>Mustela nivalis</i>	2	0	0	2
Dog <i>Canis familiaris</i>	0	1	0	1
Grey squirrel <i>Sciurus carolinensis</i>	1	2	0	3
Other rodents	3	2	0	5
Other				
Adder <i>Vipera berus</i>	0	2	0	2
Unknown	5	7	4	16

^aPartial predation is defined as a predator taking some nest contents with the remainder surviving and the nest attempt continuing