

CORRECTION

Open Access



Correction to: NSABP FB-7: a phase II randomized neoadjuvant trial with paclitaxel + trastuzumab and/or neratinib followed by chemotherapy and postoperative trastuzumab in HER2+ breast cancer

Samuel A. Jacobs^{1*}, André Robidoux^{1,2}, Jame Abraham^{1,3}, José Manuel Pérez-García^{4,5}, Nicla La Verde^{6,7}, James M. Orcutt^{1,8}, Marina E. Cazzaniga^{5,9}, Fanny Piette¹⁰, Silvia Antolín¹¹, Elena Aguirre⁵, Javier Cortes^{4,5}, Antonio Llombart-Cussac⁵, Serena Di Cosimo^{5,12}, Rim S. Kim¹, Huichen Feng¹, Corey Lipchik¹, Peter C. Lucas^{1,13}, Ashok Srinivasan¹, Ying Wang¹, Nan Song¹, Patrick G. Gavin¹, April D. Balousek¹, Soonmyung Paik^{1,14}, Carmen J. Allegra¹⁵, Norman Wolmark^{1,16} and Katherine L. Pogue-Geile¹

Correction to: Breast Cancer Res

<https://doi.org/10.1186/s13058-019-1196-y>

After the publication [1] the authors report corrections for Table 3, for the headers of columns 5 and 7, which should read as follows: Column 5 Header: *P*-value for pCR, T vs Other Arms in HR⁺, and Column 7 Header: *P*-value for pCR, T vs Other Arms in HR⁻.

The correct version of the table can be found below:

Author details

¹NSABP Foundation, Inc., Nova Tower 2, Two Allegheny Center – Ste 1200, Pittsburgh, PA 15212, USA. ²Centre hospitalier de l'université de Montréal, Montréal, QC, Canada. ³Cleveland Clinic, Cleveland, OH, USA. ⁴QuironSalud Group, IOB Institute of Oncology, Barcelona, Madrid, Spain. ⁵Medica Scientia Innovation Research (MedSIR), Barcelona, Spain. ⁶Present address: ASST Fatebenefratelli Sacco - PO Luigi Sacco, Milan, Italy. ⁷ASST Fatebenefratelli Sacco - PO Fatebenefratelli, Milan, Italy. ⁸Roper St. Francis Healthcare, Charleston, SC, USA. ⁹Azienda Ospedaliera San Gerardo, Monza, Italy. ¹⁰International Drug Development Institute (IDDI), Louvain-la-Neuve, Belgium. ¹¹Hospital Universitario, Coruña, Spain. ¹²Fondazione IRCCS Istituto Nazionale di Tumori, Milan, Italy. ¹³Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA. ¹⁴Severance Biomedical Science Institute and Department of Medical Oncology, Yonsei University College of Medicine, Seoul, Republic of South Korea. ¹⁵Department of Medicine,

University of Florida Health, Gainesville, FL, USA. ¹⁶University of Pittsburgh, Pittsburgh, PA 15212, USA.

Published online: 22 January 2020

Reference

1. Jacobs, et al. *Breast Cancer Res.* 2019;21:133. <https://doi.org/10.1186/s13058-019-1196-y>.

The original article can be found online at <https://doi.org/10.1186/s13058-019-1196-y>

* Correspondence: samuel.jacobs@nsabp.org

¹NSABP Foundation, Inc., Nova Tower 2, Two Allegheny Center – Ste 1200, Pittsburgh, PA 15212, USA

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Table 3 NSABP FB-7: Pathologic complete response (pCR) (breast and nodes) by treatment arm and HR status

Treatment arms, <i>n</i> = 42 each arm	Total pCR	<i>P</i> -value for pCR T vs Other Arms in all patients ^a	pCR HR ⁺	<i>P</i> -value for pCR T vs Other Arms in HR ⁺ ^b	pCR HR ⁻	<i>P</i> -value for pCR T vs Other Arms in HR ⁻ ^b	<i>P</i> -value HR ⁺ vs HR ⁻
Arm 1 (T)	16/41 (39.0%)	NA	8/27 (29.6%)	NA	8/14 (57.1%)	NA	0.11
Arm 2 (N)	14/42 (33.3%)	0.63	8/29 (27.6%)	1	6/13 (46.2%)	0.71	0.30
Arm 3 (T plus N)	21/42 (50%)	0.22	7/23 (30.4%)	1	14/19 (73.7%)	0.46	0.01
All arms	51/126 (40%)		23/79 (29%)		28/46 (61%)		0.001

Abbreviations: T, trastuzumab; N, neratinib

^aCMH test stratified for the time period the patients were randomized to the study (prior/after the addition of Arm 3)

^bFisher's exact test