# CORRECTION



# **Open Access**

# Correction: The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women

Roberto Romero<sup>1,2,3\*</sup>, Sonia S Hassan<sup>1,4</sup>, Pawel Gajer<sup>5,6</sup>, Adi L Tarca<sup>1</sup>, Douglas W Fadrosh<sup>5</sup>, Lorraine Nikita<sup>1</sup>, Marisa Galuppi<sup>1,4</sup>, Ronald F Lamont<sup>1,7,8</sup>, Piya Chaemsaithong<sup>1</sup>, Jezid Miranda<sup>1</sup>, Tinnakorn Chaiworapongsa<sup>1,4</sup> and Jacques Ravel<sup>5,6\*</sup>

# Correction

We have identified an error in Additional file 1: Table S1 in this article [1], whereby the ethnicities were inverted for some of the subjects. An amended Additional file 1: Table S1 is published here, with the corrected ethnicities highlighted.

# **Additional file**

**Additional file 1: Table S1.** Taxonomic assignments, total 16S rRNA gene sequences per taxa, and metadata.

## Author details

<sup>1</sup>Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NIH, Detroit, MI, USA. <sup>2</sup>Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, MI, USA. <sup>3</sup>Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI, USA. <sup>4</sup>Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, MI, USA. <sup>5</sup>Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, MD, USA. <sup>6</sup>Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, MD, USA. <sup>7</sup>Department of Obstetrics and Gynaecology, University of Southern Denmark, Odense, Denmark. <sup>8</sup>Division of Surgery, University College, Northwick Park Institute for Medical Research Campus, London, UK.

Received: 3 April 2014 Accepted: 3 April 2014 Published: 15 April 2014

### \* Correspondence: romeror@mail.nih.gov; jravel@som.umaryland.edu <sup>1</sup>Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NIH, Bethesda, MD and, Detroit, MI, USA

Full list of author information is available at the end of the article



© 2014 Romero et al.; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

### Reference

 Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Nikita L, Galuppi M, Lamont RF, Chaemsaithong P, Miranda J, Chaiworapongsa T, Ravel J: The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome* 2014, 2:4.

### doi:10.1186/2049-2618-2-10

**Cite this article as:** Romero *et al.*: **Correction:** The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome* 2014 **2**:10.

<sup>&</sup>lt;sup>5</sup>Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, MD, USA