

CORRELATED X-RAY AND TEV EMISSION IN THE GAMMA-RAY BINARY LS I +61 303

H. ANDERHUB¹, L. A. ANTONELLI², P. ANTORANZ³, M. BACKES⁴, C. BAIXERAS⁵, S. BALESTRA³, J. A. BARRIO³, D. BASTIERI⁶, J. BECERRA GONZÁLEZ⁷, J. K. BECKER⁴, W. BEDNAREK⁸, K. BERGER⁸, E. BERNARDINI⁹, A. BILAND¹, O. BLANCH BIGAS¹⁴, R. K. BOCK^{10,6}, G. BONNOLI¹¹, P. BORDAS¹², D. BORLA TRIDON¹⁰, V. BOSCH-RAMON¹², D. BOSE³, I. BRAUN¹, T. BRETZ¹³, D. BRITZGER¹⁰, M. CAMARA³, E. CARMONA¹⁰, A. CAROSI², P. COLIN¹⁰, S. COMMICHAU¹, J. L. CONTRERAS³, J. CORTINA¹⁴, M. T. COSTADO^{7,15}, S. COVINO², F. DAZZI^{16,26}, A. DE ANGELIS¹⁶, E. DE CEA DEL POZO¹⁷, R. DE LOS REYES³, B. DE LOTTO¹⁶, M. DE MARIA¹⁶, F. DE SABATA¹⁶, C. DELGADO MENDEZ^{7,27}, A. DOMÍNGUEZ¹⁸, D. DOMINIS PRESTER¹⁹, D. DORNER¹, M. DORO⁶, D. ELSAESSER¹³, M. ERRANDO¹⁴, D. FERENC²⁰, E. FERNÁNDEZ¹⁴, R. FIRPO¹⁴, M. V. FONSECA³, L. FONT⁵, N. GALANTE¹⁰, R. J. GARCÍA LÓPEZ^{7,15}, M. GARCZARCYK¹⁴, M. GAUG⁷, N. GODINOVIC¹⁹, F. GOEBEL^{10,28}, D. HADASCH⁵, A. HERRERO^{7,15}, D. HILDEBRAND¹, D. HÖHNE-MÖNCH¹³, J. HOSE¹⁰, D. HRUPEC¹⁹, C. C. HSU¹⁰, T. JOGLER^{10,29}, S. KLEPSEK¹⁴, D. KRANICH¹, A. LA BARBERA², A. LAILLE²⁰, E. LEONARDO¹¹, E. LINDFORS²¹, S. LOMBARDI⁶, F. LONGO¹⁶, M. LÓPEZ⁶, E. LORENZ^{1,10}, P. MAJUMDAR⁹, G. MANEVA²², N. MANKUZHIYIL¹⁶, K. MANNHEIM¹³, L. MARASCHI², M. MARIOTTI⁶, M. MARTÍNEZ¹⁴, D. MAZIN¹⁴, M. MEUCCI¹¹, J. M. MIRANDA³, R. MIRZOYAN¹⁰, H. MIYAMOTO¹⁰, J. MOLDÓN¹², M. MOLES¹⁸, A. MORALEJO¹⁴, D. NIETO³, K. NILSSON²¹, J. NINKOVIC¹⁰, R. ORITO¹⁰, I. OYA³, R. PAOLETTI¹¹, J. M. PAREDES¹², M. PASANEN²¹, D. PASCOLI⁶, F. PAUSS¹, R. G. PEGNA¹¹, M. A. PEREZ-TORRES¹⁸, M. PERSIC^{16,23}, L. PERUZZO⁶, F. PRADA¹⁸, E. PRANDINI⁶, N. PUCHADES^{14,29}, I. PULJAK¹⁹, I. REICHARDT¹⁴, W. RHODE⁴, M. RIBÓ^{12,29}, J. RICO^{24,14}, M. RISSI¹, A. ROBERT⁵, S. RÜGAMER¹³, A. SAGGION⁶, T. Y. SAITO¹⁰, M. SALVATI², M. SÁNCHEZ-CONDE¹⁸, K. SATALECKA⁹, V. SCALZOTTO⁶, V. SCAPIN¹⁶, T. SCHWEIZER¹⁰, M. SHAYDUK¹⁰, S. N. SHORE²⁵, N. SIDRO¹⁴, A. SIERPOWSKA-BARTOSIK⁸, A. SILLANPÄÄ²¹, J. SITAREK^{10,8}, D. SOBCZYNSKA⁸, F. SPANIER¹³, S. SPIRO², A. STAMERRA¹¹, L. S. STARK¹, T. SURIC¹⁹, L. TAKALO²¹, F. TAVECCHIO², P. TEMNIKOV²², D. TESCARO¹⁴, M. TESHIMA¹⁰, D. F. TORRES^{24,17}, N. TURINI¹¹, H. VANKOV²², R. M. WAGNER¹⁰, V. ZABALZA¹², F. ZANDANEL¹⁸, R. ZANIN¹⁴, J. ZAPATERO⁵, A. FALCONE³⁰, L. VETERE³⁰, N. GEHRELS³¹, S. TRUSHKIN³², V. DHAWAN³³, P. REIG³⁴

¹ ETH Zurich, CH-8093 Switzerland

² INAF National Institute for Astrophysics, I-00136 Rome, Italy

³ Universidad Complutense, E-28040 Madrid, Spain

⁴ Technische Universität Dortmund, D-44221 Dortmund, Germany

⁵ Universitat Autònoma de Barcelona, E-08193 Bellaterra, Spain

⁶ Università di Padova and INFN, I-35131 Padova, Italy

⁷ Inst. de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain

⁸ University of Łódź, PL-90236 Lodz, Poland

⁹ Deutsches Elektronen-Synchrotron (DESY), D-15738 Zeuthen, Germany

¹⁰ Max-Planck-Institut für Physik, D-80805 München, Germany

¹¹ Università di Siena, and INFN Pisa, I-53100 Siena, Italy

¹² Universitat de Barcelona (ICC/IEEC), E-08028 Barcelona, Spain

¹³ Universität Würzburg, D-97074 Würzburg, Germany

¹⁴ IFAE, Edifici Cn., Campus UAB, E-08193 Bellaterra, Spain

¹⁵ Depto. de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain

¹⁶ Università di Udine, and INFN Trieste, I-33100 Udine, Italy

¹⁷ Institut de Ciències de l'Espai (IEEC-CSIC), E-08193 Bellaterra, Spain

¹⁸ Inst. de Astrofísica de Andalucía (CSIC), E-18080 Granada, Spain

¹⁹ Rudjer Boskovic Institute, Bijenicka 54, HR-10000 Zagreb, Croatia

²⁰ University of California, Davis, CA-95616-8677, USA

²¹ Tuorla Observatory, University of Turku, FI-21500 Piikkiö, Finland

²² Inst. for Nucl. Research and Nucl. Energy, BG-1784 Sofia, Bulgaria

²³ INAF/Osservatorio Astronomico and INFN, I-34143 Trieste, Italy

²⁴ ICREA, E-08010 Barcelona, Spain

²⁵ Università di Pisa, and INFN Pisa, I-56126 Pisa, Italy

²⁶ supported by INFN Padova

²⁷ now at: Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

²⁸ deceased

²⁹ Send offprint requests to T. Jogler jogler@mppmu.mpg.de, N. Puchades neus@ifae.es, M. Ribó mribo@am.ub.es

³⁰ Pennsylvania State University, PA-16802, USA

³¹ NASA/Goddard Space Flight Center, Greenbelt, MD-20771, USA

³² Special Astrophysical Observatory RAS, 369167, Russia

³³ National Radio Astronomy Observatory, Socorro, NM-87801, USA and

³⁴ University of Crete, Heraklion, G-71110, Greece

Draft version July 31, 2009

ABSTRACT

The discovery of TeV emitting X-ray binaries has triggered an intense effort to better understand the particle acceleration, absorption, and emission mechanisms in compact binary systems, which provide variable conditions along eccentric orbits. Despite this, the nature of some of these systems, and of the accelerated particles producing the TeV emission, is unclear. To answer some of these open questions we conducted a multiwavelength campaign of the TeV emitting X-ray binary LS I +61 303 including the MAGIC telescope, *XMM-Newton*, and *Swift* during 60% of an orbit in 2007 September. We detect a simultaneous outburst at X-ray

and TeV energies, with the peak at phase 0.62 and a similar shape at both wavelengths. A linear fit to the simultaneous X-ray/TeV pairs obtained during the outburst yields a correlation coefficient of $r = 0.97$, while a linear fit to all simultaneous pairs provides $r = 0.81$. The observed correlation and the lack of significant absorption towards the observer expected at TeV energies during the explored orbital phases indicate a simultaneity in the emission processes. Assuming that they are dominated by a single particle population, either hadronic or leptonic, the X-ray/TeV flux ratio favors leptonic models. This fact, together with the detected photon indexes, suggests that in LS I +61 303 the X-rays are the result of synchrotron radiation of the same VHE electrons that produce TeV emission as a result of inverse Compton scattering of stellar photons.

Subject headings: binaries: general — gamma rays: observations — stars: emission-line, Be — stars: individual (LS I +61 303) — X-rays: binaries — X-rays: individual (LS I +61 303)

1. INTRODUCTION

LS I +61 303 is one of the few X-ray binaries that have been detected at TeV energies (see e.g. Paredes 2008 for a recent review). It is a high mass X-ray binary system located at a distance of 2.0 ± 0.2 kpc (Frail & Hjellming 1991). The system contains a rapidly rotating early type B0 Ve star with a stable equatorial decretion disk and mass loss, and a compact object with a mass between 1 and $4 M_{\odot}$ orbiting it every ~ 26.5 d in an eccentric orbit (see Casares et al. 2005, Grundstrom et al. 2007, Aragona et al. 2009, and references therein). LS I +61 303 was classified as a microquasar based on structures detected from five to several tens of milliarcseconds with the EVN and MERLIN (Massi et al. 2004 and references therein), although analysis of later EVN and MERLIN datasets does not reveal the presence of such structures (Dhawan et al. 2006; Albert et al. 2008a). VLBA images with a resolution of one milliarcsecond (2 AU at the source distance) obtained during a full orbital cycle show an elongated structure that rotates as a function of the orbital phase (Dhawan et al. 2006). Later VLBA images show repeating structures at the same orbital phases, reinforcing the idea that the milliarcsecond structure depends on the orbital phase (Albert et al. 2008a). This may be consistent with a model based on the interaction between the relativistic wind of a young non-accreting pulsar and the wind/decretion disk of the stellar companion (Dubus 2006a), as occurs in PSR B1259–63 (but confining the pulsar wind may be problematic, see Bogovalov et al. 2008).

LS I +61 303 shows periodic non-thermal radio outbursts on average every $P_{\text{orb}} = 26.4960 \pm 0.0028$ d, with the peak of the radio emission shifting progressively from phase 0.45 to 0.95, using $T_0 = \text{JD } 2,443,366.775$, with a modulation period of 1667 ± 8 d (Gregory 2002). According to the most precise orbital parameters, the periastron takes place at phase 0.275 and the eccentricity of the orbit is 0.537 ± 0.034 (Aragona et al. 2009).

LS I +61 303 has been observed several times in X-rays (see Smith et al. 2009 and references therein). It generally displays quasi-periodic X-ray outbursts, with the maximum occurring between orbital phase 0.4 and 0.8 (Goldoni & Mereghetti 1995; Taylor et al. 1996; Paredes et al. 1997; Harrison et al. 2000; Esposito et al. 2007), although the lack of a sensitive long-term monitoring has prevented to search for a super-orbital modulation. The source also shows short-term variability on timescales of hours (Sidoli et al. 2006).

At very high energy (VHE) gamma rays, LS I +61 303 has been clearly detected several times both by MAGIC (Albert et al. 2006, 2008a, 2009) and VERITAS (Acciari et al. 2008, 2009). The source also displays TeV periodicity, with minima taking place near periastron, where only upper limits were found in MAGIC observations, and maxima occurring on av-

erage at phase 0.6–0.7, although the source has also shown a second peak at phase 0.84 in one of the cycles (Albert et al. 2009). There are indications of correlated X-ray/TeV emission, based on non-simultaneous data taken more than six hours apart (Albert et al. 2009), and one day apart (Albert et al. 2008a).

The lack of a systematic behavior from cycle to cycle at X-ray and TeV energies, and the occurrence of short-term variability in the X-ray flux, has hampered the definitive detection of an X-ray/TeV correlation from the comparison of non-simultaneous data. We therefore conducted a multiwavelength campaign in 2007 September, covering the epoch of maximum TeV emission. Here we report the first simultaneous TeV and X-ray observations of LS I +61 303 obtained with the MAGIC Cherenkov telescope and the *XMM-Newton* and *Swift* X-ray satellites that show correlated emission in both energy bands. We also briefly comment on radio and optical spectroscopic observations obtained during the campaign.

2. OBSERVATIONS AND DATA ANALYSIS

2.1. VHE Gamma Rays

The TeV observations were performed from 2007 September 4 to 21 using the MAGIC telescope on the Canary Island of La Palma (28.75°N , 17.86°W , 2225 m a.s.l.), from where LS I +61 303 is observable at zenith angles above 32° . The essential parameters of MAGIC are a 17 m diameter segmented mirror of parabolic shape (currently the largest dish for an air Cherenkov telescope), an f/D of 1.05 and an hexagonally shaped camera of 576 hemispherical photo multiplier tubes with a field of view of 3.5° diameter. MAGIC can detect gamma rays from 60 GeV to several TeV, and with a special setup the trigger threshold can be reduced to 25 GeV (Aliu et al. 2008). Its energy resolution is $\Delta E/E = 20\%$ above energies of 200 GeV. The current sensitivity is 1.6% of the Crab Nebula flux for a 5σ detection in 50 h of on-source time. The improvement compared to the previous sensitivity was achieved by installing new 2 GHz FADCs (Albert et al. 2008b).

The total observation time was 58.8 h, including 39.6 h in dark conditions and 19.2 h under moonlight or twilight (Britzger et al. 2009). The range of zenith angles for these observations was $[32^{\circ}, 50^{\circ}]$, with 96% of the data having zenith angles below 43° . The final effective total observation time is 54.2 h. Table 1 gives the MJD, the effective observation time, and the orbital phase for each of the MAGIC observations. The observations were carried out in wobble mode (Fomin et al. 1994), i.e. by alternately tracking two positions at 0.4° offset from the source position. This observing mode provides a reliable background estimate for pointlike sources such as LS I +61 303.

The data analysis was performed using the standard

TABLE 1
LOG OF THE TEV OBSERVATIONS

MJD ^a	Obs. Time (min)	Phase range	$N(> 300 \text{ GeV})^b$ ($10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$)
54348.148 ± 0.087	203.0	0.470–0.476	1.1 ± 1.7
54349.157 ± 0.080	210.1	0.508–0.514	−1.9 ± 2.0
54350.136 ± 0.094	214.5	0.544–0.551	0.2 ± 2.1
54351.160 ± 0.080	219.8	0.584–0.590	−0.1 ± 2.2
54352.161 ± 0.081	221.1	0.621–0.627	15.7 ± 2.4
54353.163 ± 0.082	224.2	0.659–0.665	7.7 ± 2.1
54354.166 ± 0.078	212.5	0.697–0.703	2.8 ± 2.2
54355.153 ± 0.091	172.4	0.734–0.741	0.3 ± 2.3
54356.139 ± 0.080	148.7	0.771–0.777	0.1 ± 2.5
54357.153 ± 0.092	178.1	0.809–0.816	6.4 ± 2.3
54358.155 ± 0.091	179.3	0.847–0.854	8.1 ± 2.4
54359.154 ± 0.092	184.4	0.885–0.892	2.9 ± 2.4
54360.154 ± 0.091	177.3	0.923–0.929	5.7 ± 2.5
54361.154 ± 0.094	183.1	0.960–0.967	5.3 ± 2.4
54362.153 ± 0.094	188.7	0.998–0.005	−0.6 ± 2.4
54363.156 ± 0.093	138.6	0.036–0.043	7.5 ± 2.8
54364.149 ± 0.099	195.7	0.073–0.081	3.8 ± 1.9

NOTE. — ^a The central MJD and half-span of each observation is quoted.

^b Uncertainties are given at 68% confidence level.

MAGIC analysis and reconstruction software (Albert et al. 2008c). The quality of the data was checked and data taken with anomalous event rates (very low atmospheric transmission, car light flashes, etc.) were rejected following the standard procedure, as previously done in Albert et al. (2009). From the remaining events, image parameters were calculated (Hillas 1985). In addition, the time parameters described in Aliu et al. (2009) were calculated as well. For the γ /hadron separation, a multidimensional classification procedure based on the image and timing parameters with the Random Forest method was used (Albert et al. 2008d). We optimized the signal selection cuts with contemporaneous Crab Nebula data taken at the same zenith angle range. The energy of the primary γ -ray was reconstructed from the image parameters using also a Random Forest method. The differential energy spectrum was unfolded taking into account the full instrumental energy resolution (Albert et al. 2007). We estimate the systematic uncertainty to be about 30% for the derived integral flux values and ± 0.2 for the obtained photon index. For more details on the systematic uncertainties present in the MAGIC data see Albert et al. (2008c).

2.2. X Rays

We observed LS I +61 303 with *XMM-Newton* during seven runs from 2007 September 4 to 11, lasting from 12 to 18 ks (see Table 2), amounting to a total observation time of 104.3 ks. The EPIC pn detector used the Large Window Mode, while the EPIC MOS detectors used the Small Window Mode. All detectors used the medium thickness optical blocking filter. Data were processed using version 8.0.0 of the *XMM-Newton* Science Analysis Software (SAS). Known hot or flickering pixels were removed using the standard SAS tasks. Further cleaning was necessary to remove from the dataset periods of high background corresponding to soft proton flares, reducing the net good exposure durations to 67.0 and 92.6 ks for the pn and MOS detectors, respectively.

Cleaned pn and MOS event files were extracted for spectral analysis. Source spectra were extracted from a $\sim 70''$ radius circle centered on the source (PSF of $15''$) while background spectra were taken from a number of source-free circles with $\sim 150''$ radius (three for the pn detector, four for MOS1 and five for MOS2). The extracted spectra were analyzed with

XSpec v12.3.1 (Arnaud 1996). The spectra were binned so that a minimum of 20 counts per bin were present and energy resolution was not over-sampled by more than a factor 3. An absorbed power-law function (`wabs*powerlaw`) yielded satisfactory fits for all observations. De-absorbed fluxes in the 0.3–10 keV range were computed from the spectral fits.

Additional observations of 2–5 ks each, consisting of several pointings of 0.2–1.0 ks, were obtained with the *Swift*/XRT from 2007 September 11 to 22. The total observation time was 28.5 ks. The *Swift* data were processed using the FTOOLS task `xrtpipeline` (version 0.12.1 under HEASoft 6.6). The spectral analysis procedures were the same as those used for the *XMM-Newton* data. Since the hydrogen column density was poorly constrained, with typical values of $(0.5 \pm 0.2) \times 10^{22} \text{ cm}^{-2}$, we fixed it to $0.5 \times 10^{22} \text{ cm}^{-2}$, a typical value for LS I +61 303 also found in the *XMM-Newton* fits and close to the average value for the *Swift* fits.

To search for short-term X-ray variability we also extracted 0.3–10 keV background-subtracted lightcurves for each observation, binned to 100 s for *XMM-Newton* and at half-time of the sparse 0.2 to 1.0 ks *Swift* pointings within each observation. For *XMM-Newton* we considered the sum of the count-rate in the three detectors. From these lightcurves we computed the degree of variability as the standard deviation with respect to the mean of the count rate divided by this mean, and the significance of this variability computed from the χ^2 probability of the count rate being constant. We also computed hardness ratios as the fraction between the count rates above and below 2 keV.

3. RESULTS

3.1. VHE Gamma Rays

The measured fluxes of LS I +61 303 at energies above 300 GeV are listed in Table 1, and the corresponding lightcurve is shown in Fig. 1-top. The periodically modulated peak emission (Albert et al. 2009) is prominently seen as the highest flux value of these observations at phase 0.62 (followed by a smaller flux at phase 0.66). In addition to this established feature we measured, as in 2006 December, significant flux on a nightly basis at phase 0.85. We note that there is significant flux in the phase range 0.8–1.0 at the level of $(5.2 \pm 1.0) \times 10^{-12} \text{ cm}^{-2} \text{ s}^{-1}$, which is compatible with the 2 σ upper limit we obtained for the shorter 2006 observations (Albert et al. 2009).

We also determined the differential energy spectrum from the ~ 10 h of observations conducted from orbital phase 0.6 to 0.7. A power-law fit yields:

$$\frac{dN}{dE} = \frac{(2.0 \pm 0.3_{\text{stat}} \pm 0.6_{\text{sys}}) \times 10^{-12}}{\text{cm}^2 \text{ s TeV}} \times \left(\frac{E}{1 \text{ TeV}} \right)^{-2.7 \pm 0.3_{\text{stat}} \pm 0.2_{\text{sys}}},$$

compatible within errors with our previous measurements (Albert et al. 2006, 2009).

3.2. X Rays

We summarize in Table 2 the parameters of the spectral fits and variability obtained for both the *XMM-Newton* and *Swift*/XRT data sets. All X-ray fluxes quoted hereafter are de-absorbed. Since there is significant X-ray count-rate variability within each observation, we have to consider it to provide

TABLE 2
LOG OF THE *XMM-Newton* (TOP) AND *Swift* (BOTTOM) X-RAY OBSERVATIONS

MJD ^a	Exposure time			Phase range	$F(0.3-10\text{ keV})^b$ ($10^{-12}\text{ erg cm}^{-2}\text{ s}^{-1}$)	Γ	$N_{\text{H}}(\text{wabs})$ (10^{22} cm^{-2})	$\chi^2/\text{d.o.f.}$	Variab./Signif. (%)/(σ)
	Total (ks)	pn (ks)	MOS (ks)						
54347.202 ± 0.072	12.4	10.8	12.4	0.434–0.440	13.8 ± 0.2 (±2.4)	1.87 ± 0.03	0.517 ± 0.014	446.2/382	17.1/22.5
54349.140 ± 0.076	13.1	5.6	9.1	0.507–0.513	12.4 ± 0.2 (±2.1)	1.66 ± 0.03	0.504 ± 0.019	379.8/352	16.8/15.1
54350.199 ± 0.106	18.3	10.5	17.1	0.546–0.554	13.3 ± 0.2 (±1.6)	1.66 ± 0.02	0.514 ± 0.013	536.5/422	11.8/13.8
54351.160 ± 0.099	17.1	10.9	12.1	0.583–0.590	13.4 ± 0.2 (±1.6)	1.68 ± 0.03	0.525 ± 0.015	431.6/400	12.1/12.8
54352.146 ± 0.085	14.7	5.8	13.2	0.621–0.627	22.9 ± 0.2 (±1.6)	1.54 ± 0.02	0.538 ± 0.012	496.7/447	6.8/ 5.9
54353.167 ± 0.084	14.5	11.9	14.5	0.659–0.665	18.6 ± 0.2 (±1.1)	1.58 ± 0.02	0.529 ± 0.013	467.5/429	5.9/ 1.9
54354.144 ± 0.082	14.2	11.5	14.2	0.696–0.702	12.6 ± 0.2 (±1.3)	1.65 ± 0.03	0.520 ± 0.015	439.5/407	10.0/ 8.0
54354.663 ± 0.073	3.3	0.716–0.722	10.3 ± 0.7 (±2.0)	1.71 ± 0.16	0.5 (fixed)	5.38/14	18.4/ 1.1
54355.670 ± 0.070	2.7	0.754–0.759	12.4 ± 0.9 (±1.6)	1.38 ± 0.14	0.5 (fixed)	8.83/15	10.7/ 0.3
54356.671 ± 0.073	3.3	0.792–0.797	17.6 ± 1.0 (±2.0)	1.36 ± 0.10	0.5 (fixed)	9.83/26	9.9/ 0.4
54357.674 ± 0.073	3.3	0.830–0.835	13.9 ± 0.9 (±2.3)	1.49 ± 0.14	0.5 (fixed)	13.07/20	15.3/ 1.2
54358.178 ± 0.103	3.8	0.848–0.855	17.7 ± 0.9 (±1.5)	1.47 ± 0.09	0.5 (fixed)	20.46/32	6.3/ 0.1
54359.951 ± 0.205	5.3	0.911–0.926	19.5 ± 0.8 (±2.6)	1.54 ± 0.07	0.5 (fixed)	34.52/50	12.7/ 0.4
54362.247 ± 0.237	2.1	0.996–0.014	13.7 ± 1.1 (±3.6)	1.78 ± 0.19	0.5 (fixed)	5.45/11	25.0/ 1.7
54363.121 ± 0.102	2.5	0.034–0.042	12.3 ± 1.0 (±1.5)	1.53 ± 0.16	0.5 (fixed)	9.71/13	9.3/ 0.1
54365.530 ± 0.040	2.2	0.127–0.130	13.3 ± 1.1 (±1.7)	1.46 ± 0.15	0.5 (fixed)	10.14/12	10.1/ 0.5

NOTE. — ^a The central MJD and half-span of each observation is quoted. ^b Fluxes are de-absorbed. The uncertainties in parentheses include the uncertainties due to intrinsic X-ray variability during the observations.




FIG. 1.— TeV and X-ray lightcurves of LS I +61 303 during the multi-wavelength campaign of 2007 September. *Top*: Flux above 300 GeV versus the observation time in MJD and the orbital phase. The horizontal dashed line indicates 0 flux. The vertical arrows mark the times of simultaneous TeV and X-ray observations. *Bottom*: De-absorbed flux in the 0.3–10 keV energy range for the seven *XMM-Newton* observations (filled circles) and the nine *Swift* ones (open circles). Error bars correspond to a 1σ confidence level in all cases. Dotted lines join consecutive data points to help following the main trends of the lightcurves. The sizes of the symbols are larger than the time span of individual observations.

more realistic values for the flux uncertainties than those directly obtained from the spectral fits. We note that there is no significant hardness ratio change within each of the observations, in contrast to what was found in the *XMM-Newton* observations reported by Sidoli et al. (2006). In principle this implies that, for each observation, the flux and corresponding uncertainty obtained from the spectral fit is a good estimate of the flux during the whole observation. However, moderate count-rate variability is present in most observations, ranging from 6 to 17% for the *XMM-Newton* data and from 9 to

25% for the *Swift* data (see Table 2), yielding to additional uncertainties that should be considered when providing a measurement spanning several ks. We converted this count-rate variability into flux variability by multiplying the degree of variability defined in Sect. 2.2 with the fluxes coming from the spectral fits. Since no spectral change is detected within each observation, we added this flux variability (as an estimate of flux uncertainty) in quadrature to the spectral fits flux errors. This procedure provides the more realistic total flux uncertainties quoted in parentheses in Table 2, and used hereafter.

We show in Fig. 1-bottom the 0.3–10 keV lightcurve of LS I +61 303. The source displays a steady flux during the first four observations and shows a steep increase at phase 0.62, which is followed by a slower decay up to phase 0.69 (*XMM-Newton*) and probably up to phase 0.72 (*Swift*). The behavior is very similar to the one seen at TeV energies, although at X-ray energies the baseline has a significant flux. Later on there is a significant increase of the X-ray flux up to phase 0.8. This high flux is detected with a sparse sampling up to phase 0.9, and the source goes back to its baseline flux at phase 1.0. This high X-ray flux between phases 0.8 and 1.0 occurs when the source is also detected at TeV energies.

3.3. X-Ray/TeV Correlation

A clear correlation between the X-ray and TeV emissions is seen during the outburst, with a simultaneous peak at phase 0.62 (see Fig. 1). To study the significance of this correlation, we selected the X-ray datasets that overlap with MAGIC observations. There are six overlapping MAGIC/*XMM-Newton* datasets, for which strictly simultaneous observations range from 3.3 to 3.9 h. For the four overlapping MAGIC/*Swift* datasets the strictly simultaneous observations range from 2.2 to 4.1 h (although the *Swift* runs have gaps). We plot in Fig. 2 the X-ray fluxes against the TeV fluxes for all ten simultaneous pairs, which are marked with arrows in Fig. 1. The linear correlation coefficient for the six simultaneous MAGIC/*XMM-Newton* pairs that trace the outburst is $r = 0.97$. For the ten simultaneous pairs we find $r = 0.81$ (which for a double tail test has a probability of about 5×10^{-3} to be produced from independent X-ray and TeV fluxes). Minimizing χ^2 we obtain $\chi^2 = 7.68$ for 8 degrees of freedom and the fol-




FIG. 2.— De-absorbed X-ray fluxes as a function of TeV fluxes. Only the 10 simultaneous fluxes, marked with arrows in Fig. 1, have been considered. Error bars correspond to a 1σ confidence level in all cases. The solid line represents a χ^2 linear fit to all data points.

lowing relationship: $F(0.3-10 \text{ keV})/[10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}] = (12.2^{+0.9}_{-1.0}) + (0.71^{+0.17}_{-0.14}) \times N(E > 300 \text{ GeV})/[10^{-12} \text{ cm}^{-2} \text{ s}^{-1}]$ (non-Gaussian uncertainties). This fit is plotted as a solid line in Fig. 2.

However, as can be seen in Fig. 2, the flux uncertainties are relatively large. This calls for a test of the reliability of the correlation strength considering the errors of individual data points. To this end we use the z-Transformed Discrete Correlation Function (ZDCF), which determines 68% confidence level intervals for the correlation coefficient from unevenly sampled data as a function of time lags (see e.g. Edelson & Krolik 1988, Alexander 1997). The Fisher z-transform of the linear correlation coefficient is used to estimate the 68% confidence level interval. The X-ray and TeV lightcurves reported here lead to the highest correlation coefficient for simultaneous observations, yielding the uncertainties for the linear correlation coefficient to be $r = 0.81^{+0.06}_{-0.21}$. The sensitivity of the MAGIC telescope requires observation times of several hours to be able to get few-sigma detections for the TeV fluxes from LS I +61 303. During such long periods, the X-ray fluxes have quite large intrinsic variability (see Table 2). This results in quite large uncertainties on both the TeV and X-ray fluxes that restrict the ability to determine a possible correlation out of the outburst.

4. DISCUSSION

We have discovered an X-ray/TeV correlation in the gamma-ray binary LS I +61 303 based on simultaneous multiwavelength data obtained with MAGIC, *XMM-Newton*, and *Swift*. The correlation is mainly due to the very similar trends of the detected flux during the outburst around orbital phase 0.62, while the uncertainties prevent to be sure about the existence of the correlation outside the outburst. Given the variability in the X-ray flux (up to 25% on hour scales), it is necessary to program simultaneous observations to perform correlation studies in LS I +61 303. This conclusion has also been reached recently by the VERITAS Collaboration, when reporting the lack of X-ray/TeV correlation based on contemporaneous data with a TeV sampling that is not dense enough (Acciari et al. 2009). Although a similar X-ray/TeV corre-

lation has been obtained for the TeV emitting X-ray binary LS 5039, this result was based on non-simultaneous data acquired years apart (Aharonian et al. 2005, 2006; Takahashi et al. 2009).

The TeV emission within a binary system can suffer photon-photon absorption via pair creation, mainly with the stellar optical/ultraviolet photons. In LS I +61 303 this absorption is only expected to be significant towards the observer for $E > 300 \text{ GeV}$ just before periastron (Dubus 2006b; Bednarek 2006; Sierpowska-Bartosik & Torres 2009), a phase range not explored here. On the other hand, the quoted X-ray fluxes are already de-absorbed (and the hydrogen column densities and the associated errors are low). Therefore, the X-ray/TeV correlation we have found for LS I +61 303 indicates that the emission processes at both wavelengths occur at the same time and are probably the result of a single physical mechanism. In this context, it is reasonable to assume that the X-ray and TeV emissions are produced by a single particle population.

It is interesting to note that the MAGIC spectrum in the 0.6–0.7 phase range yields $\sim 11 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ for $E > 300 \text{ GeV}$, while the X-ray flux is $\sim 19 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$. Therefore, the total X-ray flux approximately doubles the TeV flux. However, if we subtract an apparent baseline X-ray flux of $10 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$, the resulting X-ray flux is similar to the total TeV flux in the phase range 0.6–0.7. If the radiation mechanisms are dominated by a single particle population, the X-ray/TeV correlation and the smaller/similar TeV fluxes favor leptonic models. In hadronic models, the X-ray emitting e^\pm and the TeV photons would come from the same protons (for reasonable values of the magnetic field), and the luminosity of the e^\pm radiation should be $\lesssim 1/2$ that of TeV gamma-rays (Kelner et al. 2006) unlike it is observed. In addition, the IC cooling channel is less efficient than the synchrotron channel to produce the detected X-ray emission for reasonable values of the magnetic field (see Takahashi et al. 2009 for a similar discussion for LS 5039). This clearly suggests that the X rays are the result of synchrotron radiation of the same VHE electrons that produce TeV emission as a result of inverse Compton scattering of optical/ultraviolet stellar photons.

The observed photon indexes of the simultaneous X-ray and VHE spectra are consistent with one population of electrons following a power-law energy distribution with index ~ 2.1 . These electrons would produce X-rays via synchrotron and VHE photons via IC with an interaction angle $\lesssim \pi/2$. We note that an electron index of ~ 2.1 is too hard if synchrotron cooling dominates in the X-ray range, since it implies an injected electron index of 1.1. On the other hand, dominant adiabatic cooling implies an injection index of 2.1, a more reasonable value (see for example the discussion in Takahashi et al. 2009). Although small changes in the VHE spectrum would be expected due to variations in the IC interaction angle or the electron index (as seen in X-rays), at present they are not detectable due to the large uncertainties of the VHE photon index.

Finally, we note that contemporaneous radio lightcurves obtained with RATAN, VLBA images, and $H\alpha$ spectroscopy are consistent with previous results (Gregory 2002; Dhawan et al. 2006; Zamanov et al. 1999). Details on these observations will be reported elsewhere. Therefore, the X-ray/TeV correlation occurred when the source was showing a standard behavior in both its outflow (radio) and decretion disk ($H\alpha$ line).

We would like to thank the Instituto de Astrofísica de Canarias for the excellent working conditions at the Observatorio del Roque de los Muchachos in La Palma. The support of the German BMBF and MPG, the Italian INFN and Spanish MICINN is gratefully acknowledged. This work was also

supported by ETH Research Grant TH 34/043, by the Polish MNiSzW Grant N N203 390834, and by the YIP of the Helmholtz Gemeinschaft.

Facilities: MAGIC, XMM, Swift, RATAN, VLBA, Skinas:1.3m

REFERENCES

- Acciari, V. A., et al. 2008, *ApJ*, 679, 1427
 Acciari, V. A., et al. 2009, *ApJ*, 700, 1034
 Aharonian, F., Akhperjanian, A. G., Aye, K.-M., et al. 2005, *Science*, 309, 746
 Aharonian, F., et al. 2006, *A&A*, 460, 743
 Albert, J., et al. 2006, *Science*, 312, 1771
 Albert, J., et al. 2007, *Nucl. Instrum. Methods Phys. Res. A*, 583, 494
 Albert, J., et al. 2008a, *ApJ*, 684, 1351
 Albert, J., et al. 2008b, *Nucl. Instrum. Methods Phys. Res. A*, 594, 407
 Albert, J., et al. 2008c, *ApJ*, 674, 1037
 Albert, J., et al. 2008d, *Nucl. Instrum. Methods Phys. Res. A*, 588, 424
 Albert, J., et al. 2009, *ApJ*, 693, 303
 Alexander, T. 1997, in *Astronomical Time Series*, ed. D. Maoz, A. Sternberg, & E. M. Leibowitz (Dordrecht, Kluwer), 163
 Aliu, E., et al. 2008, *Science*, 322, 1221
 Aliu, E., et al. 2009, *Aph*, 30, 293
 Aragona, C., McSwain, M. V., Grundstrom, E. D., Marsh, A. N., Roettenbacher, R. M., Hessler, K. M., Boyajian, T. S., & Ray, P. S. 2009, *ApJ*, 698, 514
 Arnaud, K. A. 1996, in *ASP Conf. Ser. 101, Astronomical Data Analysis Software and Systems V*, ed. G. Jacoby & J. Barnes (San Francisco, CA: ASP), 17
 Bednarek, M. 2006, *MNRAS*, 368, 579
 Bogovalov, S. V., Khangulyan, D. V., Koldoba, A. V., Ustyugova, G. V., & Aharonian, F. A. 2008, *MNRAS*, 387, 63
 Britzger, D., et al. (MAGIC Collaboration) 2009, *Proc. of the 31th ICRC, Łódź*, in press (arXiv:0907.0973)
 Casares, J., Ribas, I., Paredes, J. M., Martí, J., & Allende Prieto, C. 2005, *MNRAS*, 360, 1105
 Dhawan, V., Mioduszewski, A., & Rupen, M. 2006, in *Proc. VI Microquasar Workshop: Microquasars and Beyond*, ed. T. Belloni (Trieste: PoS), 52
 Dubus, G. 2006a, *A&A*, 456, 801
 Dubus, G. 2006b, *A&A*, 451, 9
 Edelson, R. A., & Krolik, J. H. 1988, *ApJ*, 333, 646
 Esposito, P., Caraveo, P. A., Pellizzoni, A., de Luca, A., Gehrels, N., & Marelli, M. A. 2007, *A&A*, 474, 575
 Fomin, V. P., Stepanian, A. A., Lamb, R. C., Lewis, D. A., Punch, M., & Weekes, T. C. 1994, *Aph*, 2, 137
 Frail, D. A., & Hjellming, R. M. 1991, *AJ*, 101, 2126
 Goldoni, P., & Mereghetti, S. 1995, *A&A*, 299, 751
 Gregory, P. C. 2002, *ApJ*, 575, 427
 Grundstrom, E. D., et al. 2007, *ApJ*, 656, 437
 Harrison, F. A., Ray, P. S., Leahy, D. A., Waltman, E. B., & Pooley, G. G. 2000, *ApJ*, 528, 454
 Hillas, A. M. 1985, *Proc. of the 19th ICRC, La Jolla*, 3, 445
 Kelner, S. R., Aharonian, F. A., & Bugayov, V. V. 2006, *Phys. Rev. D*, 74, 034018
 Massi, M., Ribó, M., Paredes, J. M., Garrington, S. T., Peracaula, M., Martí, J. 2004, *A&A*, 414, L1
 Paredes, J. M. 2008, *American Institute of Physics Conference Series*, 1085, 157
 Paredes, J. M., Martí, J., Peracaula, M., & Ribó, M. 1997, *A&A*, 320, L25
 Sidoli, L., Pellizzoni, A., Vercellone, S., Moroni, M., Mereghetti, S., & Tavani, M. 2006, *A&A*, 459, 901
 Sierpowska-Bartosik, A., & Torres, D. F. 2009, *ApJ*, 693, 1462
 Smith, A., Kaaret, P., Holder, J., Falcone, A., Maier, G., Pandel, D., & Strohm, M. 2009, *ApJ*, 693, 1621
 Takahashi, T., et al. 2009, *ApJ*, 697, 535
 Taylor, A. R., Young, G., Peracaula, M., Kenny, H. T., & Gregory, P. C. 1996, *A&A*, 305, 817
 Zamanov, R. K., Martí, J., Paredes, J. M., Fabregat, J., Ribó, M., Tarasov, A. E. 1999, *A&A*, 351, 543