

"Made available under NASA sponsorship
In the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

E 7.4 - 1 0.5 3 2.
CR - 138281

CORRELATION OF COASTAL WATER TURBIDITY
AND CIRCULATION WITH ERTS-1 AND SKYLAB IMAGERY

V. Klemas
~~M. Otley~~
College of Marine Studies
University of Delaware
Newark, DE

R. Rogers
Bendix Aerospace Corporation
Ann Arbor, MI

May 28, 1974
Report on Significant Results
NASA ERTS-1 Contract NASS-21837
Un 362 SR 9654

Prepared for
GODDARD SPACE FLIGHT CENTER
GREENBELT, MD 20771

(E74-10532) CORRELATION OF COASTAL WATER
TURBIDITY AND CIRCULATION WITH ERTS-1 AND
SKYLAB IMAGERY Report on Significant
Results (Delaware Univ.) 2 p HC \$4.00
CSCL 08C G3/13
Unclas
00532
N74-25850

Significant Results

Circulation patterns were observed by ERTS-1 during different parts of the tidal cycle, using suspended sediment as a natural tracer. The current direction in the imagery agreed well with predicted and measured currents throughout Delaware Bay. During flood tide the suspended sediment as seen from ERTS-1 also correlated with the depth profile at the mouth of the bay.

Convergent shear boundaries between different water masses have been observed from ERTS-1, with foam lines containing high concentrations of lead, mercury and other toxic substances. Several varieties of boundaries or fronts have been seen. Those near the mouth of the bay are associated with the tidal intrusion of shelf water. Fronts in the interior of the bay on the Delaware side appear to be associated with velocity shears induced by differences in bottom topography. In several ERTS-1 and Skylab-EREP frames, acid waste disposal plumes have been detected about 40 miles off Delaware's Atlantic coast.

Under atmospheric conditions encountered along the East Coast of the United States, MSS band 5 seemed to give the best representation of sediment load in the upper one meter of the water column. Band 4 was masked by haze-like noise, while band 6 did not penetrate sufficiently into the water column. Microdensitometer traces across the band 5 imagery produce a relative irradiance curve which correlates with Secchi depth and suspended sediment concentration. A more meaningful correlation between radiance and sediment concentration can be obtained by combining the data in all four bands using computer techniques. Such a method promises to yield not only a mathematically expressible relationship between concentration and radiance but also, by identification of typical spectral signatures, allows identification of sediment and certain pollutants. If this method is refined by using a greater number of bands, it may even be capable of distinguishing between different sediment types.

As expected, Skylab imagery contains better spectral and spatial resolution than that from ERTS-1. Based on high-contrast targets, such as pier clusters and street patterns, the ERTS-1 MSS seems to have a resolution of 70 to 100 meters. On the other hand, the regular eighteen-day cycle of ERTS-1 permits observation of important man-made and natural changes, and facilitates collection of ground truth.