



Corrigendum to

“Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V – heterogeneous reactions on solid substrates” published in Atmos. Chem. Phys. 10, 9059–9223, 2010

J. N. Crowley¹, M. Ammann², R. A. Cox³, R. G. Hynes⁴, M. E. Jenkin⁵, A. Mellouki⁶, M. J. Rossi⁷, J. Troe⁸, and T. J. Wallington⁹

¹Max-Planck-Institut für Chemie, Division of Atmospheric Chemistry, Hahn-Meitner-Weg 1, 55128 Mainz, Germany

²Laboratory of Radiochemistry and Environmental Chemistry, OFLB 103, Paul Scherrer Institut, 5232 Villigen, Switzerland

³Centre for Atmospheric Science, Dept. of Chemistry, University of Cambridge, Lensfield Road Cambridge CB2 1EP, UK

⁴CSIRO Energy Technology, Lucas Heights Science and Technology Centre, Building 2, PMB7, Bangor, NSW 2234, Australia

⁵Atmospheric Chemistry Services, Okehampton, Devon EX20 1FB, UK

⁶ICARE (Institut de Combustion, Aérothermique, Réactivité et Environnement) CNRS (Centre National de la Recherche Scientifique) – UPR3021, 1C, Avenue de la recherche scientifique, 45071 Orleans cedex 02, France

⁷Laboratorium für Atmosphärenchemie, OFLA 008, Paul Scherrer Institut, 5232 Villigen, Switzerland

⁸Institute of Physical Chemistry, University of Göttingen, Tammannstr. 6, 37077 Göttingen, Germany

⁹Ford Motor Company, Research and Advanced Engineering, Mail Drop RIC-2122, Dearborn, Michigan 48121-2053, USA

* The IUPAC Subcommittee on Gas Kinetic Data Evaluation for Atmospheric Chemistry

Correspondence to: J. N. Crowley (john.crowley@mpic.de)

In the paper “Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V – heterogeneous reactions on solid substrates” (published in Atmos. Chem. Phys., 12, 5223–5235, doi:10.5194/acp-10-9059-2010), the preferred value of the uptake coefficient for N₂O₅ interacting with NAT surfaces (datasheet V.A5.5) is given as 6×10^{-3} on pages 9063 and 9208. The correct value is 6×10^{-4} .

The service charges for this open access publication have been covered by the Max Planck Society.