

Counter-current chromatography for the separation of terpenoids: a comprehensive review with respect to the solvent systems employed

Krystyna Skalicka-Woźniak · Ian Garrard



Received: 15 January 2014 / Accepted: 11 March 2014 / Published online: 25 March 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract Natural products extracts are commonly highly complex mixtures of active compounds and consequently their purification becomes a particularly challenging task. The development of a purification protocol to extract a single active component from the many hundreds that are often present in the mixture is something that can take months or even years to achieve, thus it is important for the natural product chemist to have, at their disposal, a broad range of diverse purification techniques. Counter-current chromatography (CCC) is one such separation technique utilising two immiscible phases, one as the stationary phase (retained in a spinning coil by centrifugal forces) and the second as the mobile phase. The method benefits from a number of advantages when compared with the more traditional liquid–solid separation methods, such as no irreversible adsorption, total recovery of the injected sample, minimal tailing of peaks, low risk of sample denaturation, the ability to accept particulates, and a low solvent consumption.

The selection of an appropriate two-phase solvent system is critical to the running of CCC since this is both the mobile and the stationary phase of the system. However, this is also by far the most time consuming aspect of the technique and the one that most inhibits its general take-up. In recent years, numerous natural product purifications have been published using CCC from almost every country across the globe. Many of these papers are devoted to terpenoids—one of the most diverse groups. Naturally occurring terpenoids provide opportunities to discover new drugs but many of them are available at very low levels in nature and a huge number of them still remain unexplored. The collective knowledge on performing successful CCC separations of terpenoids has been gathered and reviewed by the authors, in order to create a comprehensive document that will be of great assistance in performing future purifications.

Keywords Counter-current chromatography · Terpenoids · Terpenes · Natural products · Separation · Purification

K. Skalicka-Woźniak (✉)
Department of Pharmacognosy with Medicinal Plant Unit,
Medical University of Lublin, 1 Chodzki Str., 20-093
Lublin, Poland
e-mail: kskalicka@pharmacognosy.org

I. Garrard
Advanced Bioprocessing Centre, Brunel Institute for
Bioengineering, Brunel University, Uxbridge UB8 3PH,
UK

Introduction

Terpenoids, also referred to as terpenes, are one of the largest and the most diverse group of natural products accounting for more than 40,000 individual compounds, with several new compounds being discovered every year.




Fig. 1 Biosynthetic pathways of terpenes (Wang et al. 2005)

They are synthesized from only two five-carbon isomers: isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP). Two biosynthetic routes have been characterized: the classical acetate mevalonate pathway (described in 1967) and the triose phosphate-utilizing non-mevalonate pathway characterized in 2002. Starting from the universal precursors IPP and DMAPP, thousands of enzymes are involved in the biosynthetic pathways for terpenoid chain elongation, cyclization, and functionalization of hydrocarbon chains. The active isoprene unit (IPP) is repetitively added to DMAPP or a prenyl diphosphate in sequential head-to-tail condensations catalyzed by the prenyltransferases. Through consecutive condensations a prenyltransferase can synthesize a variety of products with fixed lengths and stereochemistry (Fig. 1) (Wang et al. 2005; Ajikumar et al. 2008). Based on the number of the building blocks, terpenoids are commonly classified into hemi-, mono-, sesqui-, di-, ses-, tri- and tetraterpenoids (carotenoids) having 1, 2, 3, 4, 5, 6 and 8 isoprenoid residues respectively, and polyterpenes consisting of long chains of many isoprene units (Koch et al. 2008).

Terpenoids display a wide range of biological activities. Monoterpenes and sesquiterpenes are the

main constituents of essential oils and share responsibility for important properties like antibacterial, antiviral, antioxidant etc. Triterpene saponins—ginsenosides—significantly reduce the production of beta-amyloid which accumulates in the brain of patients with Alzheimer's disease and play a critical role in pathology by inducing neuronal death. Ginkgolides (cyclic diterpenes of labdane type commonly isolated from *Ginkgo biloba*) protect neuronal cells from synaptic damage (Yoo and Park 2012). However the antimalarial drug Artemisinin and the anticancer drug paclitaxel (TaxolR) are two renowned terpene-based drugs with established medical applications. Artemisinin, earlier known as *Qinghaosu*, is a phytoconstituent isolated from *Artemisia annua* L. and can be described as a compound which possess antimalarial activity. Clinical studies with patients infected with *Plasmodium vivax* or *P. falciparum* demonstrated that artemisinin could kill the malarial parasite very quickly at the schizont stage of the parasite's life cycle (i.e. when it infects the human red blood cell) with no obvious side effects. The molecule has a completely new antimalarial prototype structure with an endoperoxide moiety, which is necessary for activity. Since artemisinin itself has poor bioavailability limiting its

effectiveness, several semisynthetic derivatives of artemisinin have been developed (Medhi et al. 2009; Brown 2010). Taxol, a plant diterpenoid widely used as a chemotherapeutic drug against several types of cancer, is known to interact with a specific site of β -tubulin—it binds to microtubules and inhibits their disassembly. Cells treated with taxol are arrested in mitosis and eventually undergo death by apoptosis. This very important activity is strongly depended on its unusual structure. It was shown that the side chain at position C-13 and the taxane ring system are essential for this activity (Xiang et al. 2009).

The purification of natural products is a complex process requiring a comprehensive range of techniques. CCC offers the natural product scientist a different mode of operation to conventional processes. Invented in the mid 1960's (Ito et al. 1966), to many scientists it is still known as it was back then—a technique that is slow, with separations measuring in hours or days. It was also unreliable as instruments frequently broke down and furthermore had poor capacity with injection amounts measured in tens of milligrams. There was also no opportunity for scale up as the factors required to scale up were poorly understood. However, the technique has been substantially developed since those early days. Advances in engineering and the understanding of the processes involved, particularly in the past 10 years, have created instruments that are fast, robust, permit very high injection loadings and, significantly for the natural products industry, can be rapidly scaled from analytical to pilot level (Sutherland and Fisher 2004). The new generation of coil planet centrifuges operate at higher “g” fields than conventional instruments, enabling higher flow rates to be used so that separation times are measured in minutes rather than hours at the same resolution (Yuan et al. 2008). With a number of important advantages over both solid phase chromatographic techniques and current liquid–liquid extraction techniques, modern high capacity counter-current chromatography is a worthy inclusion in the array of techniques required for natural product purifications.

Theoretical background of CCC

In a counter-current chromatography centrifuge, tubing is wound on a drum which is centrifugally rotated in planetary motion (the holder rotates about its own




Fig. 2 Motion of the bobbin in the CCC centrifuge (Sutherland et al. 1998)

axis while revolving around the centrifuge axis at the same angular velocity in the same direction) (Fig. 2). A two phase solvent system is introduced into this coil. Although a simple solvent system might consist of hexane and water, a more likely system for a purification would consist of hexane, ethyl acetate, methanol and water or, for biomolecules sensitive to organic solvents, an aqueous two phase system such as aqueous PEG1000 and potassium phosphate salt solution. With the two phase solvent system inside, as the coil travels through its planetary motion cycle, zones of mixing and settling travel along the phases coincident with the low and high accelerations caused by the epicyclic motion of the coil. The mixing zones are coincident with low accelerations and take the form of wave mixing, equivalent to the “swish-swosh” motion that occurs when a tube of liquids is tilted from side to side. The settling zones are coincident with high accelerations and take the form of a smooth interfacial area. There is one mixing and one settling zone per coil loop per revolution. A typical modern analytical CCC instrument may have 40 loops and spin at 2,000 rpm. It will therefore experience 4.8 million partitioning steps per hour ($40 \times 2,000 \times 60$). Similarly, a typical preparative instrument may have 30 loops and spin at 1,200 rpm, giving 2.2 million partitioning steps per hour.

Counter-current chromatography can be achieved not only in the above mentioned hydrodynamic CCC created by two axis of rotation, but also as hydrostatic CCC, typically represented by centrifugal partition chromatography (CPC). This is a single-axis instrument, which has a series of chambers machined circumferentially around a rotor. Rotation of the rotor produces a uniform g-field, which retains the




Fig. 3 Type-J planetary motion of a multilayer coil separation column presenting that the column holder rotates about its own axis and revolves around the centrifuge axis at the same angular velocity (ω) in the same direction (Ito 2005)

stationary phase in each chamber, while the mobile phase is flowed through in a cascading manner. A CPC instrument requires rotating seals for the mobile phase flow, whereas the hydrodynamic CCC instruments do not. This is because the rotation of the coil about its own axis unwinds the twist produced by its motion around the sun gear, and thus there is no twisting of the flow tubes linking the coil to the pump and the detector (Fig. 3).

Solvent selection process

In the past, selection of a suitable two phase solvent system involved a considerable amount of experience and know-how. With aqueous-organic phase systems coming from up to six or more different solvents mixed together, the possibilities were almost limitless. A typical mid-polarity selection table is shown in Table 1 (Garrard 2005). This is a modified version of the table produced by Oka et al. (1991) and runs from moderately polar (System No1: butanol–water) to moderately nonpolar (System No28: heptane–methanol). The italicized systems in the table can be used to rapidly screen the whole table first, allowing the operator to focus into the correct area of interest. Being all multiples of 0.5 ml when 4 ml solvent system is made, these particular systems are quick to make up and test. It is also possible to create the solvent systems on a micro-scale in 96 well plates if the crude sample is in short supply.

In order to achieve an efficient resolution of the target compounds, the K values, which express the solute concentration in the stationary phase divided by

Table 1 Table for selecting a suitable moderately polar two-phase solvent system, graded from polar (No1) to nonpolar (No28)

No	Heptane	EtOAc	MeOH	Butanol	Water
1	0	0	0	2	2
2	0	0.4	0	1.6	2
3	0	0.8	0	1.2	2
4	0	1.2	0	0.8	2
5	0	1.6	0	0.4	2
6	0	2	0	0	2
7	0.1	1.9	0.1	0	1.9
8	0.2	1.8	0.2	0	1.8
9	0.29	1.71	0.29	0	1.71
10	0.33	1.67	0.33	0	1.67
11	0.4	1.6	0.4	0	1.6
12	0.5	1.5	0.5	0	1.5
13	0.57	1.43	0.57	0	1.43
14	0.67	1.33	0.67	0	1.33
15	0.8	1.2	0.8	0	1.2
16	0.91	1.09	0.91	0	1.09
17	1	1	1	0	1
18	1.09	0.91	1.09	0	0.91
19	1.2	0.8	1.2	0	0.8
20	1.33	0.67	1.33	0	0.67
21	1.43	0.57	1.43	0	0.57
22	1.5	0.5	1.5	0	0.5
23	1.6	0.4	1.6	0	0.4
24	1.67	0.33	1.67	0	0.33
25	1.71	0.29	1.71	0	0.29
26	1.8	0.2	1.8	0	0.2
27	1.9	0.1	1.9	0	0.1
28	2	0	2	0	0

Quantities (in ml) required to make 4 ml of system using a liquid-handling robot. EtOAc = ethyl acetate, MeOH = methanol. Hexane may be used instead of heptane (Garrard 2005)

that in the mobile phase, should be calculated. The partition coefficient (K) should lie within the approximate range $0.5 < K < 2.0$. A smaller K value results in a loss of peak resolution, whilst a larger value produces excessive band broadening.

Solvent systems employed in CCC for terpenoid purification

In order to fully assess the use of two phase systems in CCC of terpenoids, approximately 3,500 scientific

papers were studied, published in the last 30 years, that related to all aspects of CCC or CPC. Papers that contained an application example of terpenoids, i.e. a purification performed by CCC or CPC were noted, together with the compounds purified and the solvent system used for the purification. Papers which gave examples of separations reported elsewhere were ignored, as were all symposium abstracts. Only papers which gave specific details of the solvent system and solute were recorded and only natural product secondary metabolites were noted e.g. no synthetic compounds, dyes or chemicals.

In total therefore, 150 solvent systems were listed in Table 2 together with the corresponding solutes that they separated. Some of the solvent systems corresponded to more than one solute, and some of the solutes corresponded to more than one solvent system, but if the same solute and the same solvent system were listed, this was simply a duplicate entry and was therefore removed.

The solvent systems tables mentioned above are presented here, sorted according to class of terpenoid separated. That table is presented for the benefit of CCC users by suggesting possible suitable solvent systems, which can act as a starting point for further refinement and optimization in the composition of the system.

Applications of high speed and high performance CCC

Being a liquid–liquid chromatography system, CCC can select from an almost infinite range of possible two-phase solvent systems for a purification. Most reported purifications with the technique have understandably concentrated on compounds of intermediate polarity. For example, a review of Chinese herbal medicines purified by CCC found a total of 214 different compounds in 198 published papers with a LogP polarity range from -4 to $+12$ (Sutherland and Fisher 2009). However, more than 60 % of those compounds fell in the narrow intermediate polarity range of $0-4$. Nevertheless, CCC and its sister technique, CPC can be particularly useful for purifications in the extreme polar and non-polar range and some impressive examples have been published. In 1995, Gasper and co-workers managed to purify C_{60} and C_{70} fullerenes using the non-polar and non-

aqueous solvent system of isooctane, dimethylformamide, 1,2-dichlorobenzene (4:2:1) plus 1 % *tert*-butylmethyl ether (Gasper et al. 1995). Still at the non-polar end of the spectrum, the carotenoid lycopene was isolated from tomato paste using a non-aqueous phase system of *n*-hexane, dichloromethane and acetonitrile (10:3.5:6.5) (Wei et al. 2001). This separation was performed in a single step from the crude material with 100 mg of crude extract injected onto a 230 ml capacity centrifuge. The purity of the final product was measured as >98 % by HPLC peak area. Lycopene has a calculated LogP value of approximately 17.6. Beyond the carotenoids, non-polar terpenoids can also be challenging to purify with solid phase chromatography, and again CCC offers an alternative approach. In one example, Liu et al. (2013a) successfully separated cycloartenyl ferulate and 24-methylene cycloartanyl ferulate. Due to the weak polarity, a series of low-polar solvent systems, including: *n*-hexane–ethyl acetate–ethanol (methanol)–water, *n*-hexane–ethanol (methanol)–water, *n*-hexane–methanol and *n*-hexane–ethyl acetate–*n*-butanol ethanol (methanol)–water were tested, but all without success. Since the HPLC mobile phase for the analysis of those two target compounds consisted of methanol, acetonitrile and isopropanol, these three solvents were further studied and a mixture of hexane and acetonitrile (1:1) was used for the purification. Another application of non-aqueous solvent systems is the purification of shionone—also a low-polar compound. Wang et al. (2012a) tested several hydrophobic two-phase solvent systems. Among these *n*-hexane–methanol (2:1) and heptane–dichloromethane–acetonitrile (20:7:13) were suitable for the separation.

Turning to the extreme polar end of the spectrum, CCC has been used with aqueous-organic solvent systems for the purification of peptides. For example, the peptide antibiotic colistin (LogP about -4.7) was isolated from a commercial microbial preparation using the polar two-phase system consisting of *n*-butanol and 0.04 M aqueous trifluoroacetic acid (1:1) (Ikai et al. 1998). Using the salt-based solvent system consisting of 1-propanol, acetonitrile, saturated ammonium sulphate and water (1:0.5:1.2:1) the highly polar glucosinolate glucoraphanin was purified to a purity >98 % from a crude broccoli extract in a single step (Fisher et al. 2005).

Sometimes modifications are necessary. The separation of three closely related triterpenes: sericic acid,

Table 2 Solvent systems in CCC for terpenoids separations

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
Monoterpenoids	Paeoniflorin	Huang et al. (2013)	CCC/ethyl acetate– <i>n</i> -butanol–water (3:2.5:5)
	Albiflorin		
	Rosiridin	Mudge et al. (2013)	CCC/ethyl acetate–butanol–water (3:2:5)
	Geranyl 1- <i>O</i> - α -L-arabinopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside		
	(<i>S</i>)-3,7-Dimethyl-5-octene-1,7-diol	Knapp et al. (1998)	CCC/chloroform–methanol–water (7:13:8)
	Thymol	Puertas Mejia et al. (2002)	CCC/ <i>n</i> -hexane– <i>tert</i> -butylmethyl ether–acetonitrile (1:0.1:1)
	Carvacrol		
	Eugenol	Geng et al. (2007)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:0.5:1:0.5)
	Chavibetol	dos Santos et al. (2009)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–methanol–water (12:4:4:3)
	Methyleugenol		
	α -Cyperone	Shi et al. (2009)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:0.2:1.1:0.2)
	1,8-Cineole	Dang et al. (2010)	CPC/petroleum ether–acetonitrile–acetone (4:3:1)
	Paeoniflorin	Chen et al. (2004)	CCC/ <i>n</i> -butanol–ethyl acetate–water (1:4:5)
	Cuminaldehyde	Chen et al. (2011)	CCC/ <i>n</i> -hexane–methanol–water (5:4:1)
	<i>p</i> -Menta-1,4-dien-7-al		
	Linalol	Skalicka-Woźniak et al. (2013)	CCC/heptane–ethyl acetate–methanol–water (5:2:5:2)
	Terpinen-4-ol		
	α -Terpineol		
	Anethole	Skalicka-Woźniak et al. (2013)	CCC/heptane–methanol (1:1)
	Foeniculin		
Sesquiterpenoids	Parthenolide	Fischedick et al. (2012)	CPC/heptane–ethyl acetate–methanol–water (1:1:1:1)
	11,13-Dihydroparthenolide		
	Anhydroverlotrin		
	3 β -Hydroxycostunolide		
	Costunolide diepoxide		
	3-Hydroxyparthenolide		
	Artemorin		
	Santamarine		
	Reynosin		
	Artecanin		
	Tanaparthin- β -peroxide		
	Artemisinin	Acton et al. (1986)	CCC/ <i>iso</i> -octane–ethyl acetate–methanol–water (7:3:6:4)
	Costunolide	Li et al. (2005)	CCC/light petroleum–methanol–water (5:6.5:3.5)
Dehydrocostuslactone			
Lactucopicrin	Wu et al. (2007)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1.5:5:2.75:5)	

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
	11 β ,13-Dihydrolactucin Lactucin	Wu et al. (2007)	CCC/ethyl acetate–methanol–water (20:1:20)
	Peroxyferolide Lipiferolide	Graziose et al. (2011)	CPC/ <i>n</i> -hexane–ethyl acetate–methanol–water (2:1:2:1)
	14-(3-Methylpentanoyl)-6-deoxybritannilactone 14-(3-Methylbutanoyl)-6-deoxybritannilactone 14-(2-Methylpropanoyl)-6-deoxybritannilactone	Fischedick et al. (2013a)	CPC/heptane–ethyl acetate–methanol–water (4:6:4:6)
	1,3-Epi-granilin 11,13-Dihydro-inuchinenolide B Pulchellin C 6-Deacetylbritanin 4H-Tomentosin Gaillardin Britannin		
	3 β -Hydroxy-8 β -[4'-hydroxytigloyloxy]-costunolide Eupalinolide A Eupalinolide B	Yan et al. (2012)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:4:2:3)
	Xanthathin 4-Epi-xanthanol 4-Epi-isoxanthanol	Pinel et al. (2007)	CPC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:1:1:1)
	β -Caryophyllene Rupestonic acid	Xie et al. (2008) Ma et al. (2005)	CCC/ <i>n</i> -hexane–dichloromethane–acetonitrile (10:3:7) CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (6:4:3.5:6.5) with 0.5 % acetic acid in stationary-phase
	Rupestonic acid Germacrone Curdione	Yang et al. (2010) Yan et al. (2005)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (3:5:3:5) CCC/light petroleum ether–ethanol–diethyl ether–water (5:4:0.5:1)
	Atractylon Atractylenolide III Nootkatone	Zhao and He (2006) Xie et al. (2009)	CCC/light petroleum–ethyl acetate–ethanol–water (4:1:4:1) CCC/ <i>n</i> -hexane–methanol–water (5:4:1)
	Caryophyllene oxide β -Farnesene Caryophyllene	Wei et al. (2012)	CCC/ <i>n</i> -hexane–acetonitrile–ethanol (5:4:3)
	(<i>S</i>)-Dehydrovomifoliol Curdione Curcumol Germacrone Curzerene β -Elemene Patchoulol	Yang et al. (2013) Dang et al. (2010) Li et al. (2011)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:5:1:5) CPC/light petroleum ether–acetonitrile–acetone (4:3:1) CPC/light petroleum ether–acetonitrile (1:1)

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
	Tussilagone	Wang et al. (2011a)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:0.5:1.1:0.3)
	14-Acetoxy-7β-(3'-ethyl <i>cis</i> -crotonoyloxy)-1α-(2'-methyl butyryloxy)-notonipetranone		
	7aβ-(3'-Ethyl <i>cis</i> -crotonoyloxy)-1α-(2'-methyl butyryloxy)-3,14-dehydro- <i>Z</i> -notonipetranone		
	Blumenol C	Roscher and Winterhalter (1993)	CCC/chloroform–methanol–water (7:13:8)
	Solanesol	Hu et al. (2007)	CCC/ <i>n</i> -hexane–methanol (10:7)
	Solanesol	Du et al. (2006)	CCC/light petroleum ether–ethanol–methanol (200:1:100)
	Kudtdiol	Rodrigues et al. (2009)	CCC/ethyl acetate–methanol–water (2:1.75:1)
	8β-Hydroxyeremophil-3,7(11)-dien-12,8α	Shi et al. (2008)	CCC/light petroleum–ethyl acetate–methanol–water (9:1:8:2)
	15,6α-Diolide and 8β-methoxyeremophil-3,7(11)-dien-12,8α;15,6α-diolide		
Diterpenoids	Ptaerobliquol	Agostinho et al. (2013)	CPC/heptane–ethyl acetate–methanol–water (6:5:6:5)
	Coniferin	Slacanin et al. (1991)	CPC/chloroform–methanol–water (7:13:8)
	Coniferaldehyde glucoside		
	Salvinorin A	Shirota et al. (2007)	CPC/ <i>n</i> -hexane–dichloromethane–methanol–water (8:8:9:2)
	Andrographolide	Du et al. (2003a)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:4:2.5:2.5)
	Neoandrographolide		
	Phytol	Xiao et al. (2013)	CCC/ <i>n</i> -hexane–acetonitrile–methanol (5:5:3)
	Carnosol	Fischer et al. (1991)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (70:30:14:8)
	15,16-Epoxy-12-hydroxy-8(17),13(16),14-labdatrien-19-oic acid	Martin et al. (2006)	CPC/chloroform–methanol–isopropanol–water (5:6:1:4)
	Imbricatolic acid		
	Isocupressic acid		
	Sandaracopimaric acid		
	Isopimaric acid		
	Kaurenoic acids	De Souza et al. (2010)	CCC/ <i>n</i> -hexane–acetonitrile–ethyl acetate (1:1:0.4)
	Polyalthic acid		

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
	Pseudolaric acid B <i>O</i> - β -D-glucopyranoside	He et al. (2012)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate–methanol–water (1:1:1:1) and (3:2:2:3) and (3.5:1:1:3.5)
	Pseudolaric acid C		
	Deacetyl pseudolaric acid A		
	Pseudolaric acid A <i>O</i> - β -D-glucopyranoside		
	Pseudolaric acid B		
	Pseudolaric acid B methyl ester		
	Pseudolaric acid A		
	Pseudolaric acid H		
	Oridonin	Lu et al. (2006)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:2:1:2)
	Oridonin	He et al. (2011)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (2.8:5:2.8:5)
	Oridonin	Lu et al. (2007)	2D-CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:5:1:5) in 1st direction and (3:5:3:5) in 2nd direction
	Ponicidin		
	Isonetriptophenolide	Peng et al. (2008a)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (3:2:3:2)
	Hypolide		
	Triptonide		
	Triptophenolide		
	Triptonoterpene methyl ether VI		
	Pseudolaric acids A and B	Han et al. (2009)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (5:5:5:5) for purification of aglicones and (1:9:4:6) for glycosides
	And their glucosides (<i>O</i> - β -D-glucopyranosides)		
	6 β -Angeloyloxykolavenic acid	Wu et al. (2008)	CCC/ <i>n</i> -hexane–ethanol–water (6:5:1)
	6 β -Tigloyloxykolavenic acid		
	Pseudolaric acid B	Han et al. (2008)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (5:5:5:5)
	lolitrem B	Grancher et al. (2004)	CPC/heptane–ethyl acetate–methanol–water (33:33:24:10)
	Carnosic acid	Fischedick et al. (2013b)	CPC/heptane–acetone–water (3:5:2)
	Carnosol		
	Carnosaldehyde		
	Epirosmanol		
	Rosmanol		
	12-Methoxy-carnosic acid		
	Sageone		
	Stevioside	Huang et al. (2010)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (1.5:3.5:5)
	Rebaudioside A		
	Rebaudioside C		

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
	Ginkgolides A Ginkgolides B Ginkgolides C Bilobalide	Liu et al. (2013b)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate– methanol–water (4:5:1:5) and (4:5:2:5)
	Cryptotanshinone Tanshinone I Tanshinone IIA	Gu et al. (2004, 2006)	CCC/stepwise gradient: <i>n</i> -hexane–ethanol–water (10:5.5:4.5) and (10:7:3)
	Cryptotanshinone Tanshinone I 1,2-Dihydrotanshinquinone Tanshinone IIA Tanshinone I Tanshinone IIA	Liang et al. (2013)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (5:5:7:3)
	Cryptotanshinone Tanshinone I Tanshinone IIA Dihydrotanshinone Cryptotanshinone Methylenetanshiquinone Tanshinone I Tanshinone II Danshenxinkun B Tanshinone I Tanshinone IIA Dihydrotanshinone I Cryptotanshinone Przewaquinone A	Wu et al. (2010)	CCC/ <i>n</i> -hexane–ethyl acetate–ethanol–water (8:2:7:3)
	Cryptotanshinone Tanshinone I Tanshinone IIA Dihydrotanshinone Cryptotanshinone Methylenetanshiquinone Tanshinone I Tanshinone II Danshenxinkun B Tanshinone I Tanshinone IIA Dihydrotanshinone I Cryptotanshinone Przewaquinone A	Tian et al. (2000)	CCC/ <i>n</i> -hexane–ethanol– water (4:2:2)
	Cryptotanshinone Tanshinone I Tanshinone IIA Dihydrotanshinone Cryptotanshinone Methylenetanshiquinone Tanshinone I Tanshinone II Danshenxinkun B Tanshinone I Tanshinone IIA Dihydrotanshinone I Cryptotanshinone Przewaquinone A	Li and Chen (2001b)	CCC/stepwise gradient: <i>n</i> -hexane–ethanol–water (10:5.5:4.5) and (10:7:3)
	Cryptotanshinone Tanshinone I Tanshinone IIA Dihydrotanshinone Cryptotanshinone Methylenetanshiquinone Tanshinone I Tanshinone II Danshenxinkun B Tanshinone I Tanshinone IIA Dihydrotanshinone I Cryptotanshinone Przewaquinone A	Tian et al. (2002)	CCC/light petroleum–ethyl acetate–methanol–water (2:3:2.5:1.7)
	Cryptotanshinone Tanshinone I Tanshinone IIA Dihydrotanshinone Cryptotanshinone Methylenetanshiquinone Tanshinone I Tanshinone II Danshenxinkun B Tanshinone I Tanshinone IIA Dihydrotanshinone I Cryptotanshinone Przewaquinone A	Han et al. (2003)	CCC/carbon tetrachloride– methanol–water– <i>n</i> -hexane (3:3:2:1)
	Cryptotanshinone Tanshinone I Tanshinone IIA Dihydrotanshinone Cryptotanshinone Methylenetanshiquinone Tanshinone I Tanshinone II Danshenxinkun B Tanshinone I Tanshinone IIA Dihydrotanshinone I Cryptotanshinone Przewaquinone A	Ye et al. (2008b)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (4:5:4:5)
	Cryptotanshinone Tanshinone I Tanshinone IIA Dihydrotanshinone Cryptotanshinone Methylenetanshiquinone Tanshinone I Tanshinone II Danshenxinkun B Tanshinone I Tanshinone IIA Dihydrotanshinone I Cryptotanshinone Przewaquinone A	Sun et al. (2011)	CCC/light petroleum–ethyl acetate–methanol–water (6:4:6.5:3.5)
	Dihydrotanshinone I 1,2,15,16-Tetrahydrotanshiquinone cryptotanshinone Tanshinone I Tanshinone IIA Neo-przewaquinone A Miltirone		

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
Triterpenoids	2 β ,3 β ,4 β -Trihydroxypregnan-16-one	Rodrigues et al. (2009)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:2:1.75:1)
	Squalene	Lu et al. (2003)	CCC/ <i>n</i> -hexane–methanol (2:1)
	Pristimerin	Gutiérrez et al. (2007)	CPC/heptane–ethyl acetate–methanol–water (8:1:6:1)
	Netzahualcoyene		
	Guyanin	Severino et al. (2009)	CCC/ <i>n</i> -hexane–ethanol–acetonitrile–water (10:8:1:1)
	Barbinervic acid	Fan and He (2006)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (3:6:4:2)
	Rotungenic acid		
	24-Hydroxy ursolic acid		
	Ursolic acid		
	Betulinic acid	Frighetto et al. (2005)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (10:5:2.5:1)
	Ursolic acid	Frighetto et al. (2008)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (10:5:2.5:1)
	Bellericagenin B	Nasser et al. (2006)	CCC/chloroform–methanol–water (43:37:20)
	Bellericaside B		
	Arjunglucoside I		
	28-Nor-17, 22-seco-2 α , 3 β , 19, 22, 23-pentahydroxy- Δ 12-Oleanane		
	2 α ,3 α ,19 β ,23 β -Tetrahydroxyurs-12-en-28-oic acid	Liu et al. (2011)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (10:5:3:1)
	2 α ,3 α ,23 β -Trihydroxyurs-12-en-28-oic acid		
	Oleanolic acid	Du et al. (1995)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (3:6:2:1)
	Ursolic acid		
	Shionone	Wang et al. (2012a)	CCC/ <i>n</i> -hexane–methanol (2:1)
	Taraxasterol acetate	Abbott et al. (1989)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–acetonitrile (5:2:4:5)
	Lupeol acetate		
	β -Amyrin acetate		
	Cycloartenyl ferulate	Liu et al. (2013a)	CCC/ <i>n</i> -hexane–acetonitrile (1:1)
	24-Methylene cycloartanyl ferulate		
	Abrusoside A, B, C, D	Fullas et al. (1990)	CCC/chloroform–methanol–water (7:13:8)
	No compound names given	Marston et al. (1988)	CCC/chloroform–methanol–water (7:13:8)
Inotodiol	Du et al. (2011)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:0.4:1:0.4)	
Trametenolic acid			
Ursolic acid	Maurya and Srivastava (2011)	CPC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:2:1.5:1) with 2 % ammonia solution in lower aqueous mobile phase (pH 9.5)	
Ursolic acid lactone			
Asiaticoside	Diallo et al. (1991)	CCC/chloroform– <i>n</i> -butanol–methanol–water (7:3:6:4)	
Madecassoside			

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
	Taraxeryl acetate	Yang et al. (1995)	CCC/chloroform–methanol–water (2:2:1)
	Triterpenic acid	Ito et al. (1990)	CCC/ <i>n</i> -hexane–ethanol–water (6:5:2)
	Acetyl-triterpenic acid		
	Celastrol	Wu et al. (2004)	CCC/light petroleum–ethyl acetate–tetrachloromethane–methanol–water (1:1:8:6:1)
	Asiatic acid	Du et al. (2004)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–0.05 M NaOH (5:1:6)
	Madecassic acid		
	Asciaticoside		
	Madecassoside		
	Euscaphic acid	Rocha et al. (2007)	CCC/gradient: <i>n</i> -hexane–ethyl acetate–methanol–water (1:2:1.25:2) and (1:2:1.5:2) and (1:2:1.75:2)
	Tormentic acid		
	2 α -Acetyl tormentic acid		
	3 β -Acetyl tormentic acid		
	Alisol B	Yoon et al. (2009)	CPC/ <i>n</i> -hexane–ethyl acetate–methanol–water (10:2:10:7)
	Alisol B 23-acetate		
	Ganoderic acids A, B, C6, D, E, F, G	Cheng et al. (2012)	CCC/stepwise gradient: light petroleum ether–ethyl acetate–methanol–water (3:5:3:5) and (4:5:4:5)
	Ganoderenic acid D		chloroform–methanol–water (13:7:4) + ammonia (22 mM) in aqueous phase and TFA (11 mM) in organic phase for further purification
	Dehydrosulphurenic acid	Zhang et al. (2013a)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:1.5:1.2:1)
	3-Ketodehydrosulphurenic		
	24-Methylene cycloartanol	Yao et al. (2007)	CCC/ <i>n</i> -hexane–ethyl acetate–acetonitrile (5:1:4)
	Lupenone	Yao et al. (2007)	CCC/ <i>n</i> -hexane–ethyl acetate–acetonitrile (5:2:5)
Triterpenoid saponins (ginsenosides)	Rc, Rb1 and Re	Wang et al. (2010)	CPC/ethyl acetate– <i>n</i> -butanol–water (1:1:2)
	Rg3, Rk1, Rg5 and F4	Ha et al. (2007)	CCC/methylene chloride–methanol–water–isopropanol (6:6:4:1)
	Rf, Rd, Re and Rb1	Qi et al. (2010)	CCC/methylene chloride–methanol–5 mM aqueous ammonium acetate–isopropanol (6:2:4:3)
	Re	Engelberth et al. (2010)	CPC/heptane– <i>n</i> -butanol–water (3:4:7)
	Rg1, Re, Rf, Rh1, Rb1, Rc, Rb2 and Rd	Shehzad et al. (2011)	CCC/chloroform–methanol–water–isopropanol (4:3:2:1)
	Re, Rb1, Rc and Rb2	Cheng et al. (2011)	CCC/methylene chloride–methanol–water–isopropanol (6:2:4:3, v/v) further purification of Rb1, Rc and Rb2 in <i>n</i> -hexane– <i>n</i> -butanol–0.1 % formic acid (0.7:3:4)
	Rb1, Rb2, Rc, Rd, Re, Rg1, Rf and Rh 1	Shehzad et al. (2012)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate–methanol–water (5:6:1:4) and (4:3:1:2) and (3:3:1:2)
	Re and Rg1	Chen et al. (2012)	CCC/ethyl acetate– <i>n</i> -butanol–water (4:1:6)
	Rh1, Rf, Rd, Rg1, Re, Rc, Rb2 and Rb1	Shehzad et al. (2013)	CCC/methylene chloride–methanol–water–isopropanol (1:1:2:1)

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
Triterpenoid saponins	Rg6, Rg5, Rk1, F4, Rg3, Rg2, Rf, Rd, Rg1, Re, Rc, Rb2, Rb1	Shehzad et al. (2013)	CCC/stepwise gradient: methylene chloride–methanol–water–isopropanol (5:4:1:3) and (2:2:1:2)
	Rb1 and Rb2	Wang et al. (2013)	CPC/ethyl acetate– <i>n</i> -butanol–water (0.8:1.2:2)
	Rg, Rd, Re and Rb	Cao et al. (2003)	CCC/chloroform–2-butanol–methanol–water (5:1:6:4)
	Notoginsenoside R	Cao et al. (2003)	CCC/ethyl acetate– <i>n</i> -butanol–water (1:1:2)
	Rg, Rd, Re and Rb	Cao et al. (2003)	CCC/ethyl acetate– <i>n</i> -butanol–water (1:1:2)
	Notoginsenoside R	Cao et al. (2003)	CCC/ethyl acetate– <i>n</i> -butanol–water (1:1:2)
	Rg, Re and Rb	Du et al. (2003b)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (3:4:7)
	Notoginsenoside R	Du et al. (2003b)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (3:4:7)
	Ro	Cheng et al. (2010)	CCC/ethyl acetate–isopropanol–0.1 % formic acid (3:1:5)
	Rg1, Re and Rb1	Wang et al. (2011b)	CPC/ethyl acetate– <i>n</i> -butanol–water (1:1:2)
	Notoginsenoside R1	Wang et al. (2011b)	CPC/ethyl acetate– <i>n</i> -butanol–water (1:1:2)
	Lucyoside Q	Du and Gao (2006)	CCC/chloroform–methanol–water (13:7:8)
	Lucyoside H	Du and Gao (2006)	CCC/chloroform–methanol–water (13:7:8)
	Astragaloside I	Han et al. (2007)	CCC/stepwise gradient: ethyl acetate–2-propanol–water (5:1:5) and (50:1:50)
	Astragaloside II	Han et al. (2007)	CCC/stepwise gradient: ethyl acetate–2-propanol–water (5:1:5) and (50:1:50)
	Lancemaside A	Shirota et al. (2008)	CPC/ <i>n</i> -hexane– <i>n</i> -butanol–methanol–0.1 % aqueous formic acid (3:4:1:6)
	Foetidissimoside A	Shirota et al. (2008)	CPC/ <i>n</i> -hexane– <i>n</i> -butanol–methanol–0.1 % aqueous formic acid (3:4:1:6)
	Astersaponin Hb	Shirota et al. (2008)	CPC/ <i>n</i> -hexane– <i>n</i> -butanol–methanol–0.1 % aqueous formic acid (3:4:1:6)
	Saikosaponins-A	Yoon and Kim (2009)	CPC/ethyl acetate– <i>n</i> -butanol–methanol–water (15:1:3:1)
	Saikosaponins-C	Yoon and Kim (2009)	CPC/ethyl acetate– <i>n</i> -butanol–methanol–water (15:1:3:1)
	Platycoside E	Han et al. (2009)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (1:40:20)
	Deapio-platycoside E	Han et al. (2009)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (1:40:20)
	Platycodin D3	Han et al. (2009)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (1:10:5)
	Deapio-platycodin D3	Han et al. (2009)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (1:10:5)
	Platycodin D	Han et al. (2009)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (1:10:5)
	Deapio-platycodin D	Han et al. (2009)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–water (1:10:5)
	2''-O-Acetylplatycodin D	Ha et al. (2011)	CCC/chloroform–methanol–isopropanol–water (3:2:2:3)
3''-O-Acetylpolygalacin D	Ha et al. (2011)	CCC/chloroform–methanol–isopropanol–water (3:2:2:3)	
2''-O-Acetylpolygalacin	Ha et al. (2011)	CCC/chloroform–methanol–isopropanol–water (3:2:2:3)	
3''-O-Acetylplatycodin D	Ha et al. (2011)	CCC/chloroform–methanol–isopropanol–water (3:2:2:3)	
Polygalacin D	Ha et al. (2011)	CCC/chloroform–methanol–isopropanol–water (3:2:2:3)	
No compound names given	Shi et al. (2007)	CCC/ethyl acetate– <i>n</i> -butanol–ethanol–0.05 % TFA (5:10:2:20)	
Astragaloside IV	Peng et al. (2008b)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate–ethanol–water (1:0.6:0.6:1) and (1:1:1:1)	
Astragaloside II	Peng et al. (2008b)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate–ethanol–water (1:0.6:0.6:1) and (1:1:1:1)	
Astragaloside I	Peng et al. (2008b)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate–ethanol–water (1:0.6:0.6:1) and (1:1:1:1)	
Acetylastragaloside I	Peng et al. (2008b)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate–ethanol–water (1:0.6:0.6:1) and (1:1:1:1)	
Gypsogenin derivatives	Yao et al. (2008)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–methanol–0.02 % TFA (1:9:1:9)	
Dianoside C	Yao et al. (2008)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–methanol–0.02 % TFA (1:9:1:9)	

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
	Hederagenin	Xin et al. (2009)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (7:3:5:5)
	3- <i>O</i> -[β-D-xylopyranosyl-(1→3)-α-L-rhamnopyranosyl-(1→2)-L-arabinopyranosyl]-hederagenin		
	23- <i>O</i> -Acetylshengmanol-3- <i>O</i> -D-xylopyranoside	Cicek et al. (2010)	CCC/ <i>n</i> -hexane–acetone–ethyl acetate–isopropanol–ethanol–water (3.5:1:2:1:0.5:2)
	Cimiracemoside D 25- <i>O</i> -acetylcimigenol-3- <i>O</i> -D-Xylopyranoside		
	Cimigenol		
	Esculentosides A, B, C, and D	Ma et al. (2010)	CCC/chloroform–methanol–water (4:4:2)
	Hederagenin		CCC/heptane–acetone–methanol (5:1:4)
	Hederagenin	He et al. (2002)	CPC/ <i>n</i> -hexane–ethyl acetate–methanol–water (7:8:5:3)
	Elatoside F	Lee et al. (2009)	CCC/chloroform–methanol–water–isopropanol (4:3:3:1)
	Tormentic acid	Wang et al. (2012b)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (4:5:4:5)
	Arganine A, C, D	Gosse et al. (2002)	CCC/ <i>tert</i> -butylmethyl ether– <i>n</i> -butanol–acetonitrile–0.5 % TFA (1:3:3:5)
	Tieghemelin		
	Triacetyl soyasaponin Ab, Aa, Ab, Ae, Ba, Af, Bb, Be and conjugated groups, αg, βg and γg	Zhao et al. (2012)	CCC/ <i>n</i> -butanol–acetic acid–water (5:0.05:5)–acid in <i>n</i> -butanol as stationary phase
	Licorice-saponin A3	Xu et al. (2013)	CCC/ethyl acetate– <i>n</i> -butanol–water (2:3:5) with 10 mM TFA in the upper organic
	Glycyrrhizic acid, 3- <i>O</i> -[β-D-glucuronopyranosyl-(1→2)-β-D-Galactopyranosyl]glycyrrhetic acid		Stationary phase and 10 mM ammonia in the lower aqueous mobile phase
	Glycyrrhizin	Jiang et al. (2004)	CCC/ethyl acetate–methanol–water (5:2:5)
	Goyaglycoside-E	Du and Yuan (2005)	CCC/ <i>tert</i> -butylmethyl ether– <i>n</i> -butanol–methanol–water (1:2:1:5) or (1:3:1:5)
	Momordicoside L		
	Goyaglycoside-a		
	Momordicoside K		
Tetranortriterpenoids	Azadirachtin A	Silva et al. (2007)	CCC/ <i>n</i> -hexane– <i>n</i> -butanol–methanol–water (1:0.9:1:0.9)
	Azadirachtin B		
	Azadirachtin H		
	Desacetylnimbin		
	Desacetylsalannin		
	Nimbin		
	Salannin		
	Methyl angolensate	da Silva et al. (2009)	CCC/stepwise gradient: <i>n</i> -hexane–ethyl acetate–methanol–water (2:1:1.5:1) and (2:1:1.75:1)
	7-Desacetoxy-7-oxogedunin		
	Deacetylgedunin		
	6α-Acetoxygedunin		
	Gedunin		
	Andirobin		

Table 2 continued

Classes of compounds	Purified compounds	References	Type of apparatus/solvent system
Tetraterpenoids (Carotenoids)	Cochloxanthin	Diallo and Vanhaelen (1988)	CCC/tetrachloromethane–methanol–water (5:4:1)
	Dihydrocochloxanthin		
	Zeaxanthin	Chen et al. (2005)	CCC/ <i>n</i> -hexane–ethyl acetate–ethanol–water (8:2:7:3)
	Crocin	Jiang et al. (2011)	CCC/ <i>tert</i> -butylmethyl ether– <i>n</i> -butanol–acetonitrile–water (2:2.5:1:5)
	Fucoxanthin	Kim et al. (2011)	CPC/ <i>n</i> -hexane–ethyl acetate–ethanol–water (5:5:7:3)
	Lutein	Wei et al. (2003)	CCC/ <i>n</i> -heptane–chloroform–acetonitrile (10:3:7)
	Zeaxanthin	Aman et al. (2005)	CCC/ <i>n</i> -hexane–ethanol–water (6:5:1.3)
	Lutein		
	Lutein	Tsao and Yang (2006)	CCC/ <i>n</i> -hexane–ethanol–water (6:4.5:1.5)
	Canthaxanthin	Li et al. (2006)	CCC/ <i>n</i> -hexane–ethanol–water (10:9:1)
	9'- <i>Cis</i> -neoxanthin	Baldermann et al. (2007)	CCC/ <i>n</i> -hexane–ethanol–water (5:5:4.5)
	Lycopene	Baldermann et al. (2008)	CCC/ <i>n</i> -hexane–dichloromethane–acetonitrile (30:11:18) with 85 mg/L of 3- <i>tert</i> -butyl-4-hydroxyanisole and 2- <i>tert</i> -butyl- <i>p</i> -cresol
	Crocins 1, 2, 5	Lechtenberg et al. (2008)	CPC/ <i>n</i> -hexane–ethyl acetate–ethanol–water (1:3:4:7)
	Picrocrocin		
	Fucoxanthin	Xiao et al. (2012)	CCC/ <i>n</i> -hexane–ethyl acetate–ethanol–water (5:5:6:4)
	Lutein	Li et al. (2001)	CCC/ <i>n</i> -hexane–ethanol–water (4:3:1)
	Lycopene	Wei et al. (2001)	CCC/ <i>n</i> -hexane–dichloromethane–acetonitrile (10:3.5:6.5)
	Astaxanthin	Li and Chen (2001a)	CCC/ <i>n</i> -hexane–ethyl acetate–ethanol–water (5:5:6.5:3)
	Tephrosin	Ye et al. (2008a)	CCC/ <i>n</i> -hexane–ethyl acetate–methanol–water (1:0.8:1:0.6)
	Deguelin		
6a,12a-Dehydrodeguelin			

CCC counter-current chromatography, CPC centrifugal partition chromatography

trachelosperogenin E and sericoside was achieved using a three-phase solvent system composed of *n*-heptane, *tert*-butylmethyl ether, acetonitrile and water. After mixing all solvents in equal volume, the upper phase was separated and *tert*-butylmethyl ether was shaken with the remaining two-phase solvent system in order to decrease the polarity. The lower phase was used as a stationary phase. Sericic acid was identified in the fractions collected when the upper phase was used as the mobile phase, while the elution of the middle phase led to the separation of trachelosperogenin E and sericoside (Hamzaoui et al. 2013).

Zhao and Du (2007) proposed a novel non-aqueous two-phase solvent system composed of sunflower oil and ethanol to separate solanesol—a non-cyclic terpene alcohol used as a food additive for preventing cardiac arrest and cancer. Before, this target molecule was purified from tobacco leaves extract with petroleum and methanol, both solvents being unsuitable for the food industry.

Sometimes the addition of a small amount of acid can significantly improve the purification. Shirota et al. (2008) purified the saponins lancemaside A, foetidissimoside A and astersaponin Hb from a hot

water extract of *Codonopsis lanceolata* roots used CPC with mixture of *n*-hexane–*n*-butanol–methanol–0.1 % aqueous formic acid (3:4:1:6) as the two-phase solvent system. Because the crude extract contained large amounts of sugars, an aqueous phase was chosen as the stationary phase so that these sugars would be retained in the partition cells, whereas an organic phase was chosen as mobile phase for elution of the target compound. Addition of formic acid to the two-phase solvent system retained the carboxy group in the non-ionised form and allowed the separation of the lincemasides from the large amounts of water-soluble sugars present. The addition of acetic acid to the ethyl acetate–*n*-butanol–water mixture improved the resolution of soyasaponins, naturally occurring triterpenoid glycosides. A suitable mixture *n*-butanol, acetic acid, water (5:0.05:5) was taken but acetic acid was added only in *n*-butanol as the stationary phase (Zhao et al. 2012). During the separation, with the movement of the mobile phase the acetic acid was gradually diluted with water and the polarity of the stationary phase changed as a gradient. It is worth mentioning that the main difficulties with soyasaponin isolation and purification are that the soyasaponins coexist in soybeans with the isoflavone glycosides and they share overlapping polarities. There is also the structural similarity of soyasaponin compounds.

CCC is popularly applied for purification of ginsenosides, an important group of terpenoids due to their biological activity. They generally fall in the moderate to polar category and therefore require similar solvent systems for purification (Qi et al. 2010; Shehzad et al. 2013). A good solubility for saponin-like compounds can be provided by the addition of chloroform or other chlorinated solvents and a gradual polarity change between phases can be achieved by varying the methanol ratio in the system or by adding another alcohol like isopropanol (Qi et al. 2010). Shehzad et al. (2013) proposed an efficient CCC separation method in which a flow-rate gradient technique was coupled with a new solvent gradient dilution strategy for the isolation of ginsenosides from Korean red ginseng (steam-treated *P. ginseng*). The column was first entirely filled with the upper stationary phase mixture of methylene chloride–methanol–isopropanol–water (5:4:1:3) in a reversed phase system. After 300 min, when five ginsenosides had eluted, the flow rate was increased from 1 to 1.2 ml/min and also the dilution of the lower phase was initiated and was changed to

100 % of lower phase composed of the mixture mentioned above in a ratio (2:2:1:2). Overall, 13 ginsenosides including Rg1, Re, Rf, Rg2, Rb1, Rb2, Rc, Rd, Rg3, Rk1, Rg5, Rg6, and F4 were purified (Shehzad et al. 2013). Qi et al. (2010) for the purification of ginsenosides Rf, Rd, Re and Rb1, applied mixtures of methylene chloride, methanol, water and isopropanol (6:2:4:3).

Emulsification can often be a problem in the CCC purification of ginsenosides. Generally, adding an electrolyte, such as salt or acid, can help to eliminate such emulsification. Salt is not recommended because an additional desalting process will be necessary after the separation. Therefore, formic acid is often chosen to prevent emulsification (Cheng et al. 2010). However, Qi et al. (2010) preferred to avoid an acidic environment, which would lead to the decomposition of ginsenosides. Instead, the inorganic salt ammonium acetate was chosen, because it is volatile and can be precipitated in warm acetone for sample recovery. The addition of this salt resulted in a very slight decrease in *K* values and thus a shorter separation time. In the end, the proposed solvent system was methylene chloride–methanol–5 mg/mL aqueous ammonium acetate–isopropanol (6:2:4:3).

As the solvent system contained methylene chloride and methanol, both toxic to humans and the environment, Wang et al. (2010) chose low toxic solvents such as ethyl acetate and *n*-butanol. Since the ginsenosides in *P. quinquefolium* L. have a comparatively larger polarity with solubility in hydrophilic solvents, both phases require a certain hydrophilicity to get a good separation. Initially, the hexane–*n*-butanol–water solvent systems were scanned at several volume ratios, but all the systems had a poor retention in the CPC column. Then ethyl acetate was then employed instead of hexane in order to enhance the retention of the stationary phase in the column. Because ginsenosides are easily dissolved in *n*-butanol, and the viscosity of *n*-butanol is comparatively large, so in an ethyl acetate–*n*-butanol–water solvent system, the addition of *n*-butanol could delay the peak elution time and affect the separation. Overall, the system of ethyl acetate–*n*-butanol–water (1:1:2) was used for successful separation of three ginsenosides Rc, Rb1, and Re.

Counter-current chromatography has also been used for the separation of minor and structurally similar compounds. Fan and He (2006) using

HSCCC with *n*-hexane–ethyl acetate–methanol–water (3:6:4:2) to separate not only ursolic acid, which is a very small content of the leaves of *Diospyros kaki*, but also two other pentacyclic triterpenes: barbinervic acid and its epimer rotungenic acid, differing only with the configuration of a hydroxyl group at position C3: one contains an axial and the other an equatorial hydroxyl group. HSCCC was also employed for the separation and purification of minor constituents in *Platycodi Radix*. Platycosides, the saponins that are the major active constituents, are typically composed of oleanene backbones with two side chains; one a glucose unit attached through an ether linkage at the C-3 position of a triterpene, and the other 28-*O*-arabinoserhamnose-xylose-apiose linked by an ester bond. They also have different substituents at the C-4 position. Because the content of these compounds is very low, conventional methods are frequently not suitable for the separation. Ha et al. (2011) used a two-phase solvent system consisting of chloroform–methanol–isopropanol–water (3:2:2:3) for purification of minor saponins 2''-*O*-acetylplatycodin D, 3''-*O*-acetyl polygalacin D, 2''-*O*-acetyl polygalacin and a mixture of 3''-*O*-acetylplatycodin D and polygalacin D, which were further successfully purified using prep-HPLC (Ha et al. 2011). Ha and Kim (2009) also used HSCCC for the separation of three pairs of platycosides and their deapiose forms. They used an interesting modification: the column was first filled with a mixture of the two phases, thus reducing the amount of time for hydrodynamic equilibrium to be established. The ratio of two phases was optimised at 70:30 (stationary phase–mobile phase) based on the amount of time required to reach hydrodynamic equilibrium. A series of solvent systems were tested. In the gradient elution mode, the retention of the stationary phase was extremely low. The authors decided to perform the separation in two stages. First, platycoside E, deapioplatycoside E, a mixed fraction containing platyodin D and deapiose form, and a fraction containing platycodin D3 and deapiose form were separated using the *n*-hexane–butanol–water (1:40:20) in reversed-mode. Then mixed fractions I and II were further purified in the normal elution mode with the above mentioned solvent system in the ratio 1:10:5.

CCC/CPC seems to be one of the few efficient approaches for the separation of xanthanolides, a bicyclic subtype of sesquiterpenic lactones characterized by a 5,7-fused system containing a γ -lactone

moiety. Because of some well documented and promising activity purification of these compounds is important. The presence of several chiral centres probably explain the difficulty obtaining enantiomerically pure xanthanolides by total synthesis, thus purification from natural source seems to be the best solution. Xanthanolides are coextracted with chlorophyll and lipids. The pigment crystallization and the delipidation in order to clean the extract is not selective and cause the partial loss of target compounds. Purification of several xanthanolides (xanthanin, 4-*epi*-xanthanol and 4-*epi*-isoxanthanol) was realized in one step, directly from the crude chloroformic extract of the leaves of *X. macrocarpum* with a mixture of heptane–ethyl acetate–methanol–water (1:1:1:1) (Pinel et al. 2007).

High-speed counter-current chromatography can be applied as a method suitable for fingerprinting. Gu et al. (2006) used it for quality control of TCMs and identification of the active compounds of *Salvia miltiorrhiza* Bunge, a popular traditional Chinese medicine. In order to purify a series of tanshinones a stepwise elution with solvent systems composed of *n*-hexane–ethanol–water (10:5.5:4.5) and (10:7:3) was used. The method was compared with more conventional approaches, such as high performance liquid chromatography (HPLC), high performance capillary electrophoresis (HPCE), and thin-layer chromatography scan (TLCS). In the HSCCC separation, 12 components were separated, with good resolution and precision, within 13 h. HSCCC showed better performance in the analysis of tanshinones, which produced a fingerprint which contained more chemical information than that of e.g. TLCS.

Lu et al. (2007) proposed this effective two-dimensional counter-current chromatographic method for the simultaneous isolation and purification of oridonin and ponocidin from a crude extract of *Rabdosia rubescens* with a pair of two-phase solvent systems composed of *n*-hexane–ethyl acetate–methanol–water (1:5:1:5 and 3:5:3:5, v/v). A combination of stepwise CCC and pH-zone-refining is also possible. Cheng et al. (2012) reported its successful combination in the separation of the main components from *G. lucidum*. In the first step, a two-phase solvent system composed of petroleum ether–ethyl acetate–methanol–water (3:5:3:5 and 4:5:4:5) led to the separation of ganoderic acids GE, GC6 and GF with high purity in one run. Also two peaks containing GG, GB, GA and

GED, GD, respectively, were collected and their separation was followed by pH-zone-refining CCC. Chloroform–methanol–water (13:7:4) with NH_4OH in upper aqueous stationary phase and trifluoroacetic acid as the eluter acid was the most suitable. In another example of two-dimensional CCC, a three-step gradient elution and two-step flow-rate gradient elution was applied to separate 8 diterpene compounds within 80 min in a single run from the alcohol extract of *Pseudolarix kaempferi*. Pseudolaric acid B *O*- β -D-glucopyranoside, pseudolaric acid C, deacetyl pseudolaric acid A, pseudolaric acid A *O*- β -D-glucopyranoside, pseudolaric acid B, pseudolaric acid B methyl ester, pseudolaric acid A and pseudolaric acid H were obtained with very high purity (He et al. 2012). The separation was performed in normal phase system. In the first step, the mobile phase, composed of hexane–ethyl acetate–methanol–water (1:1:1:1), was used with a flow rate 0.5 ml/min. While the mobile phase (upper) of (2:3:2:3) and (1:3.5:1:3.5) were used in the second and third step, and the flow rate of mobile phase was set at 1 ml/min. When the separation was performed in preparative conditions, the initial flow rate of mobile phase was 25 ml/min in the first step and then increased to 50 ml/min in the second and third step.

Principle advantages and drawbacks of CCC in the purification of natural products

Counter-current chromatography has a number of key advantages in the purification of natural products. For example, the solvent usage is generally far lower than that of solid phase chromatography systems operating at the same scale (about 25 %) (Graham et al. 2001). Furthermore, since the process is frequently an isocratic one, a simple analysis of solvent composition allows the recycling of the solvents, reducing the usage still further (Garrard et al. 2007). The technique also allows for 100 % recovery of the sample components. In other words, the target compound can always be retrieved since there is no solid phase and therefore no possibility of losses arising from irreversible adsorption onto the solid matrix. This is a significant advantage in every purification process. As a solid-free and therefore relatively gentle chromatographic technique, CCC can be used for the isolation of unstable natural compounds. Baldermann et al. (2007)

presented the successful purification of 9'-*cis*-neoxanthin, the predominant isomer of neoxanthin in green vegetables. Major problems during its isolation include isomerization and oxidation, mainly caused by higher temperatures, light or oxygen exposure. When typical solid stationary phase techniques are applied, the rearrangement products can be detected or complete isomerization can be observed. A solid-free technique is a very practical solution for avoiding the above mentioned complications.

Particulates, such as cell debris, are generally well tolerated in CCC, particularly when performed at large scale where the tubing bore may be up to 10 mm in diameter. This is another major advantage over solid phase chromatography systems. Thus filtering a sample is frequently not necessary, depending on the scale of CCC employed, and even if it is required, it is usually a simple filter paper filtration. With processing times similar to that of other purification methods, scale up is also possible with modern instruments with examples existing running from milligram to kilogram levels (Garrard et al. 2008). The technique can be operated in normal batch injection mode, or as a continuous extraction process for better throughput. A wide range of polarities can be processed due to the range of solvents that may be used (the literature reports examples with a logP range from -4.7 (colistin peptide antibiotic) (Ikai et al. 1998) to $+17.6$ (lycopene) (Wei et al. 2001). Also, the separation of compounds with vastly different polarities from a single extract is possible. For example, in order to purify a wide range of polarity of triterpene saponins from *Panax notoginseng*, Zhang et al. (2013b) successfully coupled accelerated solvent extraction (ASE) and high-performance counter-current chromatography (HPCCC). First the upper phase of the solvent system ethyl acetate–*n*-butanol–water (1:1:2 or 1.2:1:2) or ethyl acetate–*n*-butanol–methanol–water (3:5:1.5:6) was used as both the ASE solvent and HPCCC stationary phase. The polar saponins were eluted. In order to separate fractions with moderate polarity, the upper phase of system ethyl acetate–*n*-butanol–methanol–water (6:3:2:6 or 7:3:2:7) was used. Finally, the upper phase of the solvent system of *n*-hexane–*n*-butanol–methanol–water (8:2:2:8) or *n*-hexane–ethyl acetate–*n*-butanol–methanol–water (0.2:10:0.5:1.5:8) was used as both the ASE solvent and HPCCC stationary phase to elute the low polar compounds. This combination of methods allowed the

purification of notoginsenosides R6, R1, Spt1 and ginsenosides Rb1, F4, Rh3, Rg3, Rs3 and Rk1 with a wide range of polarity (Zhang et al. 2013b).

Compared to the early instruments, the quality of modern CCC apparatus is very good. In most cases the coils are tough and the machinery robust. A set of coils would be expected to last the lifetime of the centrifuge and maintenance and running costs are low. Unlike solid phase chromatography, there is no change to component retention over time (no column aging effects) as a freshly-filled coil of solvents is used each run. This makes it easier to consistently satisfy current regulatory requirements when performing purifications under a good laboratory practice (GLP) or good manufacturing practice (GMP) environment.

As has been seen in a number of quoted examples, a large advantage of CCC is that the technique can be operated in a number of different modes, since both the mobile and the stationary phase are liquid. Either normal or reverse phase chromatography can be chosen, depending on which solvent phase is selected to be the mobile one. However, it is even possible to switch from normal phase elution to reverse phase (or vice versa) in the middle of a run. Intermittent counter-current extraction (ICCE) is a continuous process where the operation alternates between normal and reverse phase mode at regular intervals, with the sample continuously introduced in the middle of the coil (Hewitson et al. 2009). In the reference quoted, this technique was successfully used to purify a diterpenoid, triptolide, a high value target compound, from a Chinese herbal plant, from 2 % in the crude extract to over 98 % purity. This was achieved by retaining and enriching the target compound within the CCC coil while washing away all the other components of the crude material. Alternatively, components can be recovered by eluting the liquid stationary phase without any compound losses whilst maintaining resolution (Berthod et al. 2003), a technique known as elution-extrusion. This technique is frequently adopted at the end of standard CCC purification runs simply to ensure that no loss of target compound has occurred within the liquid stationary phase. Another possibility is co-current chromatography (Berthod and Hassoun 2006) where both phases are pumped in the same direction. All of these options use a conventional CCC centrifuge and are thus easy to implement in the laboratory or pilot plant. However, with modifications to the CCC coils,

continuous counter-current extraction (CCCE) is possible, where one phase is pumped in the opposite direction to the other and the sample is introduced continuously into the centre of the coil (Ito et al. 2006; Van den Heuvel 2007).

The method also allows a two dimensional procedure to be applied. The purification of very similar terpene lactones from *G. biloba* L, such as bilobalide, ginkgolides A, B, C, and J, is an example. The partitioning experiment assisted the design of a 2D procedure using a pair of orthogonal solvent systems: chloroform–methanol–water (10:7:3) and hexane–ethyl acetate–methanol–water (4:6:4:6) with addition of 0.5 % DMSO to increase the resolution of ginkgolides A and B. This approach separated these almost equipolar lactones (Qiu et al. 2012).

The main drawbacks of the technique, particularly when compared to preparative HPLC and other solid phase chromatography techniques, include a lower efficiency. When measured in terms of theoretical plates, the efficiency of a “good” CCC apparatus is in the low thousand plate range. This figure cannot be directly compared to HPLC plate counts due to the much higher percentage of stationary phase (around 80 % compared to perhaps 5 % active stationary phase sites in HPLC) and the resolution of CCC can be extremely good. However, a low efficiency results in broad peaks, making the technique far more suited to a preparative application as opposed to an analytical one. Also the narrow polarity range within each run should be emphasised. As mentioned above, a wide range of polarities can be processed by CCC by using different solvent systems. However, a single CCC run operates over a relatively narrow polarity window. Although some examples of gradient elution have been reported (see Table 2), these do not give as wide a polarity range as that achievable with HPLC. The narrow polarity window can be used to advantage when it is desired to pluck a single target compound from a complex mixture. However, it is a disadvantage when a dozen pure compounds are required from the mixture in a single purification run. The CCC apparatus does not inherently lend itself to easy automation and thus its operation can appear labour-intensive. In addition, the instruments have not received the same intensive commercial development that has made modern HPLC and GC equipment so sophisticated. This situation will undoubtedly improve in the future, with advances in CCC machine design. Finally, the

solvent system selection is undoubtedly time-consuming. As the two phases are liquid, changes in one phase directly affects the other. Analysis of a range of possible solvent systems can be performed by an automated liquid-handling robot but this still requires a number of hours to complete. This is a current area of research in the field and no doubt large improvements in solvent system selection time can be expected over the next few years.

Conclusions

Although initially dismissed by many chemists and purification scientists as slow, unreliable and temperamental, steady development of the technique of CCC on both the engineering and the application side has transformed it into a technique worthy of inclusion in the natural product scientist's arsenal. On one side, engineering developments have produced machines that are robust, capable of fast, efficient separations and able to accept high injection loadings. On the other side, developments in the application protocols have produced modes of operation and solvent systems to purify out compounds from the full polarity spectrum. Combined, these have produced a technique that is wonderfully suited to natural product purifications, particularly on a large preparative scale, with advantages over solid phase techniques such as the ability to accept particulates and to always recover all components, and advantages over the old-style liquid–liquid techniques such as high speed, high loading and high resolution. To assist users of CCC and those wishing to experiment with the technique, a comprehensive phase selection table has been generated by distilling the solvent system information presented from CCC terpenoids purifications over the last 30 years.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Abbott T, Peterson R, McAlpine J et al (1989) Comparing centrifugal countercurrent chromatography, nonaqueous reversed phase HPLC and AG ion exchange HPLC for the separation and characterization of triterpene acetates. *J Liq Chromatogr Relat Technol* 12:2281–2301
- Acton N, Klayman DL, Rollman IJ et al (1986) Isolation of artemisinin (qinghaosu) and its separation from artemisinin using the Ito multilayer coil separator-extractor and isolation of arteannuin B. *J Chromatogr* 355:448–450
- Agostinho D, Boudesocque L, Thery-Kone I et al (2013) A new meroterpenoid isolated from roots of *Ptaeroxylon obliquum* Radlk. *Phytochem Lett* 6:560–566
- Ajikumar PK, Tyo K, Carlsen S et al (2008) Terpenoids: opportunities for biosynthesis of natural product drugs using engineered microorganisms. *Mol Pharm* 5:167–190
- Aman R, Carle R, Conrad J et al (2005) Isolation of carotenoids from plant materials and dietary supplements by high-speed counter-current chromatography. *J Chromatogr A* 1074:99–105
- Baldermann S, Reinhard A, Köhler N et al (2007) Application of high-speed counter-current chromatography for the isolation of 9'-cis-neoxanthin from fresh spinach. *J Chromatogr A* 1151:183–186
- Baldermann S, Ropeter K, Köhler N et al (2008) Isolation of all-trans lycopene by high-speed counter-current chromatography using a temperature-controlled solvent system. *J Chromatogr A* 192:191–193
- Berthod A, Hassoun M (2006) Using the liquid nature of the stationary phase in countercurrent chromatography: IV. The co-current CCC method. *J Chromatogr A* 1116:143–148
- Berthod A, Ruiz-Angel MJ, Carda-Broch S (2003) Elution-extrusion countercurrent chromatography: use of the liquid nature of the stationary phase to extend the hydrophobicity window. *Anal Chem* 75:5508–5517
- Brown GD (2010) The biosynthesis of artemisinin (Qinghaosu) and the phytochemistry of *Artemisia annua* L. (Qinghao). *Molecules* 15:7603–7698
- Cao X, Tian Y, Zhang TY, Liu QH et al (2003) Separation of dammarane-saponins from notoginseng, root of *Panax notoginseng* (Burk.) F.H.Chen by HSCCC coupled with evaporative light scattering detector. *J Liq Chromatogr Relat Technol* 26:1579–1591
- Chen F, Lu HT, Jiang Y (2004) Purification of paeoniflorin from *Paeonia lactiflora* Pall. by high speed counter current chromatography. *J Chromatogr A* 1040:205–208
- Chen F, Li HB, Wong R et al (2005) Isolation and purification of the bioactive carotenoid zeaxanthin from the microalga *Microcystis aeruginosa* by high-speed counter-current chromatography. *J Chromatogr A* 1064:183–186
- Chen Q, Hu X, Li J et al (2011) Preparative isolation and purification of cuminaldehyde and p-mentha-1,4-dien-7-ol from the essential oil of *Cuminum cyminum* L. by high-speed counter-current chromatography. *Anal Chim Acta* 689:149–154
- Chen F, Luo J, Kong L (2012) Fast isolation of ginsenosides Re and Rg1 from the roots of *Panax ginseng* by HSCCC-ELSD combined with MCI gel CC guided by HPLC-MS. *J Liq Chromatogr Relat Technol* 35:912–923
- Cheng Y, Liang Q, Hu P et al (2010) Combination of normal-phase medium-pressure liquid chromatography and high-performance counter-current chromatography for preparation of ginsenoside-Ro from *Panax ginseng* with high recovery and efficiency. *Sep Purif Technol* 73:397–402
- Cheng Y, Zhang M, Liang Q et al (2011) Two-step preparation of ginsenoside-Re, Rb1, Rc and Rb2 from the root of *Panax*

- ginseng by high-performance counter-current chromatography. *Sep Purif Technol* 77:347–354
- Cheng CR, Li YF, Xu PP (2012) Preparative isolation of triterpenoids from *Ganoderma lucidum* by counter-current chromatography combined with pH-zone-refining. *Food Chem* 130:1010–1016
- Cicek SS, Schwaiger S, Ellmerer EP et al (2010) Development of a fast and convenient method for the isolation of triterpene saponins from *Actaea racemosa* by high-speed countercurrent chromatography coupled with evaporative light scattering detection. *Planta Med* 76:467–473
- da Silva VP, Oliveira RR, Figueiredo MR (2009) Isolation of limonoids from seeds of *Carapa guianensis* Aublet (Meliaceae) by high-speed countercurrent chromatography. *Phytochem Anal* 20:77–81
- Dang YY, Li XC, Zhang QW et al (2010) Preparative isolation and purification of six volatile compounds from essential oil of *Curcuma wenyujin* using high-performance centrifugal partition chromatography. *J Sep Sci* 33:1658–1664
- De Souza PA, Rangel LP, Oigman SS et al (2010) Isolation of two bioactive diterpenic acids from *Copaifera glycyarpa* oleoresin by high-speed counter-current chromatography. *Phytochem Anal* 21:539–543
- Diallo B, Vanhaelen M (1988) Large scale purification of apocarotenoids from *Cocleosperrum tinctorium* by counter-current chromatography. *J Liq Chromatogr* 11:227–231
- Diallo B, Vanhaelen Fastre R, Vanhaelen M (1991) Direct coupling of high speed counter current chromatography to thin layer chromatography. Application to the separation of asiaticoside and madecassoside from *Centella asiatica*. *J Chromatogr* 558:446–450
- dos Santos BCB, da Silva JCT, Guerrero PG Jr et al (2009) Isolation of chavibetol from essential oil of *Pimenta pseudocaryophyllus* leaf by high-speed counter-current chromatography. *J Chromatogr A* 1216:4303–4306
- Du Q, Gao S (2006) Preparative separation of saponins from the *Luffa cylindrica* (L.) Roem. by slow rotary countercurrent chromatography. *J Liq Chromatogr Relat Technol* 29:2451–2456
- Du Q, Yuan J (2005) Preparation of triterpene saponins from the fruit of *L.* by high speed countercurrent chromatography (HSCCC). *J Liq Chromatogr Relat Technol* 28:1717–1724
- Du Q, Xiong X, Ito Y (1995) Separation of bioactive quadri-terpenic acids from the fruit of *Ligustrum lucidum* Ait by high-speed countercurrent chromatography. *J Liq Chromatogr* 18:1997–2004
- Du Q, Jerz G, Waibel R et al (2003a) Isolation of dammarane saponins from *Panax notoginseng* by high speed counter current chromatography. *J Chromatogr A* 1008:173–180
- Du Q, Jerz G, Winterhalter P (2003b) Separation of andrographolide and neoandrographolide from the leaves of *Andrographis paniculata* using high speed counter current chromatography. *J Chromatogr A* 984:147–151
- Du Q, Jerz G, Chen P et al (2004) Preparation of ursane triterpenoids from *Centella asiatica* using high speed countercurrent chromatography with step-gradient elution. *J Liq Chromatogr Relat Technol* 27:2201–2215
- Du Q, Daijie W, Ito Y (2006) Preparation of solanesol from a tobacco leaf extract using high speed countercurrent chromatography. *J Liq Chromatogr Relat Technol* 29:2587–2592
- Du D, Zhu F, Chen X et al (2011) Rapid isolation and purification of inotodiol and trametenolic acid from *Inonotus obliquus* by high-speed counter-current chromatography with evaporative light scattering detection. *Phytochem Anal* 22:419–423
- Engelberth AS, Clausen EC, Carrier DJ (2010) Comparing extraction methods to recover ginseng saponins from American ginseng (*Panax quinquefolium*), followed by purification using fast centrifugal partition chromatography with HPLC verification. *Sep Purif Technol* 72:1–6
- Fan JP, He CH (2006) Single-step preparative separation of barbinervic acid and its epimer (rotungenic acid), along with two other pentacyclic triterpene acids from the leaves of *Diospyros kaki* using HSCCC. *J Liq Chromatogr Relat Technol* 29:815–826
- Fischedick JT, Standiford M, Johnson DA et al (2012) Activation of antioxidant response element in mouse primary cortical cultures with sesquiterpene lactones isolated from *Tanacetum parthenium*. *Planta Med* 78:1725–1730
- Fischedick JT, Pesic M, Podolski-Renic A et al (2013a) Cytotoxic activity of sesquiterpene lactones from *Inula britannica* on human cancer cell lines. *Phytochem Lett* 6:246–252
- Fischedick JT, Standiford M, Johnson DA et al (2013b) Structure activity relationship of phenolic diterpenes from *Salvia officinalis* as activators of the nuclear factor E2-related factor 2 pathway. *Bioorgan Med Chem* 21:2618–2622
- Fischer N, Weinreich B, Nitz S et al (1991) Applications of high speed counter current chromatography for the separation and isolation of natural products. *J Chromatogr* 538:193–202
- Fisher D, Garrard IJ, van den Heuvel R et al (2005) Technology transfer and scale up of a potential cancer preventative plant dynamic extraction of glucoraphanin. *J Liq Chromatogr Relat Technol* 28:1913–1922
- Frighetto N, Welendorf RM, Pereira da Silva AM et al (2005) Purification of betulinic acid from *Eugenia florida* (Myrtaceae) by high-speed counter-current chromatography. *Phytochem Anal* 16:411–414
- Frighetto RTS, Welendorf RM, Nigro EN et al (2008) Isolation of ursolic acid from apple peels by high speed counter-current chromatography. *Food Chem* 106:767–771
- Fullas F, Choi YH, Kinghorn D et al (1990) Sweet-tasting triterpene glycoside constituents of *Abrus fruticulosus*. *Planta Med* 56:332–333
- Garrard IJ (2005) Simple approach to the development of a CCC solvent selection protocol suitable for automation. *J Liq Chromatogr Relat Technol* 28:1923–1935
- Garrard IJ, Janaway L, Fisher D (2007) Minimising solvent usage in high speed, high loading and high resolution isocratic dynamic extraction. *J Liq Chromatogr Relat Technol* 30:151–163
- Garrard IJ, Fisher D, Sutherland IA (2008) Dynamic extraction: a high speed, high capacity purification process that is rapidly scaleable. *LC/GC N Am* 26:2–7
- Gasper MP, Berthod A, Talabardon K et al (1995) An evaluation of the differential partitioning and separation of C60 and C70 fullerenes in a biphasic system using centrifugal partition chromatography. *J Liq Chromatogr Relat Technol* 18:1019–1034

- Geng Y, Liu J, Lv R et al (2007) An efficient method for extraction, separation and purification of eugenol from *Eugenia caryophyllata* by supercritical fluid extraction and high-speed counter-current chromatography. *Sep Purif Technol* 57:237–241
- Gosse BK, Gnabre JN, Ito Y et al (2002) Isolation of saponins with viral entry inhibitory activity by combined chromatographic methods. *J Liq Chromatogr Relat Technol* 25:3199–3211
- Graham AS, McConvey IF, Shering P (2001) An evaluation of the performance of a preparative CCC machind for the separation of an active pharmaceutical ingredient. *J Liq Chromatogr Relat Technol* 24:1811–1825
- Grancher D, Jaussaud P, Durix A et al (2004) Countercurrent chromatographic isolation of lolitrem B from endophyte infected ryegrass (*Lolium perenne* L) seed. *J Chromatogr A* 159:73–81
- Graziose R, Rathinasabapathy T, Lategan C et al (2011) Antiplasmodial activity of aporphine alkaloids and sesquiterpene lactones from *Liriodendron tulipifera* L. *J Ethnopharmacol* 133:26–30
- Gu M, Zhang G, Su Z et al (2004) Identification of major active constituents in the fingerprint of *Salvia miltiorrhiza* Bunge developed by high speed counter current chromatography. *J Chromatogr A* 1041:239–243
- Gu M, Su Z, Ouyang F (2006) Fingerprinting of *Salvia miltiorrhiza* Bunge by thin-layer chromatography scan compared with high speed countercurrent chromatography. *J Liq Chromatogr Relat Technol* 29:1503–1514
- Gutiérrez F, Estévez-Braun A, Ravelo AG et al (2007) Terpenoids from the medicinal plant *Maytenus ilicifolia*. *J Nat Prod* 70:1049–1052
- Ha YW, Kim YS (2009) Preparative isolation of six major saponins from *Platycodi radix* by high-speed counter-current chromatography. *Phytochem Anal* 20:207–213
- Ha YW, Lim SS, Ha JJ et al (2007) Preparative isolation of four ginsenosides from Korean red ginseng (steam-treated *Panax ginseng* C. A. Meyer), by high-speed counter-current chromatography coupled with evaporative light scattering detection. *J Chromatogr A* 1151:37–44
- Ha JJ, Kang M, Na YC et al (2011) Preparative separation of minor saponins from *Platycodi Radix* by high-speed counter-current chromatography. *J Sep Sci* 34:2559–2565
- Hamzaoui M, Renault JH, Nuzillard JM et al (2013) Stepwise elution of a three-phase solvent system in centrifugal partition extraction: a new strategy for the fractionation and phytochemical screening of a crude bark extract. *Phytochem Anal* 24:367–373
- Han X, Zhang T, Ito Y (2003) Separation of high purity przewaquinone A by high speed countercurrent chromatography. *J Liq Chromatogr Relat Technol* 26:1267–1274
- Han QB, Song JZ, Qiao CF et al (2007) Preparative isolation of cyclolanostane-type saponins from *Astragalus membranaceus* Bge. var. *mongholicus* (Bge.) Hsiao by TLC-MS/MS guided high-speed counter-current chromatography. *J Sep Sci* 30:135–140
- Han QB, Wong L, Yang NY et al (2008) A simple method to optimize the HSCCC two-phase solvent system by predicting the partition coefficient for target compound. *J Sep Sci* 31:1189–1194
- Han QB, Wong L, Lai F et al (2009) Preparative isolation of pseudolaric acids A and B and their glucosides from the root bark of *Pseudolarix kaempferi* using high-speed counter-current chromatography. *J Sep Sci* 32:309–313
- He W, Van Puyvelde L, Maes L et al (2002) Antitrichomonas in vitro activity of *Cussonia holstii* Engl. *Nat Prod Res* 17:127–133
- He F, Bai Y, Wang J et al (2011) Isolation and purification of oridonin from the whole plant of *Isodon rubescens* by high-speed counter-current chromatography. *Molecules* 16:7949–7957
- He S, Li S, Yang J et al (2012) Application of step-wise gradient high-performance counter-current chromatography for rapid preparative separation and purification of diterpene components from *Pseudolarix kaempferi* Gordon. *J Chromatogr A* 1235:34–38
- Hewitson P, Ignatova S, Ye H et al (2009) Intermittent counter-current extraction as an alternative approach to purification of Chinese herbal medicines. *J Chromatogr A* 1216:4187–4192
- Hu J, Liang Y, Xie Y et al (2007) Isolation and purification of solanesol from potato leaves by high-speed counter-current chromatography and identification by atmospheric pressure chemical ionization mass spectrometry. *Chin J Chromatogr* 25:528–531
- Huang XY, Fu JF, Di DL (2010) Preparative isolation and purification of steviol glycosides from *Stevia rebaudiana* Bertoni using high-speed counter-current chromatography. *Sep Purif Technol* 71:220–224
- Huang J, Xu X, Xie C et al (2013) Isolation and purification of paeoniflorin and albiflorin from radix *Paeoniae rubra* by high-speed counter-current chromatography. *J Liq Chromatogr Relat Technol* 36:419–427
- Ikai Y, Oka H, Hayakawa J et al (1998) Isolation of colistin C and B using high-speed countercurrent chromatography. *J Liq Chromatogr Relat Technol* 21:143–155
- Ito Y (2005) Golden rules and pitfalls in selecting optimum conditions for high-speed counter-current chromatography. *J Chromatogr A* 1065:145–168
- Ito Y, Weinstein M, Aoki I et al (1966) The coil planet centrifuge. *Nature* 212:985–987
- Ito Y, Oka H, Lee YW (1990) Improved high speed counter current chromatograph with three multilayer coils connected in series. II. Separation of various biological samples with a semipreparative column. *J Chromatogr* 498:169–178
- Ito Y, Goto T, Yamada S et al (2006) Application of dual counter-current chromatography for rapid sample preparation of N-methylcarbamate pesticides in vegetable oil and citrus fruit. *J Chromatogr A* 1108:20–25
- Jiang Y, Lu HT, Chen F (2004) Preparative purification of glycyrrhizin extracted from the root of liquorice using high-speed counter-current chromatography. *J Chromatogr A* 1033:183–186
- Jiang Z, Chen W, Liu S et al (2011) Preparation of crocin from *Gardenia yellow pigment* by slow rotary countercurrent chromatography. *Chin J Chromatogr* 29:277–280
- Kim SM, Shang YF, Um BH (2011) A preparative method for isolation of fucoxanthin from *Eisenia bicyclis* by centrifugal partition chromatography. *Phytochem Anal* 22:322–329

- Knapp H, Straubinger M, Fornari S et al (1998) {S}-3,7-dimethyl-5-octene-1,7-diol and related oxygenated monoterpenoids from petals of *Rosa damascena* Mill. *J Agric Food Chem* 46:1966–1970
- Koch A, Basar S, Richter R (2008) TLC of mono- and sesquiterpenes. In: Waksmundzka-Hajnos M, Sherma J, Kowalska T (eds) *Chromatography science series*, vol 99. CRC Press, Boca Raton, pp 451–480
- Lechtenberg M, Schepmann D, Niehues M et al (2008) Quality and functionality of saffron: quality control, species assortment and affinity of extract and isolated saffron compounds to NMDA and σ_1 (Sigma-1) receptors. *Planta Med* 74:764–772
- Lee JH, Ha YW, Jeong CS et al (2009) Isolation and tandem mass fragmentations of an anti-inflammatory compound from *Aralia elata*. *Arch Pharmacol Res* 32:831–840
- Li HB, Chen F (2001a) Preparative isolation and purification of astaxanthin from the microalga *Chlorococcum* sp. by high-speed counter-current chromatography. *J Chromatogr* 925:133–137
- Li HB, Chen F (2001b) Preparative isolation and purification of six diterpenoids from the Chinese medicinal plant *Salvia miltiorrhiza* by high-speed counter-current chromatography. *J Chromatogr A* 925:109–114
- Li HB, Chen F, Zhang TY et al (2001) Preparative isolation and purification of lutein from the microalga *Chlorella vulgaris* by high-speed counter-current chromatography. *J Chromatogr* 905:151–155
- Li A, Sun A, Liu R (2005) Preparative isolation and purification of costunolide and dehydrocostuslactone from *Aucklandia lappa* Decne by high-speed counter-current chromatography. *J Chromatogr A* 1076:193–197
- Li HB, Fan KW, Chen F (2006) Isolation and purification of canthaxanthin from the microalga *Chlorella zofingiensis* by high-speed counter-current chromatography. *J Sep Sci* 29:699–703
- Li XC, Zhang QW, Yin ZQ et al (2011) Preparative separation of patchouli alcohol from patchouli oil using high performance centrifugal partition chromatography. *J Essent Oil Res* 23:19–24
- Liang J, Meng J, Guo M et al (2013) Conical coils counter-current chromatography for preparative isolation and purification of tanshinones from *Salvia miltiorrhiza* Bunge. *J Chromatogr A* 1288:35–39
- Liu S, Cao M, Li D et al (2011) Purification and anticancer activity investigation of pentacyclic triterpenoids from the leaves of *Sinojackia sarcocarpa* L.Q. Luo by high-speed counter-current chromatography. *Nat Prod Res* 25:1600–1606
- Liu J, Liu R, Sun A et al (2013a) Connection of high-speed counter-current chromatography with evaporative light scattering detector by flow injection and its application for preparative isolation and purification of ginkgolide compounds from *Ginkgo biloba* L. *J Liq Chromatogr Relat Technol* 36:2317–2329
- Liu M, Yang F, Shi H et al (2013b) Preparative separation of triterpene alcohol ferulates from rice bran oil using a high performance counter-current chromatography. *Food Chem* 139:919–924
- Lu HT, Jiang Y, Chen F (2003) Preparative separation and purification of squalene from the microalga *Thraustochytrium* ATCC 26185 by high speed counter current chromatography. *J Chromatogr* 994:37–43
- Lu Y, Sun C, Pan Y (2006) Isolation and purification of oridonin from *Rabdosia rubescens* using upright counter-current chromatography. *J Sep Sci* 29:314–318
- Lu Y, Sun C, Liu R, Pan Y (2007) Effective two-dimensional counter-current chromatographic method for simultaneous isolation and purification of oridonin and ponocidin from the crude extract of *Rabdosia rubescens*. *J Chromatogr A* 1146:125–130
- Ma Y, Aisha HA, Liao L et al (2005) Preparative isolation and purification of rupestonic acid from the Chinese medicinal plant *Artemisia rupestris* L. by high-speed counter-current chromatography. *J Chromatogr A* 1076:198–201
- Ma J, Chen Q, Lai D et al (2010) Separation and purification of triterpene saponins from roots of *Radix Phytolaccae* by high-speed countercurrent chromatography coupled with evaporative light scattering detection. *J Liq Chromatogr Relat Technol* 33:563–571
- Marston A, Borel C, Hostettmann K (1988) Separation of natural products by centrifugal partition chromatography. *J Chromatogr* 450:91–99
- Martin AM, Queiroz EF, Marston A et al (2006) Labdane diterpenes from *Juniperus communis* L. berries. *Phytochem Anal* 17:32–35
- Maurya A, Srivastava SK (2011) Preparative-scale separation of anticancer triterpenes from *Eucalyptus hybrid* by centrifugal partition chromatography. *Sep Sci Technol* 46:1189–1194
- Medhi B, Patyar S, Rao RS et al (2009) Pharmacokinetic and toxicological profile of artemisinin compounds: an update. *Pharmacology* 84:323–332
- Mudge E, Lopes-Lutz D, Brown PN et al (2013) Purification of phenylalkanoids and monoterpene glycosides from *Rhodiola rosea* L. roots by high-speed counter-current chromatography. *Phytochem Anal* 24:129–134
- Nasser ALM, Mazzolin LP, Hiruma-Lima CA et al (2006) Preparative droplet counter-current chromatography for the separation of the new nor-seco-triterpene and pentacyclic triterpenoids from *Qualea parviflora*. *Chromatographia* 64:695–699
- Oka F, Oka H, Ito Y (1991) Systematic search for suitable two-phase solvent systems for high speed counter-current chromatography. *J Chromatogr* 538:99–108
- Peng A, Li R, Hu J et al (2008a) Flow rate gradient high-speed counter-current chromatography separation of five diterpenoids from *Triperygium wilfordii* and scale-up. *J Chromatogr A* 1200:129–135
- Peng J, Dong F, Qi Y et al (2008b) Preparative separation of four triterpene saponins from *Radix Astragali* by high-speed counter-current chromatography coupled with evaporative light scattering detection. *Phytochem Anal* 19:212–217
- Pinel B, Audo G, Mallet S et al (2007) Multi-grams scale purification of xanthanolides from *Xanthium macrocarpum*. Centrifugal partition chromatography versus silica gel chromatography. *J Chromatogr A* 1151:14–19
- Puertas Mejia M, Hillebrand S, Stashenko E et al (2002) In vitro radical scavenging activity of essential oils from Columbian plants and fractions from oregano (*Origanum vulgare* L.) essential oil. *Flavour Fragr J* 17:380–384

- Qi X, Ignatova S, Luo G et al (2010) Preparative isolation and purification of ginsenosides Rf, Re, Rd and Rb1 from the roots of *Panax ginseng* with a salt-containing solvent system and flow step-gradient by high performance counter-current chromatography coupled with an evaporative light scattering detector. *J Chromatogr A* 1217:1995–2001
- Qiu F, Friesen J, McAlpine JB et al (2012) Design of counter-current separation of *Ginkgo biloba* terpene lactones by nuclear magnetic resonance. *J Chromatogr A* 1242:26–34
- Rocha GG, Simões M, Lúcio KA et al (2007) Natural triterpenoids from *Cecropia lyratiloba* are cytotoxic to both sensitive and multidrug resistant leukemia cell lines. *Bioorgan Med Chem* 15:7355–7360
- Rodrigues VF, Carmo HM, Oliveira RR et al (2009) Isolation of terpenoids from *Trichilia quadrijuga* (Meliaceae) by droplet counter-current chromatography. *Chromatographia* 70:1191–1195
- Roscher R, Winterhalter P (1993) 1,1,6-trimethyl-1,2-dihydro-naphthalene (TDN) formation in wine. Application of multilayer coil counter-current chromatography for the study of *Vitis vinifera* cv. Riesling leaf glycosides. *J Agric Food Chem* 41:1452–1457
- Severino VGP, Cazal CM, Forim MR et al (2009) Isolation of secondary metabolites from *Hortia oreadica* (Rutaceae) leaves through high-speed counter-current chromatography. *J Chromatogr A* 1216:4275–4281
- Shehzad O, Jin Ha I et al (2011) Development of a rapid and convenient method to separate eight ginsenosides from *Panax ginseng* by high-speed counter-current chromatography coupled with evaporative light scattering detection. *J Sep Sci* 34:1116–1122
- Shehzad O, Khan S, Ha IJ et al (2012) Rational development of a selection model for solvent gradients in single-step separation of ginsenosides from *Panax ginseng* using high-speed counter-current chromatography. *J Sep Sci* 35:1462–1469
- Shehzad O, Kim HP, Kim YS (2013) State-of-the-art separation of ginsenosides from Korean white and red ginseng by countercurrent chromatography. *Anal Bioanal Chem* 405:4523–4530
- Shi S, Jiang D, Zhao M et al (2007) Preparative isolation and purification of triterpene saponins from *Clematis mandshurica* by high-speed counter-current chromatography coupled with evaporative light scattering detection. *J Chromatogr B* 852:679–683
- Shi SY, Zhou HH, Huang KL et al (2008) Application of high-speed counter-current chromatography for the isolation of antiviral eremophilanolides from *Ligularia atroviolacea*. *Biomed Chromatogr* 22:985–991
- Shi X, Wang X, Wang D et al (2009) Separation and purification of α -cyperone from *Cyperus rotundus* with supercritical fluid extraction and high-speed counter-current chromatography. *Sep Sci Technol* 44:712–721
- Shirota O, Nagamatsu K, Sekita S (2007) Simple preparative isolation of salvinorin A from the hallucinogenic sage, *Salvia divinorum*, by centrifugal partition chromatography. *J Liq Chromatogr Relat Technol* 30:1105–1114
- Shirota O, Nagamatsu K, Sekita S et al (2008) Preparative separation of the saponin lancemaside A from *Codonopsis lanceolata* by centrifugal partition chromatography. *Phytochem Anal* 19:403–410
- Silva JCT, Jham GN, Oliveira RDL et al (2007) Purification of the seven tetranortriterpenoids in neem (*Azadirachta indica*) seed by counter-current chromatography sequentially followed by isocratic preparative reversed-phase high-performance liquid chromatography. *J Chromatogr A* 1151:203–210
- Skalicka-Woźniak K, Walasek M, Ludwiczuk A et al (2013) Isolation of terpenoids from *Pimpinella anisum* essential oil by high-performance counter-current chromatography. *J Sep Sci* 36:2611–2614
- Slacanin I, Marston A, Hostettmann K et al (1991) The isolation of *Eleutherococcus senticosus* constituents by centrifugal partition chromatography and their quantitative determination by high performance liquid chromatography. *Phytochem Anal* 2:137–142
- Sun A, Zhang Y, Li A et al (2011) Extraction and preparative purification of tanshinones from *Salvia miltiorrhiza* Bunge by high-speed counter-current chromatography. *J Chromatogr B Anal Technol Biomed Life Sci* 879:1899–1904
- Sutherland IA, Fisher D (2004) Dynamic extraction technology. *Innov Pharm Technol* 10:68–71
- Sutherland IA, Fisher D (2009) Role of counter-current chromatography in the modernisation of Chinese herbal medicines. *J Chromatogr A* 1216:740–753
- Sutherland IA, Brown L, Forbes S et al (1998) Countercurrent chromatography (CCC) and its versatile application as an industrial purification and production process. *J Liq Chromatogr Relat Technol* 21:279–298
- Tian G, Zhang Y, Zhang T et al (2000) Separation of tanshinones from *Salvia miltiorrhiza* Bunge by high-speed counter-current chromatography. *J Chromatogr* 904:107–111
- Tian G, Zhang T, Zhang Y et al (2002) Separation of tanshinones from *Salvia miltiorrhiza* Bunge by multidimensional counter current chromatography. *J Chromatogr A* 945:281–285
- Tsao R, Yang R (2006) Lutein in selected Canadian crops and agri-food processing by-products and purification by high-speed counter-current chromatography. *J Chromatogr A* 1112:202–208
- Van den Heuvel R (2007) Investigation into the mechanics and feasibility of continuous counter-current extraction. PhD Thesis, Brunel University, London, UK
- Wang G, Tang W, Bidigare RR (2005) Terpenoids as therapeutic drugs and pharmaceutical agents. In: Zhang L, Demain AL (eds) *Natural products: drug discovery and therapeutic medicine*. Humana Press Inc., Totowa, pp 197–227
- Wang J, Bai HL, Liu CM et al (2010) Isolation and purification of ginsenosides from plant extract of *Panax quinquefolium* L. by high performance centrifugal partition chromatography coupled with ELSD. *Chromatographia* 71:267–271
- Wang D, Fang L, Wang X et al (2011a) Preparative separation and purification of sesquiterpenoids from *Tussilago farfara* L. by high-speed counter-current chromatography. *Quim Nova* 34:804–807
- Wang J, Liu CM, Li L et al (2011b) Isolation of four high-purity dammarane saponins from extract of *Panax notoginseng* by centrifugal partition chromatography coupled with evaporative light scattering detection in one operations. *Phytochem Anal* 22:263–267

- Wang D, Bai A, Lin X et al (2012a) Efficient method for extraction and isolation of shionone from *Aster tataricus* L. f. by supercritical fluid extraction and high-speed counter-current chromatography. *Acta Chromatogr* 24:615–625
- Wang Y, Liu M, Zheng L et al (2012b) Preparative purification of five bioactive components from *Agrimonia pilosa* Ledeb by high-speed counter-current chromatography. *J Sep Sci* 35:1977–1984
- Wang J, Yu J, Li L et al (2013) Isolation of high purity ginsenosides from plant extract of *Panax ginseng* by high performance centrifugal partition chromatography coupled with evaporative light scattering detection. *J Liq Chromatogr Relat Technol* 36:583–590
- Wei Y, Zhang T, Xu G et al (2001) Application of analytical and preparative high-speed counter-current chromatography for separation of lycopene from crude extract of tomato paste. *J Chromatogr A* 929:169–173
- Wei Y, Zhang T, Xu G et al (2003) Application of CCC for the separation of lutein from a crude extract of marigold flower petals. *J Liq Chromatogr Relat Technol* 26:1659–1669
- Wei Y, Du J, Lu Y (2012) Preparative separation of bioactive compounds from essential oil of *Flaveria bidentis* (L.) Kuntze using steam distillation extraction and one step high-speed counter-current chromatography. *J Sep Sci* 35:2608–2614
- Wu S, Sun C, Wang K et al (2004) Preparative isolation and purification of celastrol from *Celastrus orbicularis* Thunb. by a new counter current chromatography method with an upright coil planet centrifuge. *J Chromatogr A* 1028:171–174
- Wu H, Su Z, Yang Y et al (2007) Isolation of three sesquiterpene lactones from the roots of *Cichorium glandulosum* Boiss. et Huet. by high-speed counter-current chromatography. *J Chromatogr A* 1176:217–222
- Wu S, Yang L, Gao Y et al (2008) Multi-channel counter-current chromatography for high-throughput fractionation of natural products for drug discovery. *J Chromatogr A* 1180:99–107
- Wu D, Jiang X, Wu S (2010) Direct purification of tanshinones from *Salvia miltiorrhiza* Bunge by high-speed counter-current chromatography without presaturation of the two-phase solvent mixture. *J Sep Sci* 33:67–73
- Xiang F, Yu J, Yin R et al (2009) Structure-activity relationship of Taxol inferring from docking Taxol analogues to microtubule binding site. *Z Naturforsch* 64c:551–556
- Xiao X, Si X, Yuan Z et al (2012) Isolation of fucoxanthin from edible brown algae by microwave-assisted extraction coupled with high-speed countercurrent chromatography. *J Sep Sci* 35:2313–2317
- Xiao XH, Yuan ZQ, Li GK (2013) Preparation of phytosterols and phytol from edible marine algae by microwave-assisted extraction and high-speed counter-current chromatography. *Sep Purif Technol* 104:284–289
- Xie J, Wang S, Sun B et al (2008) Preparative separation and purification of β -caryophyllene from leaf oil of *Vitex negundo* L. var. *heterophylla* (Franch.) Rehd. by high speed countercurrent chromatography. *J Liq Chromatogr Relat Technol* 31:2621–2631
- Xie J, Sun B, Wang S et al (2009) Isolation and purification of nootkatone from the essential oil of fruits of *Alpinia oxyphylla* Miquel by high-speed counter-current chromatography. *Food Chem* 117:375–380
- Xin X, Yang Y, Zhong J et al (2009) Preparative isolation and purification of isobenzofuranone derivatives and saponins from seeds of *Nigella glandulifera* Freyn by high-speed counter-current chromatography combined with gel filtration. *J Chromatogr A* 1216:4258–4262
- Xu J, Luo J, Kong L (2013) Simultaneous separation of triterpenoid saponins and flavonoid glycosides from the roots of *Glycyrrhiza uralensis* Fisch by pH-zone-refining counter-current chromatography. *J Sep Sci* 36:3295–3301
- Yan J, Chen G, Tong S et al (2005) Preparative isolation and purification of germacrone and curdione from the essential oil of the rhizomes of *Curcuma wenyujin* by high-speed counter-current chromatography. *J Chromatogr A* 1070:207–210
- Yan G, Ji L, Luo Y et al (2012) Preparative isolation and purification of three sesquiterpenoid lactones from *Eupatorium lindleyanum* DC. by high-speed counter-current chromatography. *Molecules* 17:9002–9009
- Yang F, Ou Q, Yu W (1995) Semi-preparative separation of taraxeryl-acetate and coumarins from *Artemisia dalailamae* Kraschen by high-speed countercurrent chromatography. *J Liq Chromatogr* 18:395–403
- Yang Y, Gu D, Yili A et al (2010) One-step separation and purification of rupestonic acid and chrysosptertin B from *Artemisia rupestris* L. by high-speed counter-current chromatography. *Phytochem Anal* 21:205–209
- Yang Y, Bakri M, Gu D et al (2013) Separation of (S)-dehydrovomifoliol from leaves of *Nitraria sibirica* Pall. by high-speed counter-current chromatography. *J Liq Chromatogr Relat Technol* 36:573–582
- Yao S, Liu R, Huang X et al (2007) Preparative isolation and purification of chemical constituents from the root of *Adenophora tetraphylla* by high-speed counter-current chromatography with evaporative light scattering detection. *J Chromatogr A* 1139:254–262
- Yao S, Luo J, Huang X et al (2008) Application of preparative high-speed counter-current chromatography/preparative high-performance liquid chromatography mode in rapid separation of saponins. *J Chromatogr B* 864:69–77
- Ye H, Chen L, Li Y et al (2008a) Preparative isolation and purification of three rotenoids and one isoflavone from the seeds of *Millettia pachycarpa* Benth by high-speed counter-current chromatography. *J Chromatogr A* 1178:101–107
- Ye H, Ignatova S, Luo H et al (2008b) Preparative separation of a terpenoid and alkaloids from *Tripterygium wilfordii* Hook. using high-performance counter-current chromatography. Comparison of various elution and operating strategies. *J Chromatogr A* 1213:145–153
- Yoo KY, Park SY (2012) Terpenoids as potential anti-Alzheimer's disease therapeutics. *Molecules* 17:3524–3538
- Yoon KD, Kim J (2009) Application of centrifugal partition chromatography coupled with evaporative light scattering detection for the isolation of saikosaponins-a and -c from *Bupleurum falcatum* roots. *J Sep Sci* 32:74–78
- Yoon KD, Chin YW, Ahn MJ et al (2009) Preparative isolation of alisol B and alisol b 23-acetate from *Alismatis rhizoma* by centrifugal partition chromatography coupled with ELSD. *Chromatographia* 69:791–793

- Yuan Y, Wang B, Chen L et al (2008) How to realize the linear scale-up process for rapid purification using high-performance counter-current chromatography. *J Chromatogr A* 1194:192–198
- Zhang H, Feng B, Liu K et al (2013a) Isolation and purification of two triterpenoids from the Chinese medicinal plant *Fomes officinalis* Ames. *Asian J Chem* 25:6130–6132
- Zhang Y, Liu C, Qi Y et al (2013b) Application of accelerated solvent extraction coupled with counter-current chromatography to extraction and online isolation of saponins with a broad range of polarity from *Panax notoginseng*. *Sep Purif Technol* 106:82–89
- Zhao Y, Du Q (2007) Separation of solanesol in tobacco leaves extract by slow rotary counter-current chromatography using a novel non-aqueous two-phase solvent system. *J Chromatogr A* 1151:193–196
- Zhao C, He C (2006) Preparative isolation and purification of atractylon and atractylenolide III from the Chinese medicinal plant *Atractylodes macrocephala* by high-speed counter-current chromatography. *J Sep Sci* 29:1630–1636
- Zhao D, Yan M, Huang Y et al (2012) Efficient protocol for isolation and purification of different soyasaponins from soy hypocotyls. *J Sep Sci* 35:3281–3292