

## COUNTEREXAMPLES CONCERNING OBSERVATION OPERATORS FOR $C_0$ -SEMIGROUPS\*

BIRGIT JACOB<sup>†</sup> AND HANS ZWART<sup>‡</sup>

**Abstract.** This paper concerns systems of the form  $\dot{x}(t) = Ax(t)$ ,  $y(t) = Cx(t)$ , where  $A$  generates a  $C_0$ -semigroup. Two conjectures which were posed in 1991 and 1994 are shown not to hold. The first conjecture (by G. Weiss) states that if the range of  $C$  is one-dimensional, then  $C$  is admissible if and only if a certain resolvent estimate holds. The second conjecture (by D. Russell and G. Weiss) states that a system is exactly observable if and only if a test similar to the Hautus test for finite-dimensional systems holds. The  $C_0$ -semigroup in both counterexamples is analytic and possesses a basis of eigenfunctions. Using the  $(A, C)$ -pair from the second counterexample, we construct a generator  $A_e$  on a Hilbert space such that  $(sI - A_e)$  is uniformly left-invertible, but its semigroup does not have this property.

**Key words.** infinite-dimensional system, admissible observation operator, exact observability, conditional basis,  $C_0$ -semigroup, left-invertibility

**AMS subject classifications.** 93C25, 93A05, 93B07, 47D60

**DOI.** 10.1137/S0363012903423235

**1. Introduction.** Consider the abstract system

$$(1.1) \quad \dot{x}(t) = Ax(t), \quad y(t) = Cx(t), \quad x(0) = x_0$$

with  $x(t) \in H$  and  $y(t) \in Y$ , where  $H$  and  $Y$  are Hilbert spaces. For this abstract differential equation one would like to obtain conditions in terms of  $A$  and  $C$  such that it has a solution with certain properties. If one only considers the differential equation  $\dot{x}(t) = Ax(t)$ , then it is well known that it has a unique (weak) solution which is strongly continuous and depends continuously on the initial state  $x(0) = x_0 \in H$  if and only if  $A$  satisfies the estimates of the Hille–Yosida theorem (see, e.g., [4, Theorem 2.1.12]). Since  $\dot{x}(t) = Ax(t)$  is a part of (1.1) we have to assume that  $A$  satisfies the estimates of the Hille–Yosida theorem, or equivalently, that  $A$  generates a  $C_0$ -semigroup. If in addition  $C$  is a bounded linear operator from  $H$  to  $Y$ , then it is straightforward to see that  $y(\cdot)$  in (1.1) is well defined and continuous. However, many PDEs rewritten in the form (1.1) do not have a bounded operator  $C$ , although the output is a well-defined square integrable function. We assume that  $C$  is a bounded operator from  $D(A)$  (with the graph norm) to a Hilbert space  $Y$ . If the output is locally square integrable, then  $C$  is called an *admissible observation operator* (see Weiss [20] and the survey article by Jacob and Partington [7]). In other words,  $C$  is an admissible observation operator if and only if for some  $t_0 > 0$  (and hence any  $t_0 > 0$ ) there exists a constant  $L > 0$  such that

$$\int_0^{t_0} \|CT(t)x\|^2 dt \leq L\|x\|^2, \quad x \in D(A).$$

---

\*Received by the editors February 17, 2003; accepted for publication (in revised form) October 23, 2003; published electronically June 4, 2004.

<http://www.siam.org/journals/sicon/43-1/42323.html>

<sup>†</sup>Department of Mathematics, University of Dortmund, D-44221 Dortmund, Germany (birgit.jacob@math.uni-dortmund.de).

<sup>‡</sup>Department of Applied Mathematics, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands (h.j.zwart@math.utwente.nl).

Here  $(T(t))_{t \geq 0}$  is the  $C_0$ -semigroup generated by  $A$ . If the  $C_0$ -semigroup is exponentially stable, then  $t_0$  can be replaced by  $\infty$ . Now an interesting question is if there are simple conditions on  $C$  (and  $A$ ) such that  $C$  is an admissible observation operator.

Dual to the concept of admissible observation operator is the concept of admissible control operator. An operator  $B$  is said to be an admissible control operator if  $\dot{x}(t) = Ax(t) + Bu(t)$  has a continuous (weak) solution for every locally square integrable input  $u$ . It is well known that  $C$  is an admissible observation operator for  $A$  if and only if  $C^*$  is an admissible control operator for  $A^*$ ; see [20] for a proof of this statement. Here  $*$  denotes the adjoint operator. Because of this duality any result for admissible observation operators has an equivalent counterpart for admissible control operators, and vice versa. Hence if we refer to a paper which only deals with control operators, we trust that the reader can make the equivalent statement for observation operators. Basically, it boils down to replacing  $B$  by  $C^*$  and replacing the infinitesimal generator by its dual one.

In Weiss [21] it is shown that if  $C$  is admissible, then there exists a constant  $M > 0$  such that

$$(1.2) \quad \|C(sI - A)^{-1}\| \leq \frac{M}{\sqrt{\operatorname{Re}(s)}}$$

for all  $s$  in some right-half plane. He conjectured in [21] (see also [22]) that this condition is also sufficient. The sufficiency of condition (1.2) was proved for surjective semigroups in Weiss [21], for normal, analytic semigroups in Weiss [21, 22], for the right shift semigroup with scalar output in Partington and Weiss [15], for contraction semigroups with scalar output by Jacob and Partington [6], and for analytic contraction semigroups by Le Merdy [12]. Recently, Zwart, Jacob, and Staffans [26] and Jacob, Partington, and Pott [8] showed that in general estimate (1.2) is not sufficient. Their observation operator is infinite-dimensional. Here we use techniques similar to those in [26] to show that (1.2) is not sufficient for scalar outputs. Note that in [5] a necessary and sufficient condition has been obtained. This condition involves all powers of the resolvent, as in the Hille–Yosida theorem. Some sufficient conditions for admissibility can be found in [24].

Apart from the well-posedness of the abstract differential equation (1.1) one would like to characterize other properties in terms of the pair  $(A, C)$ . One property that has received a lot of attention is the property of exact observability. Assuming that the observation operator  $C$  is admissible, the system (1.1) is said to be exactly observable if there is a bounded mapping from the output trajectory to the initial condition, that is, for some  $t_0 > 0$  (and hence any  $t_0 > 0$ ) there exists a constant  $l > 0$  such that

$$\int_0^{t_0} \|CT(t)x\|^2 dt \geq l\|x\|^2, \quad x \in D(A).$$

If the  $C_0$ -semigroup is exponentially stable, then  $t_0$  can be replaced by  $\infty$ . Note that admissibility gives that the mapping from initial condition to output trajectory is bounded. If the state space  $H$  is finite-dimensional, and thus  $A$  and  $C$  are just matrices, then it is well known that (1.1) is exactly observable if and only if

$$\operatorname{rank} \begin{bmatrix} C \\ sI - A \end{bmatrix}$$

is full for all complex  $s$ . For infinite-dimensional systems, Russell and Weiss [17], proposed the following test for exact observability of an exponentially stable system:

$$(1.3) \quad \|(sI - A)x_0\|^2 + |\operatorname{Re}(s)|\|Cx_0\|^2 \geq m|\operatorname{Re}(s)|^2\|x_0\|^2$$

for all complex  $s$  with negative real part, for all  $x_0 \in D(A)$ , and for some positive  $m$  independent of  $s$  and  $x_0$ . In [17] they proved that this condition is always necessary, and that for  $A$  and  $C$  bounded this condition is sufficient as well. In the same paper they showed that if  $A$  has a Riesz basis of eigenfunctions and an extra condition on the eigenvalues is satisfied, then (1.3) is sufficient. In Zhou and Yamamoto [23] it was shown that (1.3) is sufficient if  $A$  is skew adjoint and  $C$  is bounded. For Riesz spectral systems with finite-dimensional output space  $Y$  inequality (1.3) is sufficient as well; see Jacob and Zwart [9, 10]. Grabowski and Callier [5] proved that if  $m$  in (1.3) is equal to one, then this estimate implies exact observability. In section 3 we show that for general  $m$  estimate (1.3) is not sufficient. Note that in our counterexample the output is one-dimensional and that  $A$  generates an analytic semigroup. In [11] we give a refined version of this conjecture.

We conclude this paper with a section on left-invertibility of  $C_0$ -semigroups. It is known that uniform left-invertibility of the semigroup implies uniform left-invertibility of the generator on the open left-half plane. We show that in general the inverse implication does not hold.

**2. General results.** Let  $H$  be a separable Hilbert space with a conditional basis  $\{\varphi_n\}_{n \in \mathbb{N}}$ . Since  $\{\varphi_n\}_{n \in \mathbb{N}}$  is a conditional basis, we have that for every  $x \in H$  there exists a unique sequence of complex numbers  $\alpha_n$  such that

$$(2.1) \quad x = \lim_{k \rightarrow \infty} \sum_{n=1}^k \alpha_n \varphi_n.$$

Hence, we can write

$$x = \sum_{n=1}^{\infty} \alpha_n \varphi_n.$$

Using (2.1) it is not hard to see that the following holds (see also Singer [18, pages 18–20]).

LEMMA 2.1. *If  $\{\varphi_n\}_{n \in \mathbb{N}}$  is a conditional basis, then the following mappings are uniformly bounded:*

$$(2.2) \quad P_n x = \sum_{k=1}^n \alpha_k \varphi_k$$

and

$$(2.3) \quad \tilde{P}_n x = \alpha_n \varphi_n,$$

where  $x = \sum_{n=1}^{\infty} \alpha_n \varphi_n$ .

Furthermore, if  $\inf_{n \in \mathbb{N}} \|\varphi_n\| > 0$ , then

$$(2.4) \quad \sup_{n \in \mathbb{N}} |\alpha_n| \leq \kappa \|x\|$$

for some  $\kappa > 0$  independent of  $x$ .

The following two properties of a conditional basis are important for the construction of our counterexamples.

DEFINITION 2.2. *Let  $\{\varphi_n\}_{n \in \mathbb{N}}$  be a conditional basis.*

1.  $\{\varphi_n\}_{n \in \mathbb{N}}$  is Besselian if there exists a constant  $c > 0$  such that

$$\sum_{k=1}^n |a_k|^2 \leq c \left\| \sum_{k=1}^n a_k \varphi_k \right\|^2$$

for all finite sequences of scalars  $a_1, \dots, a_n$ .

2.  $\{\varphi_n\}_{n \in \mathbb{N}}$  is Hilbertian if there exists a constant  $c > 0$  such that

$$\left\| \sum_{k=1}^n a_k \varphi_k \right\|^2 \leq c \sum_{k=1}^n |a_k|^2$$

for all finite sequences of scalars  $a_1, \dots, a_n$ .

Equivalently,  $\{\varphi_n\}_{n \in \mathbb{N}}$  is Besselian if and only if there exists a bounded linear operator  $S$  such that  $v_n := S\varphi_n$  is an orthonormal basis for  $H$ . More information on conditional bases can be found in Singer [18].

For diagonal operators on a conditional basis of  $H$  there is the following nice result, which can be found in Benamara and Nikolski [1, Lemma 3.2.5].

LEMMA 2.3. *Let  $\{\varphi_n\}_n$  be a conditional basis of  $H$ . If  $Q$  is defined as*

$$Q\varphi_n = q_n \varphi_n$$

with  $\{q_n\}_{n \in \mathbb{N}} \subset \mathbb{C}$ , and the total variation of the sequence  $\{q_n\}$  is finite, i.e.,

$$\text{Var}(q_n) := \sum_{n=1}^{\infty} |q_{n+1} - q_n| < \infty,$$

then  $Q$  can be extended to a linear bounded operator on  $H$ , and

$$(2.5) \quad \|Q\| \leq K(\text{Var}(q_n) + \limsup |q_n|),$$

where  $K$  is the supremum of  $\|P_n\|$ ; see Lemma 2.1.

In order to calculate the total variation, the following observation is useful. If  $f$  is a continuous function which is nondecreasing or nonincreasing on the interval  $(a, b)$ , and if the sequence  $\{q_n\}_n \subset (a, b)$  is nondecreasing or nonincreasing, then

$$\text{Var}(f(q_n)) \leq |f(a) - f(b)|.$$

In [26] the following useful result can be found.

LEMMA 2.4. *Let  $\{\mu_n\}_n \subset (-\infty, -1]$  be a monotonically decreasing sequence with  $\lim_{n \rightarrow \infty} \mu_n = -\infty$ . Furthermore, let  $\{\varphi_n\}_{n \in \mathbb{N}}$  be a conditional basis for the Hilbert space  $H$ .*

*For  $t \geq 0$ , we define  $(T(t))_{t \geq 0}$  by*

$$(2.6) \quad T(t)\varphi_n := e^{\mu_n t} \varphi_n, \quad n \in \mathbb{N}.$$

*The operator valued function  $(T(t))_{t \geq 0}$  defines an analytic, exponentially stable  $C_0$ -semigroup on  $H$ .*

**3. Counterexample on admissibility.** In this section we show that the conjecture of George Weiss for admissibility of scalar observation operators (see [21, 22]) does not hold. That means we construct an exponentially stable  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  on  $H$  with infinitesimal generator  $A$  and an operator  $C \in \mathcal{L}(D(A), \mathbb{C})$  such that

$$\|C(sI - A)^{-1}\| \leq \frac{M}{\sqrt{\operatorname{Re}(s)}}$$

for all  $s$  in some right-half plane and some constant  $M > 0$ , but  $C$  is not an admissible observation operator for  $(T(t))_{t \geq 0}$ .

Let  $\{e_n\}_{n \in \mathbb{N}}$  be a conditional basis on  $H$  which has the following properties:

1.  $\inf_{n \in \mathbb{N}} \|e_n\| > 0$ .
2.  $\{e_n\}_{n \in \mathbb{N}}$  is not Besselian.

Such Hilbert spaces and bases do exist; see, for example, Singer [18, page 351, example 11.2].

We define the sequence  $\mu_n$  as

$$(3.1) \quad \mu_n := -4^n, \quad n \in \mathbb{N},$$

and the  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  as

$$(3.2) \quad T(t)e_n = e^{\mu_n t} e_n.$$

By Lemma 2.4 we know that  $(T(t))_{t \geq 0}$  is an exponentially stable analytic semigroup. By  $A$  we denote the infinitesimal generator of  $(T(t))_{t \geq 0}$ . It is easy to see that  $A$  satisfies

$$Ae_n = \mu_n e_n, \quad n \in \mathbb{N}.$$

For  $x \in D(A)$ ,  $x = \sum_{n=1}^{\infty} x_n e_n$ , we further define

$$(3.3) \quad Cx = \sum_{n=1}^{\infty} \sqrt{-\mu_n} x_n.$$

First of all we show that  $C$  is a bounded linear operator from the domain of  $A$  into  $\mathbb{C}$ .

**PROPOSITION 3.1.** *Let  $C$  be given as in (3.3) and let  $A$  be the infinitesimal generator of the  $C_0$ -semigroup (3.2). Then we have  $C \in \mathcal{L}(D(A), \mathbb{C})$ .*

*Proof.* It is enough to show that there exists a constant  $c > 0$  such that

$$|CA^{-1}x| \leq c, \quad x \in H, \|x\| = 1.$$

Let  $x \in H$  with  $\|x\| = 1$ . Then there exist scalars  $x_n$ ,  $n \in \mathbb{N}$ , such that

$$x = \sum_{n=1}^{\infty} x_n e_n.$$

Using that  $\inf_{n \in \mathbb{N}} \|e_n\| > 0$ , we get from Lemma 2.1 that  $\sup_{n \in \mathbb{N}} |x_n| \leq \kappa < \infty$ . Note that  $\kappa$  is independent of  $x \in H$  with  $\|x\| = 1$ . Now we have

$$|CA^{-1}x| = \left| \sum_{n=1}^{\infty} \frac{x_n}{\sqrt{-\mu_n}} \right| \leq \kappa \sum_{n=1}^{\infty} 2^{-n} = \kappa.$$

Thus the proposition is proved.  $\square$

Next we show that  $C$  satisfies the estimate (1.2).

PROPOSITION 3.2. *For  $C$  given by (3.3) and  $A$  as the infinitesimal generator of the semigroup (3.2) the following holds. There exists a constant  $M > 0$  such that*

$$\|C(sI - A)^{-1}\| \leq \frac{M}{\sqrt{\operatorname{Re}(s)}}, \quad s \in \mathbb{C}_+.$$

*Proof.* Let  $s$  be an element of  $\mathbb{C}_+$ , and let  $x \in H$  have norm one. We have the following estimate:

$$\begin{aligned} \sqrt{\operatorname{Re}(s)}|C(sI - A)^{-1}x| &= \sqrt{\operatorname{Re}(s)} \left| \sum_{k=1}^{\infty} \frac{2^k}{s + 4^k} x_k \right| \\ &\leq \sqrt{\operatorname{Re}(s)} \sum_{k=1}^{\infty} \frac{2^k}{|\operatorname{Re}(s) + 4^k|} |x_k| \\ &\leq \kappa \sqrt{\operatorname{Re}(s)} \sum_{k=1}^{\infty} \frac{2^k}{\operatorname{Re}(s) + 4^k}, \end{aligned}$$

where we have used Lemma 2.1. Note that  $\kappa$  is independent of  $x$ . In order to estimate this last expression we introduce the monotonically decreasing sequence  $a_k := \frac{1}{\operatorname{Re}(s) + k^2}$ . Then for  $N \geq 2^K$  we have

$$\begin{aligned} \sum_{k=1}^N a_k &\geq a_1 + a_2 + (a_3 + a_4) + \cdots + (a_{2^{K-1}+1} + \cdots + a_{2^K}) \\ &\geq a_2 + 2a_4 + \cdots + 2^{K-1}a_{2^K} \\ &= \frac{1}{2} \sum_{k=1}^K 2^k a_{2^k}, \end{aligned}$$

and so

$$\sum_{k=1}^{\infty} \frac{2^k}{\operatorname{Re}(s) + 4^k} \leq 2 \sum_{k=1}^{\infty} \frac{1}{\operatorname{Re}(s) + k^2}.$$

Using this in our estimate of  $\sqrt{\operatorname{Re}(s)}|C(sI - A)^{-1}x|$ , we obtain that

$$\begin{aligned} \sqrt{\operatorname{Re}(s)}|C(sI - A)^{-1}x| &\leq 2\kappa \sqrt{\operatorname{Re}(s)} \sum_{k=1}^{\infty} \frac{1}{\operatorname{Re}(s) + k^2} \\ &\leq 2\kappa \sqrt{\operatorname{Re}(s)} \int_0^{\infty} \frac{1}{\operatorname{Re}(s) + t^2} dt \\ &\leq 2\kappa \sqrt{\operatorname{Re}(s)} \left( \frac{1}{\sqrt{\operatorname{Re}(s)}} \arctan \left( \frac{t}{\sqrt{\operatorname{Re}(s)}} \right) \right) \Big|_0^{\infty} \\ &\leq 2\kappa \frac{\pi}{2} = \kappa\pi, \end{aligned}$$

which proves our assertion.  $\square$

PROPOSITION 3.3. *If  $C$  given by (3.3) is an admissible observation operator for the  $C_0$ -semigroup given by (3.2), then  $\{e_n\}$  is Besselian.*

*Proof.* If  $C$  is an admissible observation operator for  $(T(t))_{t \geq 0}$ , then there would exist a constant  $L > 0$  such that

$$\int_0^\infty |CT(t)x|^2 dt \leq L\|x\|^2, \quad x \in D(A).$$

Now take a finite sequence of  $\alpha_k$ 's and consider

$$x := \sum_{k=1}^n \alpha_k e_k.$$

Then the above estimate gives

$$\int_0^\infty \left| \sum_{k=1}^n \sqrt{-\mu_k} e^{\mu_k t} \alpha_k \right|^2 dt \leq L\|x\|^2.$$

However, from Nikolski and Pavlov [14] (see also Jacob and Zwart [10]), we know that there exists a constant  $L_1 > 0$ , independent of  $x$ , such that

$$\int_0^\infty \left| \sum_{k=1}^n \sqrt{-\mu_k} e^{\mu_k t} \alpha_k \right|^2 dt \geq L_1 \sum_{k=1}^n |\alpha_k|^2.$$

Thus we have that for any finite sequence

$$\|x\|^2 \geq \frac{L_1}{L} \sum_{k=1}^n |\alpha_k|^2,$$

which shows that  $\{e_n\}$  is Besselian.  $\square$

Thus we have disproved the scalar admissibility conjecture of George Weiss.

**4. Counterexample on exact observability.** In this section we use the operators  $A$  and  $C$  constructed in section 3 with different assumptions on the basis to settle another question about operator semigroups.

We disprove the conjecture of Russell and Weiss [17] on exact observability. That means we construct an exponentially stable  $C_0$ -semigroup  $(T(t))_{t \geq 0}$  with infinitesimal generator  $A$  and an operator  $C \in \mathcal{L}(D(A), \mathbb{C})$  such that

$$\|(sI - A)x_0\|^2 + |\operatorname{Re}(s)|\|Cx_0\|^2 \geq m|\operatorname{Re}(s)|^2\|x_0\|^2, \quad s \in \mathbb{C}_-, x_0 \in D(A),$$

for some constant  $m > 0$ , but the pair  $(A, C)$  is not exactly observable.

Let  $\{e_n\}_{n \in \mathbb{N}}$  be a conditional basis on  $H$  which is Besselian, normalized—that is,  $\|e_n\| = 1$ , but not Hilbertian. Such Hilbert spaces and bases do exist; see, for example, Singer [18, page 351, example 11.2].

We define the sequence  $\mu_n$  as

$$(4.1) \quad \mu_n := -4^n, \quad n \in \mathbb{N},$$

and the  $C_0$ -semigroup as

$$(4.2) \quad T(t)e_n = e^{\mu_n t} e_n.$$

By Lemma 2.4 we know that this is an exponentially stable analytic  $C_0$ -semigroup. By  $A$  we denote the infinitesimal generator of  $(T(t))_{t \geq 0}$ . It is easy to see that  $A$  satisfies

$$Ae_n = \mu_n e_n, \quad n \in \mathbb{N}.$$

Since  $\{e_n\}_{n \in \mathbb{N}}$  is Besselian, we know that there exists a bounded linear operator  $S$  such that  $v_n := Se_n$  is an orthonormal basis for  $H$ . On this new basis we define

$$\tilde{A}v_n = \mu_n v_n.$$

It is easy to see that  $\tilde{A}$  generates a  $C_0$ -semigroup  $(\tilde{T}(t))_{t \geq 0}$ , and that

$$(4.3) \quad ST(t) = \tilde{T}(t)S.$$

Now define the operator  $\tilde{C}$  as

$$\tilde{C}v_n = \sqrt{-\mu_n}.$$

It is easy to see that we can extend  $\tilde{C}$  as a bounded operator from the domain of  $\tilde{A}$  to  $\mathbb{C}$ . We denote this extension again by  $\tilde{C}$ . We shall prove that  $\tilde{C}$  is an admissible observation operator for  $(\tilde{T}(t))_{t \geq 0}$ . Since  $(\tilde{T}(t))_{t \geq 0}$  has an orthonormal basis of eigenfunctions, we can use the result of Weiss [19], which tells us that  $\tilde{C}$  is admissible if and only if

$$\sum_{-\mu_n \in R(h, \omega)} |\mu_n| \leq \beta h,$$

where

$$R(h, \omega) := \{s \in \mathbb{C}_+ \mid \operatorname{Re}(s) \leq h, |\operatorname{Im}(s) - \omega| \leq h\}$$

and  $\beta$  is independent of  $h$ . Using the definition of  $\mu_n$  this is easy to prove. Now we define for  $x \in D(A)$ ,

$$(4.4) \quad Cx = \tilde{C}Sx.$$

From this and (4.3) we see that for  $x \in D(A)$

$$CT(t)x = \tilde{C}\tilde{T}(t)Sx.$$

Since  $S$  is bounded and since  $\tilde{C}$  is admissible for  $(\tilde{T}(t))_{t \geq 0}$ , we obtain that  $C$  is an admissible output operator for  $(T(t))_{t \geq 0}$ .

In several steps we shall prove that the pair  $(A, C)$  satisfies the estimate of Russell and Weiss, but that it is not exactly observable. In our proof we follow closely the proof of Theorem 4.4 of Russell and Weiss [17]. As in [17] we define  $N : \mathbb{C}_- \rightarrow \mathbb{N}$  as the integer such that

$$(4.5) \quad |s - \mu_{N(s)}| = \min_{k \in \mathbb{N}} |s - \mu_k|.$$

This number is well defined if the real part of  $s$  is unequal to  $(\mu_k + \mu_{k+1})/2$  for all  $k$ . We define the set for which this mapping is well defined as  $\mathbb{C}_g$ .



LEMMA 4.1. *There exists a constant  $c > 0$  such that, for all  $s \in \mathbb{C}_g$ , we have that*

$$\left| \frac{\operatorname{Re}(s)}{s - \mu_k} \right| \leq c, \quad s \in \mathbb{C}_g, k \neq N(s),$$

and

$$\left| \frac{\operatorname{Re}(s)}{\operatorname{Re}(s) - \mu_k} \right| \leq c, \quad s \in \mathbb{C}_g, k \neq N(s).$$

*Proof.* In Weiss and Russell [17] it is shown that the first estimate holds. Since  $\{\mu_k\}$  is a real sequence, it is easy to see that  $N(s) = N(\operatorname{Re}(s))$ . Taking  $s$  to be real in the first inequality, and using this observation, proves the second inequality.  $\square$

For  $s \in \mathbb{C}_g$ , we define

$$(4.6) \quad V(s) := \overline{\operatorname{span}_{n \neq N(s)} \{e_n\}}.$$

Clearly,  $V(s)$  is again a Hilbert space and in Singer [18, page 26, Proposition 4.1] it is shown that  $\{e_n\}_{n \neq N(s)}$  is a conditional basis of  $V(s)$ . By  $P_{V(s)}$  we denote the projection from  $H$  onto  $V(s)$  given by

$$P_{V(s)} := I - \tilde{P}_{N(s)}.$$

Using Lemma 2.1 we see that the projections  $P_{V(s)}$  are uniformly bounded. For  $s \in \mathbb{C}_g$ , we introduce the notation

$$(4.7) \quad e_n^s := \begin{cases} e_n, & n < N(s), \\ e_{n+1}, & n \geq N(s), \end{cases}$$

and

$$(4.8) \quad \mu_n^s := \begin{cases} \mu_n, & n < N(s), \\ \mu_{n+1}, & n \geq N(s). \end{cases}$$

The constant  $K$  in Lemma 2.3 is given by  $K := \sup_{n \in \mathbb{N}} \|P_n\|$ . Let  $K(s)$  be the corresponding constant for  $V(s)$  with conditional basis  $\{e_n^s\}$ , for  $s \in \mathbb{C}_g$ . Then it follows easily that  $K(s) \leq K$ .

Let  $s \in \mathbb{C}_g$ . We denote by  $A_s$  the part of  $A$  in  $V(s)$ , that is,

$$A_s x := Ax, \quad x \in D(A_s),$$

and  $D(A_s) := D(A) \cap V(s)$ . Note that  $V(s)$  is a  $T(t)$ -invariant subspace. Thus it is easy to see that  $C_s$ , defined by

$$C_s x := Cx, \quad x \in D(A_s),$$

is an admissible observation operator for  $(T_s(t))_{t \geq 0}$ . Here  $(T_s(t))_{t \geq 0}$  is the  $C_0$ -semigroup generated by  $A_s$ . Now we shall prove two important estimates.

LEMMA 4.2. *Let  $A_s$ ,  $C_s$ , and  $V(s)$  denote the objects defined above. The following two estimates hold.*

1. *There exists a constant  $M > 0$  such that*

$$\|(sI - A_s)^{-1}\|_{V(s)} \leq \frac{M}{|\operatorname{Re}(s)|}, \quad s \in \mathbb{C}_g.$$

2. There exists a constant  $d > 0$  such that

$$\|C_s(sI - A_s)^{-1}\| \leq \frac{d}{\sqrt{|\operatorname{Re}(s)|}}, \quad s \in \mathbb{C}_g.$$

*Proof. Part 1.* Let  $s = s_r + is_i \in \mathbb{C}_g$ . Clearly,

$$(sI - A_s)^{-1}e_n^s = \frac{1}{s - \mu_n^s}e_n^s, \quad n \in \mathbb{N}.$$

This is an operator of the form as discussed in Lemma 2.3, and thus we have to show that  $1/(s - \mu_n^s)$  is of bounded variation. We begin with the following simple observation:

$$\begin{aligned} \left| \frac{1}{s - \mu_{n+1}^s} - \frac{1}{s - \mu_n^s} \right| &= \left| \frac{\mu_{n+1}^s - \mu_n^s}{(s - \mu_{n+1}^s)(s - \mu_n^s)} \right| \\ &\leq \left| \frac{\mu_{n+1}^s - \mu_n^s}{(s_r - \mu_{n+1}^s)(s_r - \mu_n^s)} \right| \\ (4.9) \qquad \qquad \qquad &= \left| \frac{1}{s_r - \mu_{n+1}^s} - \frac{1}{s_r - \mu_n^s} \right|, \end{aligned}$$

where we have used the fact that  $\mu_n^s$  is real.

Next we define

$$h : \mathbb{R}_- \setminus \{s_r\} \rightarrow \mathbb{R}, \quad h(x) := \frac{1}{s_r - x}.$$

Then we have  $h(-\infty) = 0$ ,  $h(0) = \frac{1}{s_r}$ , and  $h$  is monotonically increasing on  $(-\infty, s_r)$  and on  $(s_r, 0)$ . Combining the above results with Lemma 2.3 we get the following estimate for  $\|(sI - A_s)^{-1}\|$ :

$$\begin{aligned} &\|(sI - A_s)^{-1}\| \\ &\leq K \left( \operatorname{Var} \left( \frac{1}{s - \mu_n^s} \right) + \left| \lim_{n \rightarrow \infty} \frac{1}{s - \mu_n^s} \right| \right) = K \sum_{n=1}^{\infty} \left| \frac{1}{s - \mu_{n+1}^s} - \frac{1}{s - \mu_n^s} \right| \\ &\leq K \sum_{n=1}^{\infty} \left| \frac{1}{s_r - \mu_{n+1}^s} - \frac{1}{s_r - \mu_n^s} \right| \\ &\leq K \left[ \left[ 0 + \frac{1}{s_r - \mu_{N(s)+1}} \right] + \left[ \frac{1}{s_r - \mu_{N(s)+1}} - \frac{1}{s_r - \mu_{N(s)-1}} \right] \right. \\ &\quad \left. + \left[ \frac{1}{s_r} - \frac{1}{s_r - \mu_{N(s)-1}} \right] \right] \\ &\leq \frac{(4c+1)K}{|\operatorname{Re}(s)|}, \end{aligned}$$

where we have used Lemmas 2.3 and 4.1 and (4.9). Since  $c$  and  $K$  are independent of  $s$  we have proved the statement.

*Part 2.* In order to prove this statement we follow Lemma 4.6 of Russell and Weiss [17]. Let  $s \in \mathbb{C}_g$ . Using the resolvent identity, we have

$$C_s(sI - A_s)^{-1} = C_s(-\bar{s}I - A_s)^{-1}[I - (\bar{s} + s)(sI - A_s)^{-1}].$$

Since  $C_s$  is an admissible observation operator for  $(T_s(t))_{t \geq 0}$  there exists a constant  $\tilde{d} > 0$ , independent of  $s$ , such that

$$\|C_s(-\bar{s}I - A_s)^{-1}\| \leq \frac{\tilde{d}}{\sqrt{|\operatorname{Re}(s)|}}$$

(see, e.g., Weiss [22]). Combining this with Part 1, the statement is proved.  $\square$

Now we can prove the estimate of Russell and Weiss [17].

LEMMA 4.3. *For  $C$  defined by (4.4) and  $A$  as the infinitesimal generator of (4.2) the following holds. There exists a constant  $m > 0$  such that, for every  $s \in \mathbb{C}_-$  and every  $x \in D(A)$ , we have*

$$(4.10) \quad \frac{1}{|\operatorname{Re}(s)|^2} \|(sI - A)x\|^2 + \frac{1}{|\operatorname{Re}(s)|} \|Cx\|^2 \geq m\|x\|^2.$$

*Proof.* The proof of this lemma is divided into two steps. First, we show that the estimate holds for  $s \in \mathbb{C}_- \setminus \mathbb{C}_g$ . Second, we prove the estimate for  $s \in \mathbb{C}_g$ .

*Part 1.* If  $s$  is not in  $\mathbb{C}_g$ , then there exists an  $k_0 \in \mathbb{N}$  such that  $\operatorname{Re}(s) = (\mu_{k_0+1} + \mu_{k_0})/2$ . It is easy to see that

$$(sI - A)^{-1}e_n = \frac{1}{s - \mu_n}e_n.$$

We use Lemma 2.3 to estimate the norm of this operator. Using (4.9) we see that it is sufficient to show that  $\left\{\frac{1}{\operatorname{Re}(s) - \mu_n}\right\}$  is of bounded variation. Similar to the proof of Part 1 of Lemma 4.2, we obtain that

$$\|(sI - A)^{-1}\| \leq K \sum_{n=1}^{\infty} \left| \frac{1}{\operatorname{Re}(s) - \mu_{n+1}} - \frac{1}{\operatorname{Re}(s) - \mu_n} \right|.$$

Now we have that  $\operatorname{Re}(s) = (\mu_{k_0+1} + \mu_{k_0})/2$ , and thus we obtain

$$\begin{aligned} & \|(sI - A)^{-1}\| \\ & \leq K \left[ \left[ 0 + \frac{1}{\operatorname{Re}(s) - \mu_{k_0+1}} \right] + \left[ \frac{1}{\operatorname{Re}(s) - \mu_{k_0+1}} - \frac{1}{\operatorname{Re}(s) - \mu_{k_0}} \right] \right. \\ & \quad \left. + \left[ \frac{1}{\operatorname{Re}(s)} - \frac{1}{\operatorname{Re}(s) - \mu_{k_0}} \right] \right] \\ & \leq K \left[ \frac{8}{\mu_{k_0} - \mu_{k_0+1}} + \frac{1}{|\operatorname{Re}(s)|} \right]. \end{aligned}$$

Now the sequence  $\{\mu_n\} = \{-4^n\}$  satisfies

$$\frac{1}{\mu_n - \mu_{n+1}} = \frac{5/3}{|\mu_n + \mu_{n+1}|}.$$

So we see that

$$\|(sI - A)^{-1}\| \leq \frac{40K}{3|\mu_{k_0} + \mu_{k_0+1}|} + \frac{K}{|\operatorname{Re}(s)|} = \frac{23K}{3|\operatorname{Re}(s)|}.$$

This is equivalent to

$$|\operatorname{Re}(s)|^{-1} \|(sI - A)x\| \geq \frac{3}{23K} \|x\|,$$

and so (4.10) holds for  $s \in \mathbb{C}_- \setminus \mathbb{C}_g$ .

*Part 2.* In order to prove this statement we follow Theorem 4.4 of Russell and Weiss.

If (4.10) would not hold, then there would exist sequences  $\{s_n\}$  and  $\{z^n\}$  such that  $s_n \in \mathbb{C}_g$ ,  $z^n \in D(A)$ ,  $\|z^n\| = 1$ , and

$$(4.11) \quad \frac{1}{|\operatorname{Re}(s_n)|^2} \|(s_n I - A)z^n\|^2 + \frac{1}{|\operatorname{Re}(s_n)|} |Cz^n|^2 = \varepsilon_n^2,$$

where  $\varepsilon_n \geq 0$  and  $\varepsilon_n \rightarrow 0$ .

Now define

$$q^n := \frac{1}{|\operatorname{Re}(s_n)|} (s_n I - A_{s_n}) P_{V(s_n)} z^n$$

and the scalar  $\alpha_n$  such that

$$\alpha_n e_{N(s_n)} = \tilde{P}_{N(s_n)} z^n = (I - P_{V(s_n)}) z^n.$$

Thus we have that

$$\frac{1}{|\operatorname{Re}(s_n)|} (s_n I - A) z^n = \frac{s_n - \mu_{N(s_n)}}{|\operatorname{Re}(s_n)|} \alpha_n e_{N(s_n)} + q^n.$$

Now we have that

$$(4.12) \quad \|q^n\| = \left\| P_{V(s_n)} \frac{1}{|\operatorname{Re}(s_n)|} (s_n I - A) z^n \right\| \leq K \frac{1}{|\operatorname{Re}(s_n)|} \|(s_n I - A) z^n\| \leq K \varepsilon_n$$

by (4.11). For  $\alpha_n$ , we obtain

$$(4.13) \quad \begin{aligned} \left| \frac{s_n - \mu_{N(s_n)}}{\operatorname{Re}(s_n)} \alpha_n \right| &= \left\| \frac{s_n - \mu_{N(s_n)}}{\operatorname{Re}(s_n)} \alpha_n e_{N(s_n)} \right\| = \left\| \frac{s_n - \mu_{N(s_n)}}{\operatorname{Re}(s_n)} \tilde{P}_{N(s_n)} z^n \right\| \\ &= \frac{1}{|\operatorname{Re}(s_n)|} \|\tilde{P}_{N(s_n)} (s_n - A) z^n\| \\ &\leq 2K \frac{1}{|\operatorname{Re}(s_n)|} \|(s_n - A) z^n\| \leq 2K \varepsilon_n. \end{aligned}$$

By definition of  $q^n$ , we have that

$$P_{V(s_n)} z^n = |\operatorname{Re}(s_n)| (s_n I - A_{s_n})^{-1} q^n.$$

Using (4.12) and Lemma 4.2, we get

$$\|P_{V(s_n)} z^n\| \leq MK \varepsilon_n,$$

whence  $P_{V(s_n)} z^n \rightarrow 0$ . Since  $\|z^n\| = 1$ , it follows that  $\|(I - P_{V(s_n)}) z^n\| \rightarrow 1$ , i.e.,

$$(4.14) \quad \lim_{n \rightarrow \infty} |\alpha_n| = 1.$$

Together with (4.13) this implies that

$$\lim_{n \rightarrow \infty} \left| \frac{s_n - \mu_{N(s_n)}}{\operatorname{Re}(s_n)} \right| = 0.$$

It is now easy to see that

$$(4.15) \quad \lim_{n \rightarrow \infty} \left| \frac{\mu_{N(s_n)}}{\operatorname{Re}(s_n)} \right| = 1.$$

Now we turn our attention to the second term of (4.11). We have

$$\begin{aligned} Cz^n &= C(I - P_{V(s_n)})z^n + CP_{V(s_n)}z^n \\ &= \alpha_n Ce_{N(s_n)} + C_{s_n}(s_n I - A_{s_n})^{-1}(s_n I - A_{s_n})P_{V(s_n)}z^n \\ &= \alpha_n \sqrt{-\mu_{N(s_n)}} + |\operatorname{Re}(s_n)| C_{s_n}(s_n I - A_{s_n})^{-1}q^n. \end{aligned}$$

Thus we can estimate the norm of this number as

$$|Cz^n| \geq |\alpha_n \sqrt{-\mu_{N(s_n)}}| - |\operatorname{Re}(s_n)| |C_{s_n}(s_n I - A_{s_n})^{-1}q^n|.$$

Hence using Lemma 4.2, Part 2, we obtain that

$$(4.16) \quad \frac{1}{\sqrt{|\operatorname{Re}(s_n)|}} |Cz^n| \geq |\alpha_n| \left| \frac{\mu_{N(s_n)}}{\operatorname{Re}(s_n)} \right|^{\frac{1}{2}} - d \|q^n\|.$$

By (4.12) and (4.14)–(4.16), we conclude that there exists a positive number  $\kappa$  such that for  $n$  sufficiently large,

$$\frac{1}{|\operatorname{Re}(s_n)|} |Cz^n|^2 \geq \kappa.$$

On the other hand, (4.11) implies that for each  $n \in \mathbb{N}$ ,

$$\frac{1}{|\operatorname{Re}(s_n)|} |Cz^n|^2 \leq \varepsilon_n^2,$$

which is a contradiction. Therefore, (4.10) must be true.  $\square$

So we know that the system  $(A, C)$  as defined in the beginning of this section satisfies the estimate of Russell and Weiss. Suppose now that the pair would be exactly observable. Then there would exist a constant  $l > 0$  such that

$$\int_0^\infty |CT(t)x|^2 dt \geq l \|x\|^2, \quad x \in D(A).$$

Now take a finite sequence of  $\alpha_k$ 's and consider

$$x := \sum_{k=1}^n \alpha_k e_k.$$

Then the above estimate gives

$$\int_0^\infty \left| \sum_{k=1}^n \sqrt{-\mu_k} e^{\mu_k t} \alpha_k \right|^2 dt \geq l \|x\|^2.$$

However, from Nikolski and Pavlov [14] (see also Russell and Weiss [17]) we know

that there exists a constant  $l_1 > 0$  such that

$$\int_0^\infty \left| \sum_{k=1}^n \sqrt{-\mu_k} e^{\mu_k t} \alpha_k \right|^2 dt \leq l_1 \sum_{k=1}^n |\alpha_k|^2.$$

Thus we have that for any finite sequence,

$$\|x\|^2 \leq \frac{l_1}{l} \sum_{k=1}^n |\alpha_k|^2.$$

However, this implies that  $\{e_n\}$  is Hilbertian, providing the contradiction.

Thus we have disproved the conjecture of Russell and Weiss on exact observability.

**5. On left-invertibility of  $C_0$ -semigroups.** We consider a bounded  $C_0$ -semigroup  $(T_e(t))_{t \geq 0}$  with infinitesimal generator  $A_e$  on a separable Hilbert space  $Z$ . A natural question is whether uniform left-invertibility of the  $C_0$ -semigroup, that is,

$$(5.1) \quad \|T_e(t)x\| \geq c_1 \|x\|, \quad x \in Z,$$

for some  $c_1 > 0$ , is equivalent to uniform left-invertibility of  $sI - A_e$  on the open left-half plane, that is,

$$(5.2) \quad \|(sI - A_e)x\| \geq c_2 |\operatorname{Re}(s)| \|x\|, \quad x \in D(A_e), s \in \mathbb{C}_-,$$

for some constant  $c_2 > 0$ .

In van Neerven [13] it is shown that (5.1) implies (5.2). Van Neerven considered only the case of a semigroup of isometries, but the general case can be proved in a similar way. If  $(T_e(t))_{t \geq 0}$  can be extended to a group or if  $\mathbb{C}_-$  is contained in the resolvent set of  $A$ , then (5.2) implies (5.1); see van Casteren [2, 3] or Zwart [25].

We now show that in general (5.2) does not imply (5.1). Consider the operators  $A$  and  $C$  of section 3, and let  $(T(t))_{t \geq 0}$  denote the exponentially stable  $C_0$ -semigroup generated by  $A$ . We now define the semigroup  $(T_e(t))_{t \geq 0}$  on  $H \oplus L^2(0, \infty)$  by

$$T_e(t) \begin{pmatrix} x \\ f \end{pmatrix} := \begin{pmatrix} T(t)x \\ CT(t-\cdot)x|_{[0,t]} + f(\cdot-t)|_{[t,\infty)} \end{pmatrix}.$$

In Grabowski and Callier [5] it is shown that  $(T_e(t))_{t \geq 0}$  is a uniformly bounded  $C_0$ -semigroup on  $H \oplus L^2(0, \infty)$ , and that the infinitesimal generator  $A_e$  of  $(T_e(t))_{t \geq 0}$  is given by

$$A_e \begin{pmatrix} x \\ f \end{pmatrix} := \begin{pmatrix} Ax \\ -\dot{f} \end{pmatrix}, \quad \begin{pmatrix} x \\ f \end{pmatrix} \in D(A_e),$$

$$D(A_e) := \left\{ \begin{pmatrix} x \\ f \end{pmatrix} \mid x \in D(A), f, \dot{f} \in L^2(0, \infty), \right.$$

$$\left. f \text{ is absolutely continuous and } f(0) = Cx \right\}.$$

Next we calculate the norm of  $\|(sI - A_e)\begin{pmatrix} x \\ f \end{pmatrix}\|$ . For  $s = s_r + is_i \in \mathbb{C}_-$  we have

$$\begin{aligned} & \left\| (sI - A_e) \begin{pmatrix} x \\ f \end{pmatrix} \right\|^2 \\ &= \|(sI - A)x\|^2 + \|sf + \dot{f}\|_{L^2(0,\infty)}^2 \\ &= \|(sI - A)x\|^2 + |s|^2\|f\|_{L^2(0,\infty)}^2 + \|\dot{f}\|_{L^2(0,\infty)}^2 \\ &\quad + 2s_r \operatorname{Re}(\langle f, \dot{f} \rangle_{L^2(0,\infty)}) + is_i(\langle f, \dot{f} \rangle_{L^2(0,\infty)} - \langle \dot{f}, f \rangle_{L^2(0,\infty)}) \\ &= \|(sI - A)x\|^2 + \|is_i f + \dot{f}\|_{L^2(0,\infty)}^2 + s_r^2\|f\|_{L^2(0,\infty)}^2 + 2s_r \operatorname{Re}(\langle f, \dot{f} \rangle_{L^2(0,\infty)}) \\ &= \|(sI - A)x\|^2 + \|is_i f + \dot{f}\|_{L^2(0,\infty)}^2 + s_r^2\|f\|_{L^2(0,\infty)}^2 \\ &\quad + s_r \int_0^\infty \frac{d}{dt} \langle f(t), f(t) \rangle dt \\ &= \|(sI - A)x\|^2 + \|is_i f + \dot{f}\|_{L^2(0,\infty)}^2 + s_r^2\|f\|_{L^2(0,\infty)}^2 - s_r \|Cx\|^2, \end{aligned}$$

because  $f(0) = Cx$  and  $f, \dot{f} \in L^2(0, \infty)$ . Thus

$$\begin{aligned} & \left\| (sI - A_e) \begin{pmatrix} x \\ f \end{pmatrix} \right\|^2 \\ & \geq \|(sI - A)x\|^2 + |\operatorname{Re}(s)|^2\|f\|_{L^2(0,\infty)}^2 + |\operatorname{Re}(s)|\|Cx\|^2 \\ & \geq c_2 |\operatorname{Re}(s)|^2 \left\| \begin{pmatrix} x \\ f \end{pmatrix} \right\|^2 \quad (\text{using Lemma 4.3}), \end{aligned}$$

where  $c_2$  is independent of  $x$  and  $f$ . This shows that (5.2) holds. Assuming (5.1) holds as well, we get

$$\left\| T_e(t) \begin{pmatrix} x \\ f \end{pmatrix} \right\| \geq c_1 \left\| \begin{pmatrix} x \\ f \end{pmatrix} \right\|, \quad t \geq 0, x \in H, f \in L^2(0, \infty),$$

for some constant  $c_1 > 0$ . Thus

$$(5.3) \quad \|T(t)x\|^2 + \|CT(\cdot)x\|_{L^2(0,t)}^2 = \left\| T_e(t) \begin{pmatrix} x \\ 0 \end{pmatrix} \right\|^2 \geq c_1 \|x\|^2, \quad x \in H, t \geq 0.$$

Using that  $(T(t))_{t \geq 0}$  is exponentially stable, we get  $\lim_{t \rightarrow \infty} \|T(t)x\|^2 = 0$ , and so letting  $t$  to infinity in (5.3) gives

$$\|CT(\cdot)x\|_{L^2(0,\infty)} \geq \sqrt{c_1} \|x\|, \quad x \in H,$$

which says that the pair  $(A, C)$  is exactly observable. However, this is in contradiction with section 3, where we showed that the pair  $(A, C)$  is not exactly observable. Thus (5.2) holds, but (5.1) is not valid.

We conclude this section with a positive result; it shows that (5.2) implies (5.1) if the constant  $c_2$  satisfies  $c_2 \geq 1$ .

**PROPOSITION 5.1.** *Let  $(T_e(t))_{t \geq 0}$  be a bounded  $C_0$ -semigroup with infinitesimal generator  $A_e$  on a separable Hilbert space  $Z$ . If (5.2) holds with  $c_2 \geq 1$ , then (5.1) holds as well.*

*Proof.* If  $c_2 \geq 1$ , then it is easy to see that (5.2) implies that

$$\|(sI - A_e)x\| \geq |\operatorname{Re} s| \|x\|, \quad s \in \mathbb{C}_-,$$

for all  $x \in D(A)$ . Choosing  $s < 0$  and taking the square of the above equation gives

$$\|(sI - A_e)x\|^2 \geq s^2\|x\|^2.$$

Using the fact that  $Z$  is a Hilbert space gives that the above inequality is equivalent to

$$s^2\|x\|^2 - 2s \operatorname{Re}\langle x, A_ex \rangle + \|A_ex\|^2 \geq s^2\|x\|^2,$$

which is equivalent to

$$-2s \operatorname{Re}\langle x, A_ex \rangle + \|A_ex\|^2 \geq 0.$$

Since this must hold for all negative  $s$ , we see that

$$\operatorname{Re}\langle x, A_ex \rangle \geq 0.$$

We now consider the function  $f(t) := \|T_e(t)x\|^2$ . Taking the derivative of  $f$  gives

$$\dot{f}(t) = 2 \operatorname{Re}\langle T_e(t)x, A_e T_e(t)x \rangle \geq 0.$$

Hence  $f$  is nondecreasing, and thus

$$\|T_e(t)x\|^2 = f(t) \geq f(0) = \|x\|^2.$$

Since  $x$  was arbitrary, we have shown the result.  $\square$

#### REFERENCES

- [1] N.-E. BENAMARA AND N. NIKOLSKI, *Resolvent test for similarity to a normal operator*, Proc. London Math. Soc., 78 (1999), pp. 585–626.
- [2] J. A. VAN CASTEREN, *Operators similar to unitary or selfadjoint ones*, Pacific J. Math., 104 (1983), pp. 241–255.
- [3] J. A. VAN CASTEREN, *Boundedness properties of resolvents and semigroups of operators*, in Linear Operators (Warsaw, 1994), Banach Center Publ. 38, Polish Acad. Sci., Warsaw, 1997, pp. 59–74.
- [4] R. F. CURTAIN AND H. ZWART, *An Introduction to Infinite-Dimensional Linear Systems Theory*, Texts Appl. Math. 21, Springer-Verlag, New York, 1995.
- [5] P. GRABOWSKI AND F. M. CALLIER, *Admissible observation operators, semigroup criteria of admissibility*, Integral Equations Operator Theory, 25 (1996), pp. 182–198.
- [6] B. JACOB AND J. R. PARTINGTON, *The Weiss conjecture on admissibility of observation operators for contraction semigroups*, Integral Equations Operator Theory, 40 (2001), pp. 231–243.
- [7] B. JACOB AND J. R. PARTINGTON, *Admissibility of control and observation operators for semigroups: A survey*, in Proceedings of the IWOTA 2002, J. A. Ball, J. W. Helton, M. Klaus, and L. Rodman, eds., Birkhäuser Verlag, 2004, to appear.
- [8] B. JACOB, J. R. PARTINGTON, AND S. POTT, *Admissible and weakly admissible observation operators for the right shift semigroup*, Proc. Edinburgh Math. Soc., 45 (2002), pp. 353–362.
- [9] B. JACOB AND H. ZWART, *Exact observability of diagonal systems with a finite-dimensional output operator*, Systems Control Lett., 43 (2001), pp. 101–109.
- [10] B. JACOB AND H. ZWART, *Exact observability of diagonal systems with a one-dimensional output operator*, Appl. Math. Comput. Sci., 11 (2001), pp. 1277–1283.
- [11] B. JACOB AND H. ZWART, *A Hautus test for infinite-dimensional systems*, in Unsolved Problems in Mathematical Systems and Control Theory, V. Blondel and A. Megretski, eds., Princeton University Press, Princeton, NJ, 2004, to appear.
- [12] C. LE MERDY, *The Weiss conjecture for bounded analytic semigroups*, J. London Math. Soc., 67 (2003), pp. 715–738.



- [13] J. VAN NEERVEN, *The Asymptotic Behaviour of Semigroups of Linear Operators*, Oper. Theory Adv. Appl. 88, Birkhäuser, Basel, 1996.
- [14] N. K. NIKOL'SKIĬ AND B. S. PAVLOV, *Bases of eigenvectors of completely nonunitary contractions and the characteristic function*, Math. USSR-Izvestija, 4 (1970), pp. 91–134.
- [15] J. R. PARTINGTON AND G. WEISS, *Admissible observation operators for the right shift semigroup*, Math. Control Signals Systems, 13 (2000), pp. 179–192.
- [16] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer-Verlag, Berlin, 1983.
- [17] D. L. RUSSELL AND G. WEISS, *A general necessary condition for exact observability*, SIAM J. Control Optim., 32 (1994), pp. 1–23.
- [18] I. SINGER, *Bases in Banach Spaces I*, Springer-Verlag, Berlin, 1970.
- [19] G. WEISS, *Admissibility of input elements for diagonal semigroups on  $l^2$* , Systems Control Lett., 10 (1988), pp. 79–82.
- [20] G. WEISS, *Admissible observation operators for linear semigroups*, Israel J. Math., 65 (1989), pp. 17–43.
- [21] G. WEISS, *Two conjectures on the admissibility of control operators*, in Estimation and Control of Distributed Parameter Systems, F. Kappel and W. Desch, eds., Birkhäuser Verlag, Basel, 1991, pp. 367–378.
- [22] G. WEISS, *A powerful generalization of the Carleson measure theorem?*, in Open Problems in Mathematical Systems Theory and Control, V. Blondel, E. Sontag, M. Vidyasagar, and J. Willems, eds., Springer-Verlag, London, 1999, pp. 267–272.
- [23] Q. ZHOU AND M. YAMAMOTO, *Hautus condition on the exact controllability of conservative systems*, Internat. J. Control, 67 (1997), pp. 371–379.
- [24] H. ZWART, *Sufficient Conditions for Admissibility*, Memorandum 1547, Department of Applied Mathematics, University of Twente, 2000, available online from <http://www.math.utwente.nl/publications>.
- [25] H. ZWART, *On the invertibility and bounded extension of  $C_0$ -semigroups*, Semigroup Forum, 63 (2001), pp. 153–160.
- [26] H. ZWART, B. JACOB, AND O. STAFFANS, *Weak admissibility does not imply admissibility for analytic semigroups*, Systems Control Lett., 48 (2003), pp. 341–350.