

COVID-19: Factors Associated with Psychological Distress, Fear, and Coping Strategies among Community Members Across 17 Countries

Muhammad Aziz Rahman (✉ drazizdmc@gmail.com)

Federation University Australia <https://orcid.org/0000-0003-1665-7966>

Sheikh Mohammed Shariful Islam

Deakin University

Patraporn Tungpunkom

Chiang Mai University

Farhana Sultana

Telstra Health

Sheikh M Alif

Monash University

Biswajit Banik

Federation University Australia

Masudus Salehin

Federation University Australia

Bindu Joseph

Federation University Australia

Louisa Lam

Federation University Australia

Mimmie Claudine Watts

Federation University Australia

Sabria Jihan Khan

The University of Melbourne

Sherief Ghozy

Mayo Clinic Hospital

Sek Ying Chair

The Chinese University of Hong Kong

Wai Tong Chien

The Chinese University of Hong Kong

Carlos Schoenfeldt-Lecuona

University of Ulm: Universitat Ulm

Nashwa El-Khazragy

Ain Shams University

Ilias Mahmud

Qassim University

Adhra Hilal Al Mawali

Ministry of Health Oman

Turkiya Saleh Al Maskari

Oman College of Health Sciences

Rayan Jafnan Alharbi

Jazan University

Amr Hamza

University of Aleppo

Mohamad Ali Keblawi

University of Aleppo

Majeda Hammoud

Kuwait University

Asmaa M Elaidy

Al-Azhar University

Agus Dwi Susanto

Universitas Indonesia

Ahmed Suparno Bahar Moni

Universiti Sains Malaysia

Alaa Ashraf AlQurashi

King Fahad Medical City

Almajdoub Ali

Berga General Hospital

Amit Wazib

Enam Medical College and Hospital

Cattaliya Siripattarakul Sanluang

Chiang Mai University

Deena H Elson

Abu Dhabi University

Farhana Yasmin

Lahore Garrison University

Feni Fitriani Taufik

Universitas Indonesia

Manal Al Kloub

The Hashemite University

Mara Gerbabe Ruiz

Oman College of Health Sciences

Mohamed Elsayed

University of Ulm: Universitat Ulm

Nael Kamel Eltewacy

Beni-Suef University: Beni Suef University

Nahed Al Laham

Al Azhar University - Gaza

Natalia Oli

Kathmandu Medical College

Ramy Abdelnaby

RWTH Aachen University Medical Faculty: Rheinisch-Westfälische Technische Hochschule Aachen Medizinische Fakultät

Rania Dweik

Abu Dhabi University

Ratree Thongyu

Rangsit University

Sami Almustanyir

Saudi Arabia Ministry of Health

Shaila Rahman

Enam Medical College and Hospital

Sirirat Nitayawan

Chiang Mai University

Sondos Al-Madhoun

Al Azhar University - Gaza

Suwit Inthong

Chiang Mai University

Talal Ali Alharbi

King Saud University

Tamanna Bahar

National Institute of Cancer Research and Hospital

Tribowo Tuahta Ginting

Persahabatan Hospital

Wendy M. Cross

Federation University Australia

Research

Keywords: COVID-19, coronavirus, mental health, psychological distress, fear, coping, resilience

Posted Date: July 7th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-654989/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: The current pandemic of COVID-19 impacted the psychological wellbeing of populations globally.

Objectives: We aimed to examine the extent and identify associated factors of psychological distress, fear of COVID-19, and coping.

Methods: We conducted a cross-sectional study across 17 countries during Jun-2020 to Jan-2021. Levels of psychological distress (Kessler Psychological Distress Scale), fear of COVID-19 (Fear of COVID-19 Scale), and coping (Brief Resilient Coping Scale) were assessed.

Results: A total of 8,559 people participated; mean age (\pm SD) was 33(\pm 13) years, 64% were females and 40% self-identified as frontline workers. More than two-thirds (69%) experienced moderate-to-very high levels of psychological distress, which was 46% in Thailand and 91% in Egypt. A quarter (24%) had high levels of fear of COVID-19, which was as low as 9% in Libya and as high as 38% in Bangladesh. More than half (57%) exhibited medium to high resilient coping; the lowest prevalence (3%) was reported in Australia and the highest (72%) in Syria. Being female (AOR 1.31 [95% CIs 1.09-1.57]), perceived distress due to change of employment status (1.56 [1.29-1.90]), comorbidity with mental health conditions (3.02 [1.20-7.60]) were associated with higher levels of psychological distress and fear. Doctors had higher psychological distress (1.43 [1.04-1.97]), but low levels of fear of COVID-19 (0.55 [0.41-0.76]); nurses had medium to high resilient coping (1.30 [1.03-1.65]).

Conclusions: The extent of psychological distress, fear of COVID and coping varied by country; however, some groups were more vulnerable than others. There is an urgent need to prioritise health and well-being of these people through well-designed intervention that may need to be tailored to meet country specific requirements.

Introduction

The COVID-19 pandemic, with more than 172 million cases and 3.7 million deaths by early June-2021, has occurred in waves.[1] The first wave raised the alarm of what was imminent; the second wave identified the in-country differences in incidence, prevalence and mortality rates as well as health system gaps, notwithstanding policy failures; while the third wave further exposed varying social, financial, policy and failures in the health system management on the global scale.

COVID-19 impacted psychological wellbeing of global populations. Studies revealed that COVID-19 pandemic affected people in discrete ways across the world and exposed varying degrees of vulnerability among divergent community members. Evidence links emotional stress to disasters, quarantine and lockdown, where people in uncertain situations lose the power to predict and control their lives under conditions of threat.[2] Prevalence of psychological distress, anxiety and depression during the COVID-19 pandemic was reported as 50%, 27% and 28% respectively, in a systematic review with 398,771 participants.[3] A recent review of the psychological effects of COVID-19 related lockdown reported many negative psychological effects associated with quarantine including fear, stress, insomnia, depression, frustration, and anger and some of those persisted post quarantine period.[4]

Factors associated with psychological wellbeing during the current COVID-19 pandemic are diverse. However, the primary reasons for COVID-related stress were associated with contracting the virus, related complications, restrictions and mandated lockdowns, social isolation, financial loss, lack of income and disruption of daily routines which have been observed globally.[5] Moreover, critical incidents such as deaths of family members, pre-existing stressors, financial impact, being older and migrant, lack of access to basic needs were substantial grounds for poor mental health outcomes.[6] A recent Australian study also found that people with higher psychological distress increased smoking and alcohol consumption during the pandemic period; females and people with pre-existing mental health conditions were more likely to experience higher levels of psychological distress.[7]

COVID-19 is unpredictable. Varying degrees of lockdown or isolation measures are implemented nationally, depending on the stage of the pandemic. Most of the published studies have examined psychological impacts of COVID-19 in a single country or small communities. Evidence from multicultural communities on a global scale is lacking. As COVID-19 continues to peak in many countries, it is imperative that ongoing planning with mental health support strategies and early identification of psychological distress are realised, because people have the ability to normalise stressful situations when they have access to support networks and resources.[8] Our study aimed to examine the extent of and the factors associated with psychological distress, the level of fear of COVID-19 and coping strategies amongst a diverse range of community people in multi-country settings.

Materials And Methods

Study design and settings

We conducted a cross-sectional study across 17 countries utilizing web-based online platforms. Participating countries included Australia, Bangladesh, Egypt, China (Hong Kong), Indonesia, Jordan, Kuwait, Libya, Malaysia, Nepal, Oman, Pakistan, Palestine, Saudi Arabia, Syria, Thailand, and the United Arab Emirates (UAE).

Study population

Adults aged ≥ 18 years, living in the participating countries, able to respond to an online questionnaire in English/ Arabic/ Thai/ Nepali were eligible. Thus, study participants included general community members, healthcare professionals, patients, university students and staff. Patients were defined as individuals who attended a general practice or an allied healthcare setting in the previous four weeks at the time of data collection. Frontline or essential service workers were defined as individuals who self-identified themselves as being in contact with patients/clients during the pandemic period.

Sampling

Sample size was calculated using OpenEpi. Study population and estimated prevalence of stress varied across the participating countries. Therefore, keeping the population size as 100,000,000, assuming 50% prevalence of stress globally, 95% confidence intervals and 80% power, the estimated minimum sample size was 385. That number was the highest possible number, even if the population size and the prevalence of stress varied across countries. Therefore, careful consideration and taking into account the opinion of the cooperating countries, we agreed a minimum sample size of 385 participants for each collaborating country.

Data collection

An online link was created with a structured survey questionnaire using the Google form. Data were collected in Jun-2020 in Australia, Aug-Sep-2020 in Bangladesh and Malaysia, and during Nov-2020 to Jan-2021 for the other 14 countries. A separate link was created for each language (English, Arabic, Thai and Nepali). The plain language information statement (PLIS) and the consent form appeared on the first screen. Only participants, who provided consent and met the eligibility criteria, could move to the next screen. The subsequent seven screens contained the full study questionnaire, comprising of 39 questions. All responses were anonymous.

The English version of the PLIS, consent form and the study questionnaire were translated into other languages as mentioned above, back-translated to English, reviewed and pilot-tested by the team of local lead investigators for Arabic (Egypt, Saudi Arabia, UAE), Thai (Thailand) and Nepali (Nepal) versions. An invitation with the online survey link and QR code were shared using different social media platforms, online community networks, staff and student email databases of participating universities/hospitals. Text messages using SMS, Viber, WhatsApp were also shared. Flyers containing the QR codes of the study were also distributed and posted in university/healthcare settings. The survey was open to minimise selection bias, so anyone having the survey link could participate in the study; and no incentives were provided for participation in the study.

Study tool

The structured survey questionnaire was adapted from the previous study conducted in Australia.[7] The survey questionnaire was pre-tested across different electronic devices. Psychological distress was measured using the Kessler Psychological Distress Scale (K-10),[9] fear was measured using the Fear of COVID-19 Scale (FCV-19S),[10] and coping was measured using Brief Resilient Coping Scale (BRCS).[11] Reliability of those tools in the English version was examined in the Australian study, and it was found that they worked for migrants and non-migrants.[12]

Data analyses

The database was downloaded from the Google platform and Stata statistical software Stata/SE V.15.0 for Windows (StataCorp, College Station, USA, 2017) was used for data analyses. Descriptive statistics, including frequencies and percentages, were generated for categorical variables; means and standard deviations (SD) were generated for continuous variables. Psychological distress (based on the K-10 scoring) was categorised into low (score 10–15) and moderate to very high (score 16–50), fear of COVID-19 (based on the FCV-19S scoring) was categorised into low (score 7–21) and high (score 22–35), and coping (based on the BRCS scoring) was categorised into low (score 4–13) and medium to high (score 14–20).

Univariate and multivariate logistic regression analyses were conducted to examine the association between variables. Multivariate analyses were conducted to control potential confounders and the results are presented with odds ratios (ORs), adjusted ORs (AOR) and 95% confidence intervals (CIs). We also tested the sensitivity of analyses by excluding the non-significant association from the univariate model, but no changes were observed in the adjusted model. We investigated potential effect modification between age groups, gender and psychological distress, fear of COVID-19 and coping strategies. The additive log risk model was compared with multiplicative odds ratio model using the likelihood ratio test and Bayesian information criterion. A cut-off of $p < 0.05$ was considered as statistically significant. For the country-wise comparison, we selected the reference country based on the lowest prevalence of moderate to very high psychological distress, lowest prevalence of high level of fear of COVID-19 and lowest prevalence of medium to high resilience coping, then we organised other countries chronologically for each outcome based on the scores prior to conducting the multivariate analyses.

Ethics

Ethics approval was obtained from the Human Research Ethics Committee from each participating country. The survey was voluntary in nature and participants got the opportunity to have informed decision to participate in the study. Privacy and confidentiality of the collected data were maintained.

Results

A total of 8987 individuals from 17 countries met the eligibility criteria and consented to participate in the study. However, 8559 of them (95%) completed the questionnaire and were included for analyses. Most countries contributed 6–7% of the study population except Bangladesh (11%) and Saudi Arabia (9%). Mean age (\pm SD) of the participants was 33 (\pm 13) years and two-thirds (64%) were females. More than one-third (42%) had a source of income during the pandemic, while 51% had their jobs adversely affected by COVID-19. More than one-third (40%) self-identified as frontline or essential service workers, which included 14% doctors and 16% nurses. Only 4% reported having a history of psychiatric or mental health issues. The majority (81%) had never been smokers, and only 11% reported drinking alcohol in the last four weeks prior to data collection. One in five participants ($n = 1780$; 21%) had direct contact and 952 (11%) participants had indirect contact with known/suspected COVID-19 cases. About 6% tested positive for COVID-19, and 14% reported self-isolating before receiving negative test results. A third of the population ($n = 2752$; 33%) visited a healthcare provider (and were defined as 'patients' in this study) and one in ten participants ($n = 1081$; 13%) used healthcare service due to COVID-19 related stress in the last six months. Table 1 shows the characteristics of the study population.

Table 1
Characteristics of the study population

Characteristics	Total, n(%)
Total study participants	8559
Age (in years)	7665
Mean (\pm SD)	33.3 (12.5)
IQR (25th percentile to 75th percentile)	23–41
Age groups	7664
18–29 years	3683 (48.1)
30–59 years	3646 (47.6)
\geq 60 years	335 (4.4)
Gender	8475
Male	3016 (35.6)
Female	5459 (64.4)
Country of residence	8559
Australia	587 (6.9)
Bangladesh	962 (11.2)
Egypt	416 (4.9)
Hong Kong	555 (6.5)
Indonesia	541 (6.3)
Jordan	538 (6.3)
Kuwait	417 (4.9)
Libya	114 (1.3)
Malaysia	720 (8.4)
Nepal	311 (3.6)
Oman	437 (5.1)
Pakistan	418 (4.9)
Palestine	417 (4.9)
Saudi Arabia	803 (9.4)
Syria	408 (4.8)
Thailand	498 (5.8)
UAE	417 (4.9)
Born in the same country of residence	8463
No	1310 (15.3)
Yes	7153 (83.6)

Characteristics	Total, n(%)
Living status	8441
Live without family members	1908 (22.6)
Live with family members	6533 (77.4)
Highest educational/vocational qualification	8449
Primary/Grade 1 to 6	62 (0.7)
Secondary/Higher Secondary/Grade 7 to 12	1546 (18.3)
Certificate/Diploma/Trade qualifications	877 (10.4)
Bachelor/Masters/PhD	5964 (70.6)
Current employment condition	8206
Unemployed/Housewife/Home maker/Home duties (No source of income)	643 (7.8)
Jobs affected by COVID-19 (lost job/working hours reduced/afraid of job loss)	4148 (50.5)
Have an income source (employed/Government benefits)	3415 (41.6)
Perceived distress due to change of employment status	7268
A little to none	4712 (61.8)
Moderate to a great deal	2916 (38.2)
Improved working situation due to change of employment situation	5822
A little to none	4473 (76.8)
Moderate to a great deal	1349 (23.2)
Self-identification as a frontline or essential service worker	8476
No	5046 (59.5)
Yes	3430 (40.1)
Self-identification as a healthcare worker	6290
No	3843 (61.1)
Yes, doctor	887 (14.1)
Yes, nurse	1032 (16.4)
Yes, other healthcare worker	528 (8.4)
COVID-19 impacted financial situation	8507
No impact	3783 (44.5)
Yes, impacted positively	1017 (12.0)
Yes, impacted negatively	3707 (43.6)
Affected by the change in financial situation	6122
Not at all	1397 (22.8)
Unsure at this time	912 (14.9)

Characteristics	Total, n(%)
Somewhat	2770 (45.2)
A great extent	1043 (17.0)
Co-morbidities	8416
No	5975 (71.0)
Mental health issue	362 (4.3)
Other co-morbidity	2079 (24.7)
Co-morbidities	8416
No	5975 (71.0)
Single co-morbidity	1547 (19.3)
Multiple co-morbidities	474 (5.9)
Smoking	8507
Never smoker	6910 (81.2)
Ever smoker (Daily/Non-daily/Ex)	1597 (18.8)
Increased smoking over the last 6 months	1018
No	535 (52.6)
Yes	483 (47.4)
Current alcohol drinking (last 4 weeks)	8365
No	7435 (88.9)
Yes	930 (11.1)
Increased alcohol drinking over the last 6 months	921
No	645 (70.0)
Yes	276 (30.0)
Contact with known/suspected case of COVID-19	8341
No	4899 (58.7)
Unsure	710 (8.5)
Yes, indirect contact	952 (11.4)
Yes, provided direct care	1780 (21.3)
Experience related to COVID-19 pandemic (multiple responses possible)	8171
No known exposure to COVID-19	6337 (77.6)
Tested positive for COVID-19	494 (6.0)
Tested negative for COVID-19 by self-isolated	1135 (13.9)
Had recent overseas travel history and was in quarantine	205 (2.5)
Self-identification as a patient (visited a healthcare provider in the last 6 months)	8322

Characteristics	Total, n(%)
No	5570 (66.9)
Yes	2752 (33.1)
Healthcare service use in the last 6 months	2727
In-person visit to a healthcare provider	1896 (69.5)
Telehealth consultation/Use of national helpline	636 (23.3)
Used both services	195 (7.2)
Perceived mental health status	6290
Poor to fair	1753 (27.9)
Good to excellent	4537 (72.1)
Healthcare service use to overcome COVID-19 related stress in the last 6 months	8264
No	7183 (86.9)
Yes	1081 (13.1)
Type of healthcare service used to overcome COVID-19 related stress in the last 6 months	1041
Consulted a GP	356 (34.2)
Consulted a Psychologist	53 (5.1)
Consulted a Psychiatrist	63 (6.1)
Used specialised mental healthcare settings	26 (2.5)
Used mental health resources	93 (8.9)
Used mental health resources available through media	171 (16.4)
Used mental health support services	79 (7.6)
Used combination of services	199 (19.1)

More than two-thirds of the study participants (n = 5846; 69%) experienced moderate to very high levels of psychological distress, a quarter (n = 2066; 24%) had high levels of fear of COVID-19, and 4815 (57%) exhibited medium to high resilient coping. (Tables S.1, S.2, S.3)

Psychological distress

The univariate analyses showed reasonable evidence against the null hypothesis of no association between moderate to very high levels of psychological distress and a number of variables. (Table 2) However, when adjusted for potential confounders, being female, perceived distress due to change of employment status, self-identification as a doctor, being affected by the change of financial situation, comorbidity with mental health conditions, unsure and indirect contact with COVID-19 patient, being a patient, use of healthcare service to overcome COVID-related stress, and higher levels of fear of COVID-19 were found to be associated with moderate to very high levels of psychological distress. We did not identify any effect modification between age groups, gender, and psychological distress.

Table 2
Predictors for psychological distress among the study participants (based on the K-10 score)

Characteristics	Low (score 10–15)		Moderate to Very High (score 16–50)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Age groups	2434	32.1	5157	67.9						
18–29 years	775	21.1	2884	78.8	Ref			Ref		
30–59	1429	39.7	2170	60.3	< 0.001	0.41	0.37–0.45	< 0.001	0.50	0.41–0.61
≥ 60 years	230	69.1	103	30.9	< 0.001	0.12	0.08–0.15	< 0.001	0.15	0.09–0.23
Gender	2622	31.1	5810	68.9						
Male	1100	36.7	1898	63.3	Ref			Ref		
Female	1522	28	3912	71.9	< 0.001	1.50	1.36–1.64	0.003	1.31	1.09–1.57
Born in the same country of residence	2611	31	5807	68.9						
No	421	32.7	864	67.2	Ref			Ref		
Yes	2190	30.7	4943	69.3	0.118	1.06	0.96–1.18	0.193	1.18	0.92–1.52
Living status	2609	31.1	5790	68.9						
Live without family members	608	32.1	1289	67.9	Ref			Ref		
Live with family members	2001	30.9	4501	69.2	0.133	1.09	0.97–1.24	0.064	1.25	0.99–1.56
Highest educational/vocational qualification	2603	30.9	5803	69.03						
Primary/Grade 1 to 6	20	33.9	39	66.1	Ref			Ref		
Secondary/Higher Secondary/Grade 7 to 12	373	24.2	1168	75.8	0.100	1.61	0.91–2.83	0.375	0.53	0.13–2.14
Certificate/Diploma/Trade qualifications	269	30.9	601	68.1	0.605	1.16	0.65–2.06	0.231	0.43	0.11–1.72
Bachelor/Masters/PhD	1941	32.7	3995	67.3	0.848	1.06	0.61–1.84	0.247	0.44	0.11–1.75
Current employment condition	2565	31.4	5597	68.5						
Unemployed/Housewife/Home maker/Home duties (No source of income)	242	37.6	401	62.4	Ref			Ref		
Jobs affected by COVID-19 (lost job/working hours reduced/afraid of job loss)	1499	36.4	2623	63.6	0.481	1.06	0.89–1.26	No estimates due to small number		
Have an income source (employed/Government benefits)	824	24.3	2573	75.4	< 0.001	1.88	1.58–2.25	0.003	1.35	1.10–1.63

Characteristics	Low (score 10–15)		Moderate to Very High (score 16–50)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Perceived distress due to change of employment status	2317	30.5	5300	69.6						
A little to none	1735	36.8	2970	63.1	Ref			Ref		
Moderate to a great deal	582	19.9	2330	80.01	< 0.001	2.38	2.1–2.61	< 0.001	1.56	1.29–1.90
Improved working situation due to change of employment status	1730	29.7	4092	70.3						
A little to none	1373	30.6	3100	69.3	Ref			Ref		
Moderate to a great deal	357	26.5	992	73.5	0.022	1.23	1.07–1.41	0.723	0.97	0.80–1.18
Self-identification as a frontline or essential service worker	2621	31.1	5823	68.9						
No	1588	31.6	3437	68.4	Ref			Ref		
Yes	1033	30.2	2386	69.7	0.084	1.07	0.98–1.19	0.830	0.98	0.79–1.21
Self-identification as a healthcare worker	1874	29.8	4416	70.2						
No	1072	27.8	2771	72.1	Ref			Ref		
Yes, doctor	261	29.4	626	70.6	0.291	0.92	0.78–1.08	0.028	1.43	1.04–1.97
Yes, nurse	395	38.3	637	61.7	< 0.001	0.63	0.54–0.72	0.375	1.13	0.86–1.5
Yes, other healthcare worker	146	27.6	382	72.4	0.893	1.01	0.82–1.25	0.521	1.11	0.81–1.52
COVID-19 impacted financial situation	2634	31.1	5845	68.9						
No impact	1479	39.2	2297	60.8	Ref			Ref		
Yes, impacted positively	292	28.7	725	71.3	< 0.001	1.59	1.37–1.86	0.330	1.14	0.88–1.48
Yes, impacted negatively	863	23.4	2823	76.6	< 0.001	2.10	1.89–2.32	0.770	1.03	0.84–1.27
Affected by the change in financial situation	1814	29.6	4308	70.4						
Not at all	690	49.4	707	50.6	Ref			Ref		
Unsure	268	29.4	644	70.6	< 0.001	2.35	1.96–2.80	< 0.001	1.69	1.32–2.16
Somewhat	710	25.6	2060	74.4	< 0.001	2.83	2.47–3.24	< 0.001	1.64	1.32–2.03

Characteristics	Low (score 10–15)		Moderate to Very High (score 16–50)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
A great extent	146	14	897	86	< 0.001	5.99	4.89–7.35	< 0.001	2.36	1.72–3.23
Co-morbidities	2601	31.1	5770	68.9						
No	1926	32.3	4020	67.6	Ref			Ref		
Psychiatric/Mental health problem	31	8.7	327	91.3	< 0.001	5.04	3.47–7.32	0.019	3.02	1.20–7.60
Other co-morbidities*	644	31.2	1423	68.8	0.436	1.04	0.94–1.17	0.147	1.30	0.91–1.82
Co-morbidities	2465	30.9	5502	69.1						
No	1926	32.4	4020	67.6	Ref			Ref		
Single co-morbidity	411	26.6	1136	73.4	0.001	1.32	1.17–1.50	0.859	0.97	0.67–1.40
Multiple co-morbidities	128	27	346	73	0.114	1.30	1.05–1.60	No estimates due to small number		
Perceived status of own mental health	1874	29.8	4416	70.2						
Poor to Fair	131	7.5	1622	92.5	Ref			Ref		
Good to Excellent	1743	38.4	2794	61.6	< 0.001	0.13	0.11–0.16	< 0.001	0.17	0.13–0.22
Smoking	2634	31.1	5846	68.9						
Never smoker	2226	32.3	4668	67.6	Ref			Ref		
Ever smoker (Daily/Non-daily/Ex)	408	25.7	1178	74.3	< 0.001	1.38	1.22–1.56	0.434	1.10	0.87–1.39
Increased smoking over the last 6 months	206	20.3	808	79.7						
No	151	28.2	384	71.9	Ref			Not included in multivariate model		
Yes	55	11.5	424	88.5	< 0.000	3.03	2.16–4.25			
Current alcohol drinking (last 4 weeks)	2583	30.9	5755	69.02						
No	2314	31.2	5104	68.7	Ref			Ref		
Yes	269	29.2	651	70.7	0.199	1.10	0.95–1.28	0.069	1.29	0.99–1.68
Increased alcohol drinking over the last 6 months	266	29.2	645	70.8						
No	235	36.9	404	63.2	Ref			Not included in multivariate model		
Yes	31	11.4	241	88.6	< 0.001	4.52	3.01–6.80			

Characteristics	Low (score 10–15)		Moderate to Very High (score 16–50)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Contact with known/suspected case of COVID-19	2574	30.9	5743	69.1						
No	1754	35.9	3127	64.1	Ref			Ref		
Unsure	141	19.9	567	80.1	< 0.001	2.26	1.85–2.73	< 0.001	1.80	1.36–2.40
Yes, had indirect contact	223	23.4	729	76.5	< 0.001	1.83	1.55–2.16	0.019	1.32	1.04–1.67
Yes, provided direct care	456	25.6	1320	74.3	< 0.001	1.63	1.44–1.85	0.814	1.03	0.81–1.30
Experience related to COVID-19 pandemic	2518	30.9	5631	69.1						
No known exposure to COVID-19	2095	33.2	4224	66.8	Ref			Ref		
Tested positive for COVID-19	124	25.2	369	74.8	< 0.001	1.48	1.2–1.82	0.988	1.00	0.72–1.38
Tested negative for COVID-19 by self-isolated	256	22.6	876	77.3	< 0.001	1.69	1.45–1.97	0.086	1.24	0.97–1.58
Had recent overseas travel history and was in quarantine	43	20.9	162	79.02	0.002	1.87	1.32–2.62	0.696	1.12	0.64–1.93
Self-identification as a patient (visited a healthcare provider in the last 6 months)	2579	31.1	5719	68.9						
No	1945	35.1	3606	64.9	Ref			Ref		
Yes	634	23.1	2113	76.9	< 0.001	1.80	1.61–2.00	< 0.001	1.67	1.40–1.99
Healthcare service use in the last 6 months	646	23.7	2079	76.3						
In-person visit to a healthcare provider	493	26.1	1401	73.9	Ref			Ref		
Telehealth consultation/Use of national helpline	120	18.9	516	81.1	< 0.001	1.51	1.21–1.89	Not included in multivariate model		
Used both services	33	16.9	162	83.1	0.005	1.72	1.17–2.54			
Level of fear of COVID-19 (FCV-19S categories)	2634	31.1	5845	68.9						
Low (score 7–21)	2328	36.3	4088	63.7	Ref			Ref		
High (score 22–35)	306	14.8	1757	85.2	< 0.001	3.27	2.87–3.73	< 0.001	3.26	2.57–4.13
Level of coping (BRCS categories)	2633	31.1	5840	68.9						

Characteristics	Low (score 10–15)		Moderate to Very High (score 16–50)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Low resilient copers (score 4–13)	1011	27.6	2648	72.4	Ref			Ref		
Medium to high resilient copers (score 14–20)	1622	33.7	3192	66.3	< 0.001	0.75	0.69–0.82	0.637	0.96	0.81–1.14
Healthcare service use to overcome COVID-19 related stress in the last 6 months	2560	31	5697	69						
No	2422	33.7	4754	66.3	Ref			Ref		
Yes	138	12.8	943	69	< 0.001	3.48	2.89–4.19	< 0.001	1.99	1.45–2.72
<i>Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient</i>										

Levels of fear

Similar to psychological distress, participants from all 17 countries demonstrated significant levels of fear to COVID 19. (Table 3) After adjusting for potential confounders, high levels of fear were associated with being aged 30–59 years, being female, perceived distress due to a change of employment status, self-identification as a frontline or essential service worker, being affected by the change of financial situation, having comorbidities, drinking alcohol in the previous four weeks, unsure contact with a COVID-19 case, health service use to overcome COVID-related stress, and having moderate to very high levels of psychological distress. We did observe some effect modification with gender and fear of COVID-19 (contact with a COVID-19 patient) (data not shown).

Table 3
Predictors for fear of COVID-19 among the study participants (based on the FCV-19S score)

Characteristics	Low (score 7–21)		High (score 22–35)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Age groups	5710	75.20	1886	24.8						
18–29 years	2777	75.8	883	24.1	Ref			Ref		
30–59 years	2661	73.8	942	26.1	0.047	1.11	1.00–1.24	0.004	1.35	1.10–1.64
≥ 60 years	272	81.6	61	18.3	0.017	0.71	0.53–0.94	0.184	1.40	0.86–2.30
Gender	6383	75.6	2055	24.4						
Male	2305	76.8	695	23.20	Ref			Ref		
Female	4078	74.9	1360	25.1	0.059	1.11	0.99–1.23	0.001	1.51	1.25–1.83
Born in the same country of residence	6365	75.5	2059	24.4						
No	933	72.4	355	27.6	Ref			Ref		
Yes	5432	76.1	1704	23.8	0.005	0.82	0.72–0.94	0.001	0.66	0.51–0.85
Living status	6354	75.6	2050	24.4						
Live without family members	1322	69.6	577	30.4	Ref			Ref		
Live with family members	5032	77.4	1473	22.6	< 0.001	0.67	0.6–0.75	0.431	1.10	0.86–1.41
Highest educational/vocational qualification	6359	75.6	2052	24.4						
Primary/Grade 1 to 6	47	79.6	12	20.3	Ref			Ref		
Secondary/Higher Secondary/Grade 7 to 12	1176	76.3	366	23.7	0.547	1.22	0.64–2.32	0.569	1.41	0.44–4.55
Certificate/Diploma/Trade qualifications	626	71.8	245	28.1	0.198	1.53	0.8–2.93	0.298	1.87	0.57–6.09
Bachelor/Masters/PhD	4510	75.9	1429	24.1	0.506	1.24	0.66–2.35	0.689	1.27	0.40–4.05
Current employment condition	6174	75.6	1994	24.4						
Unemployed/Housewife/Home maker/Home duties (No source of income)	433	67.3	210	32.6	Ref			Ref		
Jobs affected by COVID-19 (lost job/working hours reduced/afraid of job loss)	3304	80.1	821	19.9	< 0.001	0.51	0.42–0.61	No estimate due to small number		
<i>Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient</i>										

Characteristics	Low (score 7–21)		High (score 22–35)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Have an income source (employed/Government benefits)	2437	71.7	963	28.3	0.026	0.81	0.68–0.98	0.588	1.05	0.87–1.27
Perceived distress due to change of employment status	5772	75.7	1847	24.2						
A little to none	3767	80.1	939	19.9	Ref			Ref		
Moderate to a great deal	2005	68.8	908	31.2	< 0.001	1.82	1.63–2.02	< 0.001	1.52	1.27–1.82
Improved working situation due to change of employment status	4570	78.5	1251	21.5						
A little to none	3566	79.7	906	20.3	Ref			Ref		
Moderate to a great deal	1004	74.4	345	25.6	< 0.001	1.35	1.17–1.56	0.401	1.08	0.9–1.32
Self-identification as a frontline or essential service worker	6398	75.7	2052	24.3						
No	3839	76.3	1191	23.7	Ref			Ref		
Yes	2559	74.8	861	25.2	0.115	1.08	0.99–1.2	0.001	1.47	1.20–1.82
Self-identification as a healthcare worker	4950	78.7	1339	21.3						
No	2990	77.8	853	22.2	Ref			Ref		
Yes, doctor	712	80.4	174	19.6	0.096	0.86	0.71–1.03	< 0.001	0.55	0.41–0.76
Yes, nurse	838	81.2	194	18.8	0.018	0.81	0.68–0.97	0.053	0.75	0.56–1.01
Yes, other healthcare worker	410	77.6	118	22.4	0.937	1.01	0.81–1.26	0.131	0.79	0.58–1.07
COVID-19 impacted financial situation	6418	75.6	2066	24.4						
No impact	3053	80.8	725	19.2	Ref			Ref		

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	Low (score 7–21)		High (score 22–35)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Yes, impacted positively	768	75.5	249	24.5	< 0.001	1.37	1.16–1.61	0.075	1.29	0.98–1.70
Yes, impacted negatively	2597	70.4	1092	29.6	< 0.001	1.77	1.6–1.97	0.004	1.36	1.11–1.68
Affected by the change in financial situation	4813	78.6	1308	21.4						
Not at all	1201	85.9	196	14	Ref			Ref		
Unsure	724	79.4	188	20.6	< 0.001	1.59	1.28–1.98	0.149	1.23	0.93–1.64
Somewhat	2169	78.3	600	21.7	< 0.001	1.69	1.42–2.02	0.033	1.32	1.02–1.08
A great extent	719	68.9	324	31.1	< 0.001	2.76	2.26–3.37	0.021	1.44	1.06–1.96
Co-morbidities	6345	75.7	2032	24.3						
No	4645	78.1	1303	21.9	Ref			Ref		
Psychiatric/Mental health problem	248	68.7	113	31.3	< 0.001	1.62	1.29–2.05	0.984	1.00	0.64–1.60
Other co-morbidities*	1452	70.2	616	29.8	< 0.001	1.51	1.35–1.7	0.001	1.71	1.25–2.32
Co-morbidities	6059	76.1	1910	23.9						
No	4645	78.1	1303	21.9	Ref			Ref		
Single co-morbidity	1096	70.9	451	29.2	< 0.001	1.47	1.29–1.66	0.021	0.69	0.51–0.95
Multiple co-morbidities	318	67.1	156	32.9	< 0.001	1.75	1.43–2.14	No estimate due to small number		
Perceived status of own mental health	4950	78.7	1339	21.3						
Poor to Fair	1190	67.9	563	32.1	Ref			Ref		
Good to Excellent	3760	82.9	776	17.1	< 0.001	0.44	0.39–0.5	< 0.001	0.72	0.60–0.86
Smoking	6420	75.6	2065	24.3						
Never smoker	5251	76.1	1647	23.8	Ref			Ref		
Ever smoker (Daily/Non-daily/Ex)	1169	73.6	418	26.3	0.039	1.14	1.01–1.30	0.708	1.04	0.84–1.31
Increased smoking over the last 6 months	758	74.7	256	25.3						

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	Low (score 7–21)		High (score 22–35)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
No	418	78.1	117	21.9	Ref			Not included in multivariate model		
Yes	340	70.9	139	29	0.009	1.46	1.1–1.94			
Current alcohol drinking (last 4 weeks)	6309	75.6	2035	24.4						
No	5646	76.1	1776	23.9	Ref			Ref		
Yes	663	71.9	259	28.1	0.006	1.24	1.07–1.45	0.038	1.33	1.02–1.73
Increased alcohol drinking over the last 6 months	658	72.1	255	27.9						
No	511	79.7	130	20.3	Ref			Not included in multivariate model		
Yes	147	54.1	125	45.9	< 0.001	3.34	2.46–4.54			
Contact with known/suspected case of COVID-19	6292	75.6	2031	24.4						
No	3769	77.1	1117	22.9	Ref			Ref		
Unsure	488	68.8	221	31.2	< 0.001	1.53	1.29–1.82	0.006	1.41	1.10–1.80
Yes, had indirect contact	722	75.8	230	24.2	0.384	1.07	0.92–1.26	0.713	1.04	0.86–1.35
Yes, provided direct care	1313	73.9	463	26.1	0.007	1.19	1.04–1.35	0.782	0.97	0.76–1.23
Experience related to COVID-19 pandemic	6155	75.5	2000	24.5						
No known exposure to COVID-19	4833	76.4	1490	23.6	Ref			Ref		
Tested positive for COVID-19	391	79.2	103	20.8	0.170	0.85	0.68–1.07	0.175	0.80	0.57–1.11
Tested negative for COVID-19 by self-isolated	791	69.8	342	30.2	< 0.001	1.40	1.22–1.61	0.336	1.12	0.89–1.41
Had recent overseas travel history and was in quarantine	140	68.3	65	31.7	0.007	1.51	1.12–2.03	0.808	0.93	0.54–1.61
Self-identification as a patient (visited a healthcare provider in the last 6 months)	6273	75.5	2031	24.5						
No	4247	76.5	1308	23.6	Ref			Ref		
Yes	2026	73.7	723	26.3	0.006	1.16	1.04–1.29	0.217	0.90	0.76–1.06

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	Low (score 7–21)		High (score 22–35)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Healthcare service use in the last 6 months	1973	72.4	754	27.6						
In-person visit to a healthcare provider	1413	74.5	483	25.5	Ref			Ref		
Telehealth consultation/Use of national helpline	426	66.9	210	33	< 0.001	1.44	1.19–1.75	Not included in multivariate model		
Used both services	134	68.7	61	31.3	0.079	1.33	0.97–1.83			
Level of psychological distress (K10 categories)	6416	75.7	2063	24.3						
Low (score 10–15)	2328	88.4	306	11.6	Ref			Ref		
Moderate to Very High (score 16–50)	4088	69.9	1757	30.1	< 0.001	3.26	2.87–3.72	< 0.001	3.36	2.67–4.23
Level of coping (BRCS categories)	6418	75.7	2061	24.3						
Low resilient copers (score 4–13)	2647	72.2	1018	27.8	Ref			Ref		
Medium to high resilient copers (score 14–20)	3771	78.3	1043	21.7	< 0.001	0.72	0.65–0.80	< 0.001	0.74	0.63–0.87
Healthcare service use to overcome COVID-19 related stress in the last 6 months	6243	75.6	2020	24.5						
No	5595	77.9	1587	22.1	Ref			Ref		
Yes	648	59.9	433	40.1	< 0.001	2.35	2.06–2.70	< 0.001	2.42	1.96–3.01
<i>Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient</i>										

Coping strategies

Table 4 shows the univariate analyses identifying significant association between medium to high resilient coping and other variables. From the multivariate analyses, we identified that participants who were ≥ 60 years old, self-identification as a nurse, whose financial situation was impacted negatively, who perceived their own mental health as good to excellent, who had indirect contact and direct contact with known or suspected cases of COVID-19, and who visited a healthcare provider in the previous six months were more likely to have medium to high resilient coping. We did not identify any effect modification between age group, gender, and coping strategies (data not shown).

Table 4
Predictors for coping among the study participants (based on the BRCS score)

Characteristics	Low (score 4–13)		Medium to High (score 14–20)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Age groups	3247	42.8	4344	57.2						
18–29 years	1581	43.3	2074	56.7	Ref			Ref		
30–59 years	1543	42.8	2060	57.2	0.711	1.02	0.93–1.12	0.329	1.08	0.92–1.28
≥ 60 years	123	36.9	210	63.1	0.026	1.30	1.03–1.64	0.011	1.66	1.12–2.44
Gender	3640	43.2	4792	56.8						
Male	1323	44.1	1675	55.9	Ref			Ref		
Female	2317	42.6	3117	57.4	0.186	1.07	0.97–1.17	0.235	0.91	0.79–1.06
Born in the same country of residence	3635	43.2	4783	56.8						
No	649	50.4	639	49.6	Ref			Ref		
Yes	2986	41.8	4144	58.1	< 0.001	1.41	1.25–1.59	0.124	0.85	0.69–1.05
Living status	3614	43	4784	56.9						
Live without family members	812	42.7	1087	57.2	Ref			Ref		
Live with family members	2802	43.1	3697	56.9	0.780	0.99	0.89–1.1	0.106	0.85	0.7–1.04
Highest educational/vocational qualification	3622	43.1	4783	56.9						
Primary/Grade 1 to 6	30	50.8	29	49.2	Ref			Ref		
Secondary/Higher Secondary/Grade 7 to 12	673	43.7	868	56.3	0.277	1.33	0.8–2.24	0.537	1.35	0.52–3.48
Certificate/Diploma/Trade qualifications	409	47.2	458	57.7	0.585	1.16	0.69–1.96	0.871	1.08	0.42–2.81
Bachelor/Masters/PhD	2510	42.3	3428	57.7	0.187	1.41	0.85–2.36	0.583	1.30	0.51–3.32
Current employment condition	3523	43.2	4639	56.8						
Unemployed/Housewife/Home maker/Home duties (No source of income)	260	40.4	383	59.5	Ref			Ref		
Jobs affected by COVID-19 (lost job/working hours reduced/afraid of job loss)	1734	42.1	2391	57.9	0.444	0.94	0.797–1.11	No estimate due to small number		
<i>Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient</i>										

Characteristics	Low (score 4–13)		Medium to High (score 14–20)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
			1865	54.9						
Have an income source (employed/Government benefits)	1529	45.1			0.031	0.84	0.69–0.99	0.354	0.93	0.8–1.09
Perceived distress due to change of employment status	3095	40.6	4522	59.4						
A little to none	1815	38.6	2889	61.4	Ref			Ref		
Moderate to a great deal	1280	43.9	1633	56.1	< 0.001	0.80	0.73–0.88	0.030	0.82	0.68–0.98
Improved working situation due to change of employment status	2291	39.4	3528	60.6						
A little to none	1753	39.2	2717	60.8	Ref			Ref		
Moderate to a great deal	538	39.8	811	60.1	0.662	0.98	0.86–1.1	0.342	1.09	0.92–1.28
Self-identification as a frontline or essential service worker	3646	43.2	4798	56.8						
No	2155	42.9	2869	57.1	Ref			Ref		
Yes	1491	43.6	1929	56.4	0.522	0.97	0.87–1.06	0.525	0.94	0.8–1.13
Self-identification as a healthcare worker	2482	39.5	3801	60.5						
No	1578	41.1	2259	58.9	Ref			Ref		
Yes, doctor	331	37.4	555	62.6	0.040	1.17	1.01–1.36	0.417	0.90	0.70–1.16
Yes, nurse	371	35.9	661	64.1	0.003	1.24	1.08–1.44	0.029	1.30	1.03–1.65
Yes, other healthcare worker	202	38.3	326	61.7	0.209	1.13	0.94–1.36	0.280	1.15	0.90–1.48
COVID-19 impacted financial situation	3663	43.2	4815	56.8						
No impact	1613	42.8	2160	57.3	Ref			Ref		

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	Low (score 4–13)		Medium to High (score 14–20)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Yes, impacted positively			603	59.4						
	413	40.7			0.229	1.10	0.95–1.26	0.851	0.98	0.80–1.23
Yes, impacted negatively	1637	44.4	2052	55.6	0.157	0.94	0.85–1.03	< 0.001	1.37	1.16–1.62
Affected by the change in financial situation	2403	39.3	3712	60.7						
Not at all	523	37.4	874	62.6	Ref			Ref		
Unsure	385	42.4	523	57.6	0.017	0.81	0.69–0.96	0.004	0.74	0.60–0.90
Somewhat	1051	37.9	1716	62	0.732	0.98	0.86–1.12	0.398	0.92	0.78–1.14
A great extent	444	42.6	599	57.4	0.010	0.81	0.69–0.95	0.151	0.83	0.66–1.07
Co-morbidities	3630	43.4	4741	56.6						
No	2458	41.4	3488	58.7	Ref			Ref		
Psychiatric/Mental health problem	223	62.5	134	37.5	< 0.001	0.42	0.33–0.52	0.431	0.85	0.57–1.27
Other co-morbidities*	949	45.9	1119	54.1	< 0.001	0.82	0.73–0.91	0.324	1.15	0.88–1.50
Co-morbidities	3321	41.7	4642	58.3						
No	2458	41.3	3488	58.7	Ref			Ref		
Single co-morbidity	674	43.6	873	56.4	0.113	0.91	0.81–1.02	0.149	0.82	0.62–1.09
Multiple co-morbidities	189	40.2	281	59.8	0.633	1.05	0.87–1.27	No estimate due to small number		
Perceived status of own mental health	2482	39.5	3801	60.5						
Poor to Fair	913	52.1	839	47.8	Ref			Ref		

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	Low (score 4–13)		Medium to High (score 14–20)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Good to Excellent	1569	34.6	2962	65.4	< 0.001	2.05	1.83–2.3	< 0.001	1.97	1.70–2.30
Smoking	3665	43.2	4814	56.8						
Never smoker	2912	42.2	3982	57.8	Ref			Ref		
Ever smoker (Daily/Non-daily/Ex)	753	47.5	832	52.5	< 0.001	0.81	0.72–0.90	0.533	1.06	0.88–1.28
Increased smoking over the last 6 months	447	44.2	565	55.8						
No	234	43.7	301	56.3	Ref			Ref		
Yes	213	44.6	264	55.4	0.770	0.96	0.75–1.23	Not included in multivariate model		
Current alcohol drinking (last 4 weeks)	3595	43.1	4743	56.8						
No	3089	41.6	4328	58.4	Ref			Ref		
Yes	506	54.9	415	45.1	< 0.001	0.59	0.50–0.66	0.532	0.93	0.74–1.17
Increased alcohol drinking over the last 6 months	499	54.7	413	45.3						
No	310	48.4	330	51.7	Ref			Not included in multivariate model		
Yes	189	69.5	83	30.5	< 0.001	0.40	0.31–0.56			
Contact with known/suspected case of COVID-19	3578	43	4739	56.9						
No	2223	45.5	2662	54.5	Ref			Ref		
Unsure	333	46.9	376	53	0.470	0.94	0.81–1.1	0.297	0.90	0.73–1.1
Yes, had indirect contact	353	37.3	594	62.7	< 0.001	1.41	1.21–1.63	0.004	1.33	1.10–1.62
Yes, provided direct care	669	37.7	1107	62.3	< 0.001	1.37	1.22–1.53	< 0.001	1.45	1.19–1.77
Experience related to COVID-19 pandemic	3497	42.9	4652	57.1						
No known exposure to COVID-19	2739	43.4	3580	56.6	Ref			Ref		
Tested positive for COVID-19	184	37.3	310	62.7	0.008	1.29	1.07–1.56	0.259	0.86	0.65–1.12

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	Low (score 4–13)		Medium to High (score 14–20)		Unadjusted analyses			Adjusted analyses		
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Tested negative for COVID-19 by self-isolated	480	42.4	651	57.6	0.571	1.03	0.91–1.18	0.012	0.78	0.64–0.95
Had recent overseas travel history and was in quarantine	94	45.8	111	54.2	0.476	0.90	0.68–1.2	0.312	0.80	0.51–1.24
Self-identification as a patient (visited a healthcare provider in the last 6 months)	3564	42.9	4734	57.1						
No	2466	44.4	3089	55.6	Ref			Ref		
Yes	1098	40.1	1645	59.9	0.001	1.20	1.09–1.31	0.012	1.20	1.04–1.28
Healthcare service use in the last 6 months	1089	40	1633	59.9						
In-person visit to a healthcare provider	730	38.5	1165	61.5	Ref			Ref		
Telehealth consultation/Use of national helpline	277	43.5	359	56.5	0.025	0.82	0.67–0.97	Not included in multivariate model		
Used both services	82	42.9	109	57.1	0.234	0.83	0.62–1.13			
Level of psychological distress (K10 categories)	3659	43.2	4814	56.8						
Low (score 10–15)	1011	38.4	1622	61.6	Ref			Ref		
Moderate to Very High (score 16–50)	2648	45.4	3192	54.6	< 0.001	0.74	0.67–0.81	0.498	0.95	0.81–1.11
Level of fear of COVID-19 (FCV-19S categories)	3665	43.2	4814	56.7						
Low (score 7–21)	2647	41.2	3771	58.8	Ref			Ref		
High (score 22–35)	1018	49.4	1043	50.6	< 0.001	0.71	0.64–0.78	< 0.001	0.72	0.61–0.85
Healthcare service use to overcome COVID-19 related stress in the last 6 months	3546	42.9	4718	57.1						
No	3049	42.4	4134	57.6	Ref			Ref		
Yes	497	45.9	584	54	0.030	0.87	0.76–0.99	0.375	0.91	0.75–1.12
<i>Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient</i>										

Country-wise findings

Country-wise analyses (Table 5) showed that moderate to very high levels of psychological distress was common in all 17 countries. The lowest prevalence (46%) was reported from Thailand and the highest (91%) from Egypt. When other countries were compared considering Thailand as the baseline, it was found that participants from 10 countries (Hong Kong, Oman, Libya, Kuwait, Saudi Arabia, UAE, Jordan, Syria, Palestine and Egypt), demonstrated statistically significant high psychological distress. Prevalence on high levels of fear of COVID-19 varied across 17 countries (Libya: 9%, Bangladesh: 38%). Participants from four countries (Oman, Indonesia, Hong Kong and Pakistan) exhibited higher levels of fear of COVID-19 compared to the participants from Libya. Finally, participants from 12 countries (Jordan, Egypt, Saudi Arabia, Kuwait, Hong Kong, UAE, Palestine, Thailand, Oman, Nepal, Indonesia and Syria) demonstrated statistically significant medium to high resilience coping compared to those from Australia.

Table 5

Country-wise analyses for high psychological distress, fear of COVID-19 and coping among the study participants

Characteristics	K-10 Score				Unadjusted analyses			Adjusted analyses		
	Low (score 10–15)		Moderate to Very High (score 16–50)		p	ORs	95% CIs	p	AORs	95% CIs
	n	%	n	%						
Country of residence	2634		5846							
Thailand	269	54.1	229	45.9	Ref			Ref		
Hong Kong	256	46.1	299	53.9	0.011	1.37	1.08–1.75	< 0.001	1.93	1.37–2.73
Indonesia	223	41.2	318	58.8	< 0.001	1.68	1.31–2.14	0.071	1.44	0.97–2.15
Oman	180	41.2	257	58.8	< 0.001	1.68	1.30–2.17	< 0.001	2.20	1.50–3.25
Nepal	119	38.3	192	61.7	< 0.001	1.90	1.42–2.52	0.253	1.28	0.84–1.95
Malaysia	273	37.9	447	62.1	< 0.001	1.92	1.53–2.42	Not included in multivariate model		
Australia	203	37.5	339	62.5	< 0.001	1.96	1.53–2.51	Not included in multivariate model		
Libya	38	33.3	76	66.7	< 0.001	2.35	1.53–3.60	< 0.001	3.54	1.91–6.56
Kuwait	132	31.6	285	68.4	< 0.001	2.54	1.93–3.33	< 0.001	3.06	2.05–4.58
Bangladesh	284	30.1	644	69.4	< 0.001	2.67	2.12–3.31	Not included in multivariate model		
Pakistan	121	28.9	297	71.1	< 0.001	2.88	2.19–3.80	0.105	1.40	0.93–2.11
Saudi Arabia	225	28	578	71.9	< 0.001	3.02	2.38–3.81	< 0.001	2.82	1.99–4.01
UAE	89	21.3	328	78.6	< 0.001	4.32	3.23–5.80	< 0.001	3.68	2.31–5.86
Jordan	80	14.9	458	85.1	< 0.001	6.72	5.01–9.04	< 0.001	6.83	4.05–11.5
Syria	53	13	355	87.0	< 0.001	7.87	5.61–11.0	< 0.001	6.05	3.59–10.2
Palestine	50	12	367	88.0	< 0.001	8.62	6.11–12.2	< 0.001	4.80	2.87–8.02
Egypt	39	9.4	377	90.6	< 0.001	11.4	7.81–16.5	< 0.001	9.43	5.33–16.7
Characteristics	FCV-19S Score				Unadjusted analyses			Adjusted analyses		
<i>Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient</i>										

Characteristics	K-10 Score				Unadjusted analyses			Adjusted analyses		
	Low (score 10–15)		Moderate to Very High (score 16–50)		p	ORs	95% CIs	p	AORs	95% CIs
	n	%	n	%						
	High (score 22–35)									
	Low (score 7–21)									
	n	%	n	%	p	ORs	95% CIs	p	AORs	95% CIs
Country of residence	6420		2066							
Libya	104	91.2	10	8.8	Ref			Ref		
Saudi Arabia	714	88.9	89	11.1	0.458	1.30	0.65–2.57	0.669	0.85	0.40–1.82
Thailand	427	85.7	71	14.3	0.123	1.73	0.86–3.46	0.937	1.03	0.47–2.28
Kuwait	347	83.2	70	16.8	0.037	2.1	1.04–4.22	0.395	1.40	0.64–3.07
Oman	351	80.3	86	19.7	0.008	2.55	1.28–5.08	0.044	2.23	1.02–4.88
Jordan	429	79.7	109	20.3	0.005	2.64	1.34–5.23	0.477	0.74	0.33–1.70
Nepal	248	79.7	63	20.3	0.007	2.64	1.31–5.35	0.057	2.16	0.98–4.80
Syria	324	79.6	83	20.4	0.006	2.67	1.33–5.32	0.455	1.35	0.62–2.93
Palestine	330	79.1	87	20.8	0.004	2.74	1.37–5.47	0.844	1.09	0.49–2.42
UAE	320	76.7	97	23.3	0.001	3.15	1.59–6.27	0.561	1.27	0.58–2.81
Indonesia	405	74.8	136	25.1	< 0.001	3.50	1.77–6.88	0.006	2.86	1.35–6.08
Malaysia	525	72.9	195	27.1	< 0.001	3.87	1.98–7.54	Not included in multivariate model		

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	K-10 Score				Unadjusted analyses			Adjusted analyses		
	Low (score 10–15)		Moderate to Very High (score 16–50)		p	ORs	95% CIs	p	AORs	95% CIs
	n	%	n	%						
Egypt	288	69.2	128	30.8	< 0.001	4.62	2.34–9.14	0.055	2.13	0.98–4.62
Hong Kong	382	68.8	173	31.2	< 0.001	4.71	2.40–9.24	0.003	3.21	1.47–7.01
Australia	374	68.1	175	31.8	< 0.001	4.87	2.49–9.54	Not included in multivariate model		
Pakistan	281	67.2	137	32.8	< 0.001	5.07	2.57–10.0	0.002	3.41	1.58–7.33
Bangladesh	571	61.5	357	38.4	< 0.001	6.50	3.35–12.6	Not included in multivariate model		
Characteristics	BRCS Score				Unadjusted analyses			Adjusted analyses		
	Low (score 4–13)		Medium to High (score 14–20)		p	ORs	95% CIs	p	AORs	95% CIs
	n	%	n	%						
Country of residence	3665		4815							
Australia	534	97.3	15	2.7	Ref			Ref		
Libya	70	61.9	43	38.1	< 0.001	21.86	11.6–41.4	Not included in multivariate model		
Pakistan	221	52.8	197	47.1	< 0.001	31.73	18.3–54.9	0.210	1.40	0.83–2.36
Jordan	252	46.8	286	53.2	< 0.001	40.40	23.5–69.4	0.014	1.99	1.15–3.43
Egypt	191	45.9	225	54.1	< 0.001	41.93	24.2–72.6	0.003	2.28	1.33–3.88
Saudi Arabia	354	44.1	448	55.8	< 0.001	45.05	26.5–76.7	0.016	1.84	1.12–3.02
Kuwait	183	43.8	234	56.1	< 0.001	45.52	26.3–78.8	0.009	2.01	1.20–3.40
Bangladesh	398	42.8	530	57.1	< 0.001	47.41	27.9–80.5	Not included in multivariate model		
Hong Kong	230	41.4	325	58.5	< 0.001	50.30	29.3–86.3	0.002	2.29	1.34–3.91
UAE	151	36.5	262	63.4	< 0.001	61.77	35.6–107	< 0.001	2.64	1.53–4.55

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Characteristics	K-10 Score				Unadjusted analyses			Adjusted analyses		
	Low (score 10–15)		Moderate to Very High (score 16–50)		p	ORs	95% CIs	p	AORs	95% CIs
	n	%	n	%						
Palestine	152	36.5	264	63.4	< 0.001	61.83	35.7–107	< 0.001	2.90	1.68–4.99
Thailand	175	35.1	323	64.9	< 0.001	65.71	38.1–113	0.004	2.18	1.29–3.70
Malaysia	251	34.8	469	65.1	< 0.001	66.52	38.9–114	Not included in multivariate model		
Oman	137	31.4	300	68.7	< 0.001	77.96	44.9–135	< 0.001	3.80	2.21–6.54
Nepal	97	31.2	214	68.8	< 0.001	78.54	44.6–138	< 0.001	3.45	1.99–5.98
Indonesia	156	28.8	385	71.2	< 0.001	87.86	50.9–152	< 0.001	4.16	2.51–6.92
Syria	113	27.7	295	72.3	< 0.001	92.93	53.2–162	< 0.001	4.94	2.89–8.46

Adjusted for: age, gender, smoking, alcohol intake, living status, place of birth, country, education, employment status, employment stress, healthcare worker, financial impact, contact with COVID-19 case, experience due to COVID-19 and self-identification as a patient

Discussion

To our knowledge, this study is one of the few large-scale global cross-sectional studies that assessed psychological distress, levels of fear, and coping strategies and their associated factors among community members, frontline workers, and patients across 17 countries during the first and second wave of the COVID-19 pandemic. We found that more than two-thirds (69%) participants experienced moderate to very high levels of psychological distress and about a quarter (24%) had a high level of fear of COVID-19. Despite having moderate to high levels of psychological distress and fear, more than half of the participants (57%) reported medium to high levels of resilient coping.

Findings from this study were consistent with the previous Australian study.[7] Similarly, the previous research found almost a third of the participants (33%) experienced high to very high levels of psychological distress; however, they found more participants experienced a high level of fear of COVID-19 (32%), while our study found only 24%. Furthermore, the Australian study found that almost all participants (97%) had low resilient coping, whereas this global study found 57% participants had medium to high resilient coping. Learning from previous successful experiences that enable people to cope better could explain this discrepancy.[13] When participants from the Australian study were faced with COVID-19 at an earlier stage, participants of this study (that included participants who were confronted with both 1st and 2nd waves) might have learned how to cope with all kinds of relevant practices from the 1st wave of the pandemic (such as social distancing, home quarantine, or lockdown, hand hygiene and wearing masks), leading them to high resilient coping and less fear of COVID-19. However, the context was interplayed with distress and fear in this study. It was found that participants who perceived distress due to change of their employment, whose financial situation was affected greatly, and had unsure contact with COVID-19 were more likely to have higher psychological distress and fear.

We found that females had higher psychological distress and fear of COVID-19. This finding is consistent with the Australian study,[7] and studies from elsewhere.[14] They also had a greater chance of loneliness, specifically for young people aged 18–29 years or those 60+.[15] Such distress and fear could also be related to ‘infodemic’ through the increased use of social media.[16] Having a history of mental illness and experience of family violence was shown to aggravate depression, anxiety and stress amongst women during the pandemic.[17] In addition, concerns of exposure to COVID-19 amongst family members could have accentuated their anxiety and distress. Women tend to have more care giving roles in a family and often prioritise health concerns of family members over their own.[7] That warrants improved awareness amongst women regarding regular health assessment and accessing resources to support their wellbeing.

Interestingly, participants who perceived their mental health as good to excellent, even though their financial situation was impacted negatively, and who had contact with COVID-19 patients indirectly or directly were more likely to have medium to high resilient coping. This was especially true for participants who self-identified themselves as nurses. This is incongruent with the Australian study, though consistent with earlier studies.[18] Our findings reflected that participants perceived mental resiliency could be the internal psychological aid that eases their reality during the pandemic despite having higher psychological distress. Enhancing resilience could be a possible intervention to enable people to cope with the mental health impact of COVID-19. Such a psychological resilience model has been developed and tested for its effectiveness in China and was found to improve the overall mental health of the target population during the COVID-19 pandemic.[19]

In our study, doctors had higher psychological distress, but low levels of fear of COVID-19; nurses had medium to high resilient coping. A recent systematic review of 24 studies with 13,731 health and social care workers showed that female nurses, comorbidities, lack of personal protective equipment, concerns about family, fear of infections and close contact with COVID-19 patients were the predictors for poor mental wellbeing amongst healthcare workers.[20] Low levels of fear amongst the frontline healthcare workers in our study were likely due to their prolonged professional exposure with COVID-19 patient management. Due to the heterogeneity of the health systems and varying availability of resources across participating countries, healthcare workers experienced catastrophic situations during the surge of pandemic period, which could have resulted in high resilience amongst the nurses.

Our findings showed that participants who had comorbidities and those who had a mental illness showed higher psychological distress and fear. These groups were more vulnerable under pandemic guidelines (such as social distancing, working from home), which potentially raised the risks of relapse, especially those who were mentally ill and who needed primary caregivers. Generally, evidence from clinical settings and literature indicated that mentally ill persons who lived alone would have more psychotic relapses than those being cared for by primary caregivers.[21] Medication adherence for this group of patients could have been challenging without caregiving provision.[22] Accessibility to the health care system was more difficult because most healthcare workers were overloaded with COVID-19 infected patients and the related tasks, therefore, managing chronic diseases was not a priority. In addition, lockdown policies impacted transportation and public facilities were closed in many instances. Previous evidence also suggested that people with stressful situations and pre-existing medical problems had higher levels of depression and anxiety.[23] Telemedicine to replace face-to-face consultations had been established in many countries including Australia during COVID-19. The effect of such an alternative healthcare delivery system needs to be evaluated further, especially its impact on people with non-communicable diseases and/or mental illness who need continuing care.

Eighty-one percent of the study population were never smokers. Those who smoked and drank alcohol, reported increased use of tobacco and alcohol (47% and 30% respectively) in the last six months. Moreover, drinking behavior was also associated with higher levels of fear of COVID-19. The findings were consistent with the previous Australian study and that risky behavior was associated with a higher impact on psychological distress.[12] A study conducted in China also found that participants who had a history of smoking could escalate the severe symptoms of COVID-19 once hospitalized and possibly required ventilator equipment.[24] A Polish study also revealed that current alcohol drinkers were less able to find

positives about the pandemic (positive reframing) and coping.[25] An effective coping strategy needs to be developed and implemented to target populations using social media to prevent unhealthy coping behaviors.

The change of employment status and an uncertain financial situation were associated with higher psychological distress and fear. In our study, 51% participants reported that their jobs were affected by COVID-19, due to losing jobs, reduced working hours, or being afraid of job loss. That was probably one of the significant indicators of mental wellbeing, impacted by COVID-19 on people's socioeconomic status around the globe and consistent with a study conducted among Israeli youths (20–35 years old).[26] The need for urgent action to support and elevate economic assistance, especially for those whose job was impacted negatively from the pandemic, is critical. While business enterprises were freezing around the globe due to restrictions related to controlling the spread of coronavirus, basic needs are essential, specifically for vulnerable groups to prevent psychological crisis which could potentially lead to suicidal attempts or even suicide.

The impact of COVID-19 on the psychological wellbeing was unprecedented and was different from country to country. Therefore, findings from 17 countries were found to be diverse. In our study, country specific results on psychological distress showed a specific trend. For example, more than two-thirds of the participants reported moderate to very high level of psychological distress who were living in countries with war/conflict (Syria, Palestine, Libya and the Middle East [Saudi Arabia, UAE, Jordan and Kuwait]) followed by South Asia (Pakistan, Nepal and Bangladesh) and least by the participants from South-East Asian countries (Thailand, Hong Kong and Indonesia). However, participants from Oman, Australia and Egypt could not be fitted into any of those categories. It can be assumed that such disparities could be related to geography, access to healthcare, having comorbidities, living in war-torn and conflicting countries.[27] It can be also assumed that uncertainties about COVID-19, its progression and rapid mutation, availability and access to varied range of evidence could also contribute to the report of diverse country-wise findings of moderate to high level of psychological distress. Similar higher levels of anxiety were reported in Hong Kong during the SARS epidemic amongst medicine students and students living in the area where there was a rapid spread of infection.[28]

Participants from the Middle East and war-torn countries reported less fear compared to the participants from South-East Asian countries and South Asia. The exact reasons for this could not be elicited from our study, however the reasons can be explained by two factors, firstly, high standard care and public health in Saudi Arabia, Kuwait and Oman, and success of early interventions, such as early lockdown reducing the transmission of COVID-19. It can be further emphasized that participants from war-torn countries already have experienced high levels of fear for prolonged periods which might cause an idiosyncratic response to the pandemic.[29] Further study on war-torn countries could provide more insights. Higher levels of fear of COVID-19 among participants from South-East Asian countries could be explained by their previous traumatic experience from SARS and H1N1 pandemics, which disproportionately affected South-East Asian countries.[28]

In our study, we found that more than half of the participants (57%) showed medium to high resilience towards the pandemic. Interestingly, participants from Australia found to struggle most, despite reports of very low levels of community transmission compared to the other 16 countries included in this study. This could be explained by the fact that Australian participants were predominantly from Victoria, the only state in Australia which was affected by the second wave of COVID-19 during the study period, which caused statewide strict lockdown, social isolation, job loss.[12] Nonetheless, despite potential lack of capacity and resources to manage pandemics, participants from war-torn countries like Palestine and Syria were found to have higher coping compared to the participants from Australia. It was beyond the scope of our study to examine the reasons for such findings. Research from Syria reported strategies to contain COVID-19, such as effective use of social media tools, community engagement, bottom-up approach from the local government, and coordinated support by the international donor communities.[30]

Limitations

We had some limitations in our study. The use of online surveys potentially introduced selection bias, as participants were limited to those who could access the internet only; therefore, the generalizability of the findings needs to be interpreted with

caution. Drawing predictive conclusions based on the differences is difficult and is a limitation of a cross sectional study design. Nevertheless, under the circumstances of movement restriction and social distancing, an online survey was the most robust available option during the pandemic to fulfill our research objectives. From the perspective of multi-country study (17 countries), the multicultural background, the difference of policies and compliance of public health actions that varied across participating countries, might also impact on the examined variables (psychological distress, fear, and ways of coping). We, therefore, adjusted the variable 'country' during the multivariate analyses to control potential confounding effects. Furthermore, the collaboration from researchers across 17 countries and the achievement of the target sample size during the crisis period of COVID-19 showed significant power to test our hypotheses and provided key information to plan interventions as needed.

Conclusions

Our study examined the extent and identified factors associated with psychological distress, fear of COVID-19 and coping amongst diverse community members across 17 countries. Females and people with existing mental health issues were the most vulnerable group of populations for adverse psychological impact of COVID-19. There is an urgent need to prioritise these vulnerable population; adequate medical and social support along with specific health promotion policies should be considered within the strategic response to the ongoing pandemic and future crises. Future studies should focus on developing strategies to enhance resilience and examining effectiveness of such interventions. Besides global strategies to address psychological impact, policy makers in each country should revisit existing support structures and enhance them during this critical period. Innovative approaches are needed to enhance effective coping and social support to alleviate impact and prevent emotional crisis for vulnerable people in the longer term.

Declarations

Acknowledgements

We would like to acknowledge the support from all the collaborators and individuals who supported us in collecting data from the participating countries. We would like to convey our gratitude to all the study participants, who donated their valuable time within the crisis period of coronavirus pandemic and kindly participated in this global study.

Authors' contributions

MAR was the lead investigator, who conceptualised the study and had the responsibility to coordinate with the study investigators for data collection in 17 countries. MAR, SMSI, FS, SMA, BB, MS and WMC had substantial contribution to the conception or design of the study. Data collection was coordinated by the respective country lead: MAR in Australia, PT in Thailand, SG in Egypt, SYC in Hong Kong, AHAM and TSAM in Oman, AH and MAK in Syria, MH in Kuwait, ADS in Indonesia, ASBM in Malaysia, AA in Libya, DHE and RD in United Arab Emirates, FY in Pakistan, MAK in Jordan, NAL in Palestine, NO in Nepal, SA in Saudi Arabia, SR and TB in Bangladesh. MAR, FS, SMA and SJK cleaned, analysed and interpreted data. MAR, PT, FS, SMA, BB, MS, BJ, LL, MCW and SJK wrote the manuscript. SG, SYC, WTC, CS-L, NE-K, IM, AHAM, TSAM, RJA, AH, MAK, MH, AME provided critical feedback on narrative structure or methods or results. MAR, SMSI, FS and WMC finalised the manuscript and revised it critically for important intellectual content. All authors had full access to all the data in the study, accepted responsibility for its validity and had final responsibility for the decision to submit for publication.

Funding

We did not receive funding for this investigation.

Availability of data and materials

All data relevant to the study are included in the article.

Ethics approval and consent to participate

Ethics approval was obtained from the Human Research Ethics Committee from each participating country: Australia (Federation University Australia, Ref: B20-036), Bangladesh (Enam Medical College, Ref: EMC/ERC/2020/08-2), Egypt (Ain Shams University, Ref: FMASU R 121/2020), Hong Kong (The Chinese University of Hong Kong, Ref: SBRE-20-172), Indonesia (Universitas Indonesia, Ref: KET-1425/UN2.F1/ETIK/PPN.00.02/2020), Jordan (The Hashemite University), Kuwait (Kuwait University, Ref: VDR/EC/3693), Libya (Al-Brega General Hospital), Malaysia (Universitas Sains Malaysia, Ref: USM/JEPeM/COVID19-40), Nepal (Kathmandu Medical College Public Ltd., Ref: 2611202004), Oman (Ministry of Health, Ref: MoH/CSR/20/24012), Pakistan (Lahore Garrison University), Saudi Arabia (Ministry of Health, Ref: 20-605E), Syria (University of Aleppo), Thailand (Chiang Mai University, Ref: AF 04-021), United Arab Emirates (Abu Dhabi University, Ref: CoHS-20-20-00024). Each study participant read the consent form along with plain language summary and ticked their consent in the online form prior to accessing the study questionnaire.

Consent for publication

Data were collected anonymously, therefore, no identifying information were collected from the study participants.

Competing interests

The authors confirm that there are no known conflicts of interest associated with this publication.

References

1. WHO Coronavirus (COVID-19) Dashboard [<https://covid19.who.int/>]
2. Van Rheenen TE, Meyer D, Neill E, Phillipou A, Tan EJ, Toh WL, Rossell SL: Mental health status of individuals with a mood-disorder during the COVID-19 pandemic in Australia: Initial results from the COLLATE project. *J Affect Disord* 2020, 275:69-77.
3. Nochaiwong S, Ruengorn C, Thavorn K, Hutton B, Awiphan R, Phosuya C, Ruanta Y, Wongpakaran N, Wongpakaran T: Global prevalence of mental health issues among the general population during the coronavirus disease-2019 pandemic: a systematic review and meta-analysis. *Sci Rep* 2021, 11(1):10173.
4. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, Rubin GJ: The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020, 395(10227):912-920.
5. Wang C, Tee M, Roy AE, Fardin MA, Srichokchatchawan W, Habib HA, Tran BX, Hussain S, Hoang MT, Le XT et al: The impact of COVID-19 pandemic on physical and mental health of Asians: A study of seven middle-income countries in Asia. *PLoS One* 2021, 16(2):e0246824.
6. Gloster AT, Lamnisis D, Lubenko J, Presti G, Squatrito V, Constantinou M, Nicolaou C, Papacostas S, Aydin G, Chong YY et al: Impact of COVID-19 pandemic on mental health: An international study. *PLoS One* 2020, 15(12):e0244809.
7. Rahman MA, Hoque N, Alif SM, Salehin M, Islam SMS, Banik B, Sharif A, Nazim NB, Sultana F, Cross W: Factors associated with psychological distress, fear and coping strategies during the COVID-19 pandemic in Australia. *Global Health* 2020, 16(1):95.
8. Pfefferbaum B, North CS: Mental Health and the Covid-19 Pandemic. *N Engl J Med* 2020, 383(6):510-512.

9. Furukawa TA, Kessler RC, Slade T, Andrews G: The performance of the K6 and K10 screening scales for psychological distress in the Australian National Survey of Mental Health and Well-Being. *Psychol Med* 2003, 33(2):357-362.
10. Ahorsu DK, Lin CY, Imani V, Saffari M, Griffiths MD, Pakpour AH: The Fear of COVID-19 Scale: Development and Initial Validation. *Int J Ment Health Addict* 2020:1-9.
11. Sinclair VG, Wallston KA: The development and psychometric evaluation of the Brief Resilient Coping Scale. *Assessment* 2004, 11(1):94-101.
12. Rahman MA, Salehin M, Islam SMS, Alif SM, Sultana F, Sharif A, Hoque N, Nazim NB, Cross WM: Reliability of the tools used to examine psychological distress, fear of COVID-19 and coping amongst migrants and non-migrants in Australia. *Int J Ment Health Nurs* 2021, 30(3):747-758.
13. Chen L, Qu L: From Stressful Experiences to Depression in Chinese Migrant Children: The Roles of Stress Mindset and Coping. *Front Psychol* 2021, 12:601732.
14. Gausman J, Langer A: Sex and Gender Disparities in the COVID-19 Pandemic. *J Womens Health (Larchmt)* 2020, 29(4):465-466.
15. Wickens CM, McDonald AJ, Elton-Marshall T, Wells S, Nigatu YT, Jankowicz D, Hamilton HA: Loneliness in the COVID-19 pandemic: Associations with age, gender and their interaction. *J Psychiatr Res* 2021, 136:103-108.
16. Stuart J, O'Donnell K, O'Donnell A, Scott R, Barber B: Online Social Connection as a Buffer of Health Anxiety and Isolation During COVID-19. *Cyberpsychol Behav Soc Netw* 2021.
17. Sediri S, Zgueb Y, Ouanes S, Ouali U, Bourgou S, Jomli R, Nacef F: Women's mental health: acute impact of COVID-19 pandemic on domestic violence. *Arch Womens Ment Health* 2020, 23(6):749-756.
18. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P: Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun* 2020, 88:901-907.
19. He Z, Chen J, Pan K, Yue Y, Cheung T, Yuan Y, Du N, Zhao Y, Feng Y, Zhou D et al: The development of the 'COVID-19 Psychological Resilience Model' and its efficacy during the COVID-19 pandemic in China. *Int J Biol Sci* 2020, 16(15):2828-2834.
20. De Kock JH, Latham HA, Leslie SJ, Grindle M, Munoz SA, Ellis L, Polson R, O'Malley CM: A rapid review of the impact of COVID-19 on the mental health of healthcare workers: implications for supporting psychological well-being. *BMC Public Health* 2021, 21(1):104.
21. Tsang HW, Fung KM, Corrigan PW: Psychosocial and socio-demographic correlates of medication compliance among people with schizophrenia. *J Behav Ther Exp Psychiatry* 2009, 40(1):3-14.
22. Clement J, Jacobi M, Greenwood BN: Patient access to chronic medications during the Covid-19 pandemic: Evidence from a comprehensive dataset of US insurance claims. *PLoS One* 2021, 16(4):e0249453.
23. Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, Roma P: A Nationwide Survey of Psychological Distress among Italian People during the COVID-19 Pandemic: Immediate Psychological Responses and Associated Factors. *Int J Environ Res Public Health* 2020, 17(9).
24. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC et al: Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020, 382(18):1708-1720.

25. Chodkiewicz J, Talarowska M, Miniszewska J, Nawrocka N, Bilinski P: Alcohol Consumption Reported during the COVID-19 Pandemic: The Initial Stage. *Int J Environ Res Public Health* 2020, 17(13).
26. Achdut N, Refaeli T: Unemployment and Psychological Distress among Young People during the COVID-19 Pandemic: Psychological Resources and Risk Factors. *Int J Environ Res Public Health* 2020, 17(19).
27. Lawrence D, Kisely S: Inequalities in healthcare provision for people with severe mental illness. *J Psychopharmacol* 2010, 24(4 Suppl):61-68.
28. Wong TW, Gao Y, Tam WWS: Anxiety among university students during the SARS epidemic in Hong Kong. *Stress Health* 2007, 23(1):31-35.
29. Al-Ashwal FY, Kubas M, Zawiah M, Bitar AN, Mukred Saeed R, Sulaiman SAS, Khan AH, Ghadzi SMS: Healthcare workers' knowledge, preparedness, counselling practices, and perceived barriers to confront COVID-19: A cross-sectional study from a war-torn country, Yemen. *PLoS One* 2020, 15(12):e0243962.
30. Ekzayez A, Al-Khalil M, Jasiem M, Al Saleh R, Alzoubi Z, Meagher K, Patel P: COVID-19 response in northwest Syria: innovation and community engagement in a complex conflict. *J Public Health (Oxf)* 2020, 42(3):504-509.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS.docx](#)