

9-3-2002

Creating nanocavities of tunable sizes: Hollow helices

Bing Gong

State University of New York at Buffalo, bgong@chem.buffalo.edu

Huaqiang Zeng

State University of New York at Buffalo

Jin Zhu

Chinese Academy of Sciences

Lihua Yuan


State University of New York at Buffalo

Yaohua Han

State University of New York at Buffalo

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/chemzeng>

 Part of the [Analytical Chemistry Commons](#), [Materials Chemistry Commons](#), and the [Physical Chemistry Commons](#)

Gong, Bing; Zeng, Huaqiang; Zhu, Jin; Yuan, Lihua; Han, Yaohua; Cheng, Shizhi; Furukawa, Mako; Parra, Ruben D.; Kovalevsky, Audrey Y.; Mills, Jeffrey L.; Skrzypczak-Jankun, Ewa; Martinovic, Suzana; Smith, Richard D.; Zheng, Chong; Szyperski, Thomas; and Zeng, Xiao Cheng, "Creating nanocavities of tunable sizes: Hollow helices" (2002). *Xiao Cheng Zeng Publications*. 130.
<http://digitalcommons.unl.edu/chemzeng/130>

This Article is brought to you for free and open access by the Published Research - Department of Chemistry at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Xiao Cheng Zeng Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Bing Gong, Huaqiang Zeng, Jin Zhu, Lihua Yuan, Yaohua Han, Shizhi Cheng, Mako Furukawa, Ruben D. Parra, Audrey Y. Kovalevsky, Jeffrey L. Mills, Ewa Skrzypczak-Jankun, Suzana Martinovic, Richard D. Smith, Chong Zheng, Thomas Szyperski, and Xiao Cheng Zeng

Creating nanocavities of tunable sizes: Hollow helices

Bing Gong^{*†}, Huaqiang Zeng^{*}, Jin Zhu^{**}, Lihua Yuan^{*}, Yaohua Han^{*}, Shizhi Cheng^{*§}, Mako Furukawa[¶], Rubén D. Parra^{||}, Andrey Y. Kovalevsky^{*}, Jeffrey L. Mills^{*}, Ewa Skrzypczak-Jankun^{**}, Suzana Martinovic^{††}, Richard D. Smith^{††}, Chong Zheng^{**}, Thomas Szyperski^{*}, and Xiao Cheng Zeng[¶]

^{*}Department of Chemistry, State University of New York, Buffalo, NY 14260; [¶]Department of Chemistry, University of Nebraska, Lincoln, NE 68588; ^{**}Department of Chemistry, University of Toledo, Toledo, OH 43606; ^{††}Environmental and Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, WA 99352; and [§]Department of Chemistry, Northern Illinois University, De Kalb, IL 60115


Edited by Peter B. Dervan, California Institute of Technology, Pasadena, CA, and approved June 12, 2002 (received for review May 8, 2002)

A general strategy for creating nanocavities with tunable sizes based on the folding of unnatural oligomers is presented. The backbones of these oligomers are rigidified by localized, three-center intramolecular hydrogen bonds, which lead to well-defined hollow helical conformations. Changing the curvature of the oligomer backbone leads to the adjustment of the interior cavity size. Helices with interior cavities of 10 Å to >30 Å across, the largest thus far formed by the folding of unnatural foldamers, are generated. Cavities of these sizes are usually seen at the tertiary and quaternary structural levels of proteins. The ability to tune molecular dimensions without altering the underlying topology is seen in few natural and unnatural foldamer systems.

Based on the folding of biopolymers, Nature has developed astonishingly efficient and sophisticated strategies for generating various nanostructures. Of particular interests is the availability of a wide variety of nanosized cavities and holes that are responsible for numerous biological processes and functions. In recent years there has been intense interest in developing folding oligomers and polymers (foldamers) with unnatural backbones that adopt well-defined structures (1–3), which may eventually lead to protein-like molecular objects with sizes in the nanometer range. Many foldamer systems have been described (4–16). Despite the progress made so far, the foldamer field is still in its infancy. One daunting challenge involves the design of foldamers with cavities and holes of tunable sizes in the nanometer range, the realization of which will have far-reaching significance for not only fundamental understanding but also important applications. While cavities and holes are mostly seen at the tertiary and quaternary structural levels of biopolymers, almost all foldamers reported so far fold into secondary structures. In this article we describe a general strategy for designing folded structures that combine the features of both secondary and tertiary (or quaternary) structures. Nanosized cavities are generated by enforcing stably folded helical structures. In addition, by adjusting the curvature of the corresponding backbones, the interior diameters of these hollow helices are easily tuned. This tunability is seen in few natural or unnatural folding systems. Thus, helices with hydrophilic interior cavities of ~10 Å and >30 Å across result, yielding cavities of tunable sizes and properties. Many novel properties may emerge given, the tunable and distinct cavities associated with the corresponding folding molecules, and numerous applications in catalysis, separation, molecular recognition, and transportation can be envisioned. Phenomena associated with nanodimensions that lately have attracted intense interest and have been probed by using the highly hydrophobic carbon nanotubes (17, 18) can also be investigated on the basis of the hydrophilic cavities of these hollow helices.

Our design is based on oligoamides represented by the general structures **1a** and **1b**. The backbone of these oligomers consists of benzene rings linked by localized, intramolecularly hydrogen-bonded amide groups. On each of the benzene rings, the two amide linkages can be placed *meta* to each other (*m*-residue), leading to backbones consisting of *m*-residues (*m*-backbones, **1a**), or with some of the residues, the two amide groups can be

placed in a *para* geometry (*p*-residue), leading to backbones consisting of *m*- and *p*-residues (*mp*-backbones, **1b**).



Our previous studies have demonstrated that short oligomers with *m*-backbones adopt a well-defined crescent conformation (12, 19, 20). Our results indicated that the three-center hydrogen-bonding system, consisting of the S(5) and S(6) type (21) hydrogen-bonded rings, was particularly stable in the solid state and in solution (19, 20). It persisted in chloroform, the highly polar dimethyl sulfoxide (DMSO), and even in water (unpublished data), and this structure was confirmed by NMR, IR, and x-ray crystallographic studies on short oligomers. Extending the crescent backbones may lead to helical conformations.

Materials and Methods

Compounds. All compounds described herein gave satisfactory NMR and electrospray ionization (ESI) MS results consistent with their structures. The short oligomer intermediates were prepared by iterative coupling steps based on similar procedures described before (12).

Compound 2a. To a solution of 4,6-bis{2-[2-(2-methoxyethoxy)ethoxy]ethoxy}-1,3-benzenedicarboxylic acid (0.049 g, 0.10 mmol) and tetramer amine **4a** was added *N,N*-diisopropylethylamine (1 ml), followed by slow addition of *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (0.076 g, 0.20 mmol) in dimethylformamide (DMF; 1 ml) during 10 min at 50°C under nitrogen atmosphere. The mixture was changed into solution, which was stirred overnight. The precipitate from the solution at –20°C was collected and washed with cold DMF and ethyl acetate to give pure **2a** as a solid (0.093 g, 37%). ¹H NMR (400 MHz, CDCl₃) δ 10.23 (s, 2H), 10.05 (s, 2H), 9.77 (s, 2H), 9.62 (s, 2H), 9.22 (s, 2H), 9.10 (s, 2H), 8.95 (s, 1H), 8.91 (s, 2H), 8.55 (s, 2H), 6.76 (d, 2H, *J* = 7.0 Hz), 6.72 (d, 2H, *J* = 7.0 Hz), 6.58 (s, 1H), 6.52 (br, 4H), 6.45 (s, 2H), 4.38(br, 4H), 4.26 (br, 8H), 3.94–4.04 (m, 40H),

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: ESI, electrospray ionization; DMF, dimethylformamide; 1D and 2D, one- and two-dimensional; NOESY, nuclear Overhauser and exchange spectroscopy; NOE, nuclear Overhauser enhancement.

[†]To whom reprint requests should be addressed. E-mail: bgong@chem.buffalo.edu.

[¶]Present address: Chengdu Institute of Organic Chemistry, Chinese Academy of Sciences, P.O. Box 415, Chengdu 610041, People's Republic of China.

[§]Present address: Institute of Materia Medica, Chinese Academy of Medical Sciences, 1 Xian Nong Tan Street, Beijing 100050, People's Republic of China.

^{||}Present address: Department of Chemistry, De Paul University, 1036 West Belden Avenue, Chicago, IL 60614.