

# Criteria for age estimation in living individuals

A. Schmeling · C. Grundmann · A. Fuhrmann ·  
H.-J. Kaatsch · B. Knell · F. Ramsthaler · W. Reisinger ·  
T. Riepert · S. Ritz-Timme · F. W. Rösing ·  
K. Rötzscher · G. Geserick

Received: 30 April 2008 / Accepted: 16 May 2008 / Published online: 12 June 2008  
© Springer-Verlag 2008

**Abstract** This paper presents updated recommendations of the Study Group on Forensic Age Diagnostics for age estimations in living individuals in criminal proceedings. In order to increase the diagnostic accuracy and to improve the identification of age-relevant developmental disorders, a physical examination, an X-ray examination of the left hand, as well as a dental examination including the determination of the dental status and an X-ray of the

dentition should be performed in each case. If the skeletal development of the hand is completed, an additional radiological examination of the clavicles should be carried out. Minimum requirements for reference studies are defined and recommendable studies are listed. Instructions for the examination and the preparation of expert reports are presented. The committee of the study group organizes annual proficiency tests for quality assurance.

A. Schmeling (✉)  
Institut für Rechtsmedizin, Universitätsklinikum Münster,  
Röntgenstraße 23,  
48149 Münster, Germany  
e-mail: andreas.schmeling@ukmuenster.de

C. Grundmann  
Gesundheitsamt der Stadt Duisburg,  
Viktoriastraße 8,  
47166 Duisburg-Hamborn, Germany

A. Fuhrmann  
Zentrum für Zahn-, Mund- und Kieferheilkunde,  
Universitätsklinikum Hamburg-Eppendorf,  
Martinistraße 52,  
20246 Hamburg, Germany

H.-J. Kaatsch  
Institut für Rechtsmedizin, Universität Kiel,  
Arnold-Heller-Straße 12,  
24105 Kiel, Germany

B. Knell  
Institut für Rechtsmedizin, Universität Zürich,  
Winterthurerstraße 190/52,  
8057 Zürich, Switzerland

F. Ramsthaler  
Zentrum der Rechtsmedizin, Universität Frankfurt/Main,  
Kennedyallee 104,  
60596 Frankfurt, Germany

W. Reisinger  
Institut für Radiologie,  
Charité—Universitätsmedizin Berlin,  
Charitéplatz 1,  
10117 Berlin, Germany

T. Riepert  
Institut für Rechtsmedizin, Universität Mainz,  
Am Pulverturm 3,  
55131 Mainz, Germany

S. Ritz-Timme  
Institut für Rechtsmedizin, Universität Düsseldorf,  
Moorenstr. 5,  
40225 Düsseldorf, Germany

F. W. Rösing  
Arbeitsgruppe Anthropologie,  
Medizinische Fakultät,  
Universität Ulm,  
Parkstraße 11,  
89070 Ulm, Germany

K. Rötzscher  
Arbeitskreis für Forensische Odonto-Stomatologie,  
Wimphelingstraße 7,  
67345 Speyer, Germany

G. Geserick  
Zähringerstraße 34,  
10707 Berlin, Germany

**Keywords** Forensic age estimation · Criminal proceedings · Recommendations

## Introduction

In many European countries, the growing migration into Europe and between European countries has led to an increase in the number of foreigners who cannot present documentary evidence for their date of birth. As a result of this development, age diagnostics of living individuals in criminal proceedings has become an essential part of forensic practice [3]. In many countries, the age limits of relevance in criminal proceedings are between 7 and 21 years [2].

A transregional analysis of the current state of forensic age estimation in living individuals in the German-speaking countries was conducted on the occasion of the “Xth Lübeck Meeting of German Forensic Physicians” in December 1999. At this meeting, it was suggested to set up a study group composed of forensic physicians, dentists, radiologists, and anthropologists. The aim was to develop recommendations for age estimations in order to standardize the, as yet rather, heterogeneous procedure in setting up expert reports and to implement quality assurance in this area. The interdisciplinary “Study Group on Forensic Age Diagnostics” was constituted in Berlin on March 10, 2000.

The present recommendations apply to age estimations in criminal proceedings for determining whether an individual is of criminally responsible age or whether adult criminal law is applicable. The first version of these recommendations was passed by the members of the Study Group on Forensic Age Diagnostics on September 15, 2000 [24]. The updated recommendations were adopted on March 14, 2008.

## Examination methods

The scientific basis of age estimation is the genetic control of ontogenesis, which delimits the temporal variation of developmental stages [8, 17]. The growth curves of monozygotic twins show a high degree of correspondence. Although a range of diagnostic procedures for age estimation is available [10, 18, 19, 27], only few of these appear to be suitable for forensic application in living individuals regarding the above-mentioned relevant age limits, if one takes ethical and medicolegal aspects into account.

There is a wide agreement about the most suitable methods currently available. They involve:

- physical examination with determination of anthropometric measures (body height and weight, constitution-
- al type), inspection of signs of sexual maturation as well as identification of any age-relevant developmental disorders
- X-ray examination of the left hand
- dental examination with determination of the dental status and X-ray examination of the dentition
- if the skeletal development of the hand is completed, an additional examination of the clavicles should be carried out, preferably by means of a conventional X-ray examination and/or a computed tomography scan [9, 28]

The methods should be used together to increase the diagnostic accuracy and to improve the identification of age-relevant developmental disorders. The legitimization of the X-ray examinations is to be checked in accordance with the regulations applicable in the respective country. Further radiological features for determining the individual maturation should only be considered, if the relevant radiographs and scans are already available [6, 23].

## Reference studies

A reference study evaluates data in a random sample using a specific method. A method in this context is the conversion of an ontogenetic process into a chronological scale.

Reference studies used for forensic age estimation should meet the following requirements:

- adequate sample size, considering the number of age groups and population groups included in the study
- proven age of the subjects
- even age distribution
- analysis separately for both sexes
- information on the time of examination
- clear definition of the examined features
- detailed description of the methods
- data on the reference population regarding genetic/geographic origin, socioeconomic status, state of health
- data on the sample size, mean value, and range of scatter for each examined feature

Examples of reference studies are [4, 5, 7, 12, 13, 15, 16, 21, 25, 29, 30].

## Examination

Before taking on an expert evaluation in an individual case, one should check whether the question(s) can be answered with a sufficiently high degree of reliability by using sound scientific methods. The examinations to be performed must be justified by a court order.

The individuals to be examined must be informed about the type of examination and its purpose. The court ordering the report must be notified that an interpreter may be required. Each part of the examination should be performed by a specialist experienced in setting up expert reports and participating in regular proficiency tests for quality assurance (see below).

When ionizing radiation is applied, the national regulations must be observed. From a scientific point of view, the so-called minimization principle is valid without restrictions. It requires to perform each examination as dose saving as possible and to dispense any exposition that is not mandatory. The coordinating expert has to give a comprehensive assessment on the basis of the different parts of the evaluation performed by the respective specialists.

### Expert reports

The central forensic aspect of an expert report is to give the most probable age of the examined individual and/or the degree of probability that the stated age is the actual age or that the individual's age is above the relevant penal age limit. The expert report has to quote the methods and reference studies the age estimation is based on. For each examined feature, the report must indicate the most probable age and the range of scatter of the reference population [20]. Furthermore, it should be noted that the range of tolerance may be increased by an empirical observer error. The age-relevant variations resulting from the application of the reference studies in an individual case such as deviating genetic/geographic origin, different socioeconomic status, and with that a possibly different degree of acceleration, developmental disorders of the individual, have to be discussed in the report including their effect on the estimated age and, if possible, a quantitative assessment of any such effect should be given [1, 11, 14, 22, 26].

The individual's most likely age is estimated on the basis of all partial diagnoses and a critical discussion of the individual case. When the age estimations based on the used methods are summarized, it can generally be assumed that the range of scatter is reduced. So far, however, this reduction could only be estimated. Depending on what is requested, it may be necessary to assess the legally relevant age limits and/or the probability of the age given in the court order verbally.

### Quality assurance

The committee of the study group organizes annual proficiency tests for continuing quality assurance. An

expert may also request an evaluation of an age estimation before the report is written.

The present recommendations are revised annually by the committee of the study group with regards to new scientific and practical results and updated if required. The updated version of the recommendations is available on the website of the study group (<http://rechtsmedizin.klinik.uni-muenster.de/agfad/index.htm>).

### References

1. Cameriere R, Flores-Mir C, Mauricio F, Ferrante L (2007) Effects of nutrition on timing of mineralization in teeth in a Peruvian sample by the Cameriere and Demirjian methods. *Ann Hum Biol* 34:547–556
2. Dünkel F, van Kalmthout A, Schüler-Springorum H (1997) Entwicklungstendenzen und Reformstrategien im Jugendstrafrecht im europäischen Vergleich. *Forum, Mönchengladbach*
3. Geserick G, Schmeling A (2000) Übersicht zum gegenwärtigen Stand der Altersschätzung Lebender im deutschsprachigen Raum. In: Oehmichen M, Geserick G (eds) *Osteologische Identifikation (Research in Legal Medicine/Rechtsmedizinische Forschungsergebnisse)*. Schmidt-Römhild, Lübeck, pp 255–261
4. Greulich WW, Pyle SI (1959) *Radiographic atlas of skeletal development of the hand and wrist*. Stanford University Press, Stanford
5. Gunst K, Mesotten K, Carbonez A, Willems G (2003) Third molar root development in relation to chronological age: a large sample sized retrospective study. *Forensic Sci Int* 136:52–57
6. Jung H (2000) Strahlenrisiken durch Röntgenuntersuchungen zur Altersschätzung im Strafverfahren. *Fortschr Röntgenstr* 172: 553–556
7. Kahl B, Schwarze CW (1988) Aktualisierung der Dentitionstabelle von I. Schour und M. Massler von 1941. *Fortschr Kieferorthop* 49:432–443
8. Knussmann R (1996) *Vergleichende Biologie des Menschen. Lehrbuch der Anthropologie und Humangenetik*. Fischer, Stuttgart
9. Kreitner K-F, Schweden F, Schild HH, Riepert T, Nafe B (1997) Die computertomographisch bestimmte Ausreifung der medialen Klavikulaepiphyse—eine additive Methode zur Altersbestimmung im Adoleszentenalter und in der dritten Lebensdekade? *Fortschr Röntgenstr* 166:481–486
10. Liversidge H, Herdeg B, Rösing FW (1998) Dental age estimation of non-adults. A review of methods and principles. In: Alt KW, Rösing FW, Teschler-Nicola M (eds) *Dental anthropology. Fundamentals, limits, and prospects*. Springer, Wien
11. Meijerman L, Maat GJR, Schulz R, Schmeling A (2007) Variables affecting the probability of complete fusion of the medial clavicular epiphysis. *Int J Leg Med* 121:463–468
12. Mincer HH, Harris EF, Berryman HE (1993) The A.B.F.O. study of third molar development and its use as an estimator of chronological age. *J Forensic Sci* 38:379–390
13. Olze A, Schmeling A, Rieger K, Kalb G, Geserick G (2003) Untersuchungen zum zeitlichen Verlauf der Weisheitszahnmineralisation bei einer deutschen Population. *Rechtsmedizin* 13:5–10
14. Olze A, Schmeling A, Taniguchi M, Maeda H, van Niekerk P, Wernecke K-D, Geserick G (2004) Forensic age estimation in living subjects: the ethnic factor in wisdom tooth mineralization. *Int J Leg Med* 118:170–173
15. Olze A, Taniguchi M, Schmeling A, Zhu B-L, Yamada Y, Maeda H, Geserick G (2004) Studies on the chronology of third molar mineralization in a Japanese population. *Legal Med* 6:73–79

16. Olze A, van Niekerk P, Schmidt S, Wernecke K-D, Rösing FW, Geserick G, Schmeling A (2006) Studies on the progress of third molar mineralization in a black African population. *Homo* 57:209–217
17. Pelsmaekers B, Loos R, Carels C, Derom C, Vlietinck R (1997) The genetic contribution to dental maturation. *J Dent Res* 76:1337–1340
18. Ritz S, Kaatsch H-J (1996) Methoden der Altersbestimmung an lebenden Personen: Möglichkeiten, Grenzen, Zulässigkeit und ethische Vertretbarkeit. *Rechtsmedizin* 6:171–176
19. Ritz-Timme S, Cattaneo C, Collins MJ, Waite ER, Schütz HW, Kaatsch HJ, Borrman HI (2000) Age estimation: the state of the art in relation to the specific demands of forensic practise. *Int J Leg Med* 113:129–136
20. Rösing FW (2000) Forensische Altersdiagnose: Statistik, Arbeitsregeln und Darstellung. In: Oehmichen M, Geserick G (eds) *Osteologische Identifikation (Research in Legal Medicine/Rechtsmedizinische Forschungsergebnisse)*. Schmidt-Römhild, Lübeck, pp 263–275
21. Ruhstaller P (2006) Zahnärztliche Altersdiagnostik durch röntgenologische Evaluation der Entwicklungsstadien des unteren Weisheitszahnes: Auswertung von 1260 Orthopantomogrammen von Jugendlichen und jungen Erwachsenen. *Med Diss, Zürich*
22. Schmeling A, Reisinger W, Loreck D, Vendura K, Markus W, Geserick G (2000) Effects of ethnicity on skeletal maturation: consequences for forensic age estimations. *Int J Leg Med* 13:252–258
23. Schmeling A, Reisinger W, Wormanns D, Geserick G (2000) Strahlenexposition bei Röntgenuntersuchungen zur forensischen Altersschätzung Lebender. *Rechtsmedizin* 10:135–137
24. Schmeling A, Kaatsch H-J, Marré B, Reisinger W, Riepert T, Ritz-Timme S, Rösing FW, Röttscher K, Geserick G (2001) Empfehlungen für die Altersdiagnostik bei Lebenden im Strafverfahren. *Rechtsmedizin* 11:1–3
25. Schmeling A, Schulz R, Reisinger W, Mühler M, Wernecke K-D, Geserick G (2004) Studies on the time frame for ossification of medial clavicular epiphyseal cartilage in conventional radiography. *Int J Leg Med* 118:5–8
26. Schmeling A, Schulz R, Danner B, Rösing F (2006) The impact of economic progress and modernization in medicine on the ossification of hand and wrist. *Int J Leg Med* 120:121–126
27. Schmeling A, Geserick G, Reisinger W, Olze A (2007) Age estimation. *Forensic Sci Int* 165:178–181
28. Schulz R, Mühler M, Reisinger W, Schmidt S, Schmeling A (2008) Radiographic staging of ossification of the medial clavicular epiphysis. *Int J Leg Med* 122:55–58
29. Tanner JM, Healy MJR, Goldstein H, Cameron N (2001) *Assessment of skeletal maturity and prediction of adult height (TW3 method)*. Saunders, London
30. Thiemann H-H, Nitz I, Schmeling A (2006) *Röntgenatlas der normalen Hand im Kindesalter*. Thieme, Stuttgart, New York