

CRITICAL PERIOD PLASTICITY IN LOCAL CORTICAL CIRCUITS

Takao K. Hensch

Abstract | Neuronal circuits in the brain are shaped by experience during ‘critical periods’ in early postnatal life. In the primary visual cortex, this activity-dependent development is triggered by the functional maturation of local inhibitory connections and driven by a specific, late-developing subset of interneurons. Ultimately, the structural consolidation of competing sensory inputs is mediated by a proteolytic reorganization of the extracellular matrix that occurs only during the critical period. The reactivation of this process, and subsequent recovery of function in conditions such as amblyopia, can now be studied with realistic circuit models that might generalize across systems.

CRITICAL PERIOD

A strict time window during which experience provides information that is essential for normal development and permanently alters performance.

SENSITIVE PERIOD

A limited time during development, during which the effect of experience on brain function is particularly strong.

AMBLYOPIA

Poor vision through an eye that is otherwise physically healthy due to little or no transmission of the visual image to the brain through circuits that are hard-wired during a developmental critical period. It affects 2–5% of the population.

RIKEN Brain Science Institute, 2-1 Hirosawa, Wako-shi, Saitama, 351-0198, Japan.
e-mail: hensch@riken.jp
doi:10.1038/nrn1787

From polyglots¹ to virtuosi^{2,3}, human performance reflects the neural circuits that are laid down by early experience. Although learning is possible throughout life, there is no doubt that those who start younger fare better, and that plasticity is enhanced during specific windows of opportunity. An understanding of the neural basis of such CRITICAL OR SENSITIVE PERIODS of brain development would inform not only classroom and educational policy, but also drug design, clinical therapy and strategies for improved learning into adulthood. Although which might be the critical periods for higher cognitive functions such as language, music or emotional control is the subject of popular debate, such sweeping questions fail to acknowledge the sequential nature of a multistage process that involves many brain regions. Evidence about the acquisition of these rich human behaviours is largely anecdotal, as the complex environments of everyday life are inherently difficult to quantify and normalize across individuals.

Clear evidence for critical periods has instead been found in the primary sensory systems of several species⁴. Animal models are now revealing, with greater resolution, the molecular, cellular and structural events that underlie experience-dependent circuit refinement. In this review, I focus on the primary visual cortex, which has been the premier model of critical period plasticity for 40 years⁵. During a brief postnatal period (of weeks to years, in proportion to the expected

lifespan of the species^{6,7}), the closure of one eye (but not both) causes a permanent loss of visual acuity through that eye. AMBLYOPIA occurs despite there being no damage to the retina or its target, the visual thalamus (dorsal lateral geniculate nucleus, dLGN) and is determined in the neocortex (in the visual primary cortex, V1), where the inputs from the two eyes first converge and compete for space^{5,6}. Most impressively, the seemingly innocuous act of covering an eye can profoundly alter the physical structure of the brain during the critical period only.

Shaping column size during the critical period

The cortical column is a fundamental unit of organization of the mammalian neocortex. Clusters of thalamocortical axon terminals that serve either the right or left eye tessellate layer 4 of the mature cortex to produce alternating OCULAR DOMINANCE domains^{8,9}. Occluding one eye during development yields an expansion of the columns serving the open eye at the expense of those responding to the deprived eye, which become reduced in size and afferent complexity^{10,11}. This physical manifestation of early postnatal experience occurs gradually, and is preceded by more rapid changes^{12,13} of intracortical circuits outside layer 4, which instruct the hard-wiring of an anatomical fingerprint that is unique to the individual. By extension, the segregation of columns by normal vision during the critical period has




Figure 1 | Local circuit control of developing columnar architecture in the neocortex. Activity-dependent models of segregation predict that cortical GABA (γ -aminobutyric acid) circuits are involved in determining the final column spacing from an initially overlapping mosaic^{17,18}. Neuronal activity from thalamic inputs serving the right or left eye is spread within the nearby cortex by local excitatory connections (red cell) but is inhibited at greater distances (blue cell), producing a ‘mexican hat’ activation profile. When this profile is modulated during development by preferentially enhancing or reducing horizontal, long-range inhibition (arrows), columns emerge that are wider or thinner than normal, respectively. This hypothesis was verified *in vivo* in kittens by modulating GABA_A (GABA type A) currents²⁰ with benzodiazepine agonists (diazepam) or inverse agonists (methyl-6,7-dimethoxy-4-ethyl- β -carboline, DMCM) throughout the critical period¹⁹.

OCULAR DOMINANCE
Relative anatomical or physiological strength of connections from either eye to individual cells in the primary visual cortex.

THALAMOCORTICAL AFFERENTS Axons from the thalamus (for example, the dLGN) that relay sensory input from the periphery (for example, the retina) to layer 4 of the neocortex.

BENZODIAZEPINES
Modulate chloride flux through GABA_A receptors that contain the $\gamma 2$ and any combination of $\alpha 1$, $\alpha 2$, $\alpha 3$ or $\alpha 5$ subunits. Benzodiazepine agonists enhance and inverse agonists decrease GABA efficacy.

been thought to result from similar activity-dependent rules acting on an initially overlapping continuum of THALAMOCORTICAL AFFERENTS.

This dogma has recently been challenged by the finding that axon terminals of thalamocortical afferents might, in part, be clustered well before the critical period¹⁴. There is also some evidence that siblings show substantial similarity of visual maps, which supports the idea that molecular cues establish columnar architecture¹⁵. Nonetheless, it is agreed that sensory experience is important for individualizing ocular dominance maps during the critical period. Even the focal patterns of deprivation that are produced by shadows of blood vessels in a single eye are embossed onto the primary visual cortex¹⁶. In computational models of self-organization, the recipient cortical circuits largely determine the final spacing of columns^{17,18}. Overlapping inputs segregate into clusters (‘neurons that fire together wire together’) through a neocortical organization that spreads excitation locally but is limited at a distance by farther-reaching inhibition.

FIGURE 1 shows how a central area of excitation surrounded by a larger concentric area of inhibition (the canonical ‘mexican hat’ profile) of intracortical activation could, in theory, influence thalamic innervation. In particular, lateral inhibition can establish narrow

or wide columns in computer simulations of column formation by adjusting the contrast between inputs at nascent border regions during this ‘winner-takes-all’ competition¹⁸. These long-standing theoretical predictions have recently been validated *in vivo* through the direct infusion of BENZODIAZEPINES into the kitten visual cortex during the critical period¹⁹. Such drugs come in three varieties, including agonists such as diazepam, inverse agonists such as the β -carbolines (for example, methyl-6,7-dimethoxy-4-ethyl- β -carboline DMCM), and antagonists that block the actions of both (flumazenil)²⁰. All bind to a particular subset of GABA_A (γ -aminobutyric acid type A) receptor, the agonists enhancing and inverse agonists reducing chloride flux. Increasing inhibition throughout the critical period with benzodiazepine agonists leads to a local 30% increase in column width, whereas inverse agonists produce column shrinkage¹⁹ (FIG. 1).

Interestingly, increased column spacing has also been reported for STRABISMUS through exotropic deviation of the eyes during the critical period²¹. Both enhanced lateral inhibition by direct intracortical infusion of diazepam and decorrelation of visual input by artificial squint are conditions that favour the maximal segregation of ocular dominance. In summary, local imbalances of neuronal activity influence columnar architecture during normal development, and this cannot be explained solely by genetic instruction. The broad range of column sizes observed in nature²² might reflect individual differences in the span of local circuit inhibition.

Detection by excitatory-inhibitory balance

Long-range, horizontal inhibitory axons therefore provide a scaffold for discriminating competing sensory input (for example, right or left eye) in the developing neocortex. This is in contrast to the maturing neuromuscular junction in the periphery, where individual motor axons may compete directly with each other for synaptic space on the same end plate²³. This competition is limited by the global resources (for example, neurotransmitter) that are available within individual presynaptic arbors, each of which might contact multiple muscle fibres²⁴. In the tangled circuits of the neocortex^{25,26}, the action of locally balanced excitation and inhibition first integrates and then detects competition.

Even small changes in the relative amounts of excitation and inhibition can markedly alter information processing²⁷. This delicate balance is dynamically adjusted by circuitry in the cortical layers²⁸⁻³⁰, where inhibitory connections are developed later than excitatory connections in the pre-critical period for ocular dominance³¹. Drastic pharmacological perturbations of neuronal activity, such as hyperexcitation (by, for example, glutamate³² or bicuculline³³) or total silencing (by, for instance, TETRODOTOXIN (TTX)³⁴, 2-amino-5-phosphonovaleric acid (APV)³⁵ or muscimol^{36,37}), not surprisingly disrupt plasticity, but fail to inform us about intrinsic network behaviour involved in this process. Gentle titration of endogenous neurotransmission by gene-targeted disruption in mice has been instrumental in dissecting the physiological role of




Figure 2 | GABA-mediated control of the critical period. **a** | Monocular deprivation produces a loss of response to the deprived eye and a gain of open-eye input, as measured by the neuronal discharge of single units from the mouse visual cortex³⁸. The ocular dominance of cells, rated on a seven-point scale of neuronal responsiveness, indicates a typical bias toward the contralateral eye (1–3) in the rodent (top left). After 3 or more days of monocular deprivation, the distribution shifts toward the open, ipsilateral eye (4–7; top right). **b** | Sensitivity to monocular deprivation is restricted to a critical period that begins, in mice, about 1 week after the eyes open (at postnatal day 13) and peaks 1 month after birth³⁸. Monocular deprivation causes amblyopia only during the same critical period. Red circles indicate the onset, peak and end of amblyopia resulting from monocular deprivation³⁹. The onset of plasticity can be delayed by directly preventing the maturation of GABA (γ -aminobutyric acid)-mediated transmission by gene-targeted deletion of *Gad65*, which encodes a GABA-synthetic enzyme⁴³, or by dark-rearing from birth (red arrow)^{50–53}. Conversely, the critical period can be brought forward by enhancing GABA transmission directly with benzodiazepines just after eye-opening^{44–46} or by promoting the rapid maturation of interneurons through excess brain-derived neurotrophic factor (BDNF) expression (blue arrow)^{47,48}.

STRABISMUS

Deviation of the two eyes due to a weakening of extraocular musculature that results in either an inward (esotropic) or outward (exotropic) rotation of one orbit and consequent amblyopia.

local circuit elements. As in other species^{5–9}, MONOCULAR DEPRIVATION (MD) shifts the spiking response of neurons in the mouse V1 towards the open eye³⁸, but, again, only during a critical period for behavioural amblyopia³⁹ (FIG. 2a; see also BOX 1).

The primary inhibitory neurotransmitter in the brain, GABA (γ -aminobutyric acid), is synthesized by glutamic acid decarboxylase produced by two distinct genes, *Gad65* and *Gad67*. Of these, *Gad65* is concentrated in axon terminals and bound to synaptic vesicles whereas *Gad67* is found throughout the cell⁴⁰. Deletion (knockout) of *Gad67* is lethal and eliminates

most cortical GABA content⁴¹, but *Gad65*-knockout mice are viable and show poor GABA release only on strong stimulation^{42,43}. Baseline receptive field properties are normal in the absence of GAD65, but ocular dominance plasticity is prevented until inhibition is acutely restored with diazepam⁴³. When infused directly into V1, the use-dependent nature and rapid breakdown of benzodiazepine agonists ensures that only those local inhibitory circuits that are normally engaged by monocular deprivation will be boosted. Remarkably, rescue of plasticity is possible at any age in *Gad65*-knockout mice, which indicates that the critical period is dependent on the proper level of inhibitory transmission (FIG. 2b)⁴⁴.


Conversely, the onset of the critical period can be accelerated by prematurely enhancing inhibition with benzodiazepines just after eye opening^{44–46}, as well as by transgenic overexpression of brain-derived neurotrophic factor (BDNF) to promote the maturation of GABA neurons^{47,48} (FIG. 2b). A close relationship between neuronal activity, BDNF release and GABA function also explains the classic effect of dark-rearing. Raising animals without visual experience from birth naturally reduces BDNF levels⁴⁹ and GABA-mediated transmission in the visual cortex^{50,51}, and delays the peak of plasticity into adulthood^{45,52,53}. Direct diazepam infusion⁴⁵, or BDNF overexpression⁵⁴ or secretion by enriched environments⁵⁵ in complete darkness abolish the expected delay of the critical period. These striking results indicate that tonic GABA release is sufficient to trigger the eventual closure of the plastic state, even in the total absence of visual input.

Specific GABA circuits for plasticity

Interestingly, not all GABA circuits are involved in critical period regulation. Several lines of evidence point towards a single class of interneuron that has the potential to mediate long-range inhibition and synchrony in the visual cortex. Among the many types of GABA-positive interneuron^{56–58} (FIG. 3b), neurochemical markers, such as calcium-binding proteins, have been used to reveal that the onset of the critical period corresponds closely to the emergence of PARVALBUMIN-positive cells⁵⁹ and both events are accelerated by BDNF overexpression⁴⁷. The specific blockade of a potassium channel (Kv3.1) that uniquely regulates the fidelity of FAST-SPIKING behaviour (and thereby GABA release) from parvalbumin-positive interneurons^{60–62} slows the rate of ocular dominance plasticity (Y.-T. Matsuda *et al.*, unpublished observations). To date, the molecular biology of benzodiazepine action has provided the deepest insight into the local circuits that underlie plasticity.

The α -subunits of GABA_A receptors determine benzodiazepine binding through a single amino acid residue in their amino terminus^{20,63} (FIG. 3a). In mice, knock-in of a point mutation at this site renders individual GABA_A receptor subtypes insensitive to benzodiazepines⁶⁴. Weak inhibition in the visual cortex in early life (as observed for *Gad65* deletion) prevents experience-dependent plasticity^{44,45}. Loss of responsiveness to an eye deprived of vision can be

Box 1 | Critical periods: gateway to lifelong plasticity



Ocular dominance is one of several thalamocortical circuit properties in the primary visual cortex (V1) that develop in an experience-dependent manner during early life^{4,186}. These might show different postnatal profiles (see panel) with, for example, orientation or direction selectivity (green) being shaped earlier than ocular dominance^{7,44,53} (orange), or slow-wave sleep oscillations¹⁶⁹ (dark blue). Surprisingly, the detailed molecular machinery underlying each receptive field property may also differ⁵³. Recent gene expression profiling supports the idea that the critical period for ocular dominance offers a specialized molecular milieu for plasticity^{170,171}, which is consistent with dendritic spine and axonal rearrangement being limited to this time in life^{10,85,86}.

Recently, monocular deprivation in adult animals has been reported to produce subthreshold synaptic plasticity^{172–174}, which has no further impact on spiking output^{38,44,175} spines^{85,86} or visuospatial acuity^{7,39}. This suggests that the closure of the critical period might reflect sequential ‘locks’ that are placed on the molecular pathway as it flows from mature GABA (γ -aminobutyric acid)-mediated detection towards structural consolidation (FIG. 6). Functional plasticity may persist throughout life in many other systems (such as the somatosensory cortex^{154,176} and the barn owl tectum¹⁴⁹). Interestingly, limited plastic states can be greatly facilitated (to near critical period levels) by the judicious activation of neuromodulatory systems in adulthood^{177,178}. A potent endogenous regulator of neuromodulation is sleep, which plays an intriguing part in enhancing adult learning¹⁷⁹.

Neuromodulators signal through second-messenger systems (FIG. 6), which can acutely reset the excitatory–inhibitory balance^{180,181}. Full reactivation of the critical period may further require the ‘undoing’ of hard-wired neuronal structures^{72,73}. The non-permissive growth environment of mature myelin is an especially promising target for rekindling plasticity, as it relates broadly to the acquisition of visual¹⁸², linguistic¹⁸³ or musical¹⁸⁴ abilities. Better still, laying down multiple anatomical traces through rich childhood experiences might also expand the capacity for plasticity in adulthood^{147,150}.

initiated prematurely by enhancing GABA-mediated transmission with zolpidem (FIG. 4a), a GABA_A α 1-, α 2- and α 3-subunit selective ligand⁴⁶. Systematic use of the mouse ‘knock-in’ mutation has further shown that only one of these subtypes, the α 1-subunit-containing circuits, drives cortical plasticity (FIG. 4b). Diazepam fails to trigger premature plasticity in α 1-subunit-mutants, although they are fully capable of undergoing plasticity at the proper age (postnatal day (P) 25) even without drugs⁴⁶, as they form normal α 1-subunit-containing GABA receptors at the appropriate time.

Importantly, GABA-releasing interneurons in the neocortex show precise connectivity^{25,58} (FIG. 3b). Synapse formation by parvalbumin-positive cells is largely mediated through molecular cues and then refined by neuronal activity^{65,66}. These cells include axon-ensheathing CHANDELIER CELLS and soma-targeting LARGE BASKET CELLS. The latter extend the wide-reaching, horizontal axonal plexus (FIG. 1), which has been shown

to span ocular dominance columns in the cat⁶⁷. The use of immuno-electron microscopy has further indicated that individual GABA_A receptor α -subunits are trafficked to discrete postsynaptic sites on the pyramidal cell axon, soma and dendrites (FIG. 3b). For example, α 2-subunits are preferentially enriched at the axon initial segment and at short-range basket cell synapses that are innervated by CHOLECYSTOKININ-positive axon terminals^{68,69}. Although α 2-subunit-containing connections do regulate neuronal firing, they have no effect on the induction of plasticity⁴⁶. This dissociation has implications not only for models of brain development, but also for the safe design of benzodiazepines for use in human infants (see below).

The α 1-subunit-containing GABA_A receptors are instead localized to receive parvalbumin-positive (but not cholecystokinin-positive) synapses on the soma^{68,69}, further implicating these large basket cell circuits in the control of the critical period. With age, large parvalbumin-positive cells are preferentially enwrapped in PERINEURONAL NETS of extracellular matrix (ECM) molecules and sugars⁷⁰. When these are disrupted, perisomatic inhibition of their targets is reduced⁷¹, and ocular dominance shifts can once again be induced by monocular deprivation, even in adulthood⁷². This might be the result of resetting and tapping the original GABA-mediated trigger⁴⁴ (FIG. 5b), as perineuronal nets probably control the extracellular ionic milieu (for example, potassium concentration⁷⁰) that surrounds parvalbumin-positive cells, allowing them to establish their fast firing efficiency^{60–62}, or might otherwise sequester molecular regulators of parvalbumin-positive cell maturation (S. Sugiyama *et al.*, unpublished observations). The identification of a cellular critical period trigger holds great promise as a therapeutic target and for the development of strategies for lifelong learning.

Expression by proteases: structural rewiring

The ECM is increasingly being recognized as a potent site for critical period plasticity⁷³. To convert physiological events (altered vision) into structural refinements, connections must ultimately be broken and neuronal wiring rerouted. Proteases are ideally suited to clear the way for growing neurites⁷⁴. TISSUE-TYPE PLASMINOGEN ACTIVATOR (tPA) is the main serine protease in the postnatal mammalian brain^{75,76}, and was originally identified there as an IMMEDIATE EARLY GENE activated by hippocampal seizures⁷⁷. Proteolysis by tPA is gradually upregulated in V1 within 2 days of monocular deprivation during the critical period (FIG. 5a), but not in adults or *Gad65*-knockout mice⁷⁸. Interestingly, at least 2 days of diazepam treatment are required to rescue plasticity in the absence of GAD65 (REF. 45). Functional ocular dominance plasticity is impaired when tPA action is blocked^{78–80} and can be rescued by exogenous tPA (but not diazepam)⁷⁸.

Permissive amounts of tPA might, therefore, couple functional to structural changes downstream of the excitatory–inhibitory balance that triggers visual cortical plasticity. The second messenger systems, which are known to be recruited during ocular dominance

TETRODOTOXIN (TTX). A voltage-dependent sodium channel blocker that can be used to silence all neural activity except spontaneous neurotransmitter release.

MONOCULAR DEPRIVATION (MD). Imbalanced visual input due to the occlusion of one eye by patching, eyelid suture or intraocular TTX injection.

PARVALBUMIN

One of three calcium-binding proteins, which, together with calretinin and calbindin, are expressed in most GABA-mediated neurons in the neocortex in largely non-overlapping groups.

FAST-SPIKING

Ability of parvalbumin-positive cells to fire non-adapting action potentials at rates of up to several hundred Hertz, due, in part, to unique potassium conductances (Kv3 class).

CHANDELIER CELL

Stereotypical GABA cell of the cerebral cortex that ensheathes the axon initial segment of up to 200 pyramidal cells with 'cartridge' synapses to directly control action potential generation.

LARGE BASKET CELL

Class of GABA cell with long, horizontally-extending axon that makes potent inhibitory contacts on the soma and proximal parts of the dendrites of target pyramidal cells.

CHOLECYSTOKININ

(CCK). A neuropeptide that is found in non-fast-spiking large basket cells that contact the soma of target pyramidal cells at synapses enriched in GABA_A receptor $\alpha 2$ -subunits.

PERINEURONAL NET

A conglomeration of chondroitin sulphate proteoglycans, extracellular matrix and cell-adhesion molecules that condenses around particular large-basket cells with age.

TISSUE-TYPE PLASMINOGEN ACTIVATOR

(tPA). The major serine protease in the brain, well known as an anti-clotting agent that works by cleaving fibrin in the blood.

IMMEDIATE EARLY GENES

Transcription factors that are induced within minutes of intense neuronal activity. Examples include *zif268* and *c-fos*.

PLASMIN

The active form of plasminogen, which is the primary target of tPA action, and itself a protease that is known to cleave extracellular matrix molecules (for example, laminin and phosphacan).




Figure 3 | Heterogeneity of local GABA circuits in the neocortex. **a** | Inhibitory synaptic transmission is mediated by GABA (γ -aminobutyric acid) synthesized by glutamic acid decarboxylase 65 (GAD65) in the presynaptic terminal during strong stimulation. The postsynaptic effect of released GABA is modulated by exogenous benzodiazepines acting on specific postsynaptic receptors. Among the 20 identified subunits that the GABA_A (GABA type A) receptor (GABA_AR) may comprise^{20,63}, the $\alpha 1$ -, $\alpha 2$ -, $\alpha 3$ - and $\alpha 5$ -subunits contain an amino acid residue, histidine, that is vital to the benzodiazepine binding site^{20,64}. Benzodiazepine (but not GABA) binding is lost when this amino terminus histidine is mutated to arginine (causing the $\alpha 1$ -, $\alpha 2$ -, $\alpha 3$ - and $\alpha 5$ -subunits to mimic the naturally benzodiazepine-insensitive $\alpha 4$ - and $\alpha 6$ -subunits). **b** | Many subtypes of GABA-releasing inhibitory interneuron can be identified in the neocortex on the basis of morphology, connectivity, expression of calcium-binding proteins or neuropeptide content^{56–58,25}. Moreover, specific contacts are preferentially enriched in specific GABA_A receptor α -subunits^{63,64}. The perisomatic localization of different subunits is shown next to the central pyramidal neuron. All subunits are found diffusely along the dendrite. ais, axon initial segment; CCK, cholecystokinin expressing; Ch, chandelier cell; CRC, Cajal–Retzius cell; DB, double bouquet cell; M, Martinotti neuron; N, neurogliaform neuron; PV+, parvalbumin positive. Lower part of panel **a** modified, with permission, from REF. 63 © (2001) Elsevier Science.

plasticity⁸¹ (FIG. 6), lie along a molecular cascade that links neuronal activity to tPA release^{4,78}, the structural consequences of which have recently been clarified. Thalamocortical axon rearrangement is a classical outcome of critical period plasticity (FIG. 1), but is much too slow to explain the rapid shift of ocular dominance that occurs within days of monocular deprivation^{10–13,82}. Morphological plasticity is initiated postsynaptically along the apical dendrites of target pyramidal cells in the cerebral cortex, where spines serve as pleomorphic sites of excitatory synaptic connection.

Spine shape has been shown to be highly dynamic using two-photon laser scanning microscopy in living transgenic mice expressing green fluorescent protein (GFP) in a subset of layer 5 cells. The motility of spines decreases with age in the visual cortex^{83,84}, but, during the critical period, can be transiently elevated by 2 days of monocular deprivation⁸⁵ (FIG. 5a, step 1). This occludes the motility that can be induced by the direct application of tPA to naive brain slices, which indicates that tPA and its substrate, plasminogen, might be the endogenous mediators of experience-dependent spine motility. Even along the same apical dendrite⁸⁵, spines

are set in motion by brief monocular deprivation only in layers 2, 3 and 5, consistent with early extragranular changes that instruct later events in layer 4 (REF. 12).

After 2 days of monocular deprivation, increased proteolysis degrades ECM and cell-adhesion proteins before any physiological ocular dominance shift is detectable^{38,78}. Consequently, spines are eliminated by 4 days of monocular deprivation, which corresponds to the rapid, complete loss of functional responsiveness⁸⁶ (FIG. 5a, step 2). After this postsynaptic pruning, axons carrying input from the deprived eye retract before afferents serving the open eye can migrate to spaces cleared by tPA–PLASMIN along the dendrite (FIG. 5a, step 3)^{10,11}. As expected for a true critical period event, spine density is not reduced by brief monocular deprivation in adulthood or in mice that lack tPA or GAD65 (REF. 86). The reduction in spine density can be restored in tPA- or *Gad65*-knockout mice by exogenous tPA or diazepam infusion, respectively. Importantly, spine motility and pruning faithfully reflect competitive interactions between the two eyes, as they fail to occur in the adjacent monocular segment that receives input solely from the contralateral eye^{85,86}.




Figure 4 | Specific GABA_A circuits for visual cortical plasticity. **a** | Ocular dominance shifts, as rated by the neuronal discharge of single units from the mouse visual cortex on a scale from 1 (contra) to 7 (ipsi) eye input, can be induced by monocular deprivation (MD) before the critical period in the presence of zolpidem⁴⁶, a benzodiazepine (BDZ) agonist that is selective for α1-, α2- or α3-subunit-containing GABA_A (γ-aminobutyric acid type A) receptors (GABA_ARs)²⁰. **b** | Conversely, premature plasticity cannot be induced in mice with a point mutation that renders the α1-subunit insensitive to the benzodiazepine diazepam⁴⁶ (α1(H101R)). Importantly, GABA_A receptor expression and localization occur normally, as the knock-in mice show robust plasticity (without drugs) at the typical critical period (postnatal day (P) 25). Figure modified, with permission, from REF. 46 © (2001) Elsevier Science.

Ultimately, the territory that represents the open eye grows⁹⁻¹¹ (FIG. 5a, step 3, right), and this requires cortical protein synthesis⁸⁷. Cleavage of secreted pro-neurotrophins by tPA-plasmin to yield active mature forms such as BDNF might contribute to the elongation of neurites^{88,89}. As proteolytic activity slowly wanes after a week of monocular deprivation⁷⁸, axons serving the open eye expand^{10,11}, spines grow out to meet them and their density largely recovers⁸⁶ (FIG. 5a, asterisks). Curiously, spine loss is most robust near the soma of layer 2/3 pyramidal cells. Competition detected by inhibitory circuits impinging on pyramidal cell somata might, therefore, be translated into structural changes mediated by a multistep proteolytic action of the extracellular tPA-plasmin cascade.

The homosynaptic view

It is tempting to speculate that the loss or gain of visual responsiveness of neurons in V1 during the critical period is simply the result of homosynaptic LONG-TERM DEPRESSION (LTP) or POTENTIATION (LTP) somewhere in the visual circuit (the homosynaptic view). On theoretical grounds, homosynaptic rules of excitatory synaptic plasticity alone are insufficient to produce a competitive outcome⁹⁰, as they require the involvement of other complex mechanisms, such as sliding thresholds

or metaplasticity. Although correlative evidence has been presented to support homosynaptic rules for plasticity *in vivo*, the fact that direct comparisons are not supportive is often overlooked.

In favour of the homosynaptic view, changes in phosphorylation state and membrane trafficking of AMPA (α-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid) receptor subunits — events that seem to underlie LTP or LTD at central synapses⁹¹ — have been observed in the primary sensory cortex after natural sensory experience *in vivo*. For example, whisker stimulation at P12–14 in rats drives a recombinant AMPA receptor subunit (GluR1) into synapses from layers 4 to 2/3 in the somatosensory barrel cortex⁹². Expression of the GluR1 cytoplasmic tail, which inhibits the synaptic delivery of endogenous receptors during LTP *in vitro*, blocks this insertion and subsequent synaptic potentiation *in vivo*. Whisker deprivation produces both an expansion of neighbouring input from spared barrels and a loss of response to the principal whisker. The latter largely occludes induction of NMDA (N-methyl-D-aspartate) receptor-dependent LTD by low-frequency stimulation (LFS) along the principal ascending pathway⁹³, which supports the idea that the mechanisms mediating changes in barrel cortex responsiveness are the same as those that mediate LTD.

Similarly, in the visual cortex, 24-h monocular deprivation (too short a time period to shift ocular dominance *in vivo*³⁸) reduces the saturation level of LTD in response to repeated LFS, and leads to numerous changes in the GluR1 phosphorylation status, which are akin to those that occur during hippocampal LTD⁹⁴. The main difficulty with correlative studies is that they can never demonstrate causality. Ostensibly similar molecular pathways (FIG. 6) might be used differently by LTP/LTD and by critical period mechanisms subserving circuit function *in vivo*. For example, the late onset and experience-dependent profile of NMDA receptor 2A (NR2A) subunits⁹⁵⁻⁹⁸ — known to determine LTP levels in the hippocampus^{99,100} — is irrelevant to critical period expression in the visual cortex⁵³. A decline in LTP/LTD magnitude and slow emergence of NR2A at thalamocortical synapses in the barrel cortex during the anatomical critical period in the first postnatal week might also be coincidental^{101,102}. Barrel shrinkage and expansion on whisker cauterization at birth is impervious to cortical NMDA receptor deletion¹⁰³ and might reflect changes at subcortical levels, before thalamic axons segregate in layer 4 (REF. 104).

In the absence of NR2A, the depolarizing action of NMDA currents is prolonged. Like *Gad65* deletion⁴³, this tips local circuit equilibrium towards excitation, weakening ocular dominance plasticity that can be restored by diazepam⁵³. Counter-intuitive to an LTP perspective^{99,105}, inhibition is required for plasticity *in vivo* when NMDA receptor function is high. Conversely, although LTD is blocked by NMDA-receptor antagonists, high doses of APV³⁵ promote a paradoxical loss of open eye input, as does strong postsynaptic activation of GABA_A receptors with muscimol^{36,37}. Although maturation of other receptive field

LONG-TERM DEPRESSION (LTD). A persistent reduction of synaptic transmission in response to weak, poorly-correlated input.

LONG-TERM POTENTIATION (LTP). A persistent strengthening of synaptic transmission in response to strong, correlated input.




Figure 5 | Structural consolidation during the critical period. a | Once functional detection of competing inputs is made possible by the maturation of relevant GABA (γ -aminobutyric acid) circuits^{43–48}, a series of structural rearrangements accompanies sensory deprivation. Shortly (2 days) after monocular deprivation (MD), the motility of spines (two-headed arrows)⁸⁵ is increased on apical dendrites of excitatory pyramidal neurons by an increase in tissue-type plasminogen activator (tPA)–plasmin proteolytic activity (blue background)⁷⁸. No shift in ocular dominance (OD) can be detected at this stage³⁸ (1). The total number of spines on pyramidal cell apical dendrites is transiently and significantly decreased owing to their elimination (asterisks), with a time course that corresponds to the loss of cellular responses of the deprived eye 4 days after monocular deprivation⁸⁶. This is followed by a retraction of thalamocortical axons^{10,11} as extracellular tPA–plasmin activity remains high⁷⁸ (2). After long-term monocular deprivation, new spines emerge (asterisks) to receive synaptic input from sprouting axons carrying input from the open eye (triangles)^{10,11,13} as tPA–plasmin activity subsides⁷⁸ (3). **b** | These results predict that structural modification is necessary to reactivate critical period plasticity in adulthood. Infusion of chondroitinases (scissor symbols) to break up perineuronal nets (PNNs) in the extracellular matrix restores ocular dominance shifts to adult rats⁷². Interestingly, these nets preferentially envelop large, parvalbumin-positive (PV⁺) basket cells⁷⁰, which control perisomatic inhibition⁷¹ and trigger the endogenous critical period. The section depicts parvalbumin-positive basket cells stained with monoclonal antibody (red) and perineuronal nets stained with *Wisteria floribunda agglutinin* (WFA; green). Note that small parvalbumin-positive cells are not ensheathed by perineuronal nets. BDNF, brain-derived neurotrophic factor; Glu, glutamate.

hallmark of critical period plasticity. One day of monocular deprivation does not occlude homosynaptic depression by single LFS⁹⁴, and these early forms of synaptic change, which persist in the presence of protein synthesis inhibitors¹⁰⁸, are insufficient to shift ocular dominance *in vivo*⁸⁷. However, multiple spaced conditioning stimuli, used to saturate LTP/LTD⁹⁴, activate distinct transcriptional pathways (for example, that of immediate early gene *Zif268*)¹⁰⁹, which are also not required for ocular dominance plasticity¹¹⁰. In fact, the role of proteases (tPA) in growth for long-lasting forms of LTP^{111,112} is opposite to that for ocular dominance plasticity (spine pruning⁸⁶).

Overall, no consistent correlation has been found between the ability to induce homosynaptic plasticity *in vitro* and amblyopic effects *in vivo*^{113–116}. The relationship is not straightforward, as the mechanisms of LTP and LTD might differ depending on cortical layer¹¹⁷. Moreover, unlike the very rapid induction of synaptic changes, no loss of visual response or acuity occurs until several days of eyelid suture have elapsed^{38,39}. Rapid phosphorylation dynamics of one GluR1 amino acid residue alone are unlikely to explain the complex functional and structural events that constitute the critical period (FIG. 5). Spine shrinkage by LFS is, in fact, mediated by cofilin and is independent of the AMPA receptor dephosphorylation that underlies LTD¹¹⁸. At present, there is only limited evidence directly connecting the coincident structural and functional plasticity at spines¹¹⁹.

Most strikingly, synaptic depression is intact whereas ocular dominance plasticity is lost in mice that conditionally overexpress the protein phosphatase, calcineurin¹²⁰. Underscoring a double dissociation, homosynaptic models based on NMDA-receptor activation stipulate that the maturation of inhibition will terminate plasticity¹⁰⁵, whereas quite the opposite is true: *in vivo*, GABA function is required to trigger the critical period⁴⁴. Given that diazepam¹²¹ and endogenous BDNF¹²² block LTD induction in the cortex, their acceleration of plasticity *in vivo*^{44–48} is not predicted by homosynaptic models, which are routinely studied *in vitro* in the presence of GABA-receptor blockers. In the end, adjusting local inhibitory circuit function has provided direct control over the critical period (FIG. 2), which LTP-based models have not.

Local circuit models of critical period plasticity

That inhibition might enable plasticity in the developing brain seems paradoxical. The identification of a particular GABA circuit that drives the onset of the critical period and the subsequent sequence of anatomical events (FIG. 5) now allows us to construct realistic models with which to target the plasticity process more precisely. Two scenarios centred on the parvalbumin-positive basket cell serve as heuristic examples for further study. One is an ‘instructive’ model (FIG. 7a), in which powerful, fast somatic inhibition edits one-by-one the action potentials that can pass into the dendritic arbor by back-propagation through the cell body. Recent SPIKE-TIMING DEPENDENT models of synaptic plasticity rely on precise millisecond

properties (such as orientation bias^{44,53}) might reflect subunit-specific coupling of NR2A to postsynaptic signalling pathways (like LTP¹⁰⁰), the yin-yang relationship of excitatory–inhibitory balance is essential for ocular dominance plasticity.

During learning, distinct plasticity mechanisms subservise sequential phases of memory¹⁰⁶. Local (synaptic and cellular) events are eventually consolidated at a systems (circuit) level by increasing protein synthesis and neurite growth. Excessive emphasis on LTP/LTD alone (which is all-or-none at single synapses¹⁰⁷) ignores the gradual, long-lasting changes that are the

SPIKE-TIMING DEPENDENT PLASTICITY
Describes physiological windows for LTP or LTD of synaptic transmission based on the arrival time of incoming action potentials with respect to back-propagated spikes in the target dendrite.




Figure 6 | Molecular mechanisms of visual cortical plasticity. Many candidate plasticity factors have been screened using the monocular deprivation protocol — by pharmacology in kittens or by gene-targeted disruption in mice (for a review, see REF. 81). Only a handful of second messenger molecules have been found to have a direct role in plasticity without perturbing global neuronal activity. These include PROTEIN KINASE A (PKA)¹⁸¹, calcium/calmodulin-dependent protein kinase II (CaMKII), calcineurin¹²⁰, extracellular signal-regulated kinase (ERK), cyclic AMP responsive element binding protein (CREB), protein synthesis machinery⁸⁷ and the plasmin system (tissue-type plasminogen activator (tPA)–plasmin), which is regulated by its inhibitors (plasminogen activator inhibitor, PAI-1)^{78–80}. Brain-derived neurotrophic factor (BDNF) has an early role in plasticity, establishing the GABA (γ-aminobutyric acid) cells that discriminate competing sensory inputs to trigger the critical period⁴⁷. Mature BDNF, which is produced from the cleavage of pro-BDNF by tPA–plasmin⁸⁸, in turn stimulates the expression and release of tPA¹⁸⁵. Both tPA and BDNF can then contribute sequentially to the final anatomical rewiring of the cortical circuit^{10,11,13,85,86,89} (FIG. 5). Plasticity might come to an end when permissive factors are gradually lost⁸⁶, or when further growth is actively suppressed by late-emerging inhibitory factors (myelin) in the extracellular matrix^{7,73,182}.

time windows to allow such postsynaptic spikes to meet presynaptic input^{123,124}. Sloppy gating by weak inhibition at the soma would prevent a competitive outcome by allowing excess back-propagation and spurious coincident activity with infrequent, deprived inputs from the retina¹²⁵. Identifying a molecular substrate that is unique to spike-timing dependent (as opposed to LFS or tetanus-induced¹²⁶) plasticity will be a prerequisite to testing this scheme *in vivo*.

Both immature (pre-critical period) and *Gad65*-knockout mice show prolonged neuronal discharge that continues well after stimuli have passed through the receptive field of individual cells^{43–45}. Chandelier cell ‘cartridge’ synapses (which use α2 GABA_A receptor subunits) can directly control this excessive spiking at the axon initial segment^{46,127}. Surprisingly, plasticity is still triggered prematurely by diazepam when hyperexcitability persists in mice whose α2-subunits can no longer bind benzodiazepines⁴⁶. Fast-spiking, feedforward inhibition mediated by α1-subunit-containing receptors on the soma is then ideally situated to suppress back-propagation of unwanted spikes¹²⁵ (FIG. 7a). The model incorporates the wide-reaching, horizontal axons of basket cells that receive input from

PROTEIN KINASE A (PKA). Phosphorylates multiple targets (including AMPA and GABA_A receptors) when cyclic AMP binds its regulatory subunits to release the catalytic domains.

one eye to inhibit targets of the other eye⁶⁷. This simple contrast enhancement circuit relieves the burden of discriminating competitors by LTP/LTD at single synapses. Alternatively, well-timed spikes in the soma itself might facilitate nuclear calcium entry for gene transcription (and growth) in response to waves of calcium arriving from synapses on dendritic spines.

A second, ‘permissive’ model emphasizes newfound knowledge of parvalbumin-positive cell biology. Even in adulthood, basket cells can be coupled electrically into groups of 40 or 50 cells^{128,129}, ending the network with the ability to detect synchrony¹³⁰ (FIG. 7b). Whereas simultaneous inputs (for example, from the same eye) rapidly co-excite cells through gap junctions, even a 2-ms input jitter (for example, between opposite eyes) is sufficient to dampen the coupling by reciprocal GABA_A synapses, which are also enriched in α1-subunits¹³¹. As a result, these neurons with long, horizontal axonal arbors⁶⁷, are maximally active on a columnar scale, time-locked to release growth or plasticity factors when strong synchronous activity arrives in the neocortex. Gap junctional coupling between interneurons also promotes synchronous oscillation among principal cells^{132,133}. Validation of the ‘permissive’ model awaits conditional deletion of connexins in cortical parvalbumin-positive cells alone, as the retina is also rich in gap junctions¹³⁴.

This model assumes an extracellular locus of competition to be quite distinct from the intracellular mechanisms of LTP and LTD. Axons and dendritic spines that are co-active with the synchronized parvalbumin-positive cell network can benefit from the rapid, focal creation of a permissive growth environment^{78,86,135}. The source and dynamics of tPA–plasmin release in the brain remain unclear owing to the lack of specific reagents. Laminar motility of spines⁸⁵ and their rapid pruning⁸⁶ by brief periods of monocular deprivation could reflect activity-dependent secretion of proteases^{136–138} from the axons of fast-spiking cells themselves, in which parvalbumin is an important contributor to presynaptic calcium signals and synaptic integration¹³⁹.

This might explain why spines nearest the soma of layer 2/3 pyramidal cells are most robustly lost during monocular deprivation⁸⁶, as they lie nearest the parvalbumin-positive cell-rich layer⁵⁹. Potential synaptic plasticity of GABA-mediated transmission might add a further layer of regulation. How a competitive outcome arises by uniformly bathing dendrites in proteases also needs to be considered. Cell-adhesion molecules can become insensitive to proteases during high levels of activity^{140,141}. Less active synapses also release fewer endogenous protease inhibitors¹⁴², tilting the overall balance nearby towards pruning. Therefore, open-eye axons can protect themselves from proteolytic removal in two ways, whereas silent deprived-eye inputs are left defenceless in a sea of tPA–plasmin during monocular deprivation.

Towards a common brain principle

How general is the idea of local circuit inhibition and critical periods across other systems? Zebra finches acquire a unique song once in life during a critical period for vocal learning that progresses through three stages⁴.




Figure 7 | Two models for inhibitory control of sensory plasticity. a | Somatic inhibition by fast-spiking parvalbumin-positive (PV⁺) cells (green) expressing the potassium channel Kv3.1 is mediated by the GABA_A (γ-aminobutyric acid type A) α1-subunit, whereas axonal inhibition is mediated by the GABA_A α2-subunit. Somatic inhibition is ideally situated for suppression, or ‘editing’ of unwanted spikes, preventing them from back-propagating through the cell body into the dendritic tree (red arrow). This is an ‘instructive’ model, as individual action potentials can produce long-term potentiation (LTP) or depression (LTD) based on precise spike-timing dependent windows at individual synapses of coincident pre- and postsynaptic activity^{123,124}. Notably, failure to regulate excess spiking at the axon initial segment can still be differentiated by fast-spiking inhibition at the cell body. **b** | Gap junctional coupling endows networks of parvalbumin-positive (PV⁺) interneurons with the ability to detect synchronous input^{128–130}. Even a slight jitter in input timing (for example, between eyes) dampens network activity through reciprocal GABA-mediated contacts (enriched with GABA_A-receptor α1-subunits¹³¹). Only synchronous open-eye input will produce maximal, activity-dependent release or uptake of ‘permissive’ factors for neurite growth (for example, tissue-type plasminogen activator^{78,86} and brain-derived neurotrophic factor). Competition is determined extracellularly.

After first memorizing the father’s template, birds enter a motor practice phase, followed by song crystallization. In register with the active singing component, a peak in the number of GABA neurons is observed in the motor nucleus robustus archistriangularis (RA), where descending auditory and forebrain memory circuits converge onto individual cells¹⁴³. Such a GABA profile is observed only in males, and not in females, which do not produce song. Local inhibitory circuits might allow the proper matching (and discrimination) of template and vocalized sub-song in the RA before consolidation. Indeed, infusion of cannabinoids, which primarily bind presynaptic CB1 receptors to reduce GABA release¹⁴⁴, alters developmental sensorimotor vocal learning but not adult song performance¹⁴⁵.

If barn owls are reared wearing prism lenses to skew their visual world, a learned shift occurs in the auditory map of interaural time differences in the tectum so that it corresponds with the visual input¹⁴⁶. Multiple maps can be stored in the same tectum if this experience occurs during a critical period¹⁴⁷. To allow discrimination between conflicting circuits, new GABA-mediated connections are formed to suppress the unused representation¹⁴⁸. Adult plasticity is limited to incremental displacement of the visual scene within the architectural constraints of axons that were hard-wired during the critical period^{149,150}.

Conversely, in the barrel cortex GABA circuits are formed and reorganized throughout life^{151–153}, and

are associated with lifelong plasticity¹⁵⁴. This is also the case in the mammalian olfactory system, where constant neurogenesis is responsible for the underlying memory of odour discrimination in adulthood^{155,156}. The newly born cells are GABA-containing granule cells, the functions of which include lateral inhibition and synchronization of neuronal activity^{157,158}. This raises the question of whether an olfactory critical period would emerge in the absence of neurogenesis, or whether visual plasticity could be maintained by prolonging cell proliferation in the neocortex.

Most interestingly, the control of visual cortical plasticity by GABA in animal models might also apply to the development of the human brain. In postnatal samples of V1 that were removed post-mortem¹⁵⁹, the maturation of NR2A to 2B is complete within the first 9 months after birth. By contrast, *Gad65* expression and the conversion of GABA_A receptors from α3- to α1-subunit containing occurs more slowly, over several years, which is consistent with the extended length of the critical period for amblyopia in humans⁶⁷. Strikingly, the levels of *GAD67* and α2-subunits are constant during the same early postnatal time period¹⁵⁹, consistent with their limited role in plasticity in animals⁴⁶. To avoid the rapid, premature induction of critical period plasticity through α1-subunit-containing receptors^{45,46}, new benzodiazepine agonists that are selective for α2-subunits should be developed for the control of status epilepticus in human infants¹⁶⁰.

In general, critical periods are a process of selecting the best neural representation available from among the many competing inputs that continually bombard the maturing nervous system. Growth and function of lateral inhibitory circuits offer a rational, cellular substrate to be compared and modelled across regions to gain broader insight into brain development and its disorders^{161,162}. Interestingly, cortical lesions or retinal scotomas in adulthood transiently reconfigure local circuit excitation–inhibition to an immature state¹⁶³. This provides a rationale for the administration of diazepam after acute

stroke¹⁶⁴ — not only might it reduce excitotoxicity, but it could also aid recovery by triggering plasticity machinery (FIG. 6). The concept of excitatory–inhibitory balance²⁶ in key neural systems during development might apply to epilepsy¹⁶², autism¹⁶¹, **Rett syndrome**¹⁶⁵ and **Tourette's syndrome**¹⁶⁶, schizophrenia¹⁶⁷ and even the encoding and retrieval of memories¹⁶⁸. Finding the keys to critical period plasticity could enable the development of novel therapies and training paradigms for education, rehabilitation, recovery from injury and lifelong learning in adulthood.

1. Kim, K. H., Reikun, N. R., Lee, K. M. & Hirsch, J. Distinct cortical areas associated with native and second languages. *Nature* **388**, 171–174 (2002).
2. Elbert, T., Pantev, C., Wienbruch, C., Rockstroh, B. & Taub, E. Increased cortical representation of the fingers of the left hand in string players. *Science* **270**, 305–307 (1995).
3. Pantev, C. *et al.* Increased auditory cortical representation in musicians. *Nature* **392**, 811–814 (1999).
4. Hensch, T. K. Critical period regulation. *Annu. Rev. Neurosci.* **27**, 549–579 (2004).
5. Wiesel, T. N. & Hubel, D. H. Single-cell responses in striate cortex of kittens deprived of vision in one eye. *J. Neurophysiol.* **26**, 1003–1017 (1963).
A classic paper for anyone interested in the fields of development, plasticity and vision.
6. Berardi, N., Pizzorusso, T. & Maffei, L. Critical periods during sensory development. *Curr. Opin. Neurobiol.* **10**, 138–145 (2000).
7. Daw, N. *Visual Development* (Plenum, New York, 1995).
8. Hubel, D. H., Wiesel, T. N. & LeVay, S. Functional architecture of area 17 in normal and monocularly deprived macaque monkeys. *Cold Spring Harb. Symp. Quant. Biol.* **40**, 581–589 (1976).
9. Shatz, C. J. & Stryker, M. P. Ocular dominance in layer IV of the cat's visual cortex and the effects of monocular deprivation. *J. Physiol. (Lond.)* **281**, 267–283 (1978).
10. Antonini, A. & Stryker, M. P. Plasticity of geniculocortical afferents following brief or prolonged monocular occlusion in the cat. *J. Comp. Neurol.* **369**, 64–82 (1996).
11. Antonini, A., Fagiolini, M. & Stryker, M. P. Anatomical correlates of functional plasticity in mouse visual cortex. *J. Neurosci.* **19**, 4388–4406 (1999).
Details the well known anatomical consequences of monocular deprivation in terms of thalamic input to the neocortex. These structural events are far slower than intracortical changes (discussed in references 12, 13, 85 and 86).
12. Trachtenberg, J. T., Trepel, C. & Stryker, M. P. Rapid extragranular plasticity in the absence of thalamocortical plasticity in the developing primary visual cortex. *Science* **287**, 2029–2032 (2000).
13. Trachtenberg, J. T. & Stryker, M. P. Rapid anatomical plasticity of horizontal connections in the developing visual cortex. *J. Neurosci.* **21**, 3476–3482 (2001).
14. Katz, L. C. & Crowley, J. C. Development of cortical circuits: lessons from ocular dominance columns. *Nature Rev. Neurosci.* **3**, 34–42 (2002).
15. Kaschube, M., Wolf, F., Geisel, T. & Lowel, S. Genetic influence on quantitative features of neocortical architecture. *J. Neurosci.* **22**, 7206–7217 (2002).
16. Adams, D. L. & Horton, J. C. Shadows cast by retinal blood vessels mapped in primary visual cortex. *Science* **298**, 572–576 (2002).
17. Willshaw, D. J. & von der Malsburg, C. How patterned neural connections can be set up by self-organization. *Proc. R. Soc. Lond. B* **194**, 431–445 (1976).
18. Miller, K. D., Keller, J. B. & Stryker, M. P. Ocular dominance column development: analysis and simulation. *Science* **245**, 605–615 (1989).
19. Hensch, T. K. & Stryker, M. P. Columnar architecture sculpted by GABA circuits in developing cat visual cortex. *Science* **303**, 1678–1681 (2004).
First bidirectional shaping of cortical column size by the direct manipulation of lateral inhibition, as predicted by earlier theoretical models (described in references 17 and 18).
20. Sieghart, W. Structure and pharmacology of γ -aminobutyric acid_A receptor subtypes. *Pharmacol. Rev.* **47**, 181–234 (1995).
21. Lowel, S. Ocular dominance column development: strabismus changes the spacing of adjacent columns in cat visual cortex. *J. Neurosci.* **14**, 7451–7468 (1994).
22. Horton, J. C. & Hocking, D. R. Intrinsic variability of ocular dominance column periodicity in normal macaque monkeys. *J. Neurosci.* **16**, 7228–7239 (1996).
23. Kasthuri, N. & Lichtman, J. W. The role of neuronal identity in synaptic competition. *Nature* **424**, 426–430 (2003).
24. Buffelli, M. *et al.* Genetic evidence that relative synaptic efficacy biases the outcome of synaptic competition. *Nature* **424**, 430–434 (2003).
An elegant genetic dissection of competition at a visible peripheral synapse.
25. Somogyi, P., Tamas, G., Lujan, R. & Buhl, E. H. Salient features of synaptic organisation in the cerebral cortex. *Brain Res. Rev.* **26**, 113–135 (1998).
26. Hensch, T. K. & Fagiolini, M. (eds) *Excitatory–Inhibitory Balance: Synapses, Circuits, Systems* (Kluwer/Plenum, New York, 2004).
27. Liu, G. Local structural balance and functional interaction of excitatory and inhibitory synapses in hippocampal dendrites. *Nature Neurosci.* **7**, 373–379 (2004).
28. Turrigiano, G. G. & Nelson, S. B. Homeostatic plasticity in the developing nervous system. *Nature Rev. Neurosci.* **5**, 97–107 (2004).
29. Desai, N. S., Cudmore, R. H., Nelson, S. B. & Turrigiano, G. G. Critical periods for experience-dependent synaptic scaling in visual cortex. *Nature Neurosci.* **5**, 783–789 (2002).
30. Maffei, A., Nelson, S. B. & Turrigiano, G. G. Selective reconfiguration of layer 4 visual cortical circuitry by visual deprivation. *Nature Neurosci.* **7**, 1353–1359 (2004).
31. Long, M. A., Cruikshank, S. J., Jutras, M. J. & Connors, B. W. Abrupt maturation of a spike-synchronizing mechanism in neocortex. *J. Neurosci.* **25**, 7309–7316 (2005).
32. Shaw, C. & Cynader, M. Disruption of cortical activity prevents ocular dominance changes in monocularly deprived kittens. *Nature* **308**, 731–734 (1984).
33. Ramoa, A. S., Paradiso, M. A. & Freeman, R. D. Blockade of intracortical inhibition in kitten striate cortex: effects on receptive field properties and associated loss of ocular dominance plasticity. *Exp. Brain Res.* **73**, 285–296 (1988).
34. Reiter, H. O., Waitzman, D. M. & Stryker, M. P. Cortical activity blockade prevents ocular dominance plasticity in the kitten visual cortex. *Exp. Brain Res.* **65**, 182–188 (1986).
35. Bear, M. F., Kleinschmidt, A., Gu, Q. A. & Singer, W. Disruption of experience-dependent synaptic modifications in striate cortex by infusion of an NMDA receptor antagonist. *J. Neurosci.* **10**, 909–925 (1990).
36. Reiter, H. O. & Stryker, M. P. Neural plasticity without postsynaptic action potentials: less-active inputs become dominant when kitten visual cortical cells are pharmacologically inhibited. *Proc. Natl Acad. Sci. USA* **85**, 3623–3627 (1988).
37. Hata, Y. & Stryker, M. P. Control of thalamocortical afferent rearrangement by postsynaptic activity in developing visual cortex. *Science* **265**, 1732–1735 (1994).
38. Gordon, J. A. & Stryker, M. P. Experience-dependent plasticity of binocular responses in the primary visual cortex of the mouse. *J. Neurosci.* **16**, 3274–3286 (1996).
39. Prusky, G. T. & Douglas, R. M. Developmental plasticity of mouse visual acuity. *Eur. J. Neurosci.* **17**, 167–173 (2003).
A behavioural study revealing a clear critical period for amblyopia that matches the shift of single-unit responses in the mouse visual cortex (detailed in reference 38), which indicates that there are no
40. Soghomonyan, J. J. & Martin, D. L. Two isoforms of glutamate decarboxylase: why? *Trends Pharmacol.* **19**, 500–505 (1998).
41. Asada, H. *et al.* Cleft palate and decreased brain γ -aminobutyric acid in mice lacking the 67-kDa isoform of glutamic acid decarboxylase. *Proc. Natl Acad. Sci. USA* **94**, 6496–6499 (1997).
42. Tian, N. *et al.* The role of the synthetic enzyme GAD65 in the control of neuronal γ -aminobutyric acid release. *Proc. Natl Acad. Sci. USA* **96**, 12911–12916 (1999).
43. Hensch, T. K. *et al.* Local GABA circuit control of experience-dependent plasticity in developing visual cortex. *Science* **282**, 1504–1508 (1998).
The first evidence that GABA-mediated transmission is required for plasticity in vivo. Lays the foundation for a series of studies confirming that critical period timing can be controlled through inhibitory interneurons (discussed further in references 44–48), quite unlike LTP models in vitro that are routinely blocked by inhibition (see references 105, 121).
44. Fagiolini, M. & Hensch, T. K. Inhibitory threshold for critical-period activation in primary visual cortex. *Nature* **404**, 183–186 (2000).
45. Iwai, Y., Fagiolini, M., Obata, K. & Hensch, T. K. Rapid critical period induction by tonic inhibition in visual cortex. *J. Neurosci.* **23**, 6695–6702 (2003).
46. Fagiolini, M. *et al.* Specific GABA_A circuits for visual cortical plasticity. *Science* **303**, 1681–1683 (2004).
Provides striking evidence that not all GABA-mediated connections are involved in critical period induction, which has strong implications for computational models and the design of specific benzodiazepines for use in human infants.
47. Huang, Z. J. *et al.* BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual cortex. *Cell* **98**, 739–755 (1999).
48. Hanover, J. L., Huang, Z. J., Tonegawa, S. & Stryker, M. P. Brain-derived neurotrophic factor overexpression induces precocious critical period in mouse visual cortex. *J. Neurosci.* **19**, RC40 (1999).
49. Castren, E., Zafra, F., Thoenen, H. & Lindholm, D. Light regulates expression of brain-derived neurotrophic factor mRNA in rat visual cortex. *Proc. Natl Acad. Sci. USA* **89**, 9444–9448 (1992).
50. Morales, B., Choi, S. Y. & Kirkwood, A. Dark rearing alters the development of GABAergic transmission in visual cortex. *J. Neurosci.* **22**, 8084–8090 (2002).
51. Chen, L., Yang, C. & Mower, G. D. Developmental changes in the expression of GABA_A receptor subunits (α_1 , α_2 , α_3) in the cat visual cortex and the effects of dark rearing. *Mol. Brain Res.* **88**, 135–143 (2001).
52. Mower, G. D. The effect of dark rearing on the time course of the critical period in cat visual cortex. *Dev. Brain Res.* **58**, 151–158 (1991).
53. Fagiolini, M. *et al.* Separable features of visual cortical plasticity revealed by *N*-methyl-D-aspartate receptor 2A signaling. *Proc. Natl Acad. Sci. USA* **100**, 2854–2859 (2003).
Debunks the 'NR2A subunit switch' hypothesis for critical period closure (see also reference 101). Also reveals distinct molecular pathways for individual receptive field properties.
54. Gianfranceschi, L. *et al.* Visual cortex is rescued from the effects of dark rearing by overexpression of BDNF. *Proc. Natl Acad. Sci. USA* **100**, 12486–12491 (2003).

55. Bartoletti, A., Medini, P., Berardi, N. & Maffei, L. Environmental enrichment prevents effects of dark-rearing in the rat visual cortex. *Nature Neurosci.* **7**, 215–216 (2004).
- References 54 and 55, along with reference 45, strikingly demonstrate that direct modulation of tonic GABA-mediated function in the cortex is sufficient to trigger the critical period, even in the absence of visual input.**
56. Kawaguchi, Y. & Kubota, Y. GABAergic cell subtypes and their synaptic connections in rat frontal cortex. *Cereb. Cortex* **7**, 476–486 (1997).
57. Markram, H. *et al.* Interneurons of the neocortical inhibitory system. *Nature Rev. Neurosci.* **5**, 793–807 (2004).
58. DeFelipe, J. Types of neurons, synaptic connections and chemical characteristics of cells immunoreactive for calbindin-D28K, parvalbumin and calretinin in the neocortex. *J. Chem. Neuroanat.* **14**, 1–19 (1997).
59. Del Rio, J. A., De Lecea, L., Ferrer, I. & Soriano, E. The development of parvalbumin-immunoreactivity in the neocortex of the mouse. *Dev. Brain Res.* **81**, 247–259 (1994).
60. Rudy, B. & McBain, C. J. Kv3 channels: voltage-gated K⁺ channels designed for high-frequency repetitive firing. *Trends Neurosci.* **24**, 517–526 (2001).
61. Erisir, A., Lau, D., Rudy, B. & Leonard, C. S. Function of specific K⁺ channels in sustained high-frequency firing of fast-spiking neocortical interneurons. *J. Neurophysiol.* **82**, 2476–2489 (1999).
62. Lien, C. C. & Jonas, P. K_v3 potassium conductance is necessary and kinetically optimized for high-frequency action potential generation in hippocampal interneurons. *J. Neurosci.* **23**, 2058–2068 (2003).
63. Cherubini, E. & Conti, F. Generating diversity at GABAergic synapses. *Trends Neurosci.* **24**, 155–162 (2001).
64. Rudolph, U., Crestani, F. & Möhler, H. GABA_A receptor subtypes: dissecting their pharmacological functions. *Trends Pharmacol. Sci.* **22**, 188–194 (2001).
65. Di Cristo, G. *et al.* Subcellular domain-restricted GABAergic innervation in primary visual cortex in the absence of sensory and thalamic inputs. *Nature Neurosci.* **7**, 1184–1186 (2004).
66. Chattopadhyaya, B. *et al.* Experience and activity-dependent maturation of perisomatic GABAergic innervation in primary visual cortex during a postnatal critical period. *J. Neurosci.* **24**, 9598–9611 (2004).
67. Buzas, P., Eysel, U. T., Adorjan, P. & Kisvarday, Z. F. Axonal topography of cortical basket cells in relation to orientation, direction, and ocular dominance maps. *J. Comp. Neurol.* **437**, 259–285 (2001).
68. Klausberger, T., Roberts, J. D. & Somogyi, P. Cell type- and input-specific differences in the number and subtypes of synaptic GABA_A receptors in the hippocampus. *J. Neurosci.* **22**, 2513–2521 (2002).
- Demonstrates remarkable subcellular sorting of individual GABA_A receptor subtypes to receive distinct inhibitory inputs based on α -subunit composition.**
69. Nusser, Z., Sieghart, W., Benke, D., Fritschy, J. M. & Somogyi, P. Differential synaptic localization of two major γ -aminobutyric acid type A receptor α subunits on hippocampal pyramidal cells. *Proc. Natl Acad. Sci. USA* **93**, 11939–11944 (1996).
70. Härtig, W. *et al.* Cortical neurons immunoreactive for the potassium channel K_v3.1b subunit are predominantly surrounded by perineuronal nets presumed as a buffering system for cations. *Brain Res.* **842**, 15–29 (1999).
71. Saghatelayan, A. K. *et al.* Reduced perisomatic inhibition, increased excitatory transmission, and impaired long-term potentiation in mice deficient for the extra-cellular matrix glycoprotein tenascin-R. *Mol. Cell. Neurosci.* **17**, 226–240 (2001).
72. Pizzorusso, T. *et al.* Reactivation of ocular dominance plasticity in the adult visual cortex. *Science* **298**, 1248–1251 (2002).
- The holy grail of critical period studies that showed it is possible to reactivate plasticity in adulthood, achieved by disrupting ECM components of perineuronal nets with chondroitinases. These structures preferentially surround and influence the function of large parvalbumin-positive basket cells (see also references 70 and 71).**
73. Berardi, N., Pizzorusso, T. & Maffei, L. Extracellular matrix and visual cortical plasticity: freeing the synapse. *Neuron* **44**, 905–908 (2004).
74. Liu, Y., Fields, R. D., Fitzgerald, S., Festoff, B. W. & Nelson, P. G. Proteolytic activity, synapse elimination, and the Hebb synapse. *J. Neurobiol.* **25**, 325–335 (1994).
75. Shiosaka, S. & Yoshida, S. Synaptic microenvironments – structural plasticity, adhesion molecules, proteases and their inhibitors. *Neurosci. Res.* **37**, 85–89 (2000).
76. Mataga, N. & Hensch, T. K. in *Proteases in Biology and Disease* Vol. 3 (eds Lendeckel, U. & Hooper, N.) Chapter 11 (Kluwer/Plenum, New York, in the press).
77. Qian, Z., Gilbert, M. E., Colicos, M. A., Kandel, E. R. & Kuhl, D. Tissue-plasminogen activator is induced as an immediate-early gene during seizure, kindling and long-term potentiation. *Nature* **361**, 453–457 (1993).
78. Mataga, N., Nagai, N. & Hensch, T. K. Permissive proteolytic activity for visual cortical plasticity. *Proc. Natl Acad. Sci. USA* **99**, 7717–7721 (2002).
- Definitive evidence that proteases are required for ocular dominance plasticity through the use of genetically targeted animals, supporting earlier pharmacological studies in cats (detailed in references 79 and 80).**
79. Mataga, N. *et al.* Enhancement of mRNA expression of tissue-type plasminogen activator by L1-threo-3,4-dihydroxyphenylserine in association with ocular dominance plasticity. *Neurosci. Lett.* **218**, 149–152 (1996).
80. Müller, C. M. & Griesinger, C. B. Tissue plasminogen activator mediates reverse occlusion plasticity in visual cortex. *Nature Neurosci.* **1**, 47–53 (1998).
81. Berardi, N., Pizzorusso, T., Ratto, G. M. & Maffei, L. Molecular basis of plasticity in the visual cortex. *Trends Neurosci.* **26**, 369–378 (2003).
82. Silver, M. A. & Stryker, M. P. Synaptic density in geniculocortical afferents remains constant after monocular deprivation in the cat. *J. Neurosci.* **19**, 10829–10842 (1999).
83. Grutzendler, J., Kasthuri, N. & Gan, W. B. Long-term dendritic spine stability in the adult cortex. *Nature* **420**, 812–816 (2002).
84. Majewska, A. & Sur, M. Motility of dendritic spines in visual cortex *in vivo*: changes during the critical period and effects of visual deprivation. *Proc. Natl Acad. Sci. USA* **100**, 16024–16029 (2003).
85. Oray, S., Majewska, A. & Sur, M. Dendritic spine dynamics are regulated by monocular deprivation and extracellular matrix degradation. *Neuron* **44**, 1021–1030 (2004).
86. Mataga, N., Mizuguchi, Y. & Hensch, T. K. Experience-dependent pruning of dendritic spines in visual cortex by tissue plasminogen activator. *Neuron* **44**, 1031–1041 (2004).
- References 85 and 86 reveal the first anatomical events that occur on monocular deprivation. Spine motility followed by transient pruning faithfully reflects competition during the critical period and is mediated by tPA-plasmin *in vivo*.**
87. Taha, S. & Stryker, M. P. Rapid ocular dominance plasticity requires cortical but not geniculate protein synthesis. *Neuron* **34**, 425–436 (2002).
88. Pang, P. *et al.* Cleavage of proBDNF by tPA/plasmin is essential for long-term hippocampal plasticity. *Science* **306**, 487–491 (2004).
89. Tyler, W. J. & Pozzo-Miller, L. Miniature synaptic transmission and BDNF modulates dendritic spine growth and form in rat CA1 neurons. *J. Physiol. (Lond.)* **553**, 497–509 (2004).
90. Miller, K. D. Synaptic economics: competition and cooperation in synaptic plasticity. *Neuron* **17**, 371–374 (1996).
- Careful consideration of the theory and cellular events behind the competitive nature of critical period plasticity in the visual cortex.**
91. Barry, M. F. & Ziff, E. B. Receptor trafficking and the plasticity of excitatory synapses. *Curr. Opin. Neurobiol.* **12**, 279–286 (2002).
92. Takahashi, T., Svoboda, K. & Malinow, R. Experience strengthening transmission by driving AMPA receptors into synapses. *Science* **299**, 1585–1588 (2003).
93. Allen, C. B., Celikel, T. & Feldman, D. E. Long-term depression induced by sensory deprivation during cortical map plasticity *in vivo*. *Nature Neurosci.* **6**, 291–299 (2003).
94. Heynen, A. J. *et al.* Molecular mechanism for loss of visual cortical responsiveness following brief monocular deprivation. *Nature Neurosci.* **6**, 854–862 (2003).
95. Flint, A. C., Maisch, U. S., Weishaupt, J. H., Kriegstein, A. R. & Monyer, H. NR2A subunit expression shortens NMDA receptor synaptic currents in developing neocortex. *J. Neurosci.* **17**, 2469–2476 (1997).
96. Nase, G., Weishaupt, J., Stern, P., Singer, W. & Monyer, H. Genetic and epigenetic regulation of NMDA receptor expression in the rat visual cortex. *Eur. J. Neurosci.* **11**, 4320–4326 (1999).
97. Quinlan, E. M., Philpot, B. D., Hagan, R. L. & Bear, M. F. Rapid, experience-dependent expression of synaptic NMDA receptors in visual cortex *in vivo*. *Nature Neurosci.* **2**, 352–357 (1999).
98. Philpot, B. D., Sekhar, A. K., Shouval, H. Z. & Bear, M. F. Visual experience and deprivation bidirectionally modify the composition and function of NMDA receptors in visual cortex. *Neuron* **29**, 157–169 (2001).
99. Tang, Y. P. *et al.* Genetic enhancement of learning and memory in mice. *Nature* **401**, 63–69 (1999).
100. Liu, L. *et al.* Role of NMDA receptor subtypes in governing the direction of hippocampal synaptic plasticity. *Science* **304**, 1021–1024 (2004).
101. Lu, H. C., Gonzalez, E. & Crair, M. C. Barrel cortex critical period plasticity is independent of changes in NMDA receptor subunit composition. *Neuron* **32**, 619–634 (2001).
102. Barth, A. L. & Malenka, R. C. NMDAR EPSC kinetics do not regulate the critical period for LTP at thalamocortical synapses. *Nature Neurosci.* **4**, 235–236 (2001).
103. Datwani, A., Iwasato, T., Itoharu, S. & Erzurumlu, R. S. Lesion-induced thalamocortical axonal plasticity in the S1 cortex is independent of NMDA receptor function in excitatory cortical neurons. *J. Neurosci.* **22**, 9171–9175 (2002).
104. Rebsam, A., Seif, I. & Gaspar, P. Dissociating barrel development and plasticity in the mouse somatosensory cortex. *J. Neurosci.* **25**, 706–710 (2005).
- Clever use of the monamine oxidase A-knockout mouse, whose barrel formation can be rescued by lowering serotonin levels with parachlorophenylalanine (PCPA) at ages beyond the critical period for barrel plasticity. Closure of the whisker cautery effect on the barrelfield is independent and probably not determined at the cortical level.**
105. Feldman, D. E. Inhibition and plasticity. *Nature Neurosci.* **3**, 303–304 (2000).
106. Dudai, Y. Molecular bases of long-term memories: a question of persistence. *Curr. Opin. Neurobiol.* **12**, 211–216 (2002).
107. O'Connor, D. H., Wittenberg, G. M. & Wang, S. S.-H. Graded bidirectional synaptic plasticity is composed of switch-like unitary events. *Proc. Natl Acad. Sci. USA* **102**, 9679–9684 (2005).
108. Sajikumar, S., Navakkode, S. & Frey, J. U. Protein synthesis-dependent long-term functional plasticity: methods and techniques. *Curr. Opin. Neurobiol.* (in the press).
109. Jones, M. W. *et al.* A requirement for the immediate early gene *Zif268* in the expression of late LTP and long-term memories. *Nature Neurosci.* **4**, 289–296 (2001).
110. Mataga, N., Fujishima, S., Condie, B. G. & Hensch, T. K. Experience-dependent plasticity of mouse visual cortex in the absence of the neuronal activity-dependent marker *egr1/zif268*. *J. Neurosci.* **21**, 9724–9732 (2001).
111. Baranes, D. *et al.* Tissue plasminogen activator contributes to the late phase of LTP and to synaptic growth in the hippocampal mossy fiber pathway. *Neuron* **21**, 813–825 (1998).
112. Neuhoff, H., Roepner, J. & Schweizer, M. Activity-dependent formation of perforated synapses in cultured hippocampal neurons. *Eur. J. Neurosci.* **11**, 4241–4250 (1999).
113. Daw, N. in *The Visual Neurosciences* Vol. 1 (eds Chalupa, L. & Werner, J. S.) 126–145 (MIT Press, Cambridge, Massachusetts, USA, 2004).
114. Hensch, T. K. Controlling the critical period. *Neurosci. Res.* **47**, 17–22 (2003).
115. Renger, J. J. *et al.* Experience-dependent plasticity without long-term depression by type 2 metabotropic glutamate receptors in developing visual cortex. *Proc. Natl Acad. Sci. USA* **99**, 1041–1046 (2002).
116. Bartoletti, A. *et al.* Heterozygous knock-out mice for brain-derived neurotrophic factor show a pathway-specific impairment of long-term potentiation but normal critical period for monocular deprivation. *J. Neurosci.* **22**, 10072–10077 (2002).
117. Daw, N., Rao, Y., Wang, X. F., Fischer, Q. & Yang, Y. LTP and LTD vary with layer in rodent visual cortex. *Vision Res.* **44**, 3377–3380 (2004).
118. Zhou, Q., Homma, K. J. & Poo, M. M. Shrinkage of dendritic spines associated with long-term depression of hippocampal synapses. *Neuron* **44**, 749–757 (2004).
- Demonstrates that spine shape changes can occur on a fast time scale, but independent of mechanisms that underlie the LTD of synaptic transmission.**
119. Hayashi, Y. & Majewska, A. K. Dendritic spine geometry: functional implication and regulation. *Neuron* **46**, 529–532 (2005).
120. Yang, Y. *et al.* Reversible blockade of experience-dependent plasticity by calcineurin in mouse visual cortex. *Nature Neurosci.* **8**, 791–796 (2005).
121. Wan, H. *et al.* Benzodiazepine impairment of perirhinal cortical plasticity and recognition memory. *Eur. J. Neurosci.* **20**, 2214–2224 (2004).
122. Jiang, B., Akaneva, Y., Hata, Y. & Tsumoto, T. Long-term depression is not induced by low-frequency stimulation in rat visual cortex *in vivo*: a possible preventing role of endogenous brain-derived neurotrophic factor. *J. Neurosci.* **23**, 3761–3770 (2003).

123. Bi, G. & Poo, M. Synaptic modification by correlated activity: Hebb's postulate revisited. *Annu. Rev. Neurosci.* **24**, 139–166 (2001).
124. Song, S., Miller, K. D. & Abbott, L. F. Competitive Hebbian learning through spike-timing-dependent synaptic plasticity. *Nature Neurosci.* **3**, 919–926 (2000).
125. Pouille, F. & Scanziani, M. Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. *Science* **293**, 1159–1163 (2001).
126. Sanes, J. R. & Lichtman, J. W. Can molecules explain long-term potentiation? *Nature Neurosci.* **2**, 597–604 (1999).
127. Zhu, Y., Stornetta, R. L. & Zhu, J. J. Chandelier cells control excessive cortical excitation: characteristics of whisker-evoked synaptic responses of layer 2/3 nonpyramidal and pyramidal neurons. *J. Neurosci.* **24**, 5101–5108 (2004).
128. Galarreta, M. & Hestrin, S. Electrical synapses between GABA-releasing interneurons. *Nature Rev. Neurosci.* **2**, 425–433 (2001).
129. Meyer, A. H., Katona, I., Blatow, M., Rozov, A. & Monyer, H. *In vivo* labeling of parvalbumin-positive interneurons and analysis of electrical coupling in identified neurons. *J. Neurosci.* **22**, 7055–7064 (2002).
130. Galarreta, M. & Hestrin, S. Spike transmission and synchrony detection in networks of GABAergic interneurons. *Science* **292**, 2295–2299 (2001).
- Paired recordings from electrically-coupled, parvalbumin-positive cells reveals a novel property of such networks, namely synchrony detection, which could operate on a columnar scale.**
131. Gao, B. & Fritschy, J. M. Selective allocation of GABA_A receptors containing the $\alpha 1$ subunit to neurochemically distinct subpopulations of rat hippocampal interneurons. *Eur. J. Neurosci.* **6**, 837–853 (1994).
132. Deans, M. R., Gibson, J. R., Sellitto, C., Connors, B. W. & Paul, D. L. Synchronous activity of inhibitory networks in neocortex requires electrical synapses containing connexin36. *Neuron* **31**, 477–485 (2001).
133. Hormuzdi, S. G. *et al.* Impaired electrical signaling disrupts gamma frequency oscillations in connexin 36-deficient mice. *Neuron* **31**, 487–495 (2001).
134. Guldenagel, M. *et al.* Expression patterns of connexin genes in mouse retina. *J. Comp. Neurol.* **425**, 193–201 (2000).
135. Dityatev, A. & Schachner, M. Extracellular matrix molecules and synaptic plasticity. *Nature Rev. Neurosci.* **4**, 456–468 (2003).
136. Lochner, J. E. *et al.* Real-time imaging of the axonal transport of granules containing a tissue plasminogen activator/green fluorescent protein hybrid. *Mol. Biol. Cell* **9**, 2463–2476 (1998).
137. Gualandris, A., Jones, T. E., Strickland, S. & Tsirka, S. E. Membrane depolarization induces calcium-dependent secretion of tissue plasminogen activator. *J. Neurosci.* **16**, 2220–2225 (1996).
138. Parmer, R. J. *et al.* Tissue plasminogen activator (tPA) is targeted to the regulated secretory pathway. *J. Biol. Chem.* **272**, 1976–1982 (1997).
139. Collin, T. *et al.* Developmental changes in parvalbumin regulate presynaptic Ca²⁺ signaling. *J. Neurosci.* **25**, 96–107 (2005).
140. Murase, S., Mosser, E. & Schuman, E. M. Depolarization drives β -catenin into neuronal spines promoting changes in synaptic structure and function. *Neuron* **35**, 91–105 (2002).
141. Tanaka, H. *et al.* Molecular modification of N-cadherin in response to synaptic activity. *Neuron* **25**, 93–107 (2000).
- Strongly suggests that synaptic adhesion is locally controlled and dynamically modulated by neuronal activity through the acquisition of protease resistance by dimerization of N-cadherin on depolarization.**
142. Wannier-Morino, P., Rager, G., Sonderegger, P. & Grabs, D. Expression of neuroserpin in the visual cortex of the mouse during the developmental critical period. *Eur. J. Neurosci.* **17**, 1853–1860 (2003).
143. Sakaguchi, H. Sex differences in the developmental changes of GABAergic neurons in zebra finch song control nuclei. *Exp. Brain Res.* **108**, 62–68 (1996).
- Intriguing evidence that the motor phase of the zebra finch critical period for song acquisition reflects maturation of GABA-containing cells in male nucleus RA.**
144. Freund, T. F., Katona, I. & Piomelli, D. Role of endogenous cannabinoids in synaptic signaling. *Physiol. Rev.* **83**, 1017–1066 (2003).
145. Soderstrom, K. & Johnson, F. Cannabinoid exposure alters learning of zebra finch vocal patterns. *Brain Res.* **142**, 215–217 (2003).
146. Knudsen, E. I., Zheng, W. & DeBello, W. M. Traces of learning in the auditory localization pathway. *Proc. Natl Acad. Sci. USA* **97**, 11815–11820 (2000).
147. Knudsen, E. I. Capacity for plasticity in the adult owl auditory system expanded by juvenile experience. *Science* **279**, 1531–1533 (1998).
- Demonstrates the power of the critical period in laying down multiple neural representations of early experienced environments that can be reactivated when re-encountered later in life. Active suppression by newly formed GABA circuits prevents confusion among multiple maps (see reference 148).**
148. Zheng, W. & Knudsen, E. I. Functional selection of adaptive auditory space map by GABA_A-mediated inhibition. *Science* **284**, 962–965 (1999).
149. Linkenhoker, B. A. & Knudsen, E. I. Incremental training increases the plasticity of the auditory space map in adult barn owls. *Nature* **419**, 293–296 (2002).
- Suggests a new strategy for adult plasticity (without drugs) based on incremental training to cumulatively overcome the anatomical constraints established during the critical period.**
150. Linkenhoker, B. A., von der Ohe, C. G. & Knudsen, E. I. Anatomical traces of juvenile learning in the auditory system of adult barn owls. *Nature Neurosci.* **8**, 93–98 (2005).
151. Fuchs, J. L. & Salazar, E. Effects of whisker trimming on GABA_A receptor binding in the barrel cortex of developing and adult rats. *J. Comp. Neurol.* **395**, 209–216 (1998).
152. Micheva, K. D. & Beaulieu, C. An anatomical substrate for experience-dependent plasticity of the rat barrel field cortex. *Proc. Natl Acad. Sci. USA* **92**, 11834–11838 (1995).
153. Knott, G. W., Quairiaux, C., Genoud, C. & Welker, E. Formation of dendritic spines with GABAergic synapses induced by whisker stimulation in adult mice. *Neuron* **34**, 265–273 (2002).
- Demonstrates plasticity of GABA-mediated connections in the adult barrel cortex.**
154. Diamond, M. E., Armstrong-James, M. & Ebner, F. F. Experience-dependent plasticity in adult rat barrel cortex. *Proc. Natl Acad. Sci. USA* **90**, 2082–2086 (1993).
155. Gheusi, G. *et al.* Importance of newly generated neurons in the adult olfactory bulb for odor discrimination. *Proc. Natl Acad. Sci. USA* **97**, 1823–1828 (2000).
156. Cecchi, G. A., Petreanu, L. T., Alvarez-Buylla, A. & Magnasco, M. O. Unsupervised learning and adaptation in a model of adult neurogenesis. *J. Comput. Neurosci.* **11**, 175–182 (2001).
157. Yokoi, M., Mori, K. & Nakanishi, S. Refinement of odor molecule tuning by dendro-dendritic synaptic inhibition in the olfactory bulb. *Proc. Natl Acad. Sci. USA* **92**, 3371–3375 (1995).
158. Lagier, S., Carleton, A. & Lledo, P. M. Interplay between local GABA-mediated interneurons and relay neurons generates gamma oscillations in the rat olfactory bulb. *J. Neurosci.* **24**, 4382–4392 (2004).
159. Murphy, K. M., Beston, B. R., Boley, P. M. & Jones, D. G. Development of human visual cortex: a balance between excitatory and inhibitory plasticity mechanisms. *Dev. Psychobiol.* **46**, 209–221 (2005).
- Crucial evidence from human autopsy samples showing slow maturation of GAD65 and GABA_A receptor $\alpha 1$ -subunits in the visual cortex that better matches the prolonged critical period for binocular vision in this species (see references 6 and 7) than does NMDA-receptor subunit switching (for further information, see references 43, 46 and 53).**
160. De Negri, M., Baglietto, M. G. & Biancheri, R. Electrical status epilepticus in childhood: treatment with short cycles of high dosage benzodiazepine. *Brain Dev.* **15**, 311–312 (1993).
161. Rubenstein, J. L. & Merzenich, M. M. Model of autism: increased ratio of excitation/inhibition in key neural systems. *Genes Brain Behav.* **2**, 255–267 (2003).
162. Möhler, H., Fritschy, J. M., Crestani, F., Hensch, T. & Rudolph, U. Specific GABA_A circuits in brain development and therapy. *Biochem. Pharmacol.* **68**, 1685–1690 (2004).
163. Arckens, L. *et al.* Cooperative changes in GABA, glutamate and activity levels: the missing link in cortical plasticity. *Eur. J. Neurosci.* **12**, 4222–4232 (2000).
164. Lodder, J., Luijckx, G., van Raak, L. & Kessels, F. Diazepam treatment to increase the cerebral GABAergic activity in acute stroke: a feasibility study in 104 patients. *Cerebrovasc. Dis.* **10**, 437–440 (2000).
165. Dani, V. S. *et al.* Reduced cortical activity due to a shift in the balance between excitation and inhibition in a mouse model of Rett syndrome. *Proc. Natl Acad. Sci. USA* **102**, 12560–12565 (2005).
166. Kalanithi, P. S. A. *et al.* Altered parvalbumin-positive neuron distribution in basal ganglia of individuals with Tourette syndrome. *Proc. Natl Acad. Sci. USA* **102**, 13307–13312 (2005).
167. Lewis, D. A., Hashimoto, T. & Volk, D. W. Cortical inhibitory neurons and schizophrenia. *Nature Rev. Neurosci.* **6**, 312–324 (2005).
168. Paulsen, O. & Moser, E. I. A model of hippocampal memory encoding and retrieval: GABAergic control of synaptic plasticity. *Trends Neurosci.* **21**, 273–378 (1998).
169. Miyamoto, H., Katagiri, H. & Hensch, T. Experience-dependent slow-wave sleep development. *Nature Neurosci.* **6**, 553–554 (2003).
170. Prasad, S. S. *et al.* Gene expression patterns during enhanced periods of visual cortex plasticity. *Neuroscience* **111**, 35–45 (2002).
171. Ossipow, V., Pellissier, F., Schaad, O. & Ballivet, M. Gene expression analysis of the critical period in the visual cortex. *Mol. Cell. Neurosci.* **27**, 70–83 (2004).
172. Sawtell, N. B. *et al.* NMDA receptor-dependent ocular dominance plasticity in adult visual cortex. *Neuron* **38**, 977–985 (2003). Erratum *Neuron* **39**, 727 (2003).
173. Pham, T. A. *et al.* A semi-persistent adult ocular dominance plasticity in visual cortex is stabilized by activated CREB. *Learn. Mem.* **11**, 738–747 (2004).
174. Tagawa, Y., Kanold, P. O., Majdan, M. & Shatz, C. J. Multiple periods of functional ocular dominance plasticity in mouse visual cortex. *Nature Neurosci.* **8**, 380–388 (2005).
175. Fagiolini, M., Pizzorusso, T., Berardi, N., Domenici, L. & Maffei, L. Functional postnatal development of the rat primary visual cortex and the role of visual experience: dark rearing and monocular deprivation. *Vision Res.* **34**, 709–720 (1994).
176. Wang, X., Merzenich, M. M., Sameshima, K. & Jenkins, W. M. Remodelling of hand representation in adult cortex determined by timing of tactile stimulation. *Nature* **378**, 71–75 (1995).
177. Kilgard, M. P. & Merzenich, M. M. Cortical map reorganization enabled by nucleus basalis activity. *Science* **279**, 1714–1718 (1998).
178. Bao, S., Chan, V. T. & Merzenich, M. M. Cortical remodelling induced by activity of ventral tegmental dopamine neurons. *Nature* **412**, 79–83 (2001).
- Two studies that demonstrate the potential for neuromodulatory systems (attention) to restore near critical period levels of plasticity to the adult brain.**
179. Miyamoto, H. & Hensch, T. K. Reciprocal interaction of sleep and synaptic plasticity. *Mol. Interv.* **3**, 404–417 (2003).
180. Hinkle, D. J. & Macdonald, R. L. β subunit phosphorylation selectively increases fast desensitization and prolongs deactivation of $\alpha 1\beta 1\gamma 2L$ and $\alpha 1\beta 3\gamma 2L$ GABA_A receptor currents. *J. Neurosci.* **23**, 11698–11710 (2003).
181. Fischer, Q. S. *et al.* Requirement for the RII β isoform of PKA, but not calcium-stimulated adenylyl cyclase, in visual cortical plasticity. *J. Neurosci.* **24**, 9049–9058 (2004).
182. McGee, A. *et al.* Experience-driven plasticity of visual cortex limited by myelin and Nogo receptor. *Science* **309**, 2222–2226 (2005).
183. Mechelli, A. *et al.* Neurolinguistics: structural plasticity in the bilingual brain. *Nature* **431**, 757 (2004).
184. Bengtsson, S. L. *et al.* Extensive piano practicing has regionally specific effects on white matter development. *Nature Neurosci.* **8**, 1148–1150 (2005).
185. Fiumelli, H., Jabaudon, D., Magistretti, P. J. & Martin, J.-L. BDNF stimulates expression, activity and release of tissue-type plasminogen activator in mouse cortical neurons. *Eur. J. Neurosci.* **11**, 1639–1646 (1999).
186. Lewis, T. L. & Maurer, D. Multiple sensitive periods in human visual development: evidence from visually deprived children. *Dev. Psychobiol.* **46**, 163–183 (2005).

Acknowledgements

I thank M. Fagiolini for comments, N. Mataga for immunostaining in figure 5b and all members of the lab for Neuronal Circuit Development (RIKEN BSI) for their dedication to understanding critical period mechanisms.

Competing interests statement

The author declares no competing financial interests.

Online links

DATABASES

The following terms in this article are linked online to:

Entrez Gene: <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene>

BDNF | [Gad65](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene) | [Gad67](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene) | [Zif268](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene)

OMIM: <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM>

Rett syndrome | Tourette's syndrome

FURTHER INFORMATION

<http://www.brain.riken.jp/>

Access to this interactive links box is free online.