

Critical points of the Moser-Trudinger functional on a disk

Andrea Malchiodi
SISSA, Trieste
malchiod@sissa.it

Luca Martinazzi
Rutgers University
martinazzi@math.rutgers.edu

March 5, 2012

Abstract

On the unit disk $B_1 \subset \mathbb{R}^2$ we study the Moser-Trudinger functional

$$E(u) = \int_{B_1} (e^{u^2} - 1) dx, \quad u \in H_0^1(B_1)$$

and its restrictions $E|_{M_\Lambda}$, where $M_\Lambda := \{u \in H_0^1(B_1) : \|u\|_{H_0^1}^2 = \Lambda\}$ for $\Lambda > 0$. We prove that if a sequence u_k of positive critical points of $E|_{M_{\Lambda_k}}$ (for some $\Lambda_k > 0$) blows up as $k \rightarrow \infty$, then $\Lambda_k \rightarrow 4\pi$, and $u_k \rightarrow 0$ weakly in $H_0^1(B_1)$ and strongly in $C_{\text{loc}}^1(\overline{B_1} \setminus \{0\})$.

Using this we also prove that when Λ is large enough, then $E|_{M_\Lambda}$ has no positive critical point, complementing previous existence results by Carleson-Chang, M. Struwe and Lamm-Robert-Struwe.

Key Words: Moser-Trudinger inequality, critical points, blow-up analysis, variational methods

AMS subject classification: 35B33, 35B44, 35A01, 34E05

1 Introduction

Let $\Omega \Subset \mathbb{R}^2$ be a smooth bounded and connected open set. It is well-known that there is a Sobolev embedding $W_0^{1,p}(\Omega) \hookrightarrow L^{\frac{2p}{2-p}}(\Omega)$ for $p \in [1, 2)$, but $H_0^1(\Omega) := W_0^{1,2}(\Omega) \not\hookrightarrow L^\infty(\Omega)$. However it was proven by N. Trudinger [29] that $e^{u^2} \in L^1(\Omega)$ whenever $u \in H_0^1(\Omega)$. This embedding was sharpened by J. Moser [23] who showed that

$$\sup_{u \in H_0^1(\Omega), \|u\|_{H_0^1}^2 \leq 4\pi} \int_{\Omega} (e^{u^2} - 1) dx \leq C|\Omega|, \quad \|u\|_{H_0^1} := \left(\int_{\Omega} |\nabla u|^2 dx \right)^{\frac{1}{2}}, \quad (1)$$

and

$$\sup_{u \in H_0^1(\Omega), \|u\|_{H_0^1}^2 \leq 4\pi + \delta} \int_{\Omega} (e^{u^2} - 1) dx = +\infty, \quad \text{for every } \delta > 0. \quad (2)$$

Since then, a formidable amount of work has been devoted to the study of the functional

$$E(u) := \int_{\Omega} (e^{u^2} - 1) dx, \quad u \in H_0^1(\Omega)$$

and in particular of its critical points. Clearly $u \equiv 0$ is the only global minimum of E , but because of (2) we cannot look for a global maximizer of E in H_0^1 . Instead one might hope to find a maximizer of $E|_{M_\Lambda}$, i.e. of E constrained to the manifold

$$M_\Lambda := \left\{ u \in H_0^1(\Omega) : \|u\|_{H_0^1}^2 = \Lambda \right\}$$

for $\Lambda \in (0, 4\pi]$, or to find other kinds of critical points (local maxima or minima, saddle points, etc.) when $\Lambda > 4\pi$. As long as $\Lambda < 4\pi$ the embedding (1) is in fact compact, so the existence of a maximizer is elementary, but when $\Lambda \geq 4\pi$ compactness is lost and also the Palais-Smale condition does not hold anymore, see [3].

In spite of these difficulties Carleson and Chang [7] proved that when $\Omega = B_1(0)$ (the unit disk in \mathbb{R}^2) $E|_{M_{4\pi}}$ has a maximizer. This result was extended by Struwe [26] who proved the existence of a maximizer in $M_{4\pi}$ when Ω is close to a ball, and finally by Flucher [13] for any bounded smooth Ω (see also [9] for a related result in higher dimension).

The existence of critical points on M_Λ in the supercritical regime, i.e. for $\Lambda > 4\pi$, is even more challenging, and to the fundamental question of the existence of critical points of $E|_{M_\Lambda}$ for Λ large only few answers have been given. Monahan [22] gave numerical evidence that when $\Omega = B_1(0)$ then for some $\Lambda^* > 4\pi$ the functional $E|_{M_\Lambda}$ has a local maximum and a mountain pass critical point for every $\Lambda \in (4\pi, \Lambda^*)$. Assuming that a local maximum of $E|_{M_{4\pi}}$ exists (which was later shown to be true for arbitrary domains by Flucher [13]) Struwe proved in [26] that for some $\Lambda^* = \Lambda^*(\Omega) > 4\pi$ and for a.e. $\Lambda \in (4\pi, \Lambda^*)$ two critical points exist. This result was then extended in [16] to all values of $\Lambda \in (4\pi, \Lambda^*)$ through the more precise information given by a parabolic flow, compared to the one given by the Palais-Smale condition.

Further, using implicit function methods, Del Pino, Musso and Ruf [11] were able to characterize some of these critical points as one-peaked bubbling functions which blow-up as $\Lambda \searrow 4\pi$. In the same paper they showed that if Ω is not contractible, then for some $\Lambda^\dagger > 8\pi$ the functional $E|_{M_\Lambda}$ has a critical point of multi-peak type for $\Lambda \in (8\pi, \Lambda^\dagger)$. When Ω is a radially symmetric annulus they also proved for any $1 \leq \ell \in \mathbb{N}$ the existence of some $\Lambda_\ell^* > 4\pi\ell$ such that $E|_{M_\Lambda}$ has a critical point when $\Lambda \in (4\pi\ell, \Lambda_\ell^*)$. We also refer to [27] and [16] for related results on domains with *small holes*, in the spirit of [8] (where the Yamabe equation was treated).

The previous results, in particular those in [11], suggest that at least when Ω is not contractible $E|_{M_\Lambda}$ might have critical points even when Λ is much larger than 4π . In this paper we will show that such a topological assumption on Ω is natural. In fact we will prove that when $\Omega = B_1(0)$, then $E|_{M_\Lambda}$ has no positive critical points for Λ large enough.

Theorem 1 *For $\Omega = B_1(0)$ there exists $\Lambda^\sharp > 4\pi$ such that the functional $E|_{M_\Lambda}$ has*

- (i) *no positive critical points for $\Lambda > \Lambda^\sharp$,*

(ii) at least 2 positive critical points for $\Lambda \in (4\pi, \Lambda^\sharp)$,

(iii) at least one positive critical point for $\Lambda \in (0, 4\pi] \cup \{\Lambda^\sharp\}$.

The proof of the non-existence part in Theorem 1 (Part (i)) will be completely self-contained. To prove Parts (ii) and (iii) we will also use Theorem 1.7 from [26], which gives the existence of some $\Lambda^* > 4\pi$ such that a positive critical point of $E|_{M_\Lambda}$ exists for $\Lambda \in (4\pi, \Lambda^*)$. Actually by [26, Theorem 1.8] and [16, Theorem 6.5], $E|_{M_\Lambda}$ has two positive critical points whenever $\Lambda \in (4\pi, \Lambda_*)$, for some $\Lambda_* \in (4\pi, \Lambda^*)$. Then Theorem 1 complements these results by showing that, at least when $\Omega = B_1$, the existence of two positive critical points of $E|_{M_\Lambda}$ for $\Lambda > 4\pi$ persists until we reach the energy threshold $\Lambda = \Lambda^\sharp$, beyond which we have non-existence. A qualitatively similar picture has also been shown in [15], [21] (as well as in several subsequent papers in the literature) for the problem $-\Delta u = \mu f(u)$ in bounded domains of \mathbb{R}^n . Differently from these results, we focus on the Dirichlet energy rather than on the parameter μ , and we deal with a faster growth of the nonlinearity.

In order to prove Theorem 1 we first notice that a critical point of $E|_{M_\Lambda}$ solves

$$\begin{cases} -\Delta u = \lambda u e^{u^2} & \text{in } \Omega; \\ u = 0 & \text{on } \partial\Omega; \\ \|u\|_{H_0^1}^2 = \Lambda, \end{cases} \quad (3)$$

for some $\lambda > 0$, and that when $\Omega = B_1$ a positive solution to (3) is radially symmetric by Theorem 1 in [14]. Then it will be crucial to understand the blow-up behavior of a sequence of symmetric positive solutions to (3), i.e. solutions u_k to

$$\begin{cases} -\Delta u_k = \lambda_k u_k e^{u_k^2} & \text{in } B_1; \\ u_k = 0 & \text{on } \partial B_1; \\ u_k > 0 & \text{in } B_1; \\ \|u_k\|_{H_0^1}^2 = \Lambda_k. \end{cases} \quad (4)$$

In this direction a lot of work has already been done. For the sake of simplicity we shall present only the radially symmetric versions of the results which we quote, referring to the original papers for the general cases. For instance O. Druet proved (see also [4] and [2] for previous related results, where the blow-up profile was identified, and [16] where the parabolic case was treated): let (u_k) be a sequence of solutions to (4) with $\Lambda_k \leq C$ and $\sup_{B_1} u_k \rightarrow \infty$. Then up to a subsequence $\lambda_k \rightarrow \lambda_\infty \in [0, 2\pi]$,¹ $u_k \rightarrow u_\infty$ strongly in $C_{\text{loc}}^1(\overline{B_1} \setminus \{0\})$ and weakly in $H_0^1(B_1)$, where

$$-\Delta u_\infty = \lambda_\infty u_\infty e^{u_\infty^2} \quad \text{in } B_1, \quad (5)$$

and $\Lambda_k \rightarrow 4\pi L + \|u_\infty\|_{H_0^1}^2$ for some integer $L \geq 1$. More precisely

$$|\nabla u_k|^2 dx \rightharpoonup 4\pi L \delta_0 + |\nabla u_\infty|^2 dx, \quad \lambda_k u_k^2 e^{u_k^2} dx \rightharpoonup 4\pi L \delta_0 + \lambda_\infty u_\infty^2 e^{u_\infty^2} dx \quad (6)$$

¹The constant 2π is the first eigenvalue of $-\Delta$ on B_1 . As proven by Adimurthi [1], Problem (3) has a positive solution if and only if $\lambda \in (0, \lambda_1(\Omega))$, where $\lambda_1(\Omega)$ is the first eigenvalue of $-\Delta$ on Ω with Dirichlet boundary condition.

weakly in the sense of measures. The questions whether u_∞ and λ_∞ can actually be non-zero, and whether L can be greater than one (i.e. whether the blow-up can be non-simple, using a terminology introduced in [24]) were left open (in fact also higher dimensional generalization of the result of Druet, see e.g. [19], [20] and [28], produced analogous open questions), but we are now able to give a negative answer to both questions, as stated in the next theorem.

Theorem 2 *Let u_k be a sequence of solutions to (4). Then up to extracting a subsequence we have for $k \rightarrow \infty$ either*

(i) $\lambda_k \rightarrow \lambda_\infty \in [0, 2\pi]$, $u_k \rightarrow u_\infty$ in $C^1(\overline{B}_1)$, where u_∞ solves (5), or

(ii) $\lambda_k \rightarrow 0$, $\|u_k\|_{H_0^1}^2 \rightarrow 4\pi$, $u_k \rightarrow 0$ weakly in $H_0^1(B_1)$ and strongly in $C_{\text{loc}}^1(\overline{B}_1 \setminus \{0\})$ and

$$|\nabla u_k|^2 dx \rightharpoonup 4\pi\delta_0, \quad \lambda_k u_k^2 e^{u_k^2} dx \rightharpoonup 4\pi\delta_0 \quad (7)$$

weakly in the sense of measures.

The proof of Theorem 2 is self-contained. In some parts we could have used previous results of [4] or [12], but these hold for general domains, and consequently their proofs are more involved and we did not want to rest on them. Our main argument is not based on a Pohozaev-type identity as the results in [12], [16], [20] and [28], but on a simpler decay estimate of u_k away from the blow-up point, which has some partial analogies with Lemma 3 of [18] (originating in [25], see also [6]). Notice that, contrary to the previous works, e.g. [12], in our Theorem 2 we do *not* assume uniform bounds on $\|u_k\|_{H_0^1}^2$, i.e. $\lambda_k \leq C$. This is crucial if we want to apply Theorem 2 to prove Theorem 1.

The final picture that we get is then much closer to the geometric situation of the Liouville equation as studied by Brezis-Merle, Li-Shafrir and Li. More precisely, and working again on B_1 for simplicity, consider a sequence (v_k) of radially symmetric solutions to

$$-\Delta v_k = V_k e^{2v_k} \text{ in } B_1 \subset \mathbb{R}^2, \quad V_k \rightarrow V_0 > 0 \text{ in } C^0(\overline{B}_1), \quad \|e^{2v_k}\|_{L^1} \leq C, \quad \sup_{B_1} v_k \rightarrow +\infty.$$

Then, as proven in [5, Theorem 3],

$$v_k \rightarrow -\infty \text{ uniformly locally in } B_1 \setminus \{0\} \quad (8)$$

and

$$V_k e^{2v_k} dx \rightharpoonup \alpha \delta_0 \text{ weakly as measures, for some } \alpha \geq 2\pi. \quad (9)$$

Here $V_k e^{2v_k}$ plays the role of the energy density $\lambda_k u_k^2 e^{u_k^2}$ from (6) and (7). Then (8) and (9) are the equivalent of $\lambda_\infty u_\infty^2 e^{u_\infty^2} dx = 0$ in (7) (compared with (6)).

Y-Y. Li and I. Shafrir, see [17],[18], complemented the result of Brezis-Merle by showing that $\alpha = 4\pi$ in (9), finally yielding $V_k e^{2v_k} \rightharpoonup 4\pi\delta_0$, in analogy with (7). On the other hand we remark that the proof of (7) is more subtle because the nonlinearity ue^{u^2} is more difficult to handle than e^{2v} . In fact, as already noticed in previous works, e.g. [4], suitable scalings η_k of

blowing-up solutions of (4) converge in $C_{\text{loc}}^1(\mathbb{R}^2)$ to a solution η_0 of $-\Delta v = 4e^{2v}$, see Lemma 3. Unfortunately this information is too weak for our purposes and we need to linearize the equation satisfied by η_k (Eq. (14) below) to better understand its asymptotics (Lemma 4), and to have a global estimate of η_k (Lemma 5).

We also point out that an immediate consequence of the proof of Theorem 1 is the existence of blowing-up solutions to (4) with bounded energies ($\Lambda_k \rightarrow 4\pi$). This has long been an open problem: Adimurthi and Prashanth [3] were only able to prove the existence of blowing-up Palais-Smale sequences, while more recently Del Pino, Musso and Ruf, with an approach technically much richer, showed that blowing-up solutions exist for any domain Ω , see [11]. Our method applies only to the unit disk, but it is on the other hand relatively elementary and explicit.

In the following the letter C denotes a large constant which may change from line to line and even within the same line.

Acknowledgements

The authors are supported by the project FIRB-Ideas *Analysis and Beyond*, and L.M. was partially supported by the Swiss National Fond Grant no. PBEZP2-129520. During the preparation of this work L.M. was hosted by SISSA, while A.M. by Centro de Giorgi in Pisa. They are grateful to these institutions for the kind hospitality.

2 Proof of Theorem 2

By [14, Theorem 1] a positive solution u to (3) is radially symmetric. With a little abuse of notation we shall write $u(x) = u(r)$ for $x \in B_1$ with $|x| = r$. Since $\Delta u \leq 0$

$$2\pi u'(r) = \int_{B_r} \Delta u dx < 0; \quad r > 0,$$

hence $u(r)$ is decreasing.

Consider now a sequence u_k as in the statement of the theorem. By elliptic estimates, if $\max_{B_1} u_k \leq C$, then we are in case (i). Let us therefore assume that, up to a subsequence

$$\mu_k := u_k(0) = \max_{B_1} u_k \rightarrow \infty \quad \text{as } k \rightarrow \infty. \quad (10)$$

Lemma 3 *Let $r_k > 0$ be such that $r_k^2 \lambda_k \mu_k^2 e^{\mu_k^2} = 4$. Then as $k \rightarrow \infty$ we have $r_k \rightarrow 0$,*

$$\eta_k(x) := \mu_k(u_k(r_k x) - \mu_k) \rightarrow \eta_0(x) := -\log(1 + |x|^2) \quad \text{in } C_{\text{loc}}^1(\mathbb{R}^2), \quad (11)$$

and

$$\lim_{R \rightarrow \infty} \lim_{k \rightarrow \infty} \int_{B_{Rr_k}} \lambda_k u_k^2 e^{u_k^2} dx = \int_{\mathbb{R}^2} 4e^{2\eta_0} dx = 4\pi. \quad (12)$$

Proof. We first prove that $\lim_{k \rightarrow \infty} r_k = 0$. Otherwise up to extracting a subsequence we have $\lambda_k \mu_k^2 e^{\mu_k^2} \leq C$. Then, using that $u'_k \leq 0$ in $[0, 1]$ we see that

$$-\mu_k \Delta u_k \leq \lambda_k \mu_k^2 e^{\mu_k^2} \leq C \quad \text{in } B_1.$$

Therefore as $k \rightarrow \infty$ we get $\Delta u_k \rightarrow 0$ uniformly and by elliptic estimates $u_k \rightarrow 0$ in $C^1(\overline{B}_1)$, contradicting (10).

Set now $v_k(x) = u_k(r_k x) - \mu_k$. We claim that $v_k \rightarrow 0$ in $C^1_{\text{loc}}(\mathbb{R}^2)$ as $k \rightarrow \infty$. Indeed

$$-\Delta v_k(x) = \frac{4}{\mu_k} \frac{u_k(r_k x)}{\mu_k} e^{(u_k(r_k x) - \mu_k)(u_k(r_k x) + \mu_k)} \rightarrow 0 \quad \text{uniformly as } k \rightarrow \infty$$

since $4/\mu_k \rightarrow 0$, $0 \leq u_k/\mu_k \leq 1$ and $(u_k(r_k x) - \mu_k)(u_k(r_k x) + \mu_k) \leq 0$. Now notice that $v_k \leq 0$ and $v_k(0) = 0$. Then the Harnack inequality implies the claim.

Therefore we have

$$-\Delta \eta_k = V_k e^{2a_k \eta_k} \quad \text{in } B_{1/r_k},$$

where

$$V_k(x) = \frac{4u_k(r_k x)}{\mu_k} \rightarrow 4, \quad a_k = \frac{1}{2} \left(\frac{u_k(r_k x)}{\mu_k} + 1 \right) \rightarrow 1 \quad \text{in } C^0_{\text{loc}}(\mathbb{R}^2).$$

Considering that $\eta_k \leq 0$, $\Delta \eta_k$ is locally bounded and $\eta_k(0) = 0$ we have $\eta_k \rightarrow \eta^*$ in $C^1_{\text{loc}}(\mathbb{R}^2)$ by the Harnack inequality, where $-\Delta \eta^* = 4e^{2\eta^*}$ and $\eta^*(0) = 0$. On the other hand

$$-\Delta \eta_0 = 4e^{2\eta_0} \quad \text{in } \mathbb{R}^2, \quad \eta_0(0) = 0, \quad (13)$$

hence it follows from the uniqueness of solutions to the Cauchy problem (recall that all functions here are radially symmetric) that $\eta^* = \eta_0$.

Finally (12) follows from Fatou's lemma. \square

Notice that

$$-\Delta \eta_k = 4 \left(1 + \frac{\eta_k}{\mu_k^2} \right) e^{\left(2 + \frac{\eta_k}{\mu_k^2} \right) \eta_k}. \quad (14)$$

Lemma 4 *Set $w_k := \mu_k^2(\eta_k - \eta_0)$. Then we have $w_k \rightarrow w$ in $C^1_{\text{loc}}(\mathbb{R}^2)$, where*

$$w(r) := \eta_0(r) + \frac{2r^2}{1+r^2} - \frac{1}{2} \eta_0^2(r) + \frac{1-r^2}{1+r^2} \int_1^{1+r^2} \frac{\log t}{1-t} dt \quad (15)$$

is the unique solution to the ODE

$$-\Delta w = 4e^{2\eta_0}(\eta_0 + \eta_0^2 + 2w), \quad w(0) = 0, \quad w'(0) = 0. \quad (16)$$

Moreover w satisfies

$$\int_{\mathbb{R}^2} \Delta w dx = -4\pi, \quad (17)$$

and

$$\sup_{r \in [0, \infty)} |w(r) - \eta_0(r)| < \infty. \quad (18)$$

Proof. Set $\varepsilon_k := \mu_k^{-2} \rightarrow 0$ as $k \rightarrow \infty$. Using (13) and (14) we compute

$$\begin{aligned} -\Delta w_k &= \frac{1}{\varepsilon_k} \left[4(1 + \varepsilon_k \eta_k) e^{(2 + \varepsilon_k \eta_k) \eta_k} - 4e^{2\eta_0} \right] \\ &= \frac{4e^{2\eta_0}}{\varepsilon_k} \left[(1 + \varepsilon_k(\eta_0 + (\eta_k - \eta_0))) e^{2(\eta_k - \eta_0) + \varepsilon_k \eta_0^2 + 2\varepsilon_k \eta_0(\eta_k - \eta_0) + \varepsilon_k(\eta_k - \eta_0)^2} - 1 \right]. \end{aligned} \quad (19)$$

By Lemma 3 for every $R > 0$ we have $\eta_k(r) - \eta_0(r) = o(1) \rightarrow 0$ as $k \rightarrow \infty$ uniformly for $r \in [0, R]$, and we can use a Taylor expansion:

$$e^{2(\eta_k - \eta_0) + \varepsilon_k \eta_0^2 + 2\varepsilon_k \eta_0(\eta_k - \eta_0) + \varepsilon_k(\eta_k - \eta_0)^2} = 1 + 2(\eta_k - \eta_0) + \varepsilon_k \eta_0^2 + o(1)\varepsilon_k + o(1)(\eta_k - \eta_0),$$

with errors $o(1) \rightarrow 0$ as $k \rightarrow \infty$ uniformly for $r \in [0, R]$. Going back to (19) we get

$$-\Delta w_k = 4e^{2\eta_0} [\eta_0 + \eta_0^2 + 2w_k + o(1) + o(1)w_k],$$

with $o(1) \rightarrow 0$ as $k \rightarrow \infty$ uniformly for $r \in [0, R]$. By ODE theory $w_k(r)$ is locally bounded, and by elliptic estimates $w_k \rightarrow \tilde{w}$ in $C_{\text{loc}}^1(\mathbb{R}^2)$, where \tilde{w} satisfies (16).

Since the solution to the Cauchy problem (16) is unique, in order to prove that $\tilde{w} = w$ (with w given in (15)) it is enough to show that w solves (16). It is easily seen that $w(0) = 0$. First computing

$$\left(-\frac{1}{2}\eta_0^2(r) \right)' + \frac{1-r^2}{1+r^2} \frac{d}{dr} \int_1^{1+r^2} \frac{\log t}{1-t} dt = -\frac{2 \log(1+r^2)}{r(1+r^2)},$$

we get

$$w'(r) = \frac{2r(1-r^2)}{(1+r^2)^2} - \frac{2 \log(1+r^2)}{r(1+r^2)} - \frac{4r}{(1+r^2)^2} \int_1^{1+r^2} \frac{\log t}{1-t} dt, \quad (20)$$

$w'(0) = 0$, and using $\Delta w(r) = w''(r) + \frac{w'(r)}{r}$ we finally get

$$\begin{aligned} -\Delta w(r) &= \frac{16r^2}{(1+r^2)^3} - \frac{12 \log(1+r^2)}{(1+r^2)^2} + \frac{8(1-r^2)}{(1+r^2)^3} \int_1^{1+r^2} \frac{\log t}{1-t} dt \\ &= 4e^{2\eta_0} \left[\frac{4r^2}{1+r^2} + 3\eta_0 + 2\frac{1-r^2}{1+r^2} \int_1^{1+r^2} \frac{\log t}{1-t} dt \right] = 4e^{2\eta_0} [\eta_0 + \eta_0^2 + 2w]. \end{aligned}$$

To prove (17) we use the divergence theorem and (20) to get

$$\int_{\mathbb{R}^2} \Delta w dx = \lim_{r \rightarrow \infty} 2\pi r w'(r) = -4\pi.$$

Similarly from (20) we bound

$$|w'(r) - \eta_0'(r)| \leq \frac{C}{1+r^2} \quad \text{for } r \in [0, \infty),$$

and integrating in r also (18) follows. \square

Lemma 3 tells us that for $R > 0$ and $k \geq k_0(R)$ we have $\eta_k \rightarrow \eta_0$ in $C^1(B_R)$. On the other hand η_k is defined on $B_{r_k^{-1}}$ with $r_k^{-1} \rightarrow \infty$ as $k \rightarrow \infty$, and Lemma 3 gives us no information on the behavior of η_k on $B_{r_k^{-1}} \setminus B_R$. We shall now use Lemma 4 to fill this gap and have a crucial estimate of η_k in all of $B_{r_k^{-1}}$.

Lemma 5 Fix $R_0 \in (0, \infty)$ such that $w \leq -1$ on $[R_0, \infty)$, where w is given by (15) and such R_0 exists thanks to (18). Then for k large enough

$$\eta_k(r) \leq \eta_0(r), \quad \text{for } r \in [R_0, r_k^{-1}], \quad (21)$$

or equivalently

$$u_k(r) \leq \mu_k - \frac{1}{\mu_k} \log \left(1 + \left(\frac{r}{r_k} \right)^2 \right), \quad \text{for } r \in [R_0 r_k, 1]. \quad (22)$$

Proof. Write $\varepsilon_k := \mu_k^{-2} = u_k^{-2}(0)$ and $\eta_k = \eta_0 + \varepsilon_k w + \phi_k$. Then (14) is equivalent to

$$-\Delta \eta_k = 4(1 + \varepsilon_k(\eta_0 + \varepsilon_k w + \phi_k)) e^{(2+\varepsilon_k(\eta_0 + \varepsilon_k w + \phi_k))(\eta_0 + \varepsilon_k w + \phi_k)},$$

and taking (13) and (16) into account we find

$$-\Delta \phi_k = \Phi_k(\phi_k),$$

where for any function ϕ

$$\Phi_k(\phi) := 4(1 + \varepsilon_k(\eta_0 + \varepsilon_k w + \phi)) e^{(2+\varepsilon_k(\eta_0 + \varepsilon_k w + \phi))(\eta_0 + \varepsilon_k w + \phi)} - 4e^{2\eta_0} - 4e^{2\eta_0} \varepsilon_k [\eta_0 + \eta_0^2 + 2w].$$

We now expand

$$\begin{aligned} (2 + \varepsilon_k(\eta_0 + \varepsilon_k w + \phi))(\eta_0 + \varepsilon_k w + \phi) &= 2\eta_0 + 2\varepsilon_k w + 2\phi + \varepsilon_k \eta_0^2 + 2\varepsilon_k^2 w \eta_0 + \varepsilon_k^3 w^2 \\ &\quad + \varepsilon_k \phi (2\eta_0 + 2\varepsilon_k w + \phi) =: 2\eta_0 + h_k(\phi). \end{aligned}$$

To avoid cumbersome notations we will also write, for a given function ϕ and any given k ,

$$\begin{aligned} h &:= h_k(\phi) = 2\varepsilon_k w + 2\phi + \varepsilon_k \eta_0^2 + 2\varepsilon_k^2 w \eta_0 + \varepsilon_k^3 w^2 + \varepsilon_k \phi (2\eta_0 + 2\varepsilon_k w + \phi), \\ \eta &:= \eta_0 + \varepsilon_k w + \phi \end{aligned}$$

so that

$$e^{(2+\varepsilon_k \eta)\eta} = e^{2\eta_0 + h}, \quad \Phi_k(\phi) = 4e^{2\eta_0} \left[(1 + \varepsilon_k \eta) e^h - 1 - \varepsilon_k \eta_0 - \varepsilon_k \eta_0^2 - 2\varepsilon_k w \right].$$

Then with a Taylor expansion we can write

$$e^{(2+\varepsilon_k \eta)\eta} = e^{2\eta_0} [1 + h + O(h^2)],$$

where $|O(h^2)| \leq Ch^2$ for a fixed positive constant C , provided $|h| \leq 1$. Then, using (18) to bound $|w(r)| \leq C(1 + \log(1 + r^2))$,

$$\begin{aligned} (1 + \varepsilon_k \eta) e^h &= 1 + \varepsilon_k \eta_0 + 2\varepsilon_k w + \varepsilon_k \eta_0^2 + 2\phi \\ &\quad + O(\phi) (O(\phi) + O(\varepsilon_k(1 + \log(1 + r^2))^2)) + O(\varepsilon_k^2(1 + \log(1 + r^2))^3) \end{aligned}$$

where $|O(s)| \leq Cs$. Then

$$\Phi_k(\phi) = 4e^{2\eta_0} [2\phi + O(\phi) (O(\phi) + O(\varepsilon_k(1 + \log(1 + r^2))^2)) + O(\varepsilon_k^2(1 + \log(1 + r^2))^3)], \quad (23)$$

as long as $|h| \leq 1$, which is true provided for some $\delta > 0$ small enough

$$|\phi| \leq \delta; \quad \varepsilon_k(1 + \log(1 + r^2))^2 \leq \delta. \quad (24)$$

Similarly if $\tilde{\phi}$ is another function with $|\tilde{\phi}(r)| \leq \delta$ one has

$$|\Phi_k(\phi) - \Phi_k(\tilde{\phi})| \leq 4e^{2\eta_0} \left[2|\phi - \tilde{\phi}| + O(\phi - \tilde{\phi}) \left(|\phi + \tilde{\phi}| + O(\varepsilon_k(1 + \log(1 + r^2))^2) \right) \right]. \quad (25)$$

We shall now use the contraction mapping theorem to bound ϕ_k . We restrict our attention to an interval $[0, s_k]$ with $s_k = o(1)e^{\mu k}$ and to functions $\phi : [0, s_k] \rightarrow \mathbb{R}$ satisfying $\phi(r) \leq O(\varepsilon_k^2)(1 + \log(1 + r^2))$, so that (23) and (24) hold for k large enough. With these restrictions (25) gives

$$|\Phi_k(\phi) - \Phi_k(\tilde{\phi})| \leq (8 + o(1))e^{2\eta_0} |\phi - \tilde{\phi}|, \quad (26)$$

with error $o(1) \rightarrow 0$ as $k \rightarrow \infty$.

By the above computations, $\eta_k = \eta_0 + \varepsilon w + \phi_k$ solves (14) if and only if ϕ_k satisfies

$$-\Delta \phi_k = \Phi_k(\phi_k); \quad \phi_k(0) = 0, \quad \phi_k'(0) = 0.$$

Setting $\phi = \phi_k$ and $\psi = r\phi'$, the last equation gives the system

$$\begin{cases} \phi' &= \frac{1}{r}\psi, \\ \psi' &= -r\Phi_k(\phi); \end{cases} \quad (\phi(0), \psi(0)) = (0, 0). \quad (27)$$

The solutions of (27) are the fixed point of some integral equation. For technical reasons, it will be convenient to integrate starting from some value $T > 0$ (to be fixed later) of the r parameter rather than from $r = 0$. If we let (27) evolve up to time T , by the smooth dependence on initial data then (for ε_k small) the solution will satisfy

$$|\phi(r)| \leq C(T)\varepsilon_k^2, \quad |\psi(r)| \leq C(T)\varepsilon_k^2, \quad \text{for } r \in [0, T], \quad (28)$$

uniformly in ε_k . Notice that $\phi(T) = \phi_k(T)$ and $\psi(T) = T\phi_k'(T)$.

We consider then the functions

$$F_{1,(\phi,\psi)}(r) := \phi(T) + \int_T^r \psi(s) \frac{ds}{s}, \quad r \geq T;$$

$$F_{2,(\phi,\psi)}(r) := \psi(T) - \int_T^r s\Phi_k(\phi)(s)ds, \quad r \geq T.$$

Fixing $S = s_k > T$, with $s_k = o(1)e^{\mu k}$, we next define the norms

$$\|f\|_1 = \sup_{r \in (T, S]} \left| \frac{f(r)}{\log r - \log T} \right|; \quad \|f\|_2 = 2 \sup_{r \in [T, S]} |f(r)|.$$

For a large constant $\tilde{C} > 0$ to be fixed later, we will work with the following set of functions

$$\mathcal{B}_{\tilde{C}} = \left\{ (\phi, \psi) : \|\phi - \phi_k(T)\|_1 \leq \tilde{C}\varepsilon_k^2, \|\psi\|_2 \leq \tilde{C}\varepsilon_k^2, \phi(T) = \phi_k(T), \psi(T) = T\phi_k'(T) \right\}.$$

We now check next that the map $(\phi, \psi) \mapsto (F_{1,(\phi,\psi)}, F_{2,(\phi,\psi)})$ sends $\mathcal{B}_{\tilde{C}}$ in itself, for suitable choices of \tilde{C} and T , and that it is a contraction. In fact, for $(\phi, \psi) \in \mathcal{B}_{\tilde{C}}$ one has that

$$|F_{1,(\phi,\psi)}(r) - \phi(T)| \leq \frac{1}{2} \tilde{C} \varepsilon_k^2 (\log r - \log T),$$

which implies $\|F_{1,(\phi,\psi)} - \phi(T)\|_1 \leq \frac{1}{2} \tilde{C} \varepsilon_k^2$, as desired.

Moreover by (23) and (28) one has that

$$\begin{aligned} |F_{2,(\phi,\psi)}(r)| &\leq |\psi(T)| + \int_T^r s 4e^{2\eta_0(s)} [2\phi(s) + O(\phi(s))(O(\phi(s)) + O(\varepsilon_k(1 + \log(1 + s^2))^2)) \\ &\quad + O(\varepsilon_k^2(1 + \log(1 + s^2))^3)] ds \\ &\leq C(T) \varepsilon_k^2 + 8 \int_T^\infty \frac{s(\phi(s)(1 + o(1)))}{(1 + s^2)^2} ds + \int_T^\infty \frac{C_0 \varepsilon_k^2 s(1 + \log(1 + s^2))^3}{(1 + s^2)^2} ds \\ &\leq C(T) \varepsilon_k^2 + 9 \int_T^\infty \frac{s \varepsilon_k^2 (C(T) + \tilde{C} \log s)}{(1 + s^2)^2} ds + C_0 \varepsilon_k^2 \int_T^\infty \frac{s(1 + \log(1 + s^2))^3}{(1 + s^2)^2} ds \\ &\leq \varepsilon_k^2 \left[C(T) \left(1 + 9 \int_T^\infty \frac{s}{(1 + s^2)^2} ds \right) \right. \\ &\quad \left. + 9 \tilde{C} \int_T^\infty \frac{s \log s}{(1 + s^2)^2} ds + C_0 \int_T^\infty \frac{s(1 + \log(1 + s^2))^3}{(1 + s^2)^2} ds \right] \end{aligned}$$

for some fixed C_0 independent of \tilde{C} and ε_k . Now first choosing $T \geq 1$ so large that

$$9 \int_T^\infty \frac{s \log s ds}{(1 + s^2)^2} < \frac{1}{2}, \quad (29)$$

and then \tilde{C} large enough compared to $C(T)$ and C_0 , we obtain

$$\|F_{2,(\phi,\psi)}\|_2 \leq \tilde{C} \varepsilon_k^2,$$

so we are done showing that $(F_{1,(\cdot,\cdot)}, F_{2,(\cdot,\cdot)})$ maps $\mathcal{B}_{\tilde{C}}$ in itself.

Let us verify that F is a contraction. We easily estimate for $(\phi, \psi), (\tilde{\phi}, \tilde{\psi}) \in \mathcal{B}_{\tilde{C}}$

$$\|F_{1,(\phi,\psi)} - F_{1,(\tilde{\phi},\tilde{\psi})}\|_1 \leq \frac{1}{2} \|\psi - \tilde{\psi}\|_2.$$

Using (26) and (29) we also find for k large enough

$$\begin{aligned} \|F_{2,(\phi,\psi)} - F_{2,(\tilde{\phi},\tilde{\psi})}\|_2 &\leq 9 \int_T^S \frac{s|\phi(s) - \tilde{\phi}(s)|}{(1 + s^2)^2} ds \\ &\leq 9 \|\phi - \tilde{\phi}\|_1 \int_T^S \frac{s(\log s - \log T)}{(1 + s^2)^2} ds \leq \frac{1}{2} \|\phi - \tilde{\phi}\|_1, \end{aligned}$$

so we have that indeed F is a contraction. In particular the map $(\phi, \psi) \mapsto (F_{1,(\phi,\psi)}, F_{2,(\phi,\psi)})$ has a fixed point in $(\bar{\phi}, \bar{\psi}) \in \mathcal{B}_{\tilde{C}}$, which satisfies (27). Then, by uniqueness for the Cauchy problem, we have $(\bar{\phi}(r), \bar{\psi}(r)) = (\phi_k(r), r\phi'_k(r))$ for $r \in [T, S]$, whence the bounds

$$\phi_k(r) \leq C(T)\varepsilon_k^2 + \tilde{C}\varepsilon_k^2(\log r - \log T), \quad \phi'_k(r) \leq \frac{\tilde{C}\varepsilon_k^2}{2r}, \quad \text{for } T \leq r \leq S = o(1)e^{\mu_k}. \quad (30)$$

For every k large enough fix now $S = s_k = o(1)e^{\mu_k}$ such that $s_k \geq 2\mu_k$. From (28), (30) and our choice of R_0 , we get for k large enough

$$\eta_k(r) \leq \eta_0(r) - \varepsilon_k + (C(T) + \tilde{C} \log r)\varepsilon_k^2 < \eta_0(r), \quad \text{for } r \in [R_0, s_k]. \quad (31)$$

We shall now prove that

$$\int_{B_{s_k}} \Delta \eta_k dx < -4\pi$$

for k large enough. Indeed we have

$$-\int_{B_{s_k}} \Delta \eta_0 dx = \int_{B_{s_k}} \frac{4}{(1+r^2)^2} dx = 4\pi \left(1 - \frac{1}{1+s_k^2}\right) = 4\pi - \frac{4\pi}{s_k^2} + \frac{o(1)}{s_k^2}.$$

From (17) we have

$$-\int_{B_{s_k}} \varepsilon_k \Delta w dx = 4\pi(1+o(1))\varepsilon_k = \frac{4\pi(1+o(1))}{\mu_k^2}.$$

Finally, using (30) and the divergence theorem,

$$\left| \int_{B_{s_k}} \Delta \phi_k dx \right| = 2\pi s_k |\phi'_k(s_k)| = O(\varepsilon_k^2) = (\mu_k^{-4}).$$

Summing up we infer

$$-\int_{B_{s_k}} \Delta \eta_k dx = 4\pi - \frac{4\pi(1+o(1))}{s_k^2} + \frac{4\pi(1+o(1))}{\mu_k^2} + O(\mu_k^{-4}) > 4\pi$$

for k large enough, by our choice of s_k . Since $\Delta \eta_k < 0$ on all of B_{1/r_k} , we infer that

$$2\pi r \eta'_k(r) = \int_{B_r} \Delta \eta_k dx < \int_{B_{s_k}} \Delta \eta_k dx < -4\pi < \int_{B_r} \Delta \eta_0 dx = 2\pi r \eta'_0(r), \quad r \in [s_k, 1/r_k].$$

This, together with (31), completes the proof of (21). \square

Proof of Theorem 2 (completed). From (22) it follows that $\lambda_k \rightarrow 0$ as $k \rightarrow \infty$. Indeed the function $-\frac{2}{\mu_k} \log\left(\frac{r}{r_k}\right) + \mu_k$ vanishes for $r = \rho_k = \frac{2}{\sqrt{\lambda_k \mu_k}}$. Since $u_k > 0$ in B_1 we must have for k large enough $\rho_k \geq 1$, hence

$$\lambda_k \leq \frac{4}{\mu_k^2} \rightarrow 0 \quad \text{as } k \rightarrow \infty. \quad (32)$$

Set $f_k := \lambda_k u_k^2 e^{u_k^2}$. We now want to prove

$$\lim_{R \rightarrow \infty} \lim_{k \rightarrow \infty} \int_{B_1 \setminus B_{Rr_k}} f_k dx = 0, \quad (33)$$

which together with (12), yields the convergence of $f_k dx$ in (7).

Using the definition of r_k and (22) we have for $r \in [R_0 r_k, 1]$ and k large enough

$$\begin{aligned} r^2 f_k(r) &= 4 \left(\frac{r}{r_k} \right)^2 \left(\frac{u_k(r)}{\mu_k} \right)^2 e^{(1+u_k(r)/\mu_k)\eta_k(r/r_k)} \\ &\leq 4 \left(\frac{r}{r_k} \right)^2 \left(\frac{u_k(r)}{\mu_k} \right)^2 \left(1 + \left(\frac{r}{r_k} \right)^2 \right)^{-1 - \frac{u_k(r)}{\mu_k}} \\ &\leq 4 \left(\frac{u_k(r)}{\mu_k} \right)^2 \left(1 + \left(\frac{r}{r_k} \right)^2 \right)^{-\frac{u_k(r)}{\mu_k}}. \end{aligned}$$

In order to further estimate the right-hand side, we notice that the function

$$\alpha_k(t) := t^2 \left(1 + \left(\frac{r}{r_k} \right)^2 \right)^{-t}$$

satisfies for any fixed $r > 0$

$$\alpha_k \geq 0, \quad \alpha_k(0) = 0, \quad \lim_{t \rightarrow \infty} \alpha_k(t) = 0 \quad \alpha'_k(t) = 0 \Leftrightarrow t = t_k := \frac{2}{\log \left(1 + \left(\frac{r}{r_k} \right)^2 \right)}.$$

Hence $\alpha_k \leq \alpha_k(t_k)$ and we conclude

$$r^2 f_k(r) \leq \frac{16}{\log^2 \left(1 + \left(\frac{r}{r_k} \right)^2 \right)} \left(1 + \left(\frac{r}{r_k} \right)^2 \right)^{-2 \log^{-1} \left(1 + \left(\frac{r}{r_k} \right)^2 \right)}, \quad \text{for } r \geq R_0 r_k, \text{ } k \text{ large,}$$

which can be weakened to

$$\log^2 \left(\frac{r}{r_k} \right)^2 r^2 f_k(r) \leq \log^2 \left(1 + \left(\frac{r}{r_k} \right)^2 \right) r^2 f_k(r) \leq 16, \quad \text{for } r \geq R_0 r_k, \text{ } k \text{ large.} \quad (34)$$

Finally, it follows from (34) that

$$\begin{aligned} \lim_{R \rightarrow \infty} \lim_{k \rightarrow \infty} \int_{B_1 \setminus B_{Rr_k}} f_k dx &\leq \lim_{R \rightarrow \infty} \lim_{k \rightarrow \infty} \int_{Rr_k}^1 2\pi r f_k(r) dr \leq \lim_{R \rightarrow \infty} \lim_{k \rightarrow \infty} \int_{Rr_k}^1 \frac{32\pi}{r [\log(r/r_k)]^2} dr \\ &= \lim_{R \rightarrow \infty} \lim_{k \rightarrow \infty} \left[-\frac{32\pi}{\log(r/r_k)} \right]_{Rr_k}^1 = 0. \end{aligned} \quad (35)$$

Then (33) is proven, and as already noticed the first part of (7) follows.

Integrating by parts we also obtain

$$\|u_k\|_{H_0^1}^2 = \int_{B_1} u_k(-\Delta u_k) dx = \int_{B_1} f_k dx \rightarrow 4\pi \quad \text{as } k \rightarrow \infty.$$

Then, up to extracting a subsequence we have $u_k \rightharpoonup u_\infty$ weakly in $H_0^1(B_1)$ for some function $u_\infty \in H_0^1(B_1)$. Moreover, using that $\lambda_k \rightarrow 0$ as $k \rightarrow \infty$, we get for any $L > 0$

$$\int_{B_1} \lambda_k u_k e^{u_k^2} dx \leq \lambda_k \int_{\{x \in B_1 : u_k(x) \leq L\}} u_k e^{u_k^2} dx + \frac{1}{L} \int_{\{x \in B_1 : u_k(x) < L\}} f_k dx \leq o(1)C(L) + \frac{4\pi + o(1)}{L},$$

with error $o(1) \rightarrow 0$ as $k \rightarrow \infty$. Then, by letting $L \rightarrow \infty$ we infer that $\lambda_k u_k e^{u_k^2} \rightarrow 0$ in $L^1(B_1)$, and it follows that for every $\varphi \in C_c^1(B_1)$

$$\int_{B_1} (-\Delta u_\infty) \varphi dx = \lim_{k \rightarrow \infty} \int_{B_1} (-\Delta u_k) \varphi dx = \lim_{k \rightarrow \infty} \int_{B_1} -\lambda_k u_k e^{u_k^2} \varphi dx = 0,$$

hence $-\Delta u_\infty = 0$ in B_1 , i.e. $u_\infty \equiv 0$.

This also implies the convergence of $|\nabla u_k|^2 dx$ in (7). Indeed, integrating by parts and using that $u_k \rightarrow 0$ in $L^2(B_1)$ by the compactness of the embedding $H_0^1(B_1) \hookrightarrow L^2(B_1)$, we have for any $\varphi \in C_c^2(B_1)$

$$\int_{B_1} |\nabla u_k|^2 \varphi dx = \int_{B_1} \frac{\Delta(u_k^2)}{2} \varphi dx + \int_{B_1} f_k \varphi dx = \int_{B_1} \frac{u_k^2}{2} \Delta \varphi dx + \int_{B_1} f_k \varphi dx = \int_{B_1} f_k \varphi dx + o(1),$$

with error $o(1) \rightarrow 0$ as $k \rightarrow \infty$. Hence $f_k dx$ and $|\nabla u_k|^2 dx$ have the same weak limit in the sense of measures.

Now, using $u_k(1) = 0$ and (7) we infer that $u_k \rightarrow 0$ in $L_{\text{loc}}^\infty(B_1 \setminus \{0\})$. Indeed for a fixed $\delta \in (0, 1)$ and any $r \in [\delta, 1]$ we have by Hölder's inequality

$$u_k(r) = u_k(r) - u_k(1) \leq \int_\delta^1 |u_k'(\rho)| d\rho \leq \|\nabla u_k\|_{L^2(B_1 \setminus B_\delta)} \left(\frac{1}{2\pi} \log \frac{1}{\delta} \right)^{\frac{1}{2}} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Then by elliptic estimates $u_k \rightarrow 0$ in $C_{\text{loc}}^1(\bar{B}_1 \setminus \{0\})$. This completes the proof of Theorem 2. \square

3 Proof of Theorem 1

Given $\mu, \lambda > 0$ let $\bar{u}_{\mu, \lambda} \in C^\infty([0, T_{\mu, \lambda}))$ be the solution to the ODE

$$-\frac{\partial^2 \bar{u}}{\partial r^2} - \frac{1}{r} \frac{\partial \bar{u}}{\partial r} = \lambda \bar{u} e^{\bar{u}^2}, \quad \bar{u}(0) = \mu, \quad \bar{u}'(0) = 0, \quad (36)$$

where $[0, T_{\mu, \lambda})$ is the maximal interval of existence for (36) (in fact $T_{\mu, \lambda} = \infty$, but we will not prove this). Then $u_{\mu, \lambda}(x) := \bar{u}_{\mu, \lambda}(|x|)$ satisfies

$$-\Delta u_{\mu, \lambda} = \lambda u_{\mu, \lambda} e^{u_{\mu, \lambda}^2} \quad \text{in } B_{T_{\mu, \lambda}}, \quad u_{\mu, \lambda}(0) = \mu.$$

Set

$$\tau(\mu) := \inf\{r \in (0, T_{\mu,1}] : \bar{u}_{\mu,1}(r) = 0\}.$$

We claim that $\tau(\mu) < \infty$ for every $\mu > 0$. To see this, fix $r_0 \in (0, \tau(\mu))$. Then for $r \in [r_0, \tau(\mu)]$ the divergence theorem yields

$$\bar{u}'_{\mu,1}(r) = \frac{1}{2\pi r} \int_{B_r} \Delta u_{\mu,1} dx \leq \frac{1}{2\pi r} \int_{B_{r_0}} \Delta u_{\mu,1} dx < -\frac{\varepsilon}{r},$$

for some positive ε . The claim easily follows from standard comparison arguments.

Now notice that, for any $\lambda, \lambda' > 0$

$$\bar{u}_{\mu,\lambda}\left(\sqrt{\frac{\lambda'}{\lambda}}r\right) = \bar{u}_{\mu,\lambda'}(r),$$

and for every $\mu > 0$ set $\bar{u}_\mu := \bar{u}_{\mu,\tau^2(\mu)} = \bar{u}_{\mu,\lambda_\mu}$, where $\lambda_\mu := \tau^2(\mu)$. Then \bar{u}_μ is positive in $[0, 1)$, it solves (36) with $\lambda = \lambda_\mu$, and $\bar{u}_\mu(1) = 0$. By ODE theory the function $\Phi : \mu \mapsto \bar{u}_\mu|_{[0,1]}$ belongs to $C^0((0, \infty), C^2([0, 1]))$.

Now given $\Lambda > 0$ every non-negative critical point of $E|_{M_\Lambda}$ is smooth and satisfies

$$-\Delta u = \lambda u e^{u^2} \text{ in } B_1, \quad u = 0 \text{ on } \partial B_1, \quad (37)$$

for some $\lambda > 0$. By [14, Theorem 1] we have that u is radially symmetric, i.e. we can write $u(x) = \bar{u}(|x|)$, where \bar{u} satisfies (36) with $\mu = u(0)$ and the additional condition that $\bar{u} > 0$ on $[0, 1)$ and $\bar{u}(1) = 0$. This is possible only if $\bar{u} = \bar{u}_\mu$. Hence we have proven that every solution $0 \leq u \not\equiv 0$ of (37) is of the form $u(x) = u_\mu(x) := \bar{u}_\mu(|x|)$ for some $\mu > 0$. Define

$$\mathcal{E}(\mu) := \|u_\mu\|_{H_0^1(B_1)}^2, \quad \Lambda^\sharp := \sup_{\mu \in (0, \infty)} \mathcal{E}(\mu).$$

We claim that $\Lambda^\sharp < \infty$. Indeed \mathcal{E} is continuous and it is clear that $u_\mu \rightarrow 0$ smoothly as $\mu \downarrow 0$, hence $\lim_{\mu \downarrow 0} \mathcal{E}(\mu) = 0$. Moreover Theorem 2 gives $\lim_{\mu \rightarrow \infty} \mathcal{E}(\mu) = 4\pi$, hence by continuity $\Lambda^\sharp < \infty$. This completes the proof of Part (i) of the theorem. By Theorem 1.7 in [26] it follows that $\Lambda^\sharp > 4\pi$, and Parts (ii) and (iii) follow at once from the continuity of \mathcal{E} . \square

References

- [1] ADIMURTHI, *Existence of positive solutions of the semilinear Dirichlet problem with critical growth for the n -Laplacian*, Ann. Sc. Norm. Sup. Pisa, Ser. IV **17** (1990), 393-413.
- [2] ADIMURTHI, O. DRUET, *Blow-up analysis in dimension 2 and a sharp form of Trudinger-Moser inequality*, Comm. Partial Differential Equations **29** (2004), 295-322.
- [3] ADIMURTHI, S. PRASHANTH, *Failure of Palais-Smale condition and blow-up analysis for the critical exponent problem in \mathbb{R}^2* , Proc. Indian Acad. Sci. Math. Sci. **107** (1997), 283-317.
- [4] ADIMURTHI, M. STRUWE, *Global compactness properties of semilinear elliptic equations with critical exponential growth*, J. Funct. Anal. **175** (2000), 125-167.

- [5] H. BRÉZIS, F. MERLE *Uniform estimates and blow-up behaviour for solutions of $-\Delta u = V(x)e^u$ in two dimensions*, Comm. Partial Differential Equations **16** (1991), 1223-1253.
- [6] H. BRÉZIS, Y.-Y. LI, I. SHAFRIR, *A sup + inf inequality for some nonlinear elliptic equations involving exponential nonlinearities*, J. Funct. Anal. **115** (1993), 344-358.
- [7] L. CARLESON, S.-Y. A. CHANG, *On the existence of an extremal function for an inequality of J. Moser*, Bull. Sci. Math. (2) **110** (1986) 113-127.
- [8] J.-M. CORON, *Topologie et cas limite des injections de Sobolev*, C.R. Math. Acad. Sci. Paris **299** (I) (1984) 209-212.
- [9] D.G. DE FIGUEIREDO, J.M. DO O, B. RUF, *On an inequality by N. Trudinger and J. Moser and related elliptic equations*, Comm. Pure Appl. Math. **55** (2002), 135-152.
- [10] M. DEL PINO, M. MUSSO, B. RUF, *New solutions for Trudinger-Moser critical equations in \mathbb{R}^2* , J. Funct. Analysis **258** (2010), 421-457.
- [11] M. DEL PINO, M. MUSSO, B. RUF, *Beyond the Trudinger-Moser Supremum*, Calc. Var. and PDEs, to appear.
- [12] O. DRUET, *Multibumps analysis in dimension 2, quantification of blow-up levels*, Duke Math. J. **132** (2006), 217-269.
- [13] M. FLUCHER, *Extremal functions for the Trudinger-Moser inequality in 2 dimensions*, Comment. Math. Helv. **67** (3) (1992) 471-497.
- [14] B. GIDAS, W.-M. NI, L. NIRENBERG, *Symmetry and related properties via the maximum principle*, Commun. Math. Phys. **68** (1979), 209-243.
- [15] D.-D. JOSEPH, T.-S. LUNDGREN, *Quasilinear Dirichlet problems driven by positive sources*, Arch. Rat. Mech. Anal. **49** (1973), 241-269.
- [16] T. LAMM, F. ROBERT, M. STRUWE, *The heat flow with a critical exponential nonlinearity*, J. Funct. Analysis **257** (2009), 2951-2998.
- [17] Y.-Y. LI, *Harnack type inequality: The method of moving planes*, Commun. Math. Phys. **200**-2, (1999), 421-444.
- [18] Y.-Y. LI, I. SHAFRIR *Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension 2*, Indiana Univ. Math. J. **43** (1994), 1255-1270.
- [19] L. MARTINAZZI *A threshold phenomenon for embeddings of H_0^m into Orlicz spaces*, Calc. Var. Partial Differential Equations **36** (2009), 493-506.
- [20] L. MARTINAZZI, M. STRUWE *Quantization for an elliptic equation of order $2m$ with critical exponential non-linearity*, Math. Z. **270** (2012), 453-487.
- [21] F. MIGNOT, J.-P. PUEL, *Sur une classe de problèmes non linéaires avec nonlinéarité positive, croissante, convexe*, Comm. Part. Diff. Eq. **5** (8) (1980) , 791-836.
- [22] J.-P. MONAHAN, *Numerical solution of a non-linear boundary value problem*, thesis, Princeton Univ., 1971.

- [23] J. MOSER, *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. J. **20** (1971), 1077-1092.
- [24] R. SCHOEN, *On the number of constant scalar curvature metrics in a conformal class*, in "Differential Geometry: A Symposium in Honor of Manfredo Do Carmo", (H.B.Lawson and K.Tenenblat Editors), (1991), 331-320, Wiley, New York.
- [25] I. SHAFRIR, *Une inégalité de type Sup + Inf pour l'équation $-\Delta u = Ve^u$* , C. R. Acad. Sci. Paris **315** (1992), 159-164.
- [26] M. STRUWE, *Critical points of embeddings of $H_0^{1,n}$ into Orlicz spaces*, Ann. Inst. H. Poincaré Anal. Non Linéaire **5** (1984) 425-464.
- [27] M. STRUWE, *Positive solutions of critical semilinear elliptic equations on non-contractible planar domains*, J. Eur. Math. Soc. (JEMS) **2** (4) (2000) 329-388.
- [28] M. STRUWE *Quantization for a fourth order equation with critical exponential growth*, Math. Z. **256** (2007), 397-424.
- [29] N.S. TRUDINGER, *On embedding into Orlicz spaces and some applications*, J. Math. Mech. **17** (1967), 473-483.