

Crum's Theorem for 'Discrete' Quantum Mechanics

Satoru ODAKE¹ and Ryu SASAKI²

¹*Department of Physics, Shinshu University, Matsumoto 390-8621, Japan*

²*Yukawa Institute for Theoretical Physics, Kyoto University,
Kyoto 606-8502, Japan*

In one-dimensional quantum mechanics, or the Sturm-Liouville theory, Crum's theorem describes the relationship between the original and the associated Hamiltonian systems, which are iso-spectral except for the lowest energy state. Its counterpart in 'discrete' quantum mechanics is formulated algebraically, elucidating the basic structure of the discrete quantum mechanics, whose Schrödinger equation is a difference equation.

§1. Introduction

In the seminal paper of 1955, Crum showed,¹⁾ if rephrased in the language of quantum mechanics, the existence of an associated Hamiltonian system for any given one-dimensional quantum mechanical Hamiltonian system under mild assumptions. The method or the technique is quite universal and is known under many different names; the Darboux transformation,²⁾ the factorisation method³⁾ or the supersymmetric quantum mechanics.⁴⁾ Crum himself presented his results in the traditional language of Sturm-Liouville systems. Since Crum's theorem elucidates the generic structure, many exactly and quasi-exactly solvable quantum mechanical examples were constructed by employing Crum's theorem and its modifications combined with shape invariance.⁵⁾ In particular, Adler's modification of Crum's theorem⁶⁾ is quite general and useful. It allows to construct an infinitely many exactly solvable potential from any exactly solvable one.

Recently 'discrete' quantum mechanics was introduced by the present authors.^{7)–9)} It is a generalisation of quantum mechanics, in which the Schrödinger equation is a difference equation instead of differential in the ordinary quantum mechanics. In other words, the differential operator in the Hamiltonian is replaced by finite difference operators, either in the pure imaginary or the real direction. This is why they are called discrete quantum mechanics. Many explicit examples of exactly^{10), 11)} and quasi-exactly solvable^{12), 13)} systems have been constructed. These exactly solvable systems have salient features; they are solvable both in the Schrödinger and the Heisenberg pictures. The main part of the exact eigenfunctions are the known hypergeometric orthogonal polynomials of the Askey scheme.^{14)–16)} On the other hand, the exact Heisenberg operator solutions define the creation and annihilation operators^{10), 11)} which, together with the Hamiltonian, form the dynamical symmetry algebra of the exactly solvable systems. The so-called q -oscillator algebra¹⁷⁾ is the most typical dynamical symmetry algebra realised in this way.¹⁸⁾ In discrete quantum mechanics, as will be shown shortly, the counterpart of Crum's theorem holds and it also elucidates the generic structure of one dimensional systems. As in the ordinary quantum mechanics, Crum's theorem and its modifications are useful for

constructing various exactly and quasi-exactly solvable discrete quantum mechanical examples. In particular, its modification a la Adler¹⁹⁾ will also provide an infinite number of exactly solvable ones based on any exactly solvable one. When applied to discrete quantum mechanics with real shifts, the modified Crum's theorem generates an infinite or finite series of exactly solvable birth and death processes based on any known exactly solvable one.²⁰⁾ The insight obtained Crum's theorems and their modification, in the ordinary and discrete quantum mechanics, is essential for the recent derivation of infinite numbers of shape invariant systems and new exceptional orthogonal polynomials.^{21),22)}

In this paper, we present the discrete quantum mechanics version of Crum's theorem. Under mild assumptions, we construct algebraically an associated Hamiltonian system for any given one-dimensional discrete quantum Hamiltonian system. That is, if a Hamiltonian together with the full discrete energy spectra and the corresponding eigenfunctions are given, then the associated Hamiltonian together with the full discrete energy spectra and the corresponding eigenfunctions are constructed as in the original Crum's paper.¹⁾ This process goes on indefinitely, since the first associated Hamiltonian system generates the second associated Hamiltonian system and so on. Due to the essential distinction between the differential and difference equations, several technical assumptions are necessary for the discrete Crum's theorem. For example, in one-dimensional ordinary quantum mechanics, the energy spectra are non-degenerate and the oscillation theorem holds. That is, the n -th excited state wavefunction has n zeros. These two properties are not necessarily shared by the generic discrete QM. The hermiticity is almost trivial in ordinary QM, but it can only be proven after the explicit form of the groundstate wavefunction is obtained in discrete QM.^{9),12)} Another distinctive feature is the wavefunction. In ordinary QM, the wavefunction is a complex valued real function, defined on the real line, or half line or on a line segment. In one dimension, the wavefunction can be chosen real. In discrete QM with pure imaginary shifts, the wavefunction undergoes shift operations $\psi(x) \rightarrow \psi(x \pm i\gamma)$, $\gamma \in \mathbb{R}$. Thus we require the *analyticity* of the wavefunction with its domain including the real axis or a part of it in which the dynamical variable x is defined. In spite of these differences, the generation of the associated Hamiltonian system goes almost parallel in the ordinary and discrete QM, since the main part is algebraic.

This paper is organised as follows. In section two, Crum's theorem is recapitulated in some detail in the language and notation of ordinary quantum mechanics. This explains the underlying logical structure of the associated Hamiltonian system, which is also shared by the discrete quantum mechanics version. The discrete version of Crum's theorem is stated and proved in section three. The final section is for a summary and comments.

§2. Ordinary Quantum Mechanics

Let us start with a generic one-dimensional quantum mechanical system having discrete semi-infinite energy levels only:

$$0 = \mathcal{E}_0 < \mathcal{E}_1 < \mathcal{E}_2 < \dots . \quad (2.1)$$

Here we have chosen the constant part of the Hamiltonian so that the groundstate energy is zero. Then the Hamiltonian is *positive semi-definite* and in one dimension all the energy levels are non-degenerate. It is well known in linear algebra that any positive semi-definite hermitian matrix can be factorised as a product of a certain matrix, say \mathcal{A} , and its hermitian conjugate \mathcal{A}^\dagger . Similar factorisation applies to the present quantum mechanical Hamiltonian \mathcal{H} , which takes a very simple form:

$$\mathcal{H} = p^2 + U(x) = p^2 + \left(\frac{d\mathcal{W}(x)}{dx} \right)^2 + \frac{d^2\mathcal{W}(x)}{dx^2}, \quad p = -i \frac{d}{dx}, \quad (2.2)$$

$$= \mathcal{A}^\dagger \mathcal{A}, \quad \mathcal{A} \stackrel{\text{def}}{=} \frac{d}{dx} - \frac{d\mathcal{W}(x)}{dx}, \quad \mathcal{A}^\dagger = -\frac{d}{dx} - \frac{d\mathcal{W}(x)}{dx}. \quad (2.3)$$

Here a real function $\mathcal{W}(x) \in \mathbb{C}^\infty$ is called a *pre-potential* and it parametrises the groundstate wavefunction $\phi_0(x)$, which has *no node* and can be chosen real and positive:

$$\phi_0(x) = e^{\mathcal{W}(x)}. \quad (2.4)$$

It is trivial to verify

$$\mathcal{A}\phi_0(x) = 0 \Rightarrow \mathcal{H}\phi_0(x) = 0. \quad (2.5)$$

All the eigenfunctions are square-integrable and orthogonal with each other and form a complete basis of the Hilbert space:

$$\mathcal{H}\phi_n(x) = \mathcal{E}_n\phi_n(x), \quad n = 0, 1, 2, \dots, \quad (2.6)$$

$$\int \phi_n(x)^* \phi_m(x) dx = h_n \delta_{nm}, \quad 0 < h_n < \infty, \quad n, m = 0, 1, 2, \dots \quad (2.7)$$

It is well-known that the n -th excited wavefunction $\phi_n(x)$ has n zeros in the interior. For simplicity we choose all the eigenfunctions to be real. Here are a few examples:

$$\mathcal{W}(x) = -\frac{1}{2}x^2, \quad U(x) = x^2 - 1, \quad -\infty < x < \infty, \quad (2.8)$$

$$\mathcal{W}(x) = -\frac{1}{2}x^2 + g \log x, \quad U(x) = x^2 + \frac{g(g-1)}{x^2} - 1 - 2g, \quad g > 1, \quad 0 < x < \infty, \quad (2.9)$$

$$\mathcal{W}(x) = g \log \sin x, \quad U(x) = \frac{g(g-1)}{\sin^2 x} - g^2, \quad g > 1, \quad 0 < x < \pi. \quad (2.10)$$

They all lead to well-known exactly solvable quantum mechanics whose eigenfunctions consist of the classical orthogonal polynomials, the Hermite, Laguerre and Jacobi polynomials, respectively. In each case, the squared groundstate wavefunction $\phi_0(x)^2$ gives the orthogonality weight function for the polynomials.

Next let us define an associated Hamiltonian $\mathcal{H}^{[1]}$ by simply changing the order of \mathcal{A} and \mathcal{A}^\dagger :

$$\mathcal{H}^{[1]} \stackrel{\text{def}}{=} \mathcal{A}\mathcal{A}^\dagger. \quad (2.11)$$

For later convenience, let us attach the superscript $^{[0]}$ to all the quantities in the original Hamiltonian system, $\mathcal{H}^{[0]} \stackrel{\text{def}}{=} \mathcal{H}$, $\phi_n^{[0]}(x) \stackrel{\text{def}}{=} \phi_n(x)$, $\mathcal{A}^{[0]} \stackrel{\text{def}}{=} \mathcal{A}$, $U^{[0]}(x) \stackrel{\text{def}}{=} U(x)$, $\mathcal{W}^{[0]}(x) \stackrel{\text{def}}{=} \mathcal{W}(x)$. By construction \mathcal{A} and \mathcal{A}^\dagger intertwine $\mathcal{H}^{[0]}$ and $\mathcal{H}^{[1]}$:

$$\mathcal{A}^{[0]}\mathcal{H}^{[0]} = \mathcal{H}^{[1]}\mathcal{A}^{[0]}, \quad \mathcal{A}^{[0]\dagger}\mathcal{H}^{[1]} = \mathcal{H}^{[0]}\mathcal{A}^{[0]\dagger}. \quad (2.12)$$

We will show that the associated Hamiltonian system $\mathcal{H}^{[1]}$ is *iso-spectral* to the original Hamiltonian system $\mathcal{H}^{[0]}$ and the eigenfunctions are in one to one correspondence, *except for* the groundstate. Thanks to the first of the above relation (2.12), it is trivial to verify that the eigenfunctions of the associated Hamiltonian system $\mathcal{H}^{[1]}$ are generated algebraically by multiplying $\mathcal{A}^{[0]}$ to the eigenfunction of the original system:

$$\phi_n^{[1]}(x) \stackrel{\text{def}}{=} \mathcal{A}^{[0]}\phi_n^{[0]}(x), \quad \int \phi_n^{[1]}(x)^* \phi_m^{[1]}(x) dx = \mathcal{E}_n h_n \delta_{nm}, \quad n, m = 1, 2, \dots, \quad (2.13)$$

$$\mathcal{H}^{[1]}\phi_n^{[1]}(x) = \mathcal{E}_n \phi_n^{[1]}(x), \quad n = 1, 2, \dots \quad (2.14)$$

Suppose the associated Hamiltonian $\mathcal{H}^{[1]}$ has an eigenfunction $\phi'(x)$ with the eigenvalue \mathcal{E}' other than those listed above:

$$\mathcal{H}^{[1]}\phi'(x) = \mathcal{E}'\phi'(x). \quad (2.15)$$

Again, thanks to the second of the relation (2.12), it is trivial to verify

$$\mathcal{H}^{[0]}\mathcal{A}^{[0]\dagger}\phi'(x) = \mathcal{E}'\mathcal{A}^{[0]\dagger}\phi'(x). \quad (2.16)$$

Due to the completeness of the spectrum of the original Hamiltonian $\mathcal{H}^{[0]}$, the provisional eigenvalue \mathcal{E}' must belong to the above spectrum (2.14) for $n = 1, 2, \dots$. In other words, \mathcal{E}' cannot be vanishing, $\mathcal{E}' \neq 0$. Suppose that is the case ($\mathcal{E}' = 0$), then ϕ' is annihilated by $\mathcal{A}^{[0]\dagger}$. Surely there exists a solution of a first order differential equation $\mathcal{A}^{[0]\dagger}\phi'(x) = 0$, $\phi'(x) = 1/\phi_0^{[0]}(x) = e^{-\mathcal{W}(x)}$. But it is obviously non square-integrable and it does not belong to the Hilbert space of the associated Hamiltonian system. After Crum,¹⁾ we can show in the following way that $\phi_n^{[1]}(x)$ has exactly $n - 1$ zeros. Since $\phi_n^{[0]}$ has exactly n zeros, the relation

$$\frac{\phi_n^{[1]}(x)}{\phi_0^{[0]}(x)} = \frac{d}{dx} \left(\frac{\phi_n^{[0]}(x)}{\phi_0^{[0]}(x)} \right)$$

tells through Rolle's theorem that $\phi_n^{[1]}$ has at least $n - 1$ zeros. From the relation

$$\frac{d}{dx} \left(\phi_0^{[0]}(x) \phi_n^{[1]}(x) \right) = -\mathcal{E}_n \phi_0^{[0]}(x) \phi_n^{[0]}(x),$$

we find that $\phi_n^{[1]}$ has at most $n - 1$ zeros. Thus we have established that the associated Hamiltonian system $\mathcal{H}^{[1]}$ is *iso-spectral* to the original Hamiltonian system $\mathcal{H}^{[0]}$ and

the eigenfunctions are in one to one correspondence, *except for* the groundstate with the wavefunction $\phi_0^{[0]}(x)$. If the groundstate energy \mathcal{E}_1 is subtracted from the associated Hamiltonian $\mathcal{H}^{[1]}$, it is again positive semi-definite and can be factorised as above:

$$\mathcal{H}^{[1]} = \mathcal{A}^{[1]\dagger} \mathcal{A}^{[1]} + \mathcal{E}_1 = p^2 + U^{[1]}(x) + \mathcal{E}_1, \quad (2.17)$$

$$\mathcal{A}^{[1]} \stackrel{\text{def}}{=} \frac{d}{dx} - \frac{d\mathcal{W}^{[1]}(x)}{dx}, \quad \mathcal{A}^{[1]\dagger} = -\frac{d}{dx} - \frac{d\mathcal{W}^{[1]}(x)}{dx}, \quad (2.18)$$

$$e^{\mathcal{W}^{[1]}(x)} \stackrel{\text{def}}{=} |\phi_1^{[1]}(x)|, \quad U^{[1]}(x) \stackrel{\text{def}}{=} \left(\frac{d\mathcal{W}^{[1]}(x)}{dx} \right)^2 + \frac{d^2\mathcal{W}^{[1]}(x)}{dx^2}, \quad (2.19)$$

$$\mathcal{A}^{[1]}\phi_1^{[1]}(x) = 0. \quad (2.20)$$

As shown above, the groundstate wavefunction $\phi_1^{[1]}(x)$ has no node. Note that (2.11), (2.17) and (2.19) imply the Riccati equation for $d\mathcal{W}^{[1]}(x)/dx$:

$$\left(\frac{d\mathcal{W}^{[1]}(x)}{dx} \right)^2 + \frac{d^2\mathcal{W}^{[1]}(x)}{dx^2} = \left(\frac{d\mathcal{W}(x)}{dx} \right)^2 - \frac{d^2\mathcal{W}(x)}{dx^2} - \mathcal{E}_1. \quad (2.21)$$

Then by reversing the order of $\mathcal{A}^{[1]\dagger}$ and $\mathcal{A}^{[1]}$ the second associated Hamiltonian system $\mathcal{H}^{[2]}$ can be defined. This process can go indefinitely.

Here we list the definition of the s -th quantities step by step for $s \geq 1$,

$$\mathcal{H}^{[s]} \stackrel{\text{def}}{=} \mathcal{A}^{[s-1]}\mathcal{A}^{[s-1]\dagger} + \mathcal{E}_{s-1}, \quad (2.22)$$

$$\phi_n^{[s]}(x) \stackrel{\text{def}}{=} \mathcal{A}^{[s-1]}\phi_n^{[s-1]}(x), \quad (n \geq s), \quad (2.23)$$

$$e^{\mathcal{W}^{[s]}(x)} \stackrel{\text{def}}{=} |\phi_s^{[s]}(x)|, \quad (2.24)$$

$$\mathcal{A}^{[s]} \stackrel{\text{def}}{=} \frac{d}{dx} - \frac{d\mathcal{W}^{[s]}(x)}{dx}, \quad \mathcal{A}^{[s]\dagger} = -\frac{d}{dx} - \frac{d\mathcal{W}^{[s]}(x)}{dx}, \quad (2.25)$$

$$U^{[s]}(x) \stackrel{\text{def}}{=} \left(\frac{d\mathcal{W}^{[s]}(x)}{dx} \right)^2 + \frac{d^2\mathcal{W}^{[s]}(x)}{dx^2}. \quad (2.26)$$

Then we can show the following for $n \geq s \geq 0$,

$$\mathcal{H}^{[s]}\phi_n^{[s]}(x) = \mathcal{E}_n\phi_n^{[s]}(x), \quad (2.27)$$

$$\phi_n^{[s]}(x) : \text{real function}, \quad (2.28)$$

$$\mathcal{A}^{[s]}\phi_s^{[s]}(x) = 0, \quad (2.29)$$

$$\mathcal{H}^{[s]} = \mathcal{A}^{[s]\dagger}\mathcal{A}^{[s]} + \mathcal{E}_s = p^2 + U^{[s]}(x) + \mathcal{E}_s. \quad (2.30)$$

We have also

$$\phi_n^{[s-1]}(x) = \frac{\mathcal{A}^{[s-1]\dagger}}{\mathcal{E}_n - \mathcal{E}_{s-1}} \phi_n^{[s]}(x), \quad (n \geq s \geq 1). \quad (2.31)$$

In terms of the determinant (Wronskian)

$$\mathbb{W}[f_1, \dots, f_n](x) \stackrel{\text{def}}{=} \det \left(\frac{d^{j-1}f_k(x)}{dx^{j-1}} \right)_{1 \leq j, k \leq n}, \quad (2.32)$$

(for $n = 0$, we set $W[\cdot](x) = 1$.), we have

$$W[\phi_0, \phi_1, \dots, \phi_{s-1}, \phi_n](x) = \phi_0(x)\phi_1^{[1]}(x) \cdots \phi_{s-1}^{[s-1]}(x)\phi_n^{[s]}(x), \quad (n \geq s \geq 0). \quad (2.33)$$

Therefore we arrive at the concise formulas due to Crum:¹⁾

$$W[\phi_0, \phi_1, \dots, \phi_{s-1}](x) = \phi_0(x)\phi_1^{[1]}(x) \cdots \phi_{s-1}^{[s-1]}(x) = \pm e^{\mathcal{W}(x) + \mathcal{W}^{[1]}(x) + \cdots + \mathcal{W}^{[s-1]}(x)}, \quad (2.34)$$

$$U^{[s]}(x) = U(x) - 2 \frac{d^2}{dx^2} \left(\log W[\phi_0, \phi_1, \dots, \phi_{s-1}](x) \right), \quad (2.35)$$

$$\phi_n^{[s]}(x) = \frac{W[\phi_0, \phi_1, \dots, \phi_{s-1}, \phi_n](x)}{W[\phi_0, \phi_1, \dots, \phi_{s-1}](x)}, \quad (n \geq s \geq 0). \quad (2.36)$$

§3. ‘Discrete’ Quantum Mechanics (pure imaginary shifts)

In discrete quantum mechanics, the dynamical variables are, as in ordinary QM, the coordinate x , which takes value in an infinite or a semi-infinite or a finite range of the real axis and the canonical momentum p , which is realised as a differential operator $p = -i\partial_x$. Since the momentum operator appears in exponentiated forms $e^{\pm\gamma p}$, $\gamma \in \mathbb{R}$, in a Hamiltonian, it causes finite pure imaginary shifts in the wavefunction $e^{\pm\gamma p}\psi(x) = \psi(x \mp i\gamma)$. This requires the wavefunction as well as other functions appearing in the Hamiltonian to be *analytic* in x within a certain domain including the physical region of the coordinate. Let us introduce the $*$ -operation on an analytic function, $*$: $f \mapsto f^*$. If $f(x) = \sum_n a_n x^n$, $a_n \in \mathbb{C}$, then $f^*(x) \stackrel{\text{def}}{=} \sum_n a_n^* x^n$, in which a_n^* is the complex conjugation of a_n . Obviously $f^{**}(x) = f(x)$ and $f(x)^* = f^*(x^*)$. If a function satisfies $f^* = f$, then it takes real values on the real line.

The starting point is again a generic one dimensional discrete quantum mechanics Hamiltonian with discrete semi-infinite energy levels only (2.1). Again we assume that the groundstate energy is chosen to be zero $\mathcal{E}_0 = 0$, so that the Hamiltonian is positive semi-definite. The generic factorised Hamiltonian reads

$$\mathcal{H} = \mathcal{A}^\dagger \mathcal{A} = \sqrt{V(x)} e^{\gamma p} \sqrt{V^*(x)} + \sqrt{V^*(x)} e^{-\gamma p} \sqrt{V(x)} - V(x) - V^*(x), \quad (3.1)$$

$$\mathcal{A} \stackrel{\text{def}}{=} i(e^{\frac{\gamma}{2}p} \sqrt{V^*(x)} - e^{-\frac{\gamma}{2}p} \sqrt{V(x)}), \quad \mathcal{A}^\dagger \stackrel{\text{def}}{=} -i(\sqrt{V(x)} e^{\frac{\gamma}{2}p} - \sqrt{V^*(x)} e^{-\frac{\gamma}{2}p}). \quad (3.2)$$

By specifying the function $V(x)$, various explicit examples are obtained.^{7),9),11)} For instance,

$$V(x) = \frac{\prod_{j=1}^4 (1 - a_j e^{ix})}{(1 - e^{2ix})(1 - q e^{2ix})}, \quad 0 < x < \pi, \quad 0 < q < 1, \\ |a_j| < 1, \quad \{a_1^*, a_2^*, a_3^*, a_4^*\} = \{a_1, a_2, a_3, a_4\} \text{ (as a set)}, \quad (3.3)$$

gives an exactly solvable dynamics whose eigenfunctions consist of the Askey-Wilson polynomials¹⁴⁾⁻¹⁶⁾ times the groundstate wavefunction $\phi_0(x)$ (3.5), which gives the orthogonality measure $\phi_0(x)^2$.

Let us emphasise that the corresponding Schrödinger equation $\mathcal{H}\psi(x) = \mathcal{E}\psi(x)$ is a difference equation

$$\begin{aligned} \sqrt{V(x)V^*(x-i\gamma)}\psi(x-i\gamma) + \sqrt{V^*(x)V(x+i\gamma)}\psi(x+i\gamma) \\ - (V(x) + V^*(x))\psi(x) = \mathcal{E}\psi(x), \end{aligned} \quad (3.4)$$

instead of differential in ordinary QM. Again the groundstate wavefunction $\phi_0(x)$ is determined as a zero mode of \mathcal{A} :

$$\mathcal{A}\phi_0(x) = 0 \Rightarrow \mathcal{H}\phi_0(x) = 0. \quad (3.5)$$

The above equation for ϕ_0 reads

$$\sqrt{V^*(x-i\frac{\gamma}{2})}\phi_0(x-i\frac{\gamma}{2}) - \sqrt{V(x+i\frac{\gamma}{2})}\phi_0(x+i\frac{\gamma}{2}) = 0. \quad (3.6)$$

This dictates how the 'phase' of the potential function V is related to that of the groundstate wavefunction ϕ_0 . Here we also assume that the groundstate wavefunction $\phi_0(x)$ has no node and chosen to be real and positive for real x .

Due to the lack of generic theorems in the theory of difference equations, let us assume that all the energy levels are non-degenerate and all the eigenfunctions are square-integrable and orthogonal with each other and form a complete basis of the Hilbert space:

$$\mathcal{H}\phi_n(x) = \mathcal{E}_n\phi_n(x), \quad n = 0, 1, 2, \dots, \quad (3.7)$$

$$\int \phi_n(x)^*\phi_m(x)dx = h_n\delta_{nm}, \quad 0 < h_n < \infty, \quad n, m = 0, 1, 2, \dots \quad (3.8)$$

In most explicit examples these statements can be verified straightforwardly. For simplicity we choose all the eigenfunctions to be real on the real axis $\phi_n^* = \phi_n$. This is possible since \mathcal{H} maps a 'real' function to a 'real' function $f^* = f \Rightarrow (\mathcal{H}f)^* = \mathcal{H}f$.

Now the procedure to generate the associated Hamiltonian system goes almost parallel with the one shown in the preceding section. We define an associated Hamiltonian $\mathcal{H}^{[1]}$ by simply changing the order of \mathcal{A} and \mathcal{A}^\dagger , $\mathcal{H}^{[1]} \stackrel{\text{def}}{=} \mathcal{A}\mathcal{A}^\dagger$, as in (2.11). Then as in (2.12), we have $\mathcal{A}^{[0]}\mathcal{H}^{[0]} = \mathcal{H}^{[1]}\mathcal{A}^{[0]}$ and $\mathcal{A}^{[0]\dagger}\mathcal{H}^{[1]} = \mathcal{H}^{[0]}\mathcal{A}^{[0]\dagger}$. Here again we have indexed the quantities of the original Hamiltonian system by the superscript $^{[0]}$, as before. The eigenfunctions and the eigenvalues of the associated Hamiltonian $\mathcal{H}^{[1]}$ are given by

$$\phi_n^{[1]}(x) \stackrel{\text{def}}{=} \mathcal{A}^{[0]}\phi_n^{[0]}(x), \quad \int \phi_n^{[1]}(x)^*\phi_m^{[1]}(x)dx = \mathcal{E}_n h_n \delta_{nm}, \quad n, m = 1, 2, \dots, \quad (3.9)$$

$$\mathcal{H}^{[1]}\phi_n^{[1]}(x) = \mathcal{E}_n\phi_n^{[1]}(x), \quad n = 1, 2, \dots \quad (3.10)$$

The same argument as before establishes the iso-spectrality and the one to one correspondence of the eigenfunctions of $\mathcal{H}^{[0]}$ and $\mathcal{H}^{[1]}$ except for the groundstate. In explicit examples given in,⁹ one can show that $\phi'(x) = \phi'(x; \boldsymbol{\lambda}) = \varphi(x)/\phi_0(x; \boldsymbol{\lambda} + \boldsymbol{\delta})$ is annihilated by \mathcal{A}^\dagger , $\mathcal{A}^\dagger\phi'(x) = 0$ as in the Darboux transformation. Such functions, as before, do not belong to the Hilbert space of the associated Hamiltonian $\mathcal{H}^{[1]}$. Here

λ and δ are parameters and their shift. For example, $\varphi \equiv 1$ for the Meixner-Pollaczek case, $\varphi(x) = x$ for the Wilson case and $\varphi(x) = \sin x$ for the Askey-Wilson case (3.3). We have used the convention to specify the discrete quantum mechanics by the name of the polynomials constituting the main part of the eigenfunctions.

Thus $\phi_1^{[1]}(x)$ is the groundstate wavefunction of $\mathcal{H}^{[1]}$ with the eigenvalue \mathcal{E}_1 . By subtracting it from $\mathcal{H}^{[1]}$, we can again factorise the positive semi-definite Hamiltonian:

$$\mathcal{H}^{[1]} = \mathcal{A}^{[1]\dagger} \mathcal{A}^{[1]} + \mathcal{E}_1, \quad (3.11)$$

$$\begin{aligned} \mathcal{A}^{[1]} &\stackrel{\text{def}}{=} i(e^{\frac{\gamma}{2}p} \sqrt{V^{[1]*}(x)} - e^{-\frac{\gamma}{2}p} \sqrt{V^{[1]}(x)}), \\ \mathcal{A}^{[1]\dagger} &\stackrel{\text{def}}{=} -i(\sqrt{V^{[1]}(x)} e^{\frac{\gamma}{2}p} - \sqrt{V^{[1]*}(x)} e^{-\frac{\gamma}{2}p}), \end{aligned} \quad (3.12)$$

with the new potential function $V^{[1]}$ to be determined now. This imposes quadratic relations between the function $V^{[0]}$ and the unknown $V^{[1]}$:

$$V^{[0]}(x - i\frac{\gamma}{2})V^{[0]*}(x - i\frac{\gamma}{2}) = V^{[1]}(x)V^{[1]*}(x - i\gamma), \quad (3.13)$$

$$V^{[0]}(x + i\frac{\gamma}{2}) + V^{[0]*}(x - i\frac{\gamma}{2}) = V^{[1]}(x) + V^{[1]*}(x) - \mathcal{E}_1, \quad (3.14)$$

which are discrete counterparts of the Riccati equation (2.21) for the pre-potential \mathcal{W} . One essential problem is that the connection between the groundstate wavefunction $\phi_0(x)$ and the function $V(x)$ in the Hamiltonian is indirect, in contrast to the one $\phi_0(x) = e^{\mathcal{W}(x)}$, (2.4) in ordinary QM. In discrete QM, the potential function V is complex (analytic). The information on the ‘absolute value’ of $V^{[1]}(x)$ can be extracted from (3.13). The ‘phase’ part of $V^{[1]}(x)$ is given by that of $\phi_1^{[1]}(x)$ through the zero mode equation of $\mathcal{A}^{[1]}$ (3.16), just as the ‘phase’ of ϕ_0 is related to that of $V^{[0]}$ through the the zero mode equation of $\mathcal{A}^{[0]}$, (3.6). Thus the following formula determining the function $V^{[1]}(x)$ in terms of the previous function $V^{[0]}(x)$ and $\phi_1^{[1]}(x)$ is the main result of the present paper:

$$V^{[1]}(x) \stackrel{\text{def}}{=} \sqrt{V^{[0]}(x - i\frac{\gamma}{2})V^{[0]*}(x - i\frac{\gamma}{2})} \frac{\phi_1^{[1]}(x - i\gamma)}{\phi_1^{[1]}(x)}, \quad (3.15)$$

$$\mathcal{A}^{[1]}\phi_1^{[1]}(x) = 0. \quad (3.16)$$

There is no ambiguity in the phase of the square root in the above expression (3.15). It is positive for $x = \alpha + i\frac{\gamma}{2}$, $\alpha \in \mathbb{R}$, for which the function inside the square root sign is positive definite. Here we have to assume that the groundstate wavefunction $\phi_1^{[1]}(x)$ has no node, so that the function $V^{[1]}(x)$ does not develop unwanted singularities in the physical region. It is rather straightforward to verify that (3.13) is actually satisfied by the $V^{[1]}$ in (3.15). In order to verify the linear relation (3.14), one has to use the eigenvalue equation for $\phi_1^{[0]}$. By reversing the order of $\mathcal{A}^{[1]\dagger}$ and $\mathcal{A}^{[1]}$, the second associated Hamiltonian system $\mathcal{H}^{[2]}$ can be defined. Again this process can go indefinitely.

Here we list the definition of the s -th quantities step by step for $s \geq 1$,

$$\mathcal{H}^{[s]} \stackrel{\text{def}}{=} \mathcal{A}^{[s-1]} \mathcal{A}^{[s-1]\dagger} + \mathcal{E}_{s-1}, \quad (3-17)$$

$$\phi_n^{[s]}(x) \stackrel{\text{def}}{=} \mathcal{A}^{[s-1]} \phi_n^{[s-1]}(x), \quad (n \geq s), \quad (3-18)$$

$$V^{[s]}(x) \stackrel{\text{def}}{=} \sqrt{V^{[s-1]}(x - i\frac{\gamma}{2}) V^{[s-1]*}(x - i\frac{\gamma}{2})} \frac{\phi_s^{[s]}(x - i\gamma)}{\phi_s^{[s]}(x)}, \quad (3-19)$$

$$\begin{aligned} \mathcal{A}^{[s]} &\stackrel{\text{def}}{=} i(e^{\frac{\gamma}{2}p} \sqrt{V^{[s]*}(x)} - e^{-\frac{\gamma}{2}p} \sqrt{V^{[s]}(x)}), \\ \mathcal{A}^{[s]\dagger} &\stackrel{\text{def}}{=} -i(\sqrt{V^{[s]}(x)} e^{\frac{\gamma}{2}p} - \sqrt{V^{[s]*}(x)} e^{-\frac{\gamma}{2}p}). \end{aligned} \quad (3-20)$$

Let us note that the phase of the square root in (3-19) has no ambiguity. Then we can show the following for $n \geq s \geq 0$,

$$\mathcal{H}^{[s]} \phi_n^{[s]}(x) = \mathcal{E}_n \phi_n^{[s]}(x), \quad (3-21)$$

$$\phi_n^{[s]*} = \phi_n^{[s]} \quad : \text{ 'real' function}, \quad (3-22)$$

$$\mathcal{A}^{[s]} \phi_s^{[s]}(x) = 0, \quad (3-23)$$

$$\mathcal{H}^{[s]} = \mathcal{A}^{[s]\dagger} \mathcal{A}^{[s]} + \mathcal{E}_s. \quad (3-24)$$

We have also

$$\phi_n^{[s-1]}(x) = \frac{\mathcal{A}^{[s-1]\dagger}}{\mathcal{E}_n - \mathcal{E}_{s-1}} \phi_n^{[s]}(x), \quad (n \geq s \geq 1). \quad (3-25)$$

The discrete counterpart of the determinant formulas of Crum (2-33)–(2-36) requires a deformation of the Wronskian, the Casorati determinant, which has a good limiting property:

$$W_\gamma[f_1, \dots, f_n](x) \stackrel{\text{def}}{=} i^{\frac{1}{2}n(n-1)} \det \left(f_k(x + i\frac{n+1-2j}{2}\gamma) \right)_{1 \leq j, k \leq n}, \quad (3-26)$$

$$\lim_{\gamma \rightarrow 0} \gamma^{-\frac{1}{2}n(n-1)} W_\gamma[f_1, f_2, \dots, f_n](x) = W[f_1, f_2, \dots, f_n](x), \quad (3-27)$$

(for $n = 0$, we set $W_\gamma[\cdot](x) = 1$). Then we have, corresponding to (2-33),

$$W_\gamma[\phi_0, \phi_1, \dots, \phi_{s-1}, \phi_n](x) = \prod_{k=0}^{s-1} \check{\phi}_k^{[k]}(x + i\frac{k-s}{2}\gamma) \cdot \check{\phi}_n^{[s]}(x), \quad (3-28)$$

$$\check{\phi}_n^{[s]}(x) \stackrel{\text{def}}{=} \frac{\phi_n^{[s]}(x)}{\prod_{l=0}^{s-1} \sqrt{V^{[l]}(x + i\frac{s-l}{2}\gamma)}}. \quad (3-29)$$

Corresponding to (2-36), we obtain

$$\phi_n^{[s]}(x) = \prod_{l=0}^{s-1} \sqrt{V^{[l]}(x + i\frac{s-l}{2}\gamma)} \cdot \frac{W_\gamma[\phi_0, \phi_1, \dots, \phi_{s-1}, \phi_n](x)}{W_\gamma[\phi_0, \phi_1, \dots, \phi_{s-1}](x - i\frac{\gamma}{2})}, \quad (n \geq s \geq 0). \quad (3-30)$$

The proof of the above statements is elementary by induction and the necessary nontrivial formulas are only

$$\phi_n^{[s]}(x) = i \frac{\sqrt{V^{[s-1]}(x + i\frac{\gamma}{2})}}{\phi_{s-1}^{[s-1]}(x - i\frac{\gamma}{2})} \begin{vmatrix} \phi_{s-1}^{[s-1]}(x + i\frac{\gamma}{2}) & \phi_n^{[s-1]}(x + i\frac{\gamma}{2}) \\ \phi_{s-1}^{[s-1]}(x - i\frac{\gamma}{2}) & \phi_n^{[s-1]}(x - i\frac{\gamma}{2}) \end{vmatrix}, \quad (n \geq s \geq 1), \quad (3.31)$$

$$\begin{vmatrix} W_\gamma[f_0, f_1, \dots, f_{s-1}, f_s](x + i\frac{\gamma}{2}) & W_\gamma[f_0, f_1, \dots, f_{s-1}, f_n](x + i\frac{\gamma}{2}) \\ W_\gamma[f_0, f_1, \dots, f_{s-1}, f_s](x - i\frac{\gamma}{2}) & W_\gamma[f_0, f_1, \dots, f_{s-1}, f_n](x - i\frac{\gamma}{2}) \end{vmatrix} \\ = -i W_\gamma[f_0, f_1, \dots, f_{s-1}](x) W_\gamma[f_0, f_1, \dots, f_{s-1}, f_s, f_n](x), \quad (n \geq s \geq 0). \quad (3.32)$$

§4. Summary and Comments

Since Crum's paper¹⁾ is crisp and elegant, the underlying logical structure is not easy to fathom for non-experts or physicists. In section two we reproduce his results by a simplest logic in the language of quantum mechanics and using the factorisation method,³⁾ or the so-called supersymmetric quantum mechanics,⁴⁾ so that the similarity and contrast with the corresponding results of the discrete QM would become clear. Due to the lack of essential theorems in the theory of difference equations, some important properties of the spectra and eigenfunctions of the generic discrete QM must be assumed for the derivation of the associated Hamiltonian systems in section three. For example, the hermiticity or the self-adjointness of a discrete QM Hamiltonian can only be demonstrated after a proper groundstate wavefunction ϕ_0 is chosen.^{9),12)} As mentioned repeatedly, these standard properties are well satisfied in explicit examples of discrete QM in.^{7),9),11)} However, to the best of our knowledge, the general structure of the solutions of the main difference equation (3.4) as well as that for the groundstate (3.5) for generic potential $V(x)$ and $V^*(x)$, has not yet been investigated in contradistinction to the ordinary QM. In this connection, let us mention two interesting examples of quasi-exactly solvable discrete quantum mechanics (example in §IIB1 of the first paper of¹²⁾ and in §3 of¹³⁾) in which the oscillation theorem does not hold.

Here are some comments on closely related topics; shape invariance, orthogonal polynomials and limiting properties to the ordinary quantum mechanics, etc.

Shape invariance: The Hamiltonian may contain several parameters $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots)$ and we write them explicitly $\mathcal{H} = \mathcal{H}(\boldsymbol{\lambda})$, $V(x) = V(x; \boldsymbol{\lambda})$, $\mathcal{A} = \mathcal{A}(\boldsymbol{\lambda})$, etc. Let us consider the case that the potential function of the first associated Hamiltonian $V^{[1]}(x) = V^{[1]}(x; \boldsymbol{\lambda})$ has the same form as the original function V with a different set of parameters and up to a multiplicative positive constant $\kappa \in \mathbb{R}_+$:

$$V^{[1]}(x; \boldsymbol{\lambda}) = \kappa V(x; \boldsymbol{\lambda}'). \quad (4.1)$$

Here the new set of parameters $\boldsymbol{\lambda}'$ is uniquely determined by $\boldsymbol{\lambda}$ (let us write $\boldsymbol{\lambda}' = \text{si}(\boldsymbol{\lambda})$). Then this system has the shape invariance,^{5),7),9)}

$$\mathcal{A}(\boldsymbol{\lambda})\mathcal{A}(\boldsymbol{\lambda})^\dagger = \kappa \mathcal{A}(\boldsymbol{\lambda}')^\dagger \mathcal{A}(\boldsymbol{\lambda}') + \mathcal{E}_1(\boldsymbol{\lambda}). \quad (4.2)$$

The shape invariance is a sufficient condition for exact solvability. The entire energy spectrum and the excited wavefunctions are expressed in terms of $\mathcal{E}_1(\boldsymbol{\lambda})$ and $\phi_0(x; \boldsymbol{\lambda})$ as follows:

$$\mathcal{E}_n(\boldsymbol{\lambda}) = \sum_{s=0}^{n-1} \kappa^s \mathcal{E}_1(\boldsymbol{\lambda}^{[s]}), \quad (4.3)$$

$$\phi_n(x; \boldsymbol{\lambda}) \propto \mathcal{A}(\boldsymbol{\lambda}^{[0]})^\dagger \mathcal{A}(\boldsymbol{\lambda}^{[1]})^\dagger \mathcal{A}(\boldsymbol{\lambda}^{[2]})^\dagger \dots \mathcal{A}(\boldsymbol{\lambda}^{[n-1]})^\dagger \phi_0(x; \boldsymbol{\lambda}^{[n]}), \quad (4.4)$$

where $\boldsymbol{\lambda}^{[n]}$ is $\boldsymbol{\lambda}^{[0]} = \boldsymbol{\lambda}$, $\boldsymbol{\lambda}^{[n]} = \text{si}(\boldsymbol{\lambda}^{[n-1]})$ ($n = 1, 2, \dots$).

Orthogonal polynomial: Here we consider a generic Hamiltonian (3.1) of the discrete QM. That is the shape invariance is not assumed. Let us define a 'real' function $\eta(x)$ ($\eta^* = \eta$) as a ratio of $\phi_1(x)$ and $\phi_0(x)$,

$$\frac{\phi_1(x)}{\phi_0(x)} = a + b\eta(x), \quad (4.5)$$

where a and b ($b \neq 0$) are real constants. Although $\eta(x)$ is not well defined without specifying a and b , this ambiguity (affine transformation of $\eta(x)$) does not affect the following discussion. Then (3.15) implies

$$V^{[1]}(x + i\frac{\gamma}{2}) = V(x) \frac{\eta(x - i\gamma) - \eta(x)}{\eta(x) - \eta(x + i\gamma)}. \quad (4.6)$$

Let us assume further that the n -th eigenfunction ϕ_n/ϕ_0 is a degree n polynomial in this $\eta(x)$ for all $n \geq 2$:

$$\phi_n(x) = \phi_0(x) P_n(\eta(x)), \quad P_n(y) = \sum_{k=0}^n a_{n,k} y^k, \quad a_{n,k} \in \mathbb{R}, \quad a_{n,n} \neq 0. \quad (4.7)$$

Obviously $a_{0,0} = 1$, $a = a_{1,0}$ and $b = a_{1,1}$. The orthogonality of the eigenfunctions $\{\phi_n\}$ implies that $\{P_n(\eta(x))\}$ are orthogonal polynomials in $\eta(x)$ with respect to the weight function $\phi_0(x)^2$. Then the ratio of the first excited state and the groundstate of the s -th associated Hamiltonian system $\mathcal{H}^{[s]}$ takes the same form as that of the first associated Hamiltonian system $\mathcal{H}^{[1]}$ (4.5),

$$\frac{\phi_{s+1}^{[s]}(x)}{\phi_s^{[s]}(x)} = \frac{a_{s+1,s}}{a_{s,s}} + \frac{a_{s+1,s+1}}{a_{s,s}} \eta^{[s]}(x), \quad \eta^{[s]}(x) \stackrel{\text{def}}{=} \sum_{k=0}^s \eta(x + i\frac{2k-s}{2}\gamma), \quad (4.8)$$

and $V^{[s]}$ is related to $V^{[s-1]}$ as in (4.6) and this goes further down to $V^{[0]}$:

$$\begin{aligned} V^{[s]}(x + i\frac{s}{2}\gamma) &= V^{[s-1]}(x + i\frac{s-1}{2}\gamma) \frac{\eta^{[s-1]}(x + i\frac{s-3}{2}\gamma) - \eta^{[s-1]}(x + i\frac{s-1}{2}\gamma)}{\eta^{[s-1]}(x + i\frac{s-1}{2}\gamma) - \eta^{[s-1]}(x + i\frac{s+1}{2}\gamma)} \\ &= V(x) \prod_{k=0}^{s-1} \frac{\eta^{[k]}(x + i\frac{k-2}{2}\gamma) - \eta^{[k]}(x + i\frac{k}{2}\gamma)}{\eta^{[k]}(x + i\frac{k}{2}\gamma) - \eta^{[k]}(x + i\frac{k+2}{2}\gamma)} \\ &= V(x) \prod_{k=0}^{s-1} \frac{\eta(x - i\gamma) - \eta(x + ik\gamma)}{\eta(x) - \eta(x + i(k+1)\gamma)}. \end{aligned} \quad (4.9)$$

In all the known examples of exactly solvable QM (ordinary and discrete), in which the eigenfunctions have the polynomial form (4.7), the function $\eta(x)$ plays a special role to define together with the Hamiltonian \mathcal{H} an algebraic sufficient condition for exact solvability. In that case the function $\eta(x)$ is called the *sinusoidal coordinate* and the sufficient condition is named the *closure relation*.^{8),9),11)} As shown above, the shape invariance and the closure relation are very closely related.

Limit from dQM to QM: Here we show that the ordinary QM is obtained from the discrete QM in a certain limit by rescaling parameters. Let us introduce a positive parameter c and rescale γ as γ/c and the parameters in $V(x)$ appropriately,

$$\mathcal{H} = \mathcal{A}^\dagger \mathcal{A} = \sqrt{V(x)} e^{\frac{\gamma}{c} p} \sqrt{V^*(x)} + \sqrt{V^*(x)} e^{-\frac{\gamma}{c} p} \sqrt{V(x)} - V(x) - V^*(x), \quad (4.10)$$

$$\mathcal{A} \stackrel{\text{def}}{=} i \left(e^{\frac{\gamma}{2c} p} \sqrt{V^*(x)} - e^{-\frac{\gamma}{2c} p} \sqrt{V(x)} \right), \quad \mathcal{A}^\dagger \stackrel{\text{def}}{=} -i \left(\sqrt{V(x)} e^{\frac{\gamma}{2c} p} - \sqrt{V^*(x)} e^{-\frac{\gamma}{2c} p} \right). \quad (4.11)$$

Assume that $V(x)$ has the following expansion for large c ,

$$V(x) = a \left(1 + \frac{i\gamma}{c} w_1(x) + O\left(\frac{1}{c^2}\right) \right), \quad (4.12)$$

where a is a positive constant. Then we have for large c :

$$\frac{c}{\sqrt{a}\gamma} \mathcal{A} = \frac{d}{dx} - \frac{d\mathcal{W}(x)}{dx} + O\left(\frac{1}{c}\right), \quad \frac{c}{\sqrt{a}\gamma} \mathcal{A}^\dagger = -\frac{d}{dx} - \frac{d\mathcal{W}(x)}{dx} + O\left(\frac{1}{c}\right), \quad (4.13)$$

$$\frac{c^2}{a\gamma^2} \mathcal{H} = p^2 + \left(\frac{d\mathcal{W}(x)}{dx} \right)^2 + \frac{d^2\mathcal{W}(x)}{dx^2} + O\left(\frac{1}{c}\right), \quad (4.14)$$

where the derivative of the pre-potential $\mathcal{W}(x)$ is defined by $\frac{d\mathcal{W}(x)}{dx} = -\text{Re } w_1(x)$. Therefore ordinary QM is obtained from discrete QM in the $c \rightarrow \infty$ limit.

Discrete QM with real shifts: Crum's theorem for the discrete QM with real shifts can also be formulated in a similar manner. In this case the Hamiltonian is a *real symmetric tri-diagonal* (Jacobi) matrix, either of finite or infinite dimensions.⁸⁾ The factorisation of the positive semi-definite Hamiltonian $\mathcal{H} = \mathcal{A}^\dagger \mathcal{A}$ also holds and \mathcal{A} consists of the diagonal and super-diagonal elements only and \mathcal{A}^\dagger being its transpose, which consists of the diagonal and sub-diagonal elements only.

Connection with integrable systems: Since the Darboux transformation²⁾ is closely related to the inverse scattering method for soliton equations, it is natural to ask if the present formulation of Crum's theorem is related to (discrete) integrable systems. At present, all the explicit examples considered in discrete QM^{7),9),11)} (pure imaginary shifts) have an infinite number of discrete eigenvalues only. In other words, the corresponding potentials are confining, that is, they grow to infinity at the boundaries or the spatial infinities. Thus the free incoming/outgoing waves at infinity do not exist and the corresponding scattering problem cannot be formulated. To sum up, we have nothing to report on possible applications of the present Crum's theorem to (discrete) integrable systems.

After completing this work, we received a recent work by Gaillard and Matveev²³⁾ which has some overlap with the present work. We thank Vladimir Matveev for sending the new results and for many useful comments.

Acknowledgements

We are supported in part by Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, No.18340061 and No.19540179.

References

- 1) M. M. Crum, Quart. J. Math. Oxford Ser. (2) **6** (1955), 121, arXiv:physics/9908019.
- 2) G. Darboux, C. R. Acad. Paris **94** (1882), 1456.
- 3) L. Infeld and T. E. Hull, Rev. Mod. Phys. **23** (1951), 21.
- 4) See, for example, a review: F. Cooper, A. Khare and U. Sukhatme, Phys. Rep. **251** (1995), 267.
- 5) L. E. Gendenshtein, JETP Lett. **38** (1983), 356.
- 6) V.É. Adler, Theor. Math. Phys. **101** (1994) 1381.
- 7) S. Odake and R. Sasaki, J. Nonlinear Math. Phys. **12** Suppl. 1 (2005), 507, arXiv:hep-th/0410102; J. Math. Phys. **46** (2005), 063513, arXiv:hep-th/0410109.
- 8) S. Odake and R. Sasaki, J. Math. Phys. **49** (2008), 053503, arXiv:0712.4106[math.CA].
- 9) S. Odake and R. Sasaki, Prog. Theor. Phys. **119** (2008), 663, arXiv:0802.1075[quant-ph].
- 10) S. Odake and R. Sasaki, Prog. Theor. Phys. **114** (2005), 1245, arXiv:hep-th/0512155.
- 11) S. Odake and R. Sasaki, J. Math. Phys. **47** (2006), 102102, arXiv:quant-ph/0605215; Phys. Lett. B **641** (2006), 112, arXiv:quant-ph/0605221.
- 12) R. Sasaki, J. Math. Phys. **48** (2007), 122104, arXiv:0708.0702[nlin.SI]; S. Odake and R. Sasaki, J. Math. Phys. **48** (2007), 122105, arXiv:0708.0716[nlin.SI].
- 13) R. Sasaki, J. Nonlinear Math. Phys. **15** Suppl. 3 (2008), 373, arXiv:0712.2616[nlin.SI].
- 14) G. E. Andrews, R. Askey and R. Roy, *Special Functions*, Encyclopedia of mathematics and its applications (Cambridge, 1999).
- 15) M. E. H. Ismail *Classical and quantum orthogonal polynomials in one variable*, Encyclopedia of mathematics and its applications (Cambridge, 2005).
- 16) R. Koekoek and R. F. Swarttouw, arXiv:math.CA/9602214.
- 17) A. J. Macfarlane, J. of Phys. **A22** (1989), 4581; L. C. Biedenharn, J. of Phys. **A22** (1989), L873; C.-P. Sun and H.-C. Fu, J. of Phys. **A22** (1989), L983.
- 18) S. Odake and R. Sasaki, Phys. Lett. B **663** (2008), 141, arXiv:0710.2209[hep-th].
- 19) R. Kobayashi, S. Odake and R. Sasaki, in preparation.
- 20) R. Sasaki, J. Math. Phys. October issue (2009) in press.
- 21) S. Odake and R. Sasaki, Phys. Lett. B **679** (2009), 414, arXiv:0906.0142[math-ph].
- 22) S. Odake and R. Sasaki, Preprint DPSU-09-4, YITP-09-52.
- 23) P. Gaillard and V. B. Matveev, "Wronskian and Casorati determinant representation for Darboux-Pöschel-Teller potentials and their difference extensions," Preprint RIMS-1653, January 2009.