

CRYPTOSPORIDIOSIS IN HIV INFECTED PATIENTS WITH DIARRHOEA IN OSUN STATE SOUTHWESTERN NIGERIA

Yemisi Olukemi Adesiji, Rofiat Omolabake Lawal, Samuel Sunday Taiwo,
Sunday Adetona Fayemiwo, Oluwaseyi Adegboyega Adeyeba

Ladoke Akintola University of Technology, College of Health Sciences, Department of Medical Microbiology and Parasitology, Osogbo, Nigeria

Aim: Although cryptosporidiosis is said to be rare among adult HIV patients in certain parts of Nigeria, there are no documented studies on the occurrence of this parasite among HIV patients in Southwestern Nigeria.

Methods: We conducted a cross sectional study involving 150 patients (100 HIV infected and 50 HIV negative) with diarrhoea from two tertiary and one secondary health institutions in Southwestern Nigeria. Concentrated stool sample from each patient was screened for *cryptosporidium* oocysts with the modified cold Ziehl Neelsen method while direct sample was screened for other enteric parasites.

Results: The overall parasite prevalence rate in the diarrhoea patients is 79.3% (119/150) with *Cryptosporidium parvum* 52.7%, *Ascaris lumbricoides* 11.3%, *Hookworm* 3.3%, *Trichuris trichura* 2% and *Entamoeba histolytica* 10%. The parasite prevalence rate in HIV infected patients is 97% while in HIV-negative patients, the rate is 44%. The rate is significantly higher among HIV-infected patients with diarrhoea than among HIV-negative patients with diarrhoea (OR=41.152, 95%CI=11.467-147.68, P<0.0001). However this difference is attributed to *Cryptosporidium* which was found exclusively among HIV-infected patients. When *Cryptosporidium* prevalence was excluded from analysis, the parasite prevalence rates between the two groups was not significantly different ($X^2=0.8002$, df=3, P=0.8494).

Conclusion: Contrary to a previous report of apparent rarity of *cryptosporidium* infections in certain parts of Nigeria, this study revealed a high prevalence of this parasitosis among HIV infected patients in Southwestern Nigeria.

Key words: Cryptosporidiosis, prevalence, Southwestern Nigeria

Eur J Gen Med 2007;4(3):119-122

INTRODUCTION

Cryptosporidium, an intracellular protozoan has changed from that of a rare largely asymptomatic disease, to an important cause of diarrhoea in animals and humans worldwide (1).

Reported Cryptosporidiosis prevalence is 3-4% in the USA (2), 3.5-22.44% in Brazil (3) and about 50% in Africa and Haiti (4). In HIV infected patients, *cryptosporidium* is the most frequent microbial cause of diarrhea, usually causing chronic bulky and intermittent diarrhoea with liquid non-bloody stools, accompanied by pain and abdominal colic, and a noticeable loss of weight (5)

Reports of cryptosporidiosis in Nigeria are

few. In North-central Nigeria, Nwabuisi(6) reported prevalence rate of 15.1% among children aged 0-14 years with diarrhoea while Bamwat *et al* (7) reported a prevalence rate of 4.8% among malnourished children 0-5years. The role of *Cryptosporidium* in diarrhoea among HIV infected patients in Nigeria has not been established. In one tertiary institution in Southeastern Nigeria (8), no cryptosporidium oocyst was detected in diarrhoea stools of 189 HIV infected and uninfected patients. In Southwestern Nigeria, there are no documented studies on the prevalence of *Cryptosporidium* among the people. Our study is therefore aimed at determining the prevalence and contribution

Correspondence: Dr O. Y. Adesiji

Department of Medical Microbiology and Parasitology,
College of Health Sciences, Ladoke Akintola University
of Technology, PMB 4400, Osogbo, Nigeria

GSM: 2348032948270

E-mail: drmradesiji@yahoo.com

Table 1. Age and sex distribution of HIV-infected patients with diarrhoea

Age group	Male	Female	Total
6-15	3	1	4
16-25	10	12	22
26-35	19	23	42
36-45	14	12	26
46-55	5	1	6
Total	51	49	100

Mean age = 28.4±11.19

of *Cryptosporidium* to diarrhoea among HIV infected patients, the knowledge of which will help in management of HIV related opportunistic infections in our environment.

MATERIALS AND METHODS

Study area

This cross sectional study was conducted in three selected health care facilities in Osun State of Nigeria; State Hospital, Ilesha; Ladoke Akintola University Teaching Hospital, Osogbo and Obafemi Awolowo University Teaching Hospital Complex, Ile-Ife between July and October 2005.

Subjects

One hundred HIV-infected individuals presenting with diarrhoea were recruited among the patients admitted to HIV or Haematology clinics of the health institutions and 50 HIV-negative individuals with diarrhoea admitted to the medical outpatient or admitted on the wards were included as controls. HIV was diagnosed using a combination of an HIV-1 ELISA screening and a Western blot confirmatory technique or where the latter is not available, using 2 different ELISA methods. Diarrhoea was defined as passage of more than 3 loose or watery stools in 24 hours and was acute if it lasts for less than 14 days and persistent/chronic if it lasts for 14 or more days(9).

Specimen collection

Stool samples were collected from each subject into a clean Universal bottle and transported to the medical microbiology laboratory of Ladoke Akintola University Teaching Hospital for analysis. Each subject was interviewed to collect relevant demographic data.

Methodology

A direct sample of the stool was made with both saline and iodine mounts on clean grease free slides and examined under the microscope first with 10x and then 40x objective lens for

Table 2. Age and sex distribution of HIV-negative patients with diarrhoea

Age group	Male	Female	Total
6-15	2	3	5
16-25	9	8	17
26-35	6	10	16
36-45	3	6	9
46-55	2	1	3
Total	22	28	50

Mean age = 28.4±11.19

ova and cysts of parasites. A sample of the stool was concentrated using formol ether method concentration technique (10).

Detection of *Cryptosporidium* oocysts in the concentrated stool was done using the modified cold Ziehl Neelsen staining technique (11). Briefly, a concentrated smear of the stool was made on a clean grease-free slide and fixed in methanol for 3 minutes. The slide was immersed in cold Carbol fuchsin and stained for 15 minutes. It was then thoroughly rinsed in tap water and decolorized in 1% HCl (v/v) in methanol for 10-15 minutes. After rinsing again in tap water, the slide was counterstained with 0.4% malachite green for 30 seconds. The slide was then air-dried and observed under the compound light microscope using 40x objective lens for the presence of *Cryptosporidium* oocysts, which was confirmed under the oil-immersion objectives, as small pink to red spherules on pale green background

Data analysis

Demographic and other data obtained were analysed with a PC containing GraphPad software (GraphPad Software Inc, San Diego, USA). Significant differences between categorical variables were determined using Chi square or Fisher exact tests and $P < 0.05$ was taken as significant value.

RESULTS

A total of 150 patients with diarrhoea were investigated for cryptosporidiosis in three health institutions in Osun State, Southwestern Nigeria; 100 (51 males, 49 females, age range 9-54 years, mean age 32.04±9.62) were HIV infected and 50 (22 males, 28 females, age range 9-56 years, mean age 28.4±11.9) were HIV negative and served as controls (Tables 1 and 2).

The overall parasite prevalence rate in the diarrhoea patients is 79.3% (119/150) with *Cryptosporidium parvum* 52.7%, *Ascaris lumbricoides* 11.3%, Hookworm 3.3%, *Trichuris trichura* 2% and *Entamoeba*

Table 3. Distribution of parasites isolated from stool of patients with diarrhoea in 3 health institutions in Southwestern Nigeria

Parasite types	HIV-infected (n:100)	HIV-negative (n:50)	Total (%) (n:150)
<i>Cryptosporidium</i>	79	0	79 (52.7)
<i>Ascaris</i>	9	8	17 (11.3)
Hookworm	2	3	5 (3.3)
<i>Trichuris</i>	1	2	3 (2)
<i>Entamoeba</i>	6	9	15 (10)
Total	97	22	119 (79.3)

Fisher Exact OR=41.152, 95%CI=11.467-147.68, $P<0.0001$
 $X^2=0.8002$, $df=3$, $P=0.8494$ (Minus *Cryptosporidium* prevalence)

histolytica 10% (Table 3). The parasite prevalence rate in HIV infected patients is 97% while in HIV-negative patients, the rate is 44%. The rate is significantly higher among HIV-infected patients with diarrhoea than among HIV-negative patients with diarrhoea (OR=41.152, 95%CI=11.467-147.68, $P<0.0001$). However this difference is attributed to *Cryptosporidium* which was found exclusively among HIV-infected patients. When *Cryptosporidium* prevalence was excluded from analysis, the parasite prevalence rates between the two groups was not significantly different ($X^2=0.8002$, $df=3$, $P=0.8494$).

DISCUSSION

Diarrhoea is a common complication of HIV infections, inducing weight loss and cachexia and occurring in almost 90% of AIDS patients in developing countries (12). *Cryptosporidium* is a well established cause of diarrhoea among HIV infected patients worldwide with prevalence of infection ranging from 3% in developed countries to 50% in developing countries (13). However, the role of this parasite in the occurrence of diarrhoea among adult patients with HIV infection in Nigeria is not established. In a study carried out in a tertiary institution in Southeastern Nigeria (8) using modified cold Ziehl Neelsen staining technique, *Cryptosporidium* was not detected in stool samples of 189 HIV infected and non-infected patients with diarrhoea. This sharply contrasts the finding in our study using the same detection method where an overall *Cryptosporidium* prevalence of 52.7% is reported and 79% prevalence rate among HIV infected patients. The reason for this large difference can be related to the known fact that HIV opportunistic infections, cryptosporidiosis inclusive, tend to vary from one locality to another and from one country to the other depending on the level

of contamination of water, foodstuff and contacts with animals, which are important factors in dissemination of the parasite (14). Cryptosporidiosis in Nigeria has been reported from tertiary institutions mainly among children in North-central (6, 7, 15) and South-south Nigeria (16). Our study on *Cryptosporidium* among diarrhoea patients appears to be the first from Southwestern Nigeria involving two tertiary and one secondary health institutions. The *Cryptosporidium* prevalence rate of 52.7% reported among these diarrhoea patients is similar to rates reported from other African countries(4) indicating that cryptosporidiosis is an important opportunistic parasitic disease causing diarrhoea among HIV infected patients in Nigeria. This is especially so as this parasite was detected in stool samples of only HIV infected patients and this further highlights the association between immunodepression and cryptosporidiosis.

Although the overall parasite prevalence rate was higher among HIV infected patients when compared to HIV negative patients with diarrhoea in this study, this was mainly due to the high *Cryptosporidium* prevalence of 79% among the HIV infected patients. When this was removed, the prevalence rates of other parasites were not significantly different among the HIV infected and HIV negative patients.

The mechanism by which *Cryptosporidium* cause diarrhoea is not well known. Inflammatory response to the infection is variable and may be modified by co-pathogens such as *Cytomegalovirus*. However, histologic evidence of gastrointestinal mucosa injury has been reported with clinical manifestations influenced in part by the anatomic distribution of the infection with extensive infections involving both small and large intestines producing the most severe illness (17)

Contrary to a previous report of apparent rarity of *Cryptosporidium* infections in certain part of Nigeria (8), our study revealed a high prevalence of this parasitosis among HIV infected patients in Southwestern Nigeria.

ACKNOWLEDGEMENT

We are grateful to Mr. Amure- the Chief technologist of Ladoko Akintola University Teaching Hospital who assisted in processing the specimens and the entire staff of the Department of Medical Microbiology of the Hospital. Also, the staff and management of Osun State Hospital, Ilesha and Obafemi Awolowo University Teaching Hospital Complex, Ile-Ife, Nigeria who facilitated collection of stool samples from the patients. The study is significant because in Southwestern Nigeria, there are no documented studies on the prevalence of Cryptosporidium among the populace. Our study is therefore aimed at determining the prevalence and contribution of Cryptosporidium to diarrhoea among HIV infected patients, the knowledge of which will help in management of HIV related opportunistic infections in our environment.

REFERENCES

1. Flanigan T, Whalen C, Turner J, et al. Cryptosporidium infection and CD4 counts. *Ann Intern Med* 1992;116:840-2
2. Mac Kenzie WR, Hoxie NJ, Proctor ME, et al. A massive outbreak in Milwaukee of Cryptosporidium infection transmitted through the public water supply. *N Engl J Med* 1994;331:161-7
3. Rodrigues JLN, Leser P, Silva TMJ, et al. Prevalência da criptosporidiose nasíndrome diarréica do paciente HIV-positivo. *Rev Assoc Med Brasil* 1991;37: 79-84
4. Colebunders R, Francis H, Mann JM, et al. Persistent diarrhea, strongly associated with HIV infection in Kinshasa, Zaire. *Am J Gastroenterol* 1987;82:859-64
5. Kotler DP. Gastrointestinal manifestations of human immunodeficiency virus infection. *Adv Intern Med* 1995;40:197-242
6. Nwabuisi C. Childhood cryptosporidiosis and intestinal parasitosis in association with diarrhoea in Kwara State, Nigeria. *West Afr J Med* 2001;20:165-8
7. Banwat EB, Egah DZ, Onile BA, Angyo IA, Audu ES. Prevalence of cryptosporidium infection among undernourished children in Jos, Central Nigeria. *Niger Postgrad Med J* 2003;10: 84-7
8. Nwokediuko SC, Bojuwoye BJ, Onyenekwe B. Apparent rarity of cryptosporidiosis in human immunodeficiency virus (HIV)-related diarrhoea in Enugu, South-Eastern Nigeria. *Niger. Postgrad Med J* 2002;9: 70-3
9. World Health Organization. Global Programme on AIDS. Guidelines for the clinical management of HIV infection in adults, 1991
10. Cheesbrough M. District Laboratory Practice in Tropical Countries. Part I. Cambridge University Press 1999:193-208
11. Smith HV. Intestinal Protozoa. In; Gillespie SH, Hawkey PM eds. *Medical Parasitology. A Practical Approach.* Oxford University Press, New York, 1995:79-118
12. Mukhopadhyaya A, Ramakrishna BS, Kang G, et al. Enteric pathogens in Southern Indian HIV infected patients with or without diarrhoea. *Indian J Med Res* 1999;109:85-9
13. Goldstein ST, Juranek DD, Ravenholt O, et al. Cryptosporidiosis: an outbreak associated with drinking water despite state-of-the art water treatment. *Ann Intern Med* 1996;124:459-68
14. Widmer G, Carraway M, Tzipori S. Water-borne Cryptosporidium: a perspective from the USA. *Parasitol Today* 1996;12: 286-90
15. Nwabuisi C. Cryptosporidiosis among diarrhoea patients in Ilorin, Nigeria. *Niger. Med Pract* 1998;35:39-41
16. Nkanginieme KEO, Chira FW, Oruamabo RS. Cryptosporidiosis in undernourished under five children with diarrhoea at the University of Port Harcourt Teaching Hospital in Nigeria. *Niger Postgrad Med J* 1996;3:5-9
17. Lumadue JA, Manabe YC, Moore RD, Belitsos PC, Sears CL, Clark DP. A clinicopathologic analysis of AIDS-related cryptosporidiosis. *AIDS* 1998; 12:2459-66