

# Crystal Base for the Basic Representation of $U_q(\widehat{\mathfrak{sl}}(n))$

Kailash C. Misra<sup>1</sup> and Tetsuji Miwa<sup>2</sup>

<sup>1</sup> Department of Mathematics, North Carolina State University, Raleigh, NC 27695-8205, USA

<sup>2</sup> Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606, Japan

Received February 9, 1990

**Abstract.** We show the existence of the crystal base for the basic representation of  $U_q(\widehat{\mathfrak{sl}}(n))$  by giving an explicit description in terms of Young diagrams.

## 0. Introduction

In [5] Kashiwara introduces the notion of crystal base for integrable representations of  $U_q(\mathfrak{g})$ , where  $\mathfrak{g}$  is any symmetrizable Kac–Moody Lie algebra. The crystal base has a simple structure at  $q = 0$ . Let  $\{e_i, f_i, t_i^\pm\}$  be a set of generators of  $U_q(\mathfrak{g})$ . Suppose  $M$  is an integrable  $U_q(\mathfrak{g})$ -module. Kashiwara [5] constructs certain operators  $\tilde{e}_i, \tilde{f}_i$  acting on  $M$ . These operators are obtained by modifying the simple root vectors  $e_i$  and  $f_i$ . When  $M$  is an irreducible highest weight module with highest weight vector  $u$ , define:

$$L = \sum A \tilde{f}_{i_1} \tilde{f}_{i_2} \cdots \tilde{f}_{i_k} u \in M \tag{0.1}$$

and

$$B = \{v = \tilde{f}_{i_1} \tilde{f}_{i_2} \cdots \tilde{f}_{i_k} u \in L/qL \mid v \neq 0\}, \tag{0.2}$$

where  $A \subset K = \mathbf{Q}(q)$  is the ring of rational functions in  $q$  without pole at  $q = 0$ . Kashiwara [5] conjectures that  $(L, B)$  satisfies the following crucial properties:

$$\tilde{e}_i L \subset L \quad \text{and} \quad \tilde{f}_i L \subset L, \quad \text{for all } i, \tag{0.3}$$

$$\tilde{e}_i B \subset B \cup \{0\} \quad \text{and} \quad \tilde{f}_i B \subset B \cup \{0\}, \quad \text{for all } i, \tag{0.4}$$

$$u = \tilde{e}_i v \quad \text{if and only if} \quad v = \tilde{f}_i u, \quad \text{for all } i \quad \text{and} \quad u, v \in B. \tag{0.5}$$

He proves his conjecture for  $\mathfrak{g} = \mathfrak{sl}(n), \mathfrak{o}(2n + 1), \mathfrak{sp}(2n)$  and  $\mathfrak{o}(2n)$  and calls  $(L, B)$  the crystal base.

In this paper we prove this conjecture for the basic representation of  $U_q(\widehat{\mathfrak{sl}}(n))$  with highest weight  $\Lambda_0$  ( $\Lambda_0(t_i^\pm) = q^{\pm 1} \delta_{i,0}$ ). We start with the Fock space representation of  $U_q(\widehat{\mathfrak{sl}}(n))$  constructed by Hayashi [3]. We identify the Fock space  $\mathcal{F}$  with the space spanned by Young diagrams [4]. Then for each  $i$ , we decompose  $\mathcal{F}$  with respect to  $U_q(\mathfrak{sl}(2))_{(i)}$  generated by  $\{e_i, f_i, t_i^\pm\}$  (see, Theorem 3.1). This leads to the

construction of crystal base  $(L(\mathcal{F}), B(\mathcal{F}))$  for  $\mathcal{F}$  (Theorem 3.2). The set  $B(\mathcal{F})$  has a structure of a colored oriented graph which Kashiwara [5] calls a crystal graph. Let  $B(\mathcal{F})_\phi$  be the connected component of  $\phi$  (the empty Young diagram) in the graph of  $B(\mathcal{F})$ . The submodule of  $\mathcal{F}$  generated by  $\phi$  is the basic  $U_q(\widehat{\mathfrak{sl}}(n))$ -module  $M(\Lambda_0)$  with highest weight  $\Lambda_0$ . We can identify  $B(\mathcal{F})_\phi$  with the set of paths  $\mathcal{P}(\Lambda_0)$  (see [2]). As shown in [1, 2], the number of  $\Lambda_0$ -paths with weight  $\mu$  is equal to  $\dim M(\Lambda_0)_\mu$ . Using these we prove Kashiwara’s conjecture for  $M(\Lambda_0)$  (Theorem 4.7).

**1. Preliminaries**

We follow the notations in [5]. We recall some essential facts. The  $q$ -analogue  $U_q(\widehat{\mathfrak{sl}}(n))$  of the enveloping algebra of the affine Lie algebra  $\widehat{\mathfrak{sl}}(n)$  is generated by  $\{e_i, f_i, t_i^\pm = q^{\pm h_i} \mid 0 \leq i \leq n-1\}$ . These generators satisfy the following important relations:

$$[e_i, f_j] = \delta_{ij} \left( \frac{t_i^+ - t_i^-}{q - q^{-1}} \right), \tag{1.1}$$

$$t_i^+ e_j t_i^- = q^{2(\alpha_i, \alpha_j)} e_j, \tag{1.2}$$

and

$$t_i^+ f_j t_i^- = q^{-2(\alpha_i, \alpha_j)} f_j, \tag{1.3}$$

where  $(\alpha_i, \alpha_i) = 1$ . We will need the algebra  $U_q(\mathfrak{gl}(\infty))$  which is generated by  $\{e_i^\infty, f_i^\infty, t_i^{\pm \infty} = q^{\pm h_i^\infty} \mid i \in \mathbf{Z}\}$  (see [3]). These generators also satisfy the corresponding relations (1.1)–(1.3).

Let  $M$  be any integrable  $U_q(\widehat{\mathfrak{sl}}(n))$ -module. For each  $i = 0, 1, \dots, n-1$ , let  $U_q(\mathfrak{sl}(2))_{(i)}$  denote the subalgebra of  $U_q(\widehat{\mathfrak{sl}}(n))$  generated by  $e_i, f_i$  and  $t_i^\pm$ . Note that  $M$  is a union of finite-dimensional representations over  $U_q(\mathfrak{sl}(2))_{(i)}$ . Kashiwara [5] defines the following operators on  $M$ :

$$\tilde{e}_i = (qt_i^+ \Delta_i)^{-1/2} e_i, \quad \text{and} \quad \tilde{f}_i = t_i^+ (qt_i^+ \Delta_i)^{-1/2} f_i, \tag{1.4}$$

for  $i = 0, 1, \dots, n-1$ , where  $\Delta_i$  is certain element in the center of  $U_q(\mathfrak{sl}(2))_{(i)}$ . The action of the operator  $(qt_i^+ \Delta_i)^{-1/2}$  is given as follows. Let  $v$  be a weight vector in an  $(l+1)$ -dimensional irreducible  $U_q(\mathfrak{sl}(2))_{(i)}$  submodule of  $M$ . Suppose  $t_i^+ v = q^{l-2k} v$ . Then

$$\Delta_i v = (q^{l+1} - 2 + q^{-l-1}) v, \tag{1.5}$$

and

$$(qt_i^+ \Delta_i)^{-1/2} v = q^k (1 - q^{l+1})^{-1} v. \tag{1.6}$$

Let  $K = \mathbf{Q}(q)$  and  $A$  be the ring of rational functions in  $q$  without pole at  $q = 0$ . Let  $L$  be a free  $A$ -module such that  $K \otimes_A L \cong M$  and let  $B$  be a base of the  $\mathbf{Q}$ -vector space  $L/qL$ . The pair  $(L, B)$  is called a *crystal base* [5] of  $M$  if it satisfies the following conditions:

$$L = \bigoplus_{\lambda} L_{\lambda}, \tag{1.7}$$

where  $L_{\lambda} = L \cap M_{\lambda}$  and  $M_{\lambda}$  is the  $\lambda$ -weight space of  $M$ ,

$$B = \bigsqcup_{\lambda} B_{\lambda} \quad \text{where} \quad B_{\lambda} = B \cap (L_{\lambda}/qL_{\lambda}), \tag{1.8}$$

$$\tilde{e}_i L \subset L \quad \text{and} \quad \tilde{f}_i L \subset L, \quad \text{for all } i, \tag{1.9}$$

$$\tilde{e}_i B \subset B \cup \{0\} \quad \text{and} \quad \tilde{f}_i B \subset B \cup \{0\}, \quad \text{for all } i, \tag{1.10}$$

$$u = \tilde{e}_i v \quad \text{if and only if} \quad v = \tilde{f}_i u, \quad \text{for all } i \quad \text{and} \quad u, v \in B. \tag{1.11}$$

As noted in [5]  $B$  has a structure of colored oriented graph. The colors are labelled by  $i$  ( $0 \leq i \leq n - 1$ ). For  $u, v \in B$ ,  $u \xrightarrow{i} v$  when  $v = \tilde{f}_i u$ . This is called the crystal graph of  $M$ .

### 2. The Fock Space Representation of $U_q(\widehat{\mathfrak{sl}}(n))$

In this section we will briefly describe the Fock space representation of  $U_q(\mathfrak{gl}(\infty))$  and  $U_q(\widehat{\mathfrak{sl}}(n))$  given in [3] with appropriate modifications. For more details we refer the reader to [3].

Consider the lattice on the fourth quadrant of the  $xy$ -plane with sites  $\{(i, j) \in \mathbb{Z}^2 \mid i \geq 0, j \leq 0\}$ . We consider edges on the lattice as oriented, starting from  $(i, j)$  and ending at  $(i + 1, j)$  or  $(i, j + 1)$ , and labelled by the integer  $i + j$ . Any oriented path on this lattice determines uniquely a Young diagram  $Y$  and conversely. For example,




Fig. 1

In other words, given by Young diagram  $Y$ , we superimpose it on the lattice with upper left corner at the site  $(0, 0)$ . Let  $\mathscr{Y}$  be the set of all Young diagrams. Let  $\mathscr{F} = \sum_{Y \in \mathscr{Y}} KY$  be the  $K$ -vector space having all the Young diagrams as base vectors. A Young diagram viewed as a lattice path has several corners. We say the corner is concave or convex depending on whether it is of the form  $i-1 \downarrow \uparrow^i$  or  $\downarrow \uparrow^i$ .

The algebra  $U_q(\mathfrak{gl}(\infty))$  acts on the Fock space  $\mathscr{F}$ . The actions of its generators  $\{e_i^\infty, f_i^\infty, t_i^{\pm \infty} = q^{\pm h_i^\infty} \mid i \in \mathbb{Z}\}$  are given as follows. For  $Y \in \mathscr{Y}$ ,

$$\begin{aligned}
 e_i^\infty Y &= Y', & \text{if } Y \text{ has the convex corner } \downarrow \uparrow^i, \\
 & & \text{then } Y' \text{ is same as } Y \text{ except this corner} \\
 & & \text{becomes concave } i-1 \downarrow \uparrow^i, \\
 &= 0, & \text{otherwise,}
 \end{aligned} \tag{2.1}$$

$$\begin{aligned}
 f_i^\infty Y &= Y'', && \text{if } Y \text{ has the concave corner } i^{-1} \begin{array}{|c} \hline \lrcorner \\ \hline \end{array}, \\
 & && \text{then } Y'' \text{ is same as } Y \text{ except this corner} \\
 & && \text{becomes convex } \begin{array}{|c} \hline \lrcorner \\ \hline \end{array} i, \\
 &= 0, && \text{otherwise,}
 \end{aligned} \tag{2.2}$$

$$\begin{aligned}
 t_i^{\pm\infty} Y &= q^{\pm 1} Y, && \text{if } Y \text{ has the concave corner } i^{-1} \begin{array}{|c} \hline \lrcorner \\ \hline \end{array}, \\
 &= q^{\pm 1} Y, && \text{if } Y \text{ has the convex corner } \begin{array}{|c} \hline \lrcorner \\ \hline \end{array} i, \\
 &= Y, && \text{otherwise.}
 \end{aligned} \tag{2.3}$$

Under the above action  $\mathcal{F}$  becomes an irreducible integrable  $U_q(\mathfrak{gl}(\infty))$ -module with highest weight  $\Lambda_0$  ( $\Lambda_0(h_i^\infty) = \delta_{i,0}$ ) and highest weight vector  $\phi$  (the empty Young diagram).

As in [3] (with suitable normalization)  $\mathcal{F}$  becomes and  $U_q(\widehat{\mathfrak{sl}}(n))$ -module where the actions of the generators  $\{e_i, f_i, t_i^\pm = q^{\pm h_i} | 0 \leq i < n\}$  are given by the following equations:

$$e_i = \sum_{j \equiv i \pmod n} \left( \prod_{k \geq 1} t_{j-kn}^{+\infty} \right) e_j^\infty, \tag{2.4}$$

$$f_i = \sum_{j \equiv i \pmod n} f_j^\infty \left( \prod_{k \geq 1} t_{j+kn}^{-\infty} \right), \tag{2.5}$$

$$t_i^\pm = \prod_{j \equiv i \pmod n} t_j^{\pm\infty}. \tag{2.6}$$

Under the above action  $\mathcal{F}$  is an integrable  $U_q(\widehat{\mathfrak{sl}}(n))$ -module. However, it is not irreducible as an  $U_q(\widehat{\mathfrak{sl}}(n))$ -module. Observe that as an  $U_q(\widehat{\mathfrak{sl}}(n))$ -module the vector  $\phi \in \mathcal{F}$  is a highest weight vector with highest weight  $\Lambda_0$ , ( $\Lambda_0(h_i) = \delta_{i,0}$ ). The space  $M(\Lambda_0) = U_q(\widehat{\mathfrak{sl}}(n))\phi$  is the irreducible integrable highest weight  $U_q(\widehat{\mathfrak{sl}}(n))$ -module with highest weight  $\Lambda_0$ .

Given any Young diagram  $Y \in \mathcal{Y}$  we color the boxes in  $Y$  with  $n$  colors  $i = 0, 1, \dots, n - 1$ , as follows. The box with the upper left corner at site  $(i, j)$  is colored with  $(i + j)'$ -color where  $(i + j)' = (i + j) \pmod n$ . Then observe that the action of  $e_i$  (respectively  $f_i$ ) on  $Y \in \mathcal{Y}$  given by (2.4) (respectively (2.5)) is just removing (respectively adding) a box of color  $i$ . For  $Y \in \mathcal{Y}$ , the weight of  $Y$  (denoted by  $wt(Y)$ ) is  $\Lambda_0 - \sum_{i=0}^{n-1} m_i \alpha_i$  if  $Y$  contains  $m_i$  boxes of color  $i, 0 \leq i < n$ .

### 3. $U_q(\widehat{\mathfrak{sl}}(2))$ Decomposition of the Fock Space

For each  $i$  ( $0 \leq i < n$ ), let  $U_q(\widehat{\mathfrak{sl}}(2))_{(i)}$  denote the subalgebra of  $U_q(\widehat{\mathfrak{sl}}(n))$  generated by  $\{e_i, f_i, t_i^\pm\}$ . We say a Young diagram is *anti  $i$ -convex* if all its convex corners are non  $i$ -color. Given any Young diagram  $Y \in \mathcal{Y}$  let  $Y(i)$  denote its maximal subdiagram which is anti  $i$ -convex. Then  $Y$  is uniquely determined by the pair  $(Y(i), \varepsilon)$ , where  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m)$ ,  $m = \#\{\text{concave corners in } Y(i) \text{ of color } i\}$ ,

and 
$$\varepsilon_j = \begin{cases} 0, & \text{if the } j^{\text{th}} \text{ (counted from left to right)} \\ & \text{concave corner of color } i \text{ is vacant in } Y, \\ 1, & \text{if the } j^{\text{th}} \text{ concave corner of color } i \\ & \text{is occupied in } Y. \end{cases}$$

For example, let  $n = 2$  and let white be color 0 and black be color 1. Then by choosing  $i = 0$ , we have:

$Y$	$Y(0)$	$\varepsilon$
	$\phi$	(1)
		(0)
		(1)
		(1, 0, 0)
		(1, 1, 0)
		(1, 0, 1)

For any fixed  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m)$  we partition the set

$$\{1, 2, \dots, m\} = J \sqcup K_1 \sqcup \dots \sqcup K_t$$

into disjoint subsets by the following inductive procedure:

- (1) If there is no  $j$  such that  $(\varepsilon_j, \varepsilon_{j+1}) = (0, 1)$  then define  $J = \{1, 2, \dots, m\}$ .
- (2) If there is some  $j$  such that  $(\varepsilon_j, \varepsilon_{j+1}) = (0, 1)$  then define  $K_1 = \{j, j + 1\}$  and apply (1) and (2) to  $\{1, 2, \dots, m\} \setminus K_1$  to choose  $J$  or  $K_2$ . Repeat this as necessary.

For example, if  $\varepsilon = (1, 0, 1, 0, 0, 1, 1, 0)$ , then  $m = 8$  and  $J = \{1, 8\}$ ,  $K_1 = \{2, 3\}$ ,  $K_2 = \{5, 6\}$ ,  $K_3 = \{4, 7\}$ . Note that this partition is unique up to rearrangements of the sets  $K_s$ ,  $1 \leq s \leq t$ .

Let  $k = \#\{j \in J \mid \varepsilon_j = 1\}$ . For any Young diagram  $Y = (Y(i), \varepsilon)$ ,  $\varepsilon = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m)$ , and partition  $\{1, 2, \dots, m\} = J \sqcup K_1 \sqcup \dots \sqcup K_t$ , we define

$$[Y_i]_i = \sum_{J = J_0 \sqcup J_1 \mid |J_1| = k} \sum_{S \subseteq \{1, 2, \dots, t\}} q^{\#(J_0, J_1)} (-q)^{|S|} (Y(i), \varepsilon(J_0, J_1, S)),$$

where

$$\begin{aligned} \#(J_0, J_1) &= \#\{(j, j') \mid j < j', j \in J_0, j' \in J_1\} \\ \varepsilon(J_0, J_1, S) &= (\tau_1, \tau_2, \dots, \tau_m) \end{aligned}$$

such that

$$\tau_j = 0 \quad \text{if } j \in J_0, \quad \tau_j = 1 \quad \text{if } j \in J_1$$

and for  $j < j'$ ,  $\{j, j'\} \in K_s$ ,

$$(\tau_j, \tau_{j'}) = (1, 0) \quad \text{if } s \in S$$

and

$$(\tau_j, \tau_{j'}) = (0, 1) \quad \text{if } s \notin S.$$

For example, if  $Y = (Y(i), (1, 0, 1, 0))$ , then

$$[Y]_i = (Y(i), (1, 0, 1, 0)) + q\{(Y(i), (0, 0, 1, 1)) - (Y(i), (1, 1, 0, 0))\} - q^2(Y(i), (0, 1, 0, 1)).$$

**Theorem 3.1.** Fix an anti  $i$ -convex Young diagram  $Y(i)$  and a partition  $\{1, 2, \dots, m\} = J \sqcup K_1 \sqcup \dots \sqcup K_l$  such that  $|J| = l$ . For each  $k = 0, 1, \dots, l$  there is a unique diagram  $Y_k$  with the data  $(Y(i), J, K_1, \dots, K_l)$  such that  $\#\{j \in J \mid \varepsilon_j = 1\} = k$ . Furthermore,  $V_l = \bigoplus_{k=0}^l K[Y_k]_i$  is the  $(l + 1)$ -dimensional irreducible integrable  $U_q(\mathfrak{sl}(2))_{(i)}$ -module with highest weight vector  $[Y_0]_i$ . Set  $L_l = \bigoplus_{k=0}^l A[Y_k]_i$  and  $B_l = \{[Y_k]_i \mid k = 0, 1, \dots, l\}$ . Then  $(L_l, B_l)$  is a crystal base for the  $U_q(\mathfrak{sl}(2))_{(i)}$ -module  $V_l$ .

*Proof.* The first assertion is clear, for if  $J = \{i_1, \dots, i_l\}$  then there are precisely  $l + 1$  choices for  $(\varepsilon_{i_1}, \varepsilon_{i_2}, \dots, \varepsilon_{i_l})$ , namely,  $(0, 0, \dots, 0), (1, 0, \dots, 0), (1, 1, 0, \dots, 0), \dots, (1, 1, \dots, 1)$ .

Now observe that if  $Y = (Y(i), (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_m))$ , then the action of  $e_i$  and  $f_i$  on  $Y$  are given by the following formulas:

$$e_i Y = \sum_{\substack{j \\ \varepsilon_j = 1}} q^{\#\{j' \mid j' < j, \varepsilon_{j'} = 0\} - \#\{j' \mid j' < j, \varepsilon_{j'} = 1\}} (Y(i), (\varepsilon_1, \dots, \varepsilon_j - 1, \dots, \varepsilon_m)), \tag{3.2}$$

$$f_i Y = \sum_{\substack{j \\ \varepsilon_j = 0}} q^{\#\{j' \mid j' > j, \varepsilon_{j'} = 1\} - \#\{j' \mid j' > j, \varepsilon_{j'} = 0\}} (Y(i), (\varepsilon_1, \dots, \varepsilon_j + 1, \dots, \varepsilon_m)). \tag{3.3}$$

It follows from (3.2) and (3.3) that when  $e_i$  or  $f_i$  act on  $[Y]_i$  (see (3.1)) the terms corresponding to  $j \in \{r, r + p\} = K_s$  for any  $s \in \{1, 2, \dots, t\}$  cancel each other. Furthermore, for any  $j \in J$  the sum of the contributions of  $j' \in \{r, r + p\} = K_s$  for any  $s \in \{1, 2, \dots, t\}$ , to the exponent of  $q$  in (3.2) or (3.3) is zero. Hence in order to compute  $e_i[Y_k]_i$  or  $f_i[Y_k]_i$  for any  $k = 0, 1, \dots, l$ , without loss of generality we can and do assume  $t = 0, m = l$  so  $J = \{1, 2, \dots, l\}$ . Then by (3.1) we have

$$[Y_k]_i = \sum_{\substack{J = J_0 \sqcup J_1 \\ |J_0| = l - k \\ |J_1| = k}} q^{\#(J_0, J_1)} (Y(i), \varepsilon(J_0, J_1)), \tag{3.4}$$

where

$$\#(J_0, J_1) = \#\{(j, j') \mid j < j', j \in J_0, j' \in J_1\}$$

and

$$\varepsilon(J_0, J_1) = (\tau_1, \tau_2, \dots, \tau_l)$$

such that  $\tau_j = 0$  if  $j \in J_0$  and  $\tau_j = 1$  if  $j \in J_1$ .

Now applying  $e_i$  (respectively  $f_i$ ) to Eq. (3.4) and using formula (3.2) (respectively (3.3)) we easily get, for  $0 \leq k \leq l$ ,

$$e_i[Y_k]_i = q^{-(k-1)}(1 + q^2 + q^4 + \dots + q^{2(l-k)})[Y_{k-1}]_i, \tag{3.5}$$

$$f_i[Y_k]_i = q^{-l+k+1}(1 + q^2 + q^4 + \dots + q^{2k})[Y_{k+1}]_i, \tag{3.6}$$

where  $[Y_{-1}]_i = 0$ , and  $[Y_{l+1}]_i = 0$ .

Observe that (see [5]) by the definitions,

$$t_i^+[Y_k]_i = q^{l-2k}[Y_k]_i, \tag{3.7}$$

and

$$(qt_i^+ \Delta_i)^{-1/2}[Y_k]_i = q^k(1 - q^{l+1})^{-1}[Y_k]_i. \tag{3.8}$$

Now it follows from (3.5)–(3.8) that for  $1 \leq k \leq l$  we have

$$\begin{aligned}\tilde{e}_i[Y_k]_i &= (qt_i^+ \Delta_i)^{-1/2} e_i[Y_k]_i \\ &= (1 - q^{l+1})^{-1} (1 + q^2 + q^4 + \cdots + q^{2(l-k)}) [Y_{k-1}]_i,\end{aligned}\quad (3.9)$$

and

$$\begin{aligned}\tilde{f}_i[Y_k]_i &= t_i^+ (qt_i^+ \Delta_i)^{-1/2} f_i[Y_k]_i \\ &= (1 - q^{l+1})^{-1} (1 + q^2 + q^4 + \cdots + q^{2k}) [Y_{k+1}]_i.\end{aligned}\quad (3.10)$$

Hence the theorem follows. ■

**Theorem 3.2.** *Let  $L(\mathcal{F}) = \bigoplus_{Y \in \mathcal{Y}} AY$  and  $B(\mathcal{F}) = \mathcal{Y}$ . Then the pair  $(L(\mathcal{F}), B(\mathcal{F}))$  is a crystal base for the integrable  $U_q(\widehat{\mathfrak{sl}}(n))$ -module  $\mathcal{F}$ .*

*Proof.* By the definition (see (3.1)), for each  $0 \leq i \leq n-1$ ,  $[Y]_i \in L(\mathcal{F})$  and  $[Y]_i = Y + q \sum_{Y'} a_{Y'} Y'$ ,  $a_{Y'} \in A$ . For any weight  $\mu$ , the  $\mu$ -weight space  $\mathcal{F}_\mu$  of  $\mathcal{F}$  is finite-dimensional. Suppose  $\dim_{\mathbf{K}}(\mathcal{F}_\mu) = n_\mu$ . Choose a basis  $\{Y_1, Y_2, \dots, Y_{n_\mu}\}$  of  $\mathcal{F}_\mu$ . Then for each  $i$ ,

$$([Y_1]_i, \dots, [Y_{n_\mu}]_i) = (Y_1, \dots, Y_{n_\mu})(I + qX_i), \quad (3.11)$$

where  $X_i$  is a  $n_\mu \times n_\mu$  matrix with coefficients in  $A$ . Since  $I + qX_i$  is invertible in  $A$ , it follows from (3.11) that for each  $i$ ,

$$Y = [Y]_i + q \sum_{Y''} b_{Y''} [Y'']_i, \quad b_{Y''} \in A. \quad (3.12)$$

So by using Theorem 3.1, we have

$$\tilde{e}_i Y = \tilde{e}_i [Y]_i + q \sum_{Y''} b_{Y''} \tilde{e}_i [Y'']_i \in L(\mathcal{F}),$$

and

$$\tilde{f}_i Y = \tilde{f}_i [Y]_i + q \sum_{Y''} b_{Y''} \tilde{f}_i [Y'']_i \in L(\mathcal{F}),$$

which gives the required result. ■

The next proposition is an immediate consequence of Theorems (3.1), (3.2) and the definition of crystal graph (see [5]).

**Proposition 3.3.** *Let  $Y, Y' \in B(\mathcal{F})$ . In the crystal graph of  $B(\mathcal{F})$ ,  $Y \xrightarrow{i} Y'$  if and only if*

- i)  $Y = (Y(i), (\varepsilon_1, \dots, \varepsilon_m))$ ,  $Y' = (Y(i), (\varepsilon'_1, \dots, \varepsilon'_m))$ ,
- ii) the partition  $\{1, 2, \dots, m\} = J \sqcup K_1 \sqcup \cdots \sqcup K_t$  is the same for both  $Y$  and  $Y'$ ,
- iii)  $\varepsilon_j = \varepsilon'_j$  for  $j \neq r$  for some  $r \in J$ ,  $\varepsilon_r = 0$ ,  $\varepsilon'_r = 1$ , and for  $j \in J$ ,  $\varepsilon_j = \varepsilon'_j = 1$  if  $j < r$ ,  $\varepsilon_j = \varepsilon'_j = 0$  if  $j > r$ .

#### 4. Crystal Base for $M(\Lambda_0)$

As in Sect. 2, let  $M(\Lambda_0) \subset \mathcal{F}$  be the irreducible highest weight  $U_q(\widehat{\mathfrak{sl}}(n))$ -module with highest weight  $\Lambda_0$  and highest weight vector  $\phi$ . For  $Y \in \mathcal{Y}$ , let  $[f_1, f_2, \dots, f_m]$  denote the signature of  $Y$ . Let  $(g_1, g_2, \dots, g_m)$  be the largest  $m$ -tuple of nonnegative

integers (in lexicographic ordering) such that:

$$(1) \quad g_1 \geq g_2 \geq \dots \geq g_m,$$

and

$$(2) \quad f_1 - ng_1 \geq f_2 - ng_2 \geq \dots \geq f_m - ng_m \geq 0.$$

Note that they are determined by the recursive formula

$$g_j = g_{j+1} + \left\lfloor \frac{f_j - f_{j+1}}{n} \right\rfloor, \quad ([x] \text{ denotes the integral part}),$$

where  $g_j = 0$  for  $j > m$ . Define  $\sigma(Y)$  to be the Young diagram with signature  $[ng_1, ng_2, \dots, ng_m]$ . We say  $\sigma(Y)$  is the  $\sigma$ -component of  $Y$ .

**Lemma 4.1.** *For  $Y, Y' \in \mathcal{Y}$ ,  $Y \xrightarrow{i} Y'$  for some  $i$ , implies that  $\sigma(Y) = \sigma(Y')$ .*

*Proof.* Recall Proposition 3.3 which gives the condition for  $Y \xrightarrow{i} Y'$ . Suppose that  $\sigma(Y)$  has signature  $[ng_1, \dots, ng_m]$  and  $\sigma(Y')$  has signature  $[ng'_1, \dots, ng'_m]$ . If  $\sigma(Y) \neq \sigma(Y')$ , then  $g_j \neq g'_j$  for some  $j$ . Let  $[f_1, f_2, \dots, f_p]$  be the signature of  $Y(i)$ . Then we must have the following situation:

(1)  $f_j - f_{j+1} \equiv (n-1) \pmod n$  and we have concave corners of color  $i$  at the end of the  $j^{\text{th}}$  and  $(j+1)^{\text{th}}$  rows of  $Y(i)$ . Suppose that these are the  $r^{\text{th}}$  and  $(r+1)^{\text{th}}$  corners in the decomposition of  $Y = (Y(i), \varepsilon)$ .

(2) (i) Either  $\varepsilon_r = \varepsilon'_r = 0$ ,  $\varepsilon_{r+1} = 0$  and  $\varepsilon'_{r+1} = 1$ ,

(ii) or  $\varepsilon_{r+1} = \varepsilon'_{r+1} = 0$ ,  $\varepsilon_r = 0$  and  $\varepsilon'_r = 1$ .

In the case of (i),  $r+1 \in J$  since  $\varepsilon_{r+1} = 0$  in  $Y$  and  $\varepsilon_{r+1} = 1$  in  $Y'$ . But in  $Y'$ ,  $(\varepsilon_r, \varepsilon_{r+1}) = (0, 1)$ , hence  $\{r, r+1\} \in K_s$  for some  $s$ , which is a contradiction. Similarly, (ii) also leads to contradiction. ■

**Lemma 4.2.** *For any  $Y \in \mathcal{Y}$ ,  $Y = \sigma(Y)$  if and only if  $Y$  is highest in the sense of crystal graph (i.e., there is no  $Y' \in \mathcal{Y}$  such that  $Y' \xrightarrow{i} Y$ ).*

*Proof.* Let  $[f_1, f_2, \dots, f_k]$  be the signature of  $Y$ . Suppose  $Y = \sigma(Y)$ . Then  $f_j - f_{j+1} \equiv 0 \pmod n$ . Hence the color of the last box of each row is the same as the color of the concave corner of the subsequent row. So for any fixed  $i$ , if  $Y = (Y(i), (\varepsilon_1, \dots, \varepsilon_m))$  with the partition  $\{1, 2, \dots, m\} = J \sqcup K_1 \sqcup \dots \sqcup K_r$ , then  $J = \{m\}$  with  $\varepsilon_m = 0$  or  $J = \phi(\text{empty})$ . In either case  $Y$  is highest in the sense of crystal graph.

Now suppose  $Y \neq \sigma(Y)$ . Let  $[ng_1, \dots, ng_k]$  be the signature of  $\sigma(Y)$ . Then  $f_j \neq ng_j$  for some  $j$ . Assume  $j$  to be the largest integer such that  $f_j \neq ng_j$ . Let  $i$  be the color of the last box in the  $j^{\text{th}}$  row of  $Y$ . Then  $Y = (Y(i), (\varepsilon_1, \dots, \varepsilon_m))$ ,  $\{1, 2, \dots, m\} = J \sqcup K_1 \sqcup \dots \sqcup K_r$  and  $\varepsilon_r = 1$ ,  $r \in J$ , where the  $r^{\text{th}}$  corner of color  $i$  which is occupied corresponds to the last box of the  $j^{\text{th}}$  row in  $Y$ . Hence by Proposition 3.3 we can find  $Y' \in \mathcal{Y}$  such that  $Y' \xrightarrow{i} Y$ . So  $Y$  cannot be highest. ■

**Proposition 4.3.** *Let  $Z \in \mathcal{Y}$  such that  $\sigma(Z) = Z$ . Let  $B(\mathcal{F})_Z$  denote the connected component of  $Z$  in the crystal graph of  $B(\mathcal{F})$ . Then  $B(\mathcal{F})_Z = \{Y \in \mathcal{Y} \mid \sigma(Y) = Z\}$ .*

*Proof.* It follows from Lemma 4.1 that  $B(\mathcal{F})_Z \subseteq \{Y \in \mathcal{Y} \mid \sigma(Y) = Z\}$ . Now suppose  $Y \in \mathcal{Y}$  and  $\sigma(Y) = Z$ . We want to show that  $Y \in B(\mathcal{F})_Z$ . If  $Y = Z$  then there is nothing



to prove. If  $Y \neq Z$ , then by Lemma 4.2, there exists  $Y_1 \in \mathcal{Y}$  such that  $Y_1 \rightarrow Y$ . Hence using induction we get  $Y \in B(\mathcal{F})_Z$  as desired. ■

Now define

$$L = \sum_{0 \leq i_1, i_2, \dots, i_k \leq n-1} A \tilde{f}_{i_1} \tilde{f}_{i_2} \cdots \tilde{f}_{i_k} \phi \quad (4.1)$$

and

$$B = \{v = \tilde{f}_{i_1} \tilde{f}_{i_2} \cdots \tilde{f}_{i_k} \phi \in L/qL \mid v \neq 0\}. \quad (4.2)$$

Let  $M(\Lambda_0) \subset \mathcal{F}$  denote the irreducible integrable  $U_q(\hat{\mathfrak{sl}}(n))$ -module with highest weight  $\Lambda_0$  (i.e.,  $\Lambda_0(t_i^\pm) = q^{\pm 1} \delta_{i,0}$ ) and highest weight vector  $\phi$ . Then  $M(\Lambda_0) = U_q(\hat{\mathfrak{sl}}(n))\phi$ .

**Proposition 4.4.**  $M(\Lambda_0) = K \otimes_A L$ .

*Proof.* By definition,  $M(\Lambda_0) \supset K \otimes_A L$ . For  $\mu = \Lambda_0 - \sum_{i=0}^{n-1} m_i \alpha_i$  let  $M(\Lambda_0)_\mu$  denote the  $\mu$  weight space of  $M(\Lambda_0)$ . By Theorem 5.4 in [2] (also see Theorem in [1])  $\dim(M(\Lambda_0)_\mu) = \#\mathcal{P}(\Lambda_0)_\mu$ , where  $\mathcal{P}(\Lambda_0)_\mu$  denotes the set of  $\Lambda_0$ -paths of weight  $\mu$ . But there is a one-to-one correspondence between  $\mathcal{P}(\Lambda_0)_\mu$  and the set  $\{Y \in B(\mathcal{F})_\phi \mid \text{wt}(Y) = \mu\}$ . (See [2]. Young diagrams in this paper and those in [2] are transposed to each other.) For any  $Y \in B(\mathcal{F})_\phi$  by Proposition 4.3 there exists some  $(i_1, i_2, \dots, i_k)$  such that  $Y = \tilde{f}_{i_1} \tilde{f}_{i_2} \cdots \tilde{f}_{i_k} \phi$  in  $L(\mathcal{F})/qL(\mathcal{F})$ . Hence  $\dim_K(M(\Lambda_0)_\mu) \leq \dim_K(K \otimes_A L)_\mu$  for each weight  $\mu$ . Therefore,  $M(\Lambda_0) = K \otimes_A L$ . ■

**Lemma 4.5.**  $(K \otimes_A L) \cap L(\mathcal{F}) = L$ .

*Proof.* Let  $L_0 = \sum_{Y \in B(\mathcal{F})_\phi} A \tilde{f}_{i_1} \tilde{f}_{i_2} \cdots \tilde{f}_{i_k} \phi$ , where for each  $Y \in B(\mathcal{F})_\phi$  we choose a sequence  $(i_1, i_2, \dots, i_k)$  such that  $\tilde{f}_{i_1} \tilde{f}_{i_2} \cdots \tilde{f}_{i_k} \phi = Y$  in  $L(\mathcal{F})/qL(\mathcal{F})$ . Then by an argument similar to the proof of Proposition 4.4 we get  $M(\Lambda_0) = K \otimes_A L_0$ . Hence  $K \otimes_A L = K \otimes_A L_0$ .

Clearly  $L \subset (K \otimes_A L) \cap L(\mathcal{F}) = (K \otimes_A L_0) \cap L(\mathcal{F})$ . Now let  $v \in (K \otimes_A L_0) \cap L(\mathcal{F})$ . Then  $v \in L(\mathcal{F})$  and  $v \in K \otimes_A L_0 = M(\Lambda_0)$ . Let  $v \in M(\Lambda_0)_\mu$  for some weight  $\mu$  and  $\dim(M(\Lambda_0)_\mu) = n_\mu$ . Then  $v = \sum_{i=1}^{n_\mu} c_i y_i$ ,  $c_i \in K$ ,  $y_i \in L_0$ . Also since  $v \in L(\mathcal{F})$ , we have

$$v = \sum_{i=1}^{n_\mu} a_i Y_i, \quad a_i \in A, \quad Y_i \in \mathcal{Y}.$$

$$(y_1, y_2, \dots, y_{n_\mu}) = (Y_1, Y_2, \dots, Y_{n_\mu})(I + qX),$$

where  $X$  is an  $n_\mu \times n_\mu$  matrix with coefficients in  $A$ . Hence

$$\begin{aligned} v &= (Y_1, Y_2, \dots, Y_{n_\mu}) \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_{n_\mu} \end{pmatrix} \\ &= (y_1, y_2, \dots, y_{n_\mu}) \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_{n_\mu} \end{pmatrix} \end{aligned}$$

$$= (Y_1, Y_2, \dots, Y_{n_\mu})(I + qX) \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_{n_\mu} \end{pmatrix}.$$

So

$$\begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_{n_\mu} \end{pmatrix} = (I + qX) \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_{n_\mu} \end{pmatrix}.$$

But since  $a_i \in A$ ,  $i = 1, 2, \dots, n_\mu$  and  $I + qX$  is invertible in  $A$ , it follows that  $c_i \in A$ . Hence  $v = \sum c_i y_i \in L$ , which completes the proof. ■

**Corollary 4.6.**  $L/qL \subset L(\mathcal{F})/qL(\mathcal{F})$ . So  $B$  is a subset of  $B(\mathcal{F})$ .

*Proof.* It is enough to show that  $qL(\mathcal{F}) \cap L = qL$ . It follows from Proposition 4.4 and Lemma 4.5 that

$$\begin{aligned} qL(\mathcal{F}) \cap L &= qL(\mathcal{F}) \cap L(\mathcal{F}) \cap M(\Lambda_0) \\ &= qL(\mathcal{F}) \cap M(\Lambda_0) \\ &= q(L(\mathcal{F}) \cap M(\Lambda_0)) = qL. \quad \blacksquare \end{aligned}$$

**Theorem 4.7.** The pair  $(L, B)$  is a crystal base for the irreducible integrable highest weight  $U_q(\hat{\mathfrak{sl}}(n))$ -module  $M(\Lambda_0)$ .

*Proof.* Let  $v \in L$ . By the definition  $\tilde{f}_i v \in L$  for all  $i = 0, 1, \dots, n - 1$ . For each  $i$ ,  $\tilde{e}_i v \in M(\Lambda_0) = K \otimes_A L$ . Since  $L \subseteq L(\mathcal{F})$ , by Theorem 3.2  $\tilde{e}_i v \in L(\mathcal{F})$ . Hence by Lemma 4.5,  $\tilde{e}_i v \in (K \otimes_A L) \cap L(\mathcal{F}) = L$ . Now the result follows from Theorem 3.2, Proposition 4.4 and Corollary 4.6. ■

*Acknowledgements.* The authors thank T. Hayashi, M. Jimbo, M. Kashiwara and M. Okado for valuable discussions. The first author acknowledges the warm hospitality at RIMS, Kyoto University during his stay in Kyoto which was supported by JSPS fellowship.

**References**

1. Date, E., Jimbo, M., Kuniba, A., Miwa, T., Okado, M.: One-dimensional configuration sums in vertex models and affine Lie algebra characters. *Lett. Math. Phys.* **17**, 69–77 (1989)
2. Date, E., Jimbo, M., Kuniba, A., Miwa, T., Okado, M.: Paths, Maya diagrams and representations of  $\hat{\mathfrak{sl}}(r, \mathbb{C})$ . In: *Integrable systems in quantum field theory and statistical mechanics*. *Adv. Stud. Pure Math.* **19**, 149–191 (1989)
3. Hayashi, T.:  $Q$ -analogues of Clifford and Weyl algebras-spinor and oscillator representations of quantum enveloping algebras.
4. Jimbo, M., Miwa, T.: Solitons and infinite dimensional Lie algebras. *Publ. RIMS, Kyoto University* **19**, 943–1001 (1983)
5. Kashiwara, M.: Crystalizing the  $q$ -analogue of universal enveloping algebras. *RIMS, Kyoto University*, preprint RIMS-676 (1989)

Communicated by H. Araki