

Crystal chemistry of the magnetite-ulvöspinel series

FERDINANDO BOSI,^{1,2,*} ULF HÅLENIUS,¹ AND HENRIK SKOGBY¹

¹Department of Mineralogy, Swedish Museum of Natural History, Box 50007, 10405 Stockholm, Sweden

²Dipartimento di Scienze della Terra, Sapienza Università di Roma, P.le A. Moro, 5, I-00185 Rome, Italy

ABSTRACT

Spinel single crystals of 19 compositions along the magnetite-ulvöspinel join were synthesized by use of a flux-growth method. To obtain quantitative site populations, the crystals were analyzed by single-crystal X-ray diffraction, electron-microprobe techniques, and Mössbauer spectroscopy. All results were processed by using an optimization model.

The unit-cell parameter, oxygen fractional coordinate, and tetrahedral bond length increase with increasing ulvöspinel component, whereas the octahedral bond length decreases marginally. These changes result in sigmoidal crystal-chemical relationships consistent with cation substitutions in fully occupied sites. As a first approximation, the Akimoto model ${}^T(\text{Fe}_{1-x}^{3+}\text{Fe}_x^{2+})^M(\text{Fe}^{2+}\text{Fe}_{1-x}^{3+}\text{Ti}_x)\text{O}_4$ describes the cation substitutions. Deviations from this model can be explained by an electron exchange reaction ${}^T\text{Fe}^{2+} + {}^M\text{Fe}^{3+} = {}^T\text{Fe}^{3+} + {}^M\text{Fe}^{2+}$, which causes ${}^M\text{Fe}^{2+} \neq 1$ and ${}^T\text{Fe}^{2+}/\text{Ti} \neq 1$. The resultant S-shaped trends may be related to a directional change in the electron exchange reaction at $\text{Ti} \approx 0.7$ apfu. In general, variations in structural parameters over the whole compositional range can be split into two contributions: (1) a linear variation due to the ${}^T\text{Fe}^{3+} + {}^M\text{Fe}^{3+} = {}^T\text{Fe}^{2+} + {}^M\text{Ti}^{4+}$ chemical substitution and (2) non-linear variations caused by the internal electron exchange reaction.

In accordance with bond-valence theory, strained bonds ascribable to steric effects characterize the structure of magnetite-ulvöspinel crystals. To relax the bonds and thereby minimize the internal strain under retained spinel space group symmetry, the electron exchange reaction occurs.

Keywords: Spinel, Mössbauer spectroscopy, crystal synthesis, crystal structure