



CX₃CR1 deficiency alters hippocampal-dependent plasticity phenomena blunting the effects of enriched environment

Laura Maggi^{1*}, Maria Scianni¹, Igor Branchi², Ivana D'Andrea², Clotilde Lauro¹ and Cristina Limatola^{1,3}

¹ Department of Physiology and Pharmacology, Istituto Pasteur Fondazione Cenci Bolognetti, Università di Roma, Rome Italy

² Department of Cellular Biology and Neurosciences, Istituto Superiore di Sanità, Rome, Italy

³ Istituto di Ricovero e Cura a Carattere Scientifico Neuromed, Pozzilli, Italy

Edited by:

Enrico Cherubini, International School for Advanced Studies, Italy

Reviewed by:

Nicoletta Berardi, University of Florence, Italy

Hélène Marie, Centre National de la Recherche Scientifique, France

*Correspondence:

Laura Maggi, Dipartimento di Fisiologia e Farmacologia, Università di Roma Sapienza, Piazzale Aldo Moro, 5, Rome 00185, Italy.
e-mail: laura.maggi@uniroma1.it

In recent years several evidence demonstrated that some features of hippocampal biology, like neurogenesis, synaptic transmission, learning, and memory performances are deeply modulated by social, motor, and sensorial experiences. Fractalkine/CX₃CL1 is a transmembrane chemokine abundantly expressed in the brain by neurons, where it modulates glutamatergic transmission and long-term plasticity processes regulating the intercellular communication between glia and neurons, being its specific receptor CX₃CR1 expressed by microglia. In this paper we investigated the role of CX₃CL1/CX₃CR1 signaling on experience-dependent hippocampal plasticity processes. At this aim wt and CX₃CR1^{GFP/GFP} mice were exposed to long-lasting-enriched environment (EE) and the effects on hippocampal functions were studied by electrophysiological recordings of long-term potentiation of synaptic activity, behavioral tests of learning and memory in the Morris water maze paradigm and analysis of neurogenesis in the subgranular zone of the dentate gyrus (DG). We found that CX₃CR1 deficiency increases hippocampal plasticity and spatial memory, blunting the potentiating effects of EE. In contrast, exposure to EE increased the number and migration of neural progenitors in the DG of both wt and CX₃CR1^{GFP/GFP} mice. These data indicate that CX₃CL1/CX₃CR1-mediated signaling is crucial for a normal experience-dependent modulation of hippocampal functions.

Keywords: CX₃CL1, long-term potentiation, learning and memory, neurogenesis, experience

INTRODUCTION

The importance of the environment in the regulation of brain physiology and behavior has long been recognized in biological, social, and medical science. The chance to live in conditions where social interactions, but also sensory and motor inputs are potentiated (enriched environment, EE), increases brain plasticity and could be relevant to prevent or reduce the alterations in cognitive performances occurring upon aging and in several neurodegenerative diseases (van Praag et al., 1999; Nithianantharajah and Hannan, 2006). The brain of animals maintained under EE conditions has been shown to undergo molecular and morphological changes leading to: (i) modification in neuronal structure, connections, and efficacy (Globus et al., 1973; Green and Greenough, 1986; Kempermann et al., 1997); (ii) reduction of anxiety, fear, stress, and excitability (Engellenner et al., 1982); (iii) enhanced learning and memory (Escorihuela et al., 1995; Kempermann et al., 1997; van Praag et al., 1999; Rampon and Tsien, 2000). The crucial ability of the CNS to encode and retain memories is based on activity-dependent forms of synaptic plasticity and, in the mammalian brain, hippocampal long-term potentiation (LTP) is believed to be related to the storage of declarative memory (Squire and Zola-Morgan, 1991). In particular, the CA1 area of the hippocampus is critical for information processing linked to the acquisition and consolidation of spatial memories (Tsien et al., 1996). Exploration

of spatially complex or EE elicits, in the CA1 hippocampal neurons, patterns of electrical activity similar to those of electrical stimulation used to experimentally induce LTP in hippocampal slices (Otto et al., 1991). In addition, hippocampal slices from the brain of “enriched” mice show enhanced LTP in the Schaffer collateral pathway of area CA1 (Duffy et al., 2001), suggesting that exposure to EE may modify synaptic physiology in hippocampal neurons (Foster et al., 1996; van Praag et al., 1999). It has been reported that exposure of mice to EE for several weeks modifies learning and memory abilities in the hippocampus. In particular, mice exposed to a prolonged period of EE, perform better in the Morris water maze test compared with age-matched mice reared in standard cages (van Praag et al., 1999). However, the factors mediating the effect of enrichment on behavior have been only partially identified. Among these factors, cytokines appear to exert an important function (Goshen et al., 2009) however, the role of CHEMOTactic cytoKINES (chemokines) has not been explored, yet. Beside their role in the immune system, chemokines play neuromodulatory roles on brain functions being constitutively expressed in the brain both in glial cells and neurons (Asensio and Campbell, 1999). In particular, chemokines contribute to cell-to-cell communication (Tran and Miller, 2003), modulating neurotransmitter release and plasticity (Giovannelli et al., 1998; Meucci et al., 1998; Limatola et al., 2000; Bezzi et al., 2001; Ragozzino et al., 2006; Rostène et al.,

2007; Maggi et al., 2009), and modifying the functional properties of ionic channels (Meucci et al., 1998; Lax et al., 2002; Oh et al., 2002).

Although the mouse brain is completely formed few weeks after birth, it maintains some degrees of plasticity throughout life, including axonal remodeling, synaptogenesis, neurogenesis, migration, and integration in pre-existing circuits. The dentate gyrus (DG) of the hippocampus is one of the main neurogenic niches in the adult brain. The progenitor cells in the subgranular zone (SGZ) divide, generating neuronal precursors that express doublecortin (DCX, a microtubule-associated protein required for immature neural cells migration, Brown et al., 2003) and migrate for a short distance before they differentiate and integrate into the pre-existing neuronal networks (van Praag et al., 2002). A large number of studies suggests that EE and physical exercise stimulate the appearance of new neurons in the hippocampal DG (Kempermann and Gage, 1999; van Praag et al., 1999). The neurogenesis in the DG has been correlated with the function of pattern separation, an essential step in information processing to avoid memory interferences (Deng et al., 2010). The correct migration of neural precursors generated in DG requires, among the other factors, the timely expression of the chemokine CXCL12 and its specific receptors CXCR4 and CXCR7 (Sánchez-Alcañiz et al., 2011). Much less is known on the role of CX₃CL1/CX₃CR1 signaling in DG neurogenesis. CX₃CL1 is constitutively expressed in several brain area while its specific receptor CX₃CR1 is expressed by microglia and, in the hippocampus, it has been shown to modulate synaptic transmission between Schaffer collaterals and pyramidal neurons in the CA1 region, inducing AMPA-mediated excitatory synaptic depression and negatively modulating LTP (Bertollini et al., 2006; Ragozzino et al., 2006; Maggi et al., 2009). A recent study demonstrates reduced neurogenesis in the SGZ of CX₃CR1^{-/-} mice and potentiating effects of exogenous CX₃CL1 on DG neurogenesis in aged mice, which also have lower level of hippocampal CX₃CL1 expression (Bachstetter et al., 2011). In the present study we characterized the role of CX₃CR1/CX₃CL1 signaling in modulating neural plasticity, learning abilities and DG neurogenesis in a mouse model of EE.

MATERIALS AND METHODS

ANIMALS

Three-week-old female mice (*wt*: C57BL/6J and CX₃CR1^{GFP/GFP} on the C57BL/6J background from the Jackson Laboratory, Charles River) were either housed in standard environment (SE) or EE. In CX₃CR1^{GFP/GFP} mice, CX₃CR1 gene was replaced by a green fluorescent protein (GFP) reporter gene (Jung et al., 2000). Littermates *wt* and CX₃CR1^{GFP/GFP}, obtained from CX₃CR1^{+/GFP} × CX₃CR1^{+/GFP} heterozygotes, were genotyped by PCR following the protocol and primers described by the mice supplier.

ENVIRONMENTAL ENRICHMENT

Mice exposed to SE were housed in pairs, in standard cages (30 cm × 16 cm × 11 cm). Mice exposed to EE were housed 10 for cage (36 cm × 54 cm × 19 cm or 45 cm × 25 cm × 22 cm with a labyrinth), in the presence of an assortment of objects, including climbing ladders, running wheel, balls, plastic, and wood objects

suspended from the ceiling, paper, cardboard boxes, and nesting material. Toys were changed every 2–3 days, while the bedding was changed every week. Both EE and SE groups received identical type of rodent chow and water *ad libitum* and were kept on a 12-h light/dark cycle. Mice were kept in EE or SE for 2.5 months before electrophysiological, behavioral, and immunofluorescence analyses. In particular, electrophysiological and immunofluorescence experiments were performed on two different sets of trained animals (for both *wt* and CX₃CR1^{GFP/GFP}). EE experiments were not performed on littermates due to technical limitations to obtain age- and gender-matched mice of different genotypes in sufficient number.

HIPPOCAMPAL SLICE PREPARATION

Electrophysiological experiments were performed from 1 to 5 days after enrichment. The experiments were performed in agreement with international guidelines on the ethical use of animals from the European Communities Council Directive of 24 November 1986 (86/609 EEC). Hippocampal slices were routinely obtained from 13 to 14-week-old C57BL/6J or CX₃CR1^{GFP/GFP} mice. Briefly, the animals were decapitated after being anesthetized with halothane. Whole brains were rapidly removed from the skull and immersed for 10 min in ice-cold artificial cerebrospinal fluid (ACSF) solution containing (in millimolar): NaCl 125, KCl 4.4, CaCl₂ 2.5, MgSO₄ 1.5, NaHPO₄ 1, NaHCO₃ 26, and glucose 10. The ACSF was continuously oxygenated with 95% O₂, 5% CO₂ to maintain the proper pH (7.4). Transverse 350 μm thick slices were cut at 4°C with a vibratome (DSK, Japan) and the appropriate slices were placed in a chamber containing oxygenated ACSF. After their preparation, slices were allowed to recover for 2 h. Individual slices were then transferred to the interface slice-recording chamber (BSC1, Scientific System Design Inc.) with a total fluid dead space of approximately 3 ml. Slices were maintained at 30–32°C and constantly superfused at the rate of 1.5 ml/min. Solutions were applied to the slices by a peristaltic pump.

ELECTROPHYSIOLOGICAL RECORDINGS

At the beginning of each recording, a concentric bipolar stimulating electrode (SNE-100 × 50 mm long Elektronik–Harvard Apparatus GmbH) was positioned in the stratum radiatum for stimulation of Schaffer collateral pathway projections to CA1. An ACSF-filled glass micropipette (0.5–1 MΩ) was positioned 200–600 μm from the stimulating electrode for recording orthodromically evoked fEPSPs. Stimuli consisted of 100 μs constant current square pulses, applied at 0.05 Hz. The intensity of the stimulus was adjusted in each experiment to evoke ~50% of the maximal field potential amplitude without appreciable population spike contamination. Evoked responses were monitored online and stable baseline responses were recorded for at least 10 min. Only the slices that showed stable fEPSP amplitudes were included in the experiments. To analyze the time course of fEPSP slope, the recorded fEPSP was routinely averaged over 1 min (*n* = 3). LTP experiments were performed in ACSF and the averaged fEPSP (35–45 min post-induction) was normalized to the baseline values (0–10 min) before LTP induction (HFS, 1 train, 100 Hz, 1-s duration, test strength). LTP experiments in SE were replicated in

CX₃CR1^{GFP/GFP}, wt and CX₃CR1^{GFP/+} littermates (Figure A1 in Appendix).

For the paired-pulse ratio (PPR) test, closely spaced consecutive stimuli (50 ms interval) were used, and PPR was calculated as the ratio between the fEPSP amplitude evoked by the second stimulus (A2) over the first (A1; A2/A1). Input–output (I–O) curves were measured at the beginning of recordings.

DATA ACQUISITION AND ANALYSIS

Slices were visualized with a Wild M3B (Heerbrugg, Switzerland). fEPSPs were recorded and filtered (1 kHz) with an Axopatch 200 A amplifier (Axon Instruments, CA, USA) and digitized at 10 kHz with an A/D converter (Digidata 1322A, Axon Instruments). Data were stored on a computer using pClamp 9 software (Axon Instruments) and analyzed off-line with Clamp-fit 9 program (Axon Instruments).

MORRIS WATER MAZE

The water maze apparatus consisted of a black Plexiglas circular pool 88 cm in diameter and 33 cm in height, and was placed in the middle of an experimental room (dimension 4 m × 4 m × 3 m). The pool was filled with water kept at a temperature of 26 ± 1°C. A plastic transparent platform (8 cm in diameter) was placed 0.5 cm below the water surface and 10 cm from the edge of the pool. All tests were carried out between 9:00 and 14:00 h. The entire procedure took 8 days. Mice were individually transferred from the home-cage to pool. To avoid visual orientation prior to release, mice were transferred from their cages into the pool in a non-transparent plastic cup, from which they glided into the water facing the pool wall. Release points were balanced across four symmetrical positions on the pool perimeter.

The ability of experimental subjects to identify and reach a visible platform was tested in the visual cued version of the task, lasting 2 days, which preceded the 6-days of the acquisition phase. Each testing day, mice underwent three trials during which they were allowed to freely swim either for 60 s (cut-off time) or until they found and climbed onto the platform; the inter-trial interval was at least 40 min. Platform finding was defined as staying for at least 3 s on it. Once the platform was found, the mice were given the opportunity to climb on a wire-mesh grid and placed back to a cage kept in a warm environment. During the acquisition phase, mice that did not find the platform were trained in locating it by placing them on the platform for 10 s at the end of the trial. On the fourth and sixth day of the acquisition phase, the platform was removed from the pool and each mouse was tested for memory retention in a 30-s probe trial. During the probe trial, the platform was removed from the pool and the time spent in the target quadrant of the maze (where the platform was located during the acquisition phase) was scored as a reliable measure of memory retention. The swim path of the mice was recorded by means of a computer-based video-tracking system Ethovision (Noldus, The Netherlands). For the acquisition phase, the variables recorded were: latency to reach the platform, mean swimming speed, path length and turned angle, and thigmotaxis. For the probe trial, the variables recorded were: time spent in each quadrant, number of platform crossing, mean swimming speed, and thigmotaxis.

ANIMALS AND TISSUES PREPARATION

Four mice from each different set of trained animals (both wt and CX₃CR1^{GFP/GFP}) were anesthetized using chloral-hydrate (400 mg/kg), transcardially perfused with phosphate-buffered saline (PBS 0.1 M), followed by 4% paraformaldehyde (PFA) in PBS. The brains were removed and post-fixed in 4% PFA overnight at 4°C and then cryoprotected by 24 h immersion in PBS containing 30% (w/v) sucrose at 4°C. Brains were cut into 10 μm sagittal sections using a cryostat.

IMMUNOFLUORESCENCE

Immunofluorescence staining procedures were conducted on free-floating sections containing the hippocampal DG starting at −1.46 mm and ending at −2.80 mm from the bregma. The whole region of interest was covered by 4 sequential 40× microscopic fields. The same number of sections for mouse (25) was incubated for 1 h in 5% non-fat dry milk 5 and 0.5% Triton X-100, in PBS 0.1M. Sections were then incubated in goat anti DCX (Santa Cruz Biotechnology, CA, USA) diluted in 1% non-fat dry milk and 0.2% Triton X-100, in PBS 0.1 M, for 48 h at 4°C. After washing in PBS, the sections were incubated in secondary antibody (donkey anti goat, Alexafluor, Invitrogen) for 1 h, washed in PBS and then stained with Hoechst (Invitrogen) for 5 min, washed again and mounted on a microscope slide for the analysis of fluorescence (Axioscope 2 Zeiss). DCX⁺ cells in the SGZ and the granule cell layer of the DG were counted exhaustively (at 40× magnification). For analysis of cell migration, DCX⁺ cells were scored as migrating if their cell body was, in the granular layer, fully detached from the SGZ of at least 5 μm. In the same slices analyzed for neurogenesis, DG hippocampal area was calculated with Image J software and the counting of DCX⁺ cells were normalized to DG area.

STATISTICAL ANALYSIS

The values were reported as mean ± SEM. If not specify, *n* refers to the number of mice analyzed. For comparisons among different groups, a two-way analysis of variance (ANOVA) was used with genotype and enrichment as between-subjects factors. ANOVA analyses were followed by Tukey (electrophysiology) or Holm–Sidak (neurogenesis analysis) *post hoc* analyses (Sigma Plot, 11.0 software). Mixed-model ANOVA with genotype and enrichment as between-subject factor and repeated measure as within-subject factor (e.g., day, quadrant) followed by Tukey *post hoc* comparisons was performed to analyze water maze data.

RESULTS

THE ABSENCE OF CX₃CR1 INCREASES HIPPOCAMPAL LTP BUT ABOLISHES THE EFFECT OF EE

It has been reported that hippocampal LTP, induced by a weak stimulation (1 s burst of 100 Hz, HFS) in the CA1 region of the hippocampus, is enhanced by enrichment (Duffy et al., 2001). Following the same protocol of induction, we measured fEPSP potentiation in hippocampal slices in wt and CX₃CR1^{GFP/GFP} mice following exposure to either SE or EE (Figure 1A,B). In wt mice, the mean fEPSP slope potentiation, measured 35–45 min after LTP induction, was 1.05 ± 0.04 (17 slices/8 mice) and 1.56 ± 0.08 of baseline (17 slices/9 mice) in SE and EE, respectively. In CX₃CR1^{GFP/GFP} mice, LTP induction produced fEPSP slope




FIGURE 1 | CA1 plasticity of *wt* and *CX₃CR1^{GFP/GFP}* mice exposed either to SE or EE conditions. Points represent mean \pm SEM.

Of fEPSP slopes evoked every 20 s and normalized as detailed in the Section "Materials and Methods." Arrows indicate time of application of HFS. **(A)** Enhancement of LTP in *wt* mice raised in EE: 45 min after HFS, EE *wt* mice (open circle) developed a robust LTP of fEPSP slope

compared to that of mice raised in SE (dark square, $p < 0.001$). **(B)** In *CX₃CR1^{GFP/GFP}* mice LTP is enhanced and EE produced no effect: after HFS, the amplitude of fEPSP potentiation of EE *CX₃CR1^{GFP/GFP}* mice (open circle) is not different from that obtained in SE mice (dark square). Note that that basal LTP evoked in SE is increased in *CX₃CR1^{GFP/GFP}* mice compared to *wt* [in **(A)**, $p = 0.006$].

potentiation of 1.19 ± 0.06 (17 slices/7 mice) and 1.26 ± 0.07 (27 slices/9 mice) in SE and EE, respectively.

A main effect of housing condition was found [$F_{1,27} = 10.232$, $p = 0.002$]. However a significant housing \times genotype interaction [$F_{(1,27)} = 20.035$, $p < 0.001$] showed that the housing condition affected only mice of the *wt* group. In particular, the *post hoc* analysis revealed that (i) the fEPSP potentiation was significantly higher in *CX₃CR1^{GFP/GFP}* SE compared to *wt* SE ($p = 0.042$), indicating that in control conditions *CX₃CR1^{GFP/GFP}* mice showed an enhanced plasticity, also confirmed in littermates (**Figure A1** in Appendix); (ii) the *wt* EE showed a significant enhancement compared to *wt* SE ($p < 0.001$), indicating that EE affects only *wt* mice.

SHORT-TERM PLASTICITY AND BASAL fEPSP RESPONSES TO SCHAFER COLLATERAL STIMULATION ARE NOT AFFECTED IN *CX₃CR1^{GFP/GFP}* MICE

To analyze if the lack of effect of EE on LTP in *CX₃CR1^{GFP/GFP}* mice could be due to altered responses to basal synaptic transmission, we performed studies of PPR and fEPSP I–O curves of synaptic transmission in *wt* and *CX₃CR1^{GFP/GFP}* mice. PPR, that is the ratio between the fEPSP amplitude evoked by the second stimulus over the first, represents a form of pre-synaptic short-term plasticity, whose variation is generally associated with changes in transmitter release probability (Zucker, 1989). To evaluate PPR values in *wt* and *CX₃CR1^{GFP/GFP}* mice, we stimulated Schaffer collateral pathway projections to CA1 at 50 ms intervals. As shown in **Figure 2A**, PPR was 1.22 ± 0.03 (15 slices/7 mice) and 1.22 ± 0.04 (9 slices/5 mice), in *CX₃CR1^{GFP/GFP}* and *wt* mice, respectively. These findings indicate that the transmitter release probability at Schaffer collateral input is comparable between the two genotypes. We then investigated synaptic strength in *CX₃CR1^{GFP/GFP}* mice recording fEPSP I–O curves of synaptic transmission in term of slope of the evoked potentials. Sample traces of fEPSP recorded at different intensities are illustrated in **Figure 2B**: the I–O relationships of the fEPSP slope CA1 curves of the *wt* (7 slices/5 mice) were indistinguishable from those of *CX₃CR1^{GFP/GFP}* mice (12 slices/7 mice), indicating similar basal


fEPSP responses to Schaffer collateral stimulation over a range of stimulus intensities.

CX₃CR1^{GFP/GFP} MICE LEARNED THE WATER MAZE TASK FASTER THAN *WT* MICE AND THEIR PERFORMANCE WAS NOT AFFECTED BY THE EE

In order to assess the role of CX₃CL1/CX₃CR1 signaling on learning and memory abilities, we investigated the effects of 2.5 months exposure to either SE or EE on *wt* and *CX₃CR1^{GFP/GFP}* mice in the Morris water maze test. The four experimental groups, *wt* SE ($n = 6$), *wt* EE ($n = 6$), *CX₃CR1^{GFP/GFP}* SE ($n = 8$), and *CX₃CR1^{GFP/GFP}* EE ($n = 8$) mice, showed no differences in the learning performance in the visual phase, indicating that unexpected drawbacks due to enrichment and/or to genetic manipulation did not interfere with the ability to solve the maze.

In the acquisition phase, all mice significantly reduced the latency to find the platform day after day [$F_{(2,48)} = 22.379$, $p < 0.0001$]. However, the experimental subjects exposed to EE condition displayed better learning and memory abilities than those exposed to SE, as shown by the significant main effect of the environment [$F_{(1,24)} = 9.001$, $p = 0.0062$; **Figure 3A**, left]. In line with the literature (Tremml et al., 2002; Harris et al., 2009), a reduced thigmotactic behavior in EE compared to SE mice was found [$F_{(1,24)} = 6.864$, $p = 0.0150$; data not shown]. This may, at least in part, explain the difference in learning performances among groups shown during acquisition (Wolfer et al., 1998).

The effect of the environment was also evident during the probe phase on the fourth and the sixth day [respectively, $F_{(3,72)} = 4.650$, $p = 0.0050$ and $F_{(3,72)} = 3.726$, $p = 0.0150$; **Figures 3B,C**]. In particular, in both probe phases, EE mice spent significantly longer time in the quadrant where the platform was located during the acquisition phase (acquisition quadrant) compared to SE mice. By contrast, the effect of genotype was evident only during the probe phase on the fourth day [$F_{(3,72)} = 4.201$, $p = 0.0085$] even though the analysis of genotype \times environment interaction did not reach statistical significance [$F_{(3,72)} = 2.179$, $p = 0.0979$]. However, *post hoc* analysis revealed that: (i) *CX₃CR1^{GFP/GFP}* mice learned the water maze task faster than *wt* mice, the time spent by SE *CX₃CR1^{GFP/GFP}* mice in the acquisition quadrant being



significantly longer than that spent by SE wt ($p < 0.05$); (ii) EE did not improve the water maze task performance in CX₃CR1^{GFP/GFP} mice, since EE mice spent an amount of time similar to SE mice in the acquisition quadrant. On the probe phase of the sixth day, a main effect of quadrant was found [$F_{(3,72)} = 30.069$, $p < 0.0001$]. In addition, after pooling data of the two genotypes within each house condition because of no genotype effect, the interaction house condition \times quadrant is statistically significant [$F_{(3,78)} = 4.146$, $p = 0.0088$] and *post hoc* analysis revealed that all groups spent more time in the platform quadrant ($p < 0.05$; **Figure 3C**).

With regard to the other parameters considered in the analysis, a main effect of the genotype was found for speed measured during the acquisition phase and the day 4 probe [respectively, $F_{(1,24)} = 29.284$ and 4.576 , $p < 0.001$ and $p = 0.0428$], CX₃CR1^{GFP/GFP} mice swimming faster than wt, but not in the day 6 probe. A main effect of genotype was found also for path length, CX₃CR1^{GFP/GFP} mice swimming longer paths than wt during the acquisition phase (**Figure 3A**, right), the day 4 and the day 6 probe (data not shown). However, since the genotype did not affect learning performance during the acquisition phase and speed and path length are not relevant for evaluating quadrant preference during the probe phase, the results obtained have not likely been influenced by these factors. Both during the acquisition and the probe phases, no difference in thigmotactic behavior among the experimental groups was found.


EE ENHANCES NEUROGENESIS IN THE DG OF BOTH WT AND CX₃CR1^{GFP/GFP} MICE.

CX₃CR1^{GFP/GFP} mice do not have evident alterations in brain development (Jung et al., 2000) but show decreased proliferation and neurogenesis in the SGZ and granular layer of the DG (Bachstetter et al., 2011). In order to investigate if the differences in learning and memory abilities observed in CX₃CR1^{GFP/GFP} mice could be correlated with alterations in DG neurogenesis, we analyzed the number of neuronal precursors (DCX positive cells)

present in the DG of wt and CX₃CR1^{GFP/GFP} mice grown in SE or EE ($n = 4$ for both housing conditions and genotypes) (representative image shown in **Figure 4A**). We reported that the number of neuronal precursors was affected not only by the lack of CX₃CR1, as shown by the significant effect of genotype [$F_{(1,13)} = 28.327$, $p < 0.001$], but also by the housing conditions [$F_{(1,13)} = 19.238$, $p < 0.001$] without any significant interaction between the two factors ($p = 0.760$). The mean area values of the DG analyzed for wt and CX₃CR1^{GFP/GFP} mice were not significantly different, indicating that the effect of genotype was not dependent on different DG size (wt: $0.143 \text{ mm} \pm 0.004 \text{ mm}$, CX₃CR1^{GFP/GFP}: $0.141 \text{ mm} \pm 0.004 \text{ mm}$). *Post hoc* analysis revealed that the number of DCX positive cells was: (i) significantly higher in wt compared to CX₃CR1^{GFP/GFP} ($p < 0.001$; **Figure 4B–F**, as previously described (Bachstetter et al., 2011)); (ii) increased by exposure to EE in both genotypes ($p < 0.001$; **Figures 4D–F**).

We then evaluated the rate of radial migration of neuronal precursors in the DG, the step that precedes their differentiation into granule neurons and the establishment of synaptic contacts in the pre-existing hippocampal circuitries (Cameron et al., 1993; Markakis and Gage, 1999). For this reason we measured the number of migrating neuronal precursors, defined as the DCX⁺ cells whose cell body was at least $5 \mu\text{m}$ distant from the SGZ. We found that the number of DCX⁺ migrating cells was affected by both housing [$F_{(1,12)} = 85.165$, $p < 0.001$] and genotype [$F_{(1,12)} = 27.570$, $p < 0.001$] conditions but no significant interaction of these two factors was found ($p = 0.287$). *Post hoc* analysis revealed that: (i) migrating cells were increased in both genotypes in EE compared to SE condition ($p < 0.001$; **Figures 4B–E,G**); (ii) wt mice showed a higher number of migrating cells compared to CX₃CR1^{GFP/GFP} ($p < 0.001$; **Figures 4B–E,G**).

When we measured the ratio of migrating DCX⁺/total DCX⁺ cells, we found a main effect of housing [$F_{(1,12)} = 51.011$, $p < 0.001$] and genotype condition [$F_{(1,12)} = 6.399$, $p = 0.026$] with a significant factors interaction [$F_{(1,12)} = 5.668$, $p = 0.035$]. *Post hoc* analysis revealed that the ratio between migrating DCX⁺/total




DCX⁺ cells was significantly different in the two genotypes only in SE, *CX₃CR1^{GFP/GFP}* mice showing reduced levels compared to *wt* (*p* < 0.05), but not in EE condition (Figures 4B–E,H).

DISCUSSION

The main aim of the present work was to investigate the role of CX₃CR1 in mediating the effects of prolonged EE on hippocampal function and neurogenesis. As expected according to the literature (van Praag et al., 2000; Duffy et al., 2001), EE enhanced neuronal plasticity in *wt* but, surprisingly, not in *CX₃CR1^{GFP/GFP}* mice. In particular, following exposure to long-term EE, *CX₃CR1^{GFP/GFP}* mice showed no improvement in both CA1 hippocampal LTP and in water maze performance on day 4 probe. In contrast, EE enhanced hippocampal neurogenesis and migration of adult new born neuronal precursors in the DG of both genotypes.

We also report that the lack of CX₃CR1 has several effects by itself. In particular, in *CX₃CR1^{GFP/GFP}* mice grown under SE conditions, hippocampal LTP, and spatial memory are improved whereas DG neuronal precursor generation and migration are



reduced when compared with *wt* mice (see also Bachstetter et al., 2011). Specifically, we reported that, although in SE *CX₃CR1^{GFP/GFP}* mice showed similar responses to basal synaptic transmission (measured as I–O functional curves and PPF), upon a weak stimulation (100 Hz train) they had a more robust CA1 LTP compared to *wt* mice, demonstrating microglia-driven alterations of plasticity phenomena. This is in line with our previous

results, demonstrating that treatment with exogenous CX₃CL1 inhibits CA1 LTP through the activity of adenosine system, specifically of the adenosine receptor type 3 (AR3; Maggi et al., 2009). The role of microglia in modulating neuronal circuitry functions and hippocampal plasticity is also suggested by recent findings demonstrating that, during development, CX₃CR1^{GFP/GFP} mice had a decreased density in microglia cells that were compromised in their capacity to engulf synaptic material leading to immature connectivity, augmented LTD, and a global increase in excitatory synapses with immature functional characteristics (Paolicelli et al., 2011).

It has been proposed that exposure to EE may modify synaptic physiology and plasticity in hippocampal neurons (Green and Greenough, 1986; Foster et al., 1996; van Praag et al., 1999) being able to produce an enhancement of LTP in the area CA1 (Duffy et al., 2001). In plasticity processes, signaling between perisynaptic astrocytes, microglia and neurons plays an important role. Astrocytes and microglia secrete an array of cytokines and factors which contribute to synaptic modulation, learning and memory processes in healthy and diseased brain (Volterra and Meldolesi, 2005; Hanisch and Kettenmann, 2007). Nevertheless, the exact mechanisms as to how cytokines and in particular chemokines, participate in the molecular and cellular processes thought to subserve memory formation and responsiveness to environmental stimuli remain to be clarified. Interestingly, it has been recently reported that mice lacking type 1 receptor for interleukin 1 (IL-1) display impaired hippocampal memory and LTP that are restored by EE (Goshen et al., 2009).

Here we showed that, in the absence of CX₃CR1, EE failed to affect hippocampal CA1 LTP, whereas it produced an increased plasticity in *wt* animals. These evidence suggest that the absence of microglia-driven CX₃CR1 signaling occludes the enrichment effects (although we are not at saturating LTP level) possibly because of some overlapping mechanisms that need further investigation. We speculate that this lack of response to environmental stimulation might be explained with altered microglia–neurons cross-talk, affecting the circuitry underlying synaptic plasticity. These changes could reduce the responsiveness of hippocampal CA1 region to EE stimuli that, in *wt* animals, have been shown to increase the number of synaptic contacts and the density of non-perforated synapses in this area (Rampon and Tsien, 2000).

A strong association between LTP and cognition has been widely described in previous studies illustrating LTP as prime candidate for the cellular mechanisms of learning and memory (Tsien et al., 1996; Malenka, 2003). EE has been shown to enhance learning and memory functions in various tasks (van Praag et al., 1999). In particular, enriched mice display a better performance in the Morris water maze compared to controls (Pacteau et al., 1989; Kempermann et al., 1997). Here, we confirm these findings, showing that EE improved the performance of both genotypes with regard to latency to reach the platform during the acquisition phase and the preference for the platform quadrant during both probe phases. Indeed, the preference for the platform quadrant emerged earlier in the EE group, appearing after 4 days of acquisition, while it was evident only after 6 days in the SE group. *Post hoc* analysis revealed that the effect of the housing condition at day 4 probe is mainly due to a lack of preference shown by SE *wt* while CX₃CR1^{GFP/GFP} mice, independently from

the housing conditions, showed a significant preference, indicating enhanced cognitive abilities. Interestingly, improved cognitive functions have been described in mice lacking interleukin 6 (IL-6, Braida and Sacerdote, 2004) whereas increased levels of IL-6 were associated with decline in learning and memory abilities (Monje and Toda, 2003).

The lack of differences in learning performance at day 4 probe is in accordance with the LTP data, corroborating a reduced sensitivity to environmental stimulation associated with CX₃CR1 deficiency. However, we cannot exclude the presence of a ceiling effect, being the preference for the platform quadrant already at high levels after 4 days of acquisition, no further improvement due to EE might be possible. At day 6 probe, the housing condition affected the preference for the platform quadrant also in CX₃CR1^{GFP/GFP} mice, possibly because the prolonged water maze training allowed differences associated to housing condition to clearly emerge. It is worth noticing that the procedure followed in the present work consisting in exposing mice to a probe trial (i.e., day 4 probe), which is an extinction trial, during the acquisition phase may have interfered with the learning process. However, the preference for the acquisition quadrant shown by all experimental groups in the day 6 probe trial suggests that such interference, if any, was limited. Further investigations are warranted to better characterize the effect of CX₃CR1 deficiency on learning and memory.

Considerable evidence suggest that hippocampal neurogenesis plays a role in learning and memory functions (Shors et al., 2001; Snyder et al., 2005) and that EE stimulates hippocampal neurogenesis in the DG (Kempermann et al., 1997; Kempermann and Gage, 1999; van Praag et al., 1999) but the precise mechanisms by which neurogenesis facilitates memory formation remain to be fully understood. Recent data suggest that neurogenesis may be associated with some forms of hippocampal-dependent memory but not others (Shors and Townsend, 2002; Saxe et al., 2007; Garthe et al., 2009) and that reducing neurogenesis does not affect LTP in CA1 area (Garthe et al., 2009).

CX₃CL1 has been recently shown to be involved in the regulation of DG neurogenesis, since CX₃CR1^{GFP/GFP} mice have a reduced number of neuronal precursors in the DG (Bachstetter et al., 2011). We investigated whether EE was able to affect hippocampal neurogenesis in the absence of a normal CX₃CL1–CX₃CR1 signaling. Our results confirm that mice lacking CX₃CR1 have, in SE, a reduced neurogenesis in the SGZ and granular layer of the DG, as measured by DCX staining (Bachstetter et al., 2011). We report that exposure to EE induced a significant increase of neurogenesis and migration of new born cells in the DG of both *wt* and CX₃CR1^{GFP/GFP} mice and that the differences in the percentage of new born migrating cells observed in CX₃CR1 deficient mice are lost upon EE exposure. This observation indicates that EE may modulate specific signals in the neurogenic niche, likely modulating survival and differentiation of new born cells, that are functional also in the absence of CX₃CR1 but probably do not overlap with those involved in basal regulation. Possible candidates are factors involved in new born cell survival, local network activity (Deng et al., 2010), and epigenetic factors (Covic et al., 2010; Lopez-Atalaya et al., 2011). Interestingly, hippocampal neurogenesis induced by EE was associated with the activation of microglia (Ziv et al., 2006). It has been proposed that disruption

of microglia-driven CX₃CR1 signaling decreases survival and proliferation of neuronal progenitors via mechanisms that involve indirect modification of niche environment, in particular through IL-1 β (Bachstetter et al., 2011).

Our results also suggest that learning-induced regulation of adult neurogenesis is not specifically associated with modification in CA1 hippocampal plasticity or Morris water maze performance, substantiating the current idea that these forms of hippocampal-dependent plasticity and learning processes are not directly related with DG neurogenesis. Future studies are warranted to discern if specific plasticity processes in DG area could be altered in CX₃CR1^{GFP/GFP} mice, in particular those related to adult-generated granule cells that have a potential role in spatially precise, place specific strategy (Garthe et al., 2009).

In conclusion, we provide the first experimental evidence that the absence of a constitutive CX₃CL1/CX₃CR1 signaling leads to

an increase in hippocampal plasticity and learning performance and to a decrease in hippocampal-dependent responses to environmental stimulation, demonstrating microglia-driven alterations of plasticity phenomena.

ACKNOWLEDGMENTS

The authors acknowledge the technical assistance of Giuseppina Chece and Fabrizio Cecchini for mice genotyping, Alessandro Felici for animal care and Cristina Bertolini for initial EE experiments. Maria Scianni was supported by Ph.D. program in Neurophysiology, Sapienza University, Rome, Italy. Grants: Istituto Pasteur, Fondazione Cenci Bolognetti to Cristina Limatola; Ministero della Salute (Ricerca corrente, cinque per mille, Antidoping) to Cristina Limatola; Mariani Foundation of Milan (Grant n. R-09-76) to Cristina Limatola; Fondazione Viva la Vita to Cristina Limatola.

REFERENCES

- Asensio, V. C., and Campbell, I. L. (1999). Chemokines in the CNS: plurifunctional mediators in diverse states. *Trends Neurosci.* 22, 504–512.
- Bachstetter, A. D., Morganti, J. M., Jernberg, J., Schlunk, A., Mitchell, S. H., Brewster, K. W., Hudson, C. E., Cole, M. J., Harrison, J. K., Bickford, P. C., and Gemma, C. (2011). Fractalkine and CX(3)CR1 regulate hippocampal neurogenesis in adult and aged rats. *Neurobiol. Aging* 32, 2030–2044.
- Bertolini, C., Ragozzino, D., Gross, C., Limatola, C., and Eusebi, F. (2006). Fractalkine/CX3CL1 depresses central synaptic transmission in mouse hippocampal slices. *Neuropharmacology* 51, 816–821.
- Bezzi, P., Domercq, M., Brambilla, L., Galli, R., Schols, D., De Clercq, E., Vescovi, A., Bagetta, G., Kollias, G., Meldolesi, J., and Volterra, A. (2001). CXCR4-activated astrocyte glutamate release via TNF- α : amplification by microglia triggers neurotoxicity. *Nat. Neurosci.* 4, 702–710.
- Braida, D., and Sacerdote, P. (2004). Cognitive function in young and adult IL (interleukin)-6 deficient mice. *Behav. Brain Res.* 153, 423–429.
- Brown, J. P., Couillard-Després, S., Cooper-Kuhn, C. M., Winkler, J., Aigner, L., and Kuhn, H. G. (2003). Transient expression of doublecortin during adult neurogenesis. *J. Comp. Neurol.* 467, 1–10.
- Cameron, H. A., Woolley, C. S., McEwen, B. S., and Gould, E. (1993). Differentiation of newly born neurons and glia in the dentate gyrus of the adult rat. *Neuroscience* 56, 337–344.
- Covic, M., Karaca, E., and Lie, D. C. (2010). Epigenetic regulation of neurogenesis in the adult hippocampus. *Heredity* 105, 122–134.
- Deng, W., Aimone, J. B., and Gage, F. H. (2010). New neurons and new memories: how does adult hippocampal neurogenesis affect learning and memory? *Nat. Rev. Neurosci.* 11, 339–350.
- Duffy, S. N., Craddock, K. J., Abel, T., and Nguyen, P. V. (2001). Environmental enrichment modifies the PKA-dependence of hippocampal LTP and improves hippocampus-dependent memory. *Learn. Mem.* 8, 26–34.
- Engellenner, W. J., Goodlett, C. R., Burright, R. G., and Donovick, P. J. (1982). Environmental enrichment and restriction: effects on reactivity, exploration and maze learning in mice with septal lesions. *Physiol. Behav.* 29, 885–893.
- Escorihuela, R. M., Tobena, A., and Fernandez-Tereul, A. (1995). Environmental enrichment and postnatal handling prevent spatial learning deficits in aged hypoemotional (Roman high-avoidance) and hyperemotional (Roman low-avoidance) rats. *Learn. Mem.* 2, 40–48.
- Foster, T. C., Gagne, J., and Massicotte, G. (1996). Mechanism of altered synaptic strength due to experience: relation to long-term potentiation. *Brain Res.* 736, 243–250.
- Garthe, A., Behr, J., and Kempermann, G. (2009). Adult-generated hippocampal neurons allow the flexible use of spatially precise learning strategies. *PLoS ONE* 4, e5464. doi:10.1371/journal.pone.0005464
- Giovannelli, A., Limatola, C., Ragozzino, D., Mileo, A. M., Ruggieri, A., Ciotti M. T., Mercanti, D., Santoni, A., and Eusebi F. (1998). CX3 chemokines interleukin-8 (IL-8) and growth-related gene product α (GRO α) modulate Purkinje neuron activity in mouse cerebellum. *J. Neuroimmunol.* 92, 122–132.
- Globus, A., Rosenzweig, M. R., Bennett, E. L., and Diamond, M. C. (1973). Effects of differential experience on dendritic spine counts in rat cerebral cortex. *J. Comp. Physiol. Psychol.* 82, 175–181.
- Goshen, I., Avital, A., Kreisel, T., Licht, T., Segal, M., and Yirmiya, R. (2009). Environmental enrichment restores memory functioning in mice with impaired IL-1 signaling via reinstatement of long-term potentiation and spine size enlargement. *J. Neurosci.* 29, 3395–3403.
- Green, E. J., and Greenough, W. T. (1986). Altered synaptic transmission in dentate gyrus of rats reared in complex environments: evidence from hippocampal slices maintained in vitro. *J. Neurophysiol.* 55, 739–750.
- Hanisch, U. K., and Kettenmann, H. (2007). Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat. Neurosci.* 10, 1387–1394.
- Harris, A. P., D'Eath, R. B., and Healy, S. D. (2009). Environmental enrichment enhances spatial cognition in rats by reducing thigmotaxis (wall hugging) during testing. *Anim. Behav.* 77, 1459–1464.
- Jung, S., Aliberti, J., Graemmel, P., Sunshine, M. J., Kreutzberg, G. W., Sher, A., Littman, D. R. (2000). Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. *Mol. Cell. Biol.* 20, 4106–4114.
- Kempermann, G., and Gage, F. H. (1999). Experience-dependent regulation of adult hippocampal neurogenesis: effects of long-term stimulation and stimulus withdrawal. *Hippocampus* 9, 321–332.
- Kempermann, G., Kuhn, H. G., and Gage, F. H. (1997). More hippocampal neurons in adult mice living in an enriched environment. *Nature* 386, 493–495.
- Lax, P., Limatola, C., Fucile, F., Trettel, F., Di Bartolomeo, S., Renzi, M., Ragozzino, D., and Eusebi, F. (2002). Chemokine receptor CXCR2 regulates the functional properties of AMPA-type glutamate receptor GluR1 in HEK cells. *J. Neuroimmunol.* 129, 66–73.
- Limatola, C., Giovannelli, A., Maggi, L., Ragozzino, D., Castellani, L., Ciotti, M. T., Vacca, F., Mercanti, D., Santoni, A., and Eusebi, F. (2000). SDF-1 α -mediated modulation of synaptic transmission in rat cerebellum. *Eur. J. Neurosci.* 12, 2497–2504.
- Lopez-Atalaya, J. P., Ciccarelli, A., Viosca, J., Valor, L. M., Jimenez-Minchan, M., Canals, S., Giustetto, M., and Barco, A. (2011). CBP is required for environmental enrichment-induced neurogenesis and cognitive enhancement. *EMBO J.* doi: 10.1038/emboj.2011.299. [Epub ahead of print].
- Maggi, L., Trettel, F., Scianni, M., Bertolini, C., Eusebi, F., Fredholm, B. B., and Limatola, C. (2009). LTP impairment by fractalkine/CX3CL1 in mouse hippocampus is mediated through the activity of adenosine receptor type 3 (A3R). *J. Neuroimmunol.* 215, 36–42.
- Malenka, R. C. (2003). The long-term potential of LTP. *Nat. Rev. Neurosci.* 4, 923–926.

- Markakis, E. A., and Gage, F. H. (1999). Adult-generated neurons in the dentate gyrus send axonal projections to field CA3 and are surrounded by synaptic vesicles. *J. Comp. Neurol.* 406, 449–460.
- Meucci, O., Fatatis, A., Simen, A. A., Bushell, T. J., Gray, P. W., and Miller, R. J. (1998). Chemokines regulate hippocampal neuronal signaling and gp120 neurotoxicity. *Proc. Natl. Acad. Sci. U.S.A.* 95, 14500–14505.
- Monje, M. L., and Toda, H. (2003). Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 302, 1760–1765.
- Nithianantharajah, J., and Hannan, A. J. (2006). Enriched environments, experience-dependent plasticity and disorders of the nervous system. *Nat. Rev. Neurosci.* 7, 697–709.
- Oh, S. B., Endoh, T., Simen, A. A., Ren, D., and Miller, R. J. (2002). Regulation of calcium currents by chemokines and their receptors. *J. Neuroimmunol.* 123, 66–75.
- Otto, T., Eichenbaum, H., Wiener, S. I., and Wible, C. G. (1991). Learning-related patterns of CA1 spike trains parallel stimulation parameters optimal for inducing hippocampal long-term potentiation. *Hippocampus* 1, 181–192.
- Pacteau, C., Einon, D., and Sinden, J. (1989). Early rearing environment and dorsal hippocampal ibotenic acid lesions: long-term influences on spatial learning and alternation in the rat. *Behav. Brain Res.* 34, 79–96.
- Paolicelli, R., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M., Ferreira, T. A., Guiducci, E., Dumas, L., Ragozzino, R., and Gross, C. (2011). Synaptic pruning by microglia is necessary for normal brain development. *Science* 333, 1456–1458.
- Ragozzino, D., Di Angelantonio, S., Trettel, F., Bertollini, C., Maggi, L., Gross, C., Charo, I. F., Limatola, C., and Eusebi, F. (2006). Chemokine fractalkine/CX3CL1 negatively modulates active glutamatergic synapses in rat hippocampal neurons. *J. Neurosci.* 26, 10488–10498.
- Rampon, C., and Tsien, J. Z. (2000). Genetic analysis of learning behavior-induced structural plasticity. *Hippocampus* 10, 605–609.
- Rostène, W., Kitabgi, P., and Parsadaniantz, S. M. (2007). Chemokines: a new class of neuromodulator? *Nat. Rev. Neurosci.* 8, 895–903.
- Sánchez-Alcañiz, J. A., Haegel, S., Mueller, W., Pla, R., Mackay, F., Schulz, S., López-Bendito, G., Stumm, R., and Marín, O. (2011). Cxcr7 controls neuronal migration by regulating chemokine responsiveness. *Neuron* 69, 77–90.
- Saxe, M. D., Malleret, G., Vronskaya, S., Mendez, I., Garcia, A. D., Sofroniew, M. V., Kandel, E. R., and Hen, R. (2007). Paradoxical influence of hippocampal neurogenesis on working memory. *Proc. Natl. Acad. Sci. U.S.A.* 104, 4642–4646.
- Shors, T. J., Miesegages, G., Beylin, A., Zhao, M., Rydel, T., and Gould, E. (2001). Neurogenesis in the adult is involved in the formation of trace memories. *Nature* 410, 372–376.
- Shors, T. J., and Townsend, D. A. (2002). Neurogenesis may relate to some but not all types of hippocampal-dependent learning. *Hippocampus* 12, 578–584.
- Snyder, J. S., Hong, N. S., McDonald, R. J., and Wojtowicz, J. M. (2005). A role for adult neurogenesis in spatial long-term memory. *Neuroscience* 130, 843–852.
- Squire, L. R., and Zola-Morgan, S. (1991). The medial temporal lobe memory system. *Science* 20, 1380–1386.
- Tran, P. B., and Miller, R. J. (2003). Chemokine receptors: signposts to brain development and disease. *Nat. Rev. Neurosci.* 4, 444–455.
- Tremml, P., Lipp, H. P., Müller, U., and Wolfer, D. P. (2002). Enriched early experiences of mice underexpressing the β -amyloid precursor protein restore spatial learning capabilities but not normal openfield behavior of adult animals. *Genes Brain Behav.* 1, 230–241.
- Tsien, J. Z., Huerta, P. T., and Tonegawa, S. (1996). The essential role of hippocampal CA1 NMDA receptor-dependent synaptic plasticity in spatial memory. *Cell* 87, 1327–1338.
- van Praag, H., Christie, B. R., Sejnowski, T. J., and Gage, F. H. (1999). Running enhances neurogenesis, learning, and LTP in mice. *Proc. Natl. Acad. Sci. U.S.A.* 96, 13427–13431.
- van Praag, H., Kempermann, G., and Gage, F. H. (2000). Neural consequences of environmental enrichment. *Nat. Rev. Neurosci.* 1, 191–198.
- van Praag, H., Schinder, A. F., Christie, B. R., Toni, N., Palmer, T. D., and Gage, F. H. (2002). Functional neurogenesis in the adult hippocampus. *Nature* 415, 1030–1034.
- Volterra, A., and Meldolesi, J. (2005). Astrocytes, from brain glue to communication elements: the revolution continues. *Nat. Rev. Neurosci.* 6, 626–640.
- Wolfer, D. P., Stajlar-Bozicevic, M., Errington, M. L., and Lipp, H. P. (1998). Spatial memory and learning in transgenic mice: fact or artifact? *News Physiol. Sci.* 13, 118–123.
- Ziv, Y., Ron, N., Butovsky, O., Landa, G., Sudai, E., Greenberg, N., Cohen, H., Kipnis, J., and Schwartz, M. (2006). Immune cells contribute to the maintenance of neurogenesis and spatial learning abilities in adulthood. *Nat. Neurosci.* 9, 268–275.
- Zucker, R. S. (1989). Short-term synaptic plasticity. *Annu. Rev. Neurosci.* 12, 13–31.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 22 July 2011; accepted: 29 September 2011; published online: 19 October 2011.

Citation: Maggi L, Scianni M, Branchi I, D'Andrea I, Lauro C and Limatola C (2011) CX₃CR1 deficiency alters hippocampal-dependent plasticity phenomena blunting the effects of enriched environment. *Front. Cell. Neurosci.* 5:22. doi: 10.3389/fncel.2011.00022

Copyright © 2011 Maggi, Scianni, Branchi, D'Andrea, Lauro and Limatola. This is an open-access article subject to a non-exclusive license between the authors and Frontiers Media SA, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and other Frontiers conditions are complied with.

APPENDIX

