

Cyclic AMP-Response Element Regulated Cell Cycle Arrests in Cancer Cells

Ping Wang^{1*}, Shuaishuai Huang¹, Feng Wang², Yu Ren³, Michael Hehir¹, Xue Wang¹, Jie Cai^{4*}

1 Zhejiang Provincial Key Laboratory of Pathophysiology, Medical School, Ningbo University, Ningbo, China, **2** Ningbo Medical Center, LiHuiLi Hospital, Medical School, Ningbo University, Ningbo, China, **3** Department of Urologic Surgery, Ningbo Urology and Nephrology Hospital, Ningbo University, Ningbo, China, **4** Ningbo Women and Children's Hospital, Medical School, Ningbo University, Ningbo, China

Abstract

Recently, we have demonstrated that trichosanthin (TCS), a promising agent for the treatment of cervical adenocarcinoma, inhibited HeLa cell proliferation through the PKC/MAPK/CREB signal pathway. Furthermore, TCS down-regulated Bcl-2 expression was abrogated by a decoy oligonucleotide (OGN) to the cyclic AMP-responsive element (CRE). The decoy OGN blocked the binding of CRE-binding protein (CREB) to Bcl-2. These results suggested that CRE-mediated gene expression may play a pivotal role in HeLa cell proliferation. However, little is known about the effect of TCS on cell cycle arrests, particularly, whether the genes involved in cell cycle were regulated by CRE. Our present study shows that the arrests of S, G1 and G2/M phases were accompanied by the significant down-regulation of cyclin A, D1 and CDK 2, 4 in HeLa cells, cyclin D1, E and CDK 2, 4 in Caski and C33a cells, and cyclin A, B1, E and CDK 2 in SW1990 cells. However, the cell cycle arrests were reversed via the significant up-regulation of cyclin A and D1, by the combined treatment of TCS and CRE. In conclusion, these data demonstrate for the first time that specific cell cycle arrests in cancer cells can be induced by TCS by inhibiting the binding of CREB to CRE on genes related to cell proliferation.

Citation: Wang P, Huang S, Wang F, Ren Y, Hehir M, et al. (2013) Cyclic AMP-Response Element Regulated Cell Cycle Arrests in Cancer Cells. PLoS ONE 8(6): e65661. doi:10.1371/journal.pone.0065661

Editor: Hong Wanjin, Institute of Molecular and Cell Biology, Biopolis, United States of America

Received: November 18, 2012; **Accepted:** April 25, 2013; **Published:** June 28, 2013

Copyright: © 2013 Wang et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the grants from the National Natural Science Foundation of China (81071653), Natural Science Foundation of Zhejiang Province (Y2111136), Advanced Key Scientific and Technological Programs of Ningbo (2011C51005), Natural Science Foundation of Ningbo City (2011A610050) and K. C. Wong Magna Fund in Ningbo University. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: pinoav@hotmail.com (PW); doctor_caijie@hotmail.com (JC)

Introduction

Trichosanthin (TCS), an active component isolated from the root tubers of the Chinese medicinal herb *Trichosanthes kirilowii* [1], is a promising agent for the treatment of cancer [2]. Our previous reports showed that TCS inhibited cervical adenocarcinoma HeLa cell proliferation [3] through the PKC/MAPK/CREB signal pathway [4]. Furthermore, TCS down-regulated Bcl-2 expression [5], which was abrogated by a cyclic AMP-responsive element (CRE, TGACGTC) decoy oligonucleotide (OGN), blocking the CRE-binding protein (CREB) binding site on Bcl-2 [6]. These results suggest that CRE-mediated gene expression may play a pivotal role in HeLa cell proliferation.

However, little is known about the effect of TCS on cell cycle arrest of HeLa cell, cervical squamous carcinoma (Caski and C33a cell), and human pancreatic carcinoma (SW1990 cell), especially, whether genes related to cell cycle were regulated by the CRE decoy OGN.

In the present study, we further explored the effects of TCS on the proliferation of cancer cells, cell cycle arrests in the progress of cell proliferation and the role of CRE in cell cycle regulation.

The aim of this study was to investigate the effects of TCS on cancer cell proliferation and the effect of CRE on TCS-induced cell cycle arrests. An important question was whether CRE-combined cyclins played a critical role in the regulation of cell cycle arrest in these cancer cells.

Materials and Methods

Cell lines and culture

Cervical cells (HeLa, Caski, C33a) and SW1990 cells were obtained from American Type Culture Collection (ATCC, USA). HeLa, Caski, C33a [7] and SW1990 cell [8] were grown in monolayer in RPMI 1640 medium (Gibco, NY, USA) and Dulbecco's modified Eagle's medium (DMEM), containing 10% heat-inactivated fetal bovine serum, 100 U/ml of penicillin and 100 µg/ml of streptomycin (BioWhittaker, Inc., Walkersville, MD, USA), in a CO₂ incubator (Forma Scientific, USA). The medium was replaced twice a week, and cells were passaged every 4–5 days at a ratio of 1:3.

Cell treatment

Cells were plated at 2×10^6 cells/dish in 100-mm dishes in the basal medium. At confluence, they were washed briefly with PBS and then treated with TCS (Jinshan pharmacy company, Shanghai, China). Control cells were incubated in TCS-free medium.

Cell viability assay

Cell viability was assessed with Cell Counting Kit-8 (Dojindo Laboratories, Kumamoto, Japan) using the methods described previously [3,4,9].




Figure 1. Effects of TCS on cancer cell proliferation. TCS inhibited cell proliferation in a dose-dependent manner. Data represent means \pm SD of three independent experiments (* $p < 0.05$ compared with control).

doi:10.1371/journal.pone.0065661.g001

Treatment of cells with CRE-OGNs

CRE decoy OGN (5'-TGACGTCAGAGAGCGCTCTCTGACGTCA-3') and control OGN (5'-TGACGTCATGACGT-CATGACGTCA-3') were converted to phosphorothioate OGNs (Invitrogen Carlsbad, CA, USA) as previously described [5,6,10].

Preparation of cytosolic and nuclear proteins

The preparations of cytosolic and nuclear proteins were performed as previously described [6,11,12]. Briefly, at the end of each designated treatment, cells were washed briefly with cold PBS and lysed by scrapping them in 0.5 ml cold hypotonic buffer (10 mM HEPES, 40 mM KCl, 3 mM $MgCl_2$, 1 mM dithiothreitol, 0.2% NP-40, 1 μ g/ml aprotinin, 2 μ M leupeptin, 1 mM phenylmethylsulfonyl fluoride, 40 mM p-nitrophenyl phosphate, 1 mM sodium orthovanadate and 50% glycerol). The lysates were collected and incubated on ice for 5 min, then centrifuged at 15,000 \times g for 20 s. The supernatant was collected and saved as the cytosolic fractions. The pellet (i.e. cell nuclei) was resuspended in hypertonic buffer (20 mM HEPES, 420 mM KCl, 1.5 mM $MgCl_2$, 0.2 mM EDTA, 0.5 mM dithiothreitol, 0.2% NP-40, 1 μ g/ml aprotinin, 2 μ M leupeptin, 0.5 mM phenylmethylsulfonyl fluoride, 40 mM p-nitrophenyl phosphate, 1 mM sodium orthovanadate and 25% glycerol). After incubation on ice for 60 min, the samples were centrifuged at 15,000 \times g for 20 min and the supernatants were collected and saved as nuclear extracts. The cytosolic fractions and nuclear extracts were boiled for 10 min and dissolved in 8% SDS PAGE.

Cell cycle analysis

For flow cytometry analysis of cell cycle, 1×10^6 cells were harvested by centrifugation, washed with PBS and fixed with ice-cold 70% ethanol overnight. Fixed cells were treated with 25 μ g/ml RNase A at 37°C for 30 min and then stained with propidium iodide (PI) (50 μ g/ml, Sigma) solution for 30 min in the dark. The fluorescence intensity of individual cells was measured by a flow cytometer (Beckman Coulter, Inc., Miami, FL, USA). At least 10,000 cells were counted [13].

Western blot analysis

Total cell lysates were prepared by lysis of cells with radioimmunoprecipitation assay buffer and protein concentration was determined using the protein assay kit (Bio-Rad). For Western

Table 1. Concentration producing 50% growth inhibition (IC₅₀) of TCS on cancer cells.

Cell line	IC ₅₀ (μ g/ml)	
	24 h	48 h
HeLa	98.25 \pm 1.46	64.21 \pm 2.45
Caski	51.46 \pm 4.59	22.86 \pm 4.63
C33a	49.83 \pm 4.21	21.94 \pm 2.48
SW1990	101 \pm 3.84	73.14 \pm 3.51

doi:10.1371/journal.pone.0065661.t001

blot analysis, 50 μ g of each sample was processed as described [14]. The following antibodies were used: anti-cyclin A, B1, D1, E, anti-CDK2, 4 and anti-actin (Cell signaling Technology, Inc., Beverly, MA). The secondary antibodies were coupled to horseradish peroxidase and detected by chemiluminescence (Bio-Rad, Hercules, CA). The relative amounts of immunoreactive protein in each band were determined by scanning densitometric analysis of the X-ray films.

Statistical analysis

All experiments were repeated three times. The data were expressed as means \pm SD. ANOVA was used to evaluate the differences between groups. Data reported as mean \pm SD of three independent experiments. Differences were considered significant if $p < 0.05$.

Results

Effects of TCS on the proliferation of cancer cells

TCS of 20–100 μ g/ml inhibited the proliferation of cells by 3% to 70% after treatment for 24 h (Fig. 1). The 50% inhibitory concentration (IC₅₀) of TCS on Caski and C33a cells was found to be 60 μ g/ml, lower than that on HeLa and SW1990 cells (100 μ g/ml) (Table 1).

Effects of TCS on cell cycle progress and cell cycle regulatory proteins

TCS-treated cancer cells were analyzed by flow cytometry. A dose-dependent cell number increases of S, G1, G2/M phase were shown in HeLa, Caski, and SW1990 cells respectively, after they were treated for 24 hours, (Table 2). The levels of cell cycle related proteins were determined by Western blot assay. TCS decreased the abundance of cyclin A, D1 and CDK 2, 4, while it had no marked effect on the expression of cyclin B1 and E in HeLa cell (Fig. 2). In Caski cell, cyclin D1, E and CDK 2, 4 were significantly decreased, whereas no marked changes were shown in the expression of cyclin A and B1 (Fig. 3). Similar results were found in C33a cells (data not shown). In SW1990 cell, cyclin A, B1, E and CDK 2 were significantly down-regulated, no distinct effects were observed in the expression of cyclin D1 and CDK4 (Fig. 4).

Effects of CRE-decoy on the cell cycle progress and regulatory proteins

The arrests of S, G1 and G2/M phases induced by TCS in HeLa (Fig. 5), Caski (Fig. 6) and SW1990 (Fig. 7) cells, were significantly attenuated by the combined treatment of TCS and CRE (A, B). It was found that the TCS-induced decreases of cyclin A and D1 were markedly reversed by the addition of CRE, in




Figure 2. Effects of TCS on HeLa cell cycle progress and cell cycle regulatory proteins. HeLa cells were treated with indicated doses of TCS for 24 h. Cell number of S phase increased significantly (A) and expressions of cyclin A, D1 and CDK 2, 4 significantly decreased (B). Data represent means \pm SD of three independent experiments (* $p < 0.05$ compared with control). doi:10.1371/journal.pone.0065661.g002

HeLa cells (Fig. 5, C, D), Caski cells (Fig. 6 C, D) and SW1990 cells (Fig. 7 C, D).

Discussion

This study showed, for the first time, that TCS exerted cytotoxic effects on cancer cell viability in a dose-dependent manner (Fig. 1). Caski and C33a cells were sensitive to its inhibitory effect on cell proliferation (Table 1). The results of its anticancer mechanisms clearly show that the increase of S phase in HeLa cells accompanied the decrease of G₁ phase cells, while the number of G₂ phase cells did not change. In Caski and C33a cells, significant increase of G₁ phase and decrease of S and G₂ phase cells were observed. In SW1990 cells, an increase of G₂/M phase

cells was accompanied by a decrease of G₁ phase cells, with no change found in the number of S phase cells (Table 2).

Investigating the changes in multiple regulatory molecules involved in the cell cycle showed that the expressions of cyclin A, D1 and CDK 2, 4 in HeLa cells (Fig. 2), cyclin D1, E and CDK 2, 4 in Caski and C33a cells (Fig. 3), cyclin A, B1, E and CDK 2 in SW1990 cells (Fig. 4) were significantly down-regulated. Blocking the binding site of cell cycle genes to CREB by OGN, a CRE decoy, prevented the TCS-arrested S, G₁ and G₂/M cell cycles in HeLa (Fig. 5 A, B), Caski (Fig. 6 A, B) and SW1990 (Fig. 7 A, B) cells, respectively. The TCS mediated down-regulation of cyclin A, D1 in HeLa cells (Fig. 5 C, D), cyclin D1 in Caski cells (Fig. 6 C, D) and cyclin A in SW1990 cells (Fig. 7 C, D) were reversed by the combination of CRE and TCS.

Table 2. Cell cycle effects of TCS on cancer cells. $n = 3$, Mean \pm SD.

Group	HeLa			Caski			SW1990		
	G1	S	G2/M	G1	S	G2/M	G1	S	G2/M
Control	45.85 \pm 2.12	31.79 \pm 1.51	22.36 \pm 3.04	40.99 \pm 1.34	8.97 \pm 2.51	50.04 \pm 0.38	66.68 \pm 3.21	31.12 \pm 1.11	2.2 \pm 1.24
20 μ g/ml	37.23 \pm 4.28	44.46 \pm 3.19	18.31 \pm 0.97	53.42 \pm 2.47	2.98 \pm 1.18	43.60 \pm 1.63	56.15 \pm 1.56	23.49 \pm 2.0	20.36 \pm 1.56
60 μ g/ml	26.93 \pm 3.47*	49.13 \pm 0.42*	23.94 \pm 1.15	66.15 \pm 2.19*	2.99 \pm 0.99	30.86 \pm 2.10	35.58 \pm 2.41*	21.22 \pm 1.54	43.20 \pm 1.98*
100 μ g/ml	21.12 \pm 3.42*	57.85 \pm 2.63*	21.03 \pm 2.54	80.67 \pm 3.44*	1.33 \pm 2.61*	18.0 \pm 3.51*	11.88 \pm 2.54*	22.27 \pm 1.57	65.85 \pm 1.67*

* $p < 0.05$ vs control group.
doi:10.1371/journal.pone.0065661.t002




Figure 3. Effects of TCS on Caski cell cycle progress and cell cycle regulatory proteins. Caski cells were treated with indicated doses of TCS for 24 h. Cell numbers at G1 phase increased significantly (A) and expressions of cyclin D1, E and CDK2, 4 significantly decreased (B). Data represent means \pm SD of three independent experiments (* p < 0.05 compared with control). doi:10.1371/journal.pone.0065661.g003




Figure 4. Effects of TCS on SW1990 cell cycle progress and cell cycle regulatory proteins. SW1990 cells were treated with indicated doses of TCS for 24 h. Cell numbers at G2/M phase increased significantly (A), the expressions of cyclin A, B1, E and CDK2 significantly decreased (B). Data represent means \pm SD of three independent experiments (* p < 0.05 compared with control). doi:10.1371/journal.pone.0065661.g004




Figure 5. Effects of CRE-decoy on HeLa cell cycle progress and regulatory proteins. TCS-induced increase of cell numbers in S phase was significantly attenuated by the combined treatment of TCS + CRE (A, B). The down-regulated expression of cyclin A and D1 was reversed by TCS + CRE. No effect was observed on other proteins (C, D). Data represent means \pm SD of three independent experiments (* p < 0.05 compared with control, # p < 0.05 compared with TCS). doi:10.1371/journal.pone.0065661.g005

These results confirmed and extended our earlier reports showing that TCS has a cytotoxic effect on cancer cells [3,4], and partly revealed the mechanisms underlying the activation of CREB protein [5,6]. However, it has not been previously reported whether TCS-mediated cell cycle arrest occurs through the combination of CRE and target genes. The present study shows that CRE decoy OGN attenuated the decrease of the expressions of cyclin A and D1 resulting from TCS treatment, and reversed the cell cycle arrests.

The cell cycle is tightly mediated through a complex network of positive and negative cell-cycle regulatory molecules such as CDKs, CKIs and cyclins [15,16]. It has been reported that the formation of active cyclin D/CDK4 and cyclin E/CDK2 complexes controls the progression through G1 phase (G1/S transition). Further, cyclin A binds to and activates CDK2, promoting G1/S and G2/M cell cycle transitions. In late G2, cyclin B/CDK2 is activated, allowing entry into mitosis [17]. In the present study, we found that the decreased expressions of cyclin A, D1 and CDK2, 4 were associated the S phase arrest in HeLa cells (Fig. 2). The decreased-expressions of cyclin D, E and CDK2, 4 were related to G1 phase in Caski and C33a cells (Fig. 3). The expressions of cyclin A, B1 and CDK2 were significantly down-regulated in SW1990 cells, arrested in G2/M phase (Fig. 4). We also found that CDK2 was significantly decreased in cell lines. This result accords with the notion that CDK2 has a distinct and essential function in the mammalian cell-cycle, and its mutation or inhibition causes a G1-phase block [18].

The activation of genes bearing CRE is reported to occur by phosphorylating CREB at Ser 133 and binding to CRE consensus sequences in the promoter region of genes [19,20]. CREB phosphorylation peaks during the G1-S transition and then gradually decreases from S phase to M phase [21]. Combined with our previously reported results [4–6,9], it is logical to hypothesize that cell proliferation genes, bearing the CRE site, bind to CREB and affect its phosphorylation level, thereby interfering with cell division.

CRE locates upstream of the mRNA start sites. It has a key role in the cyclin A [22,23] and D1 [24,25] expression via CREB activation [26–28]. A previous study showed that the association of CREB with vaccinia-related kinase 1 (VRK1) occurred in a cell-cycle-dependent manner from late G1 to S phase. Furthermore, VRK1 specifically enhanced the activity of CRE in cyclin D1 promoter by facilitating the recruitment of phosphorylated CREB to this locus [29]. Giampuzzi et al also found that a significant diminution of CREB protein binding to the cyclin D1 promoter led to a dramatic inhibition of cyclin D1 protein expression in lysyl oxidase (LOX)-up-regulated cells [30]. This phenomenon was also confirmed in fibroblast cells [31], endometrial cell [32], INS cell [33], hepatocyte cell [34] and other cell lines [21,35,36]. In addition, a number of observations suggested cyclin A plays a pivotal role on S and G2/M transition in mammalian cells and this role is consistent with its preferential association with CDK2, instead of CDC2, during S phase [37,38]. CRE was confirmed to be required for efficient activation of the cyclin A promoter in aortic smooth muscle cells [39], fibroblast cells [40] and human




Figure 6. Effects of CRE-decoy on Caski cell cycle progress and regulatory proteins. TCS-induced increase of cell numbers in G1 phase was significantly attenuated by TCS + CRE (A, B). The down-regulated expression of cyclin D1 was reversed by the treatment of TCS + CRE. No effect was observed on other proteins (C, D). Data represent means \pm SD of three independent experiments (* p < 0.05 compared with control, # p < 0.05 compared with TCS).

doi:10.1371/journal.pone.0065661.g006




Figure 7. Effects of CRE-decoy on SW1990 cell cycle progress and regulatory proteins. TCS-induced increase of cell number in G2/M phase was significantly attenuated by the treatment of TCS + CRE (A, B). Down-regulated expression of cyclin A was reversed by the treatment of TCS + CRE. There was no effect on other proteins (C, D). Data represent means \pm SD of three independent experiments (* p < 0.05 compared with control, # p < 0.05 compared with TCS).

doi:10.1371/journal.pone.0065661.g007

embryonal carcinoma cells [22]. The present study clearly shows that combined treatment of TCS and CRE attenuated the decreases of cyclin A and D1 expression, thereby reversing the effect of TCS on cell cycle arrest. Therefore, we suggest that transcription factor CREB is one of the upstream regulators of TCS-induced cell cycle arrest in cancer cells.

References

- Maraganore JM, Joseph M, Bailey MC (1987) Purification and characterization of trichosanthin. Homology to the ricin A chain and implications as to mechanism of abortifacient activity. *J Biol Chem* 262: 11628–11633.
- Li M, Li X, Li JC (2010) Possible mechanisms of trichosanthin-induced apoptosis of tumor cells. *Anat Rec (Hoboken)* 293: 986–992.
- Wang P, Li JC (2007) Trichosanthin-induced specific changes of cytoskeleton configuration were associated with the decreased expression level of actin and tubulin genes in apoptotic HeLa cells. *Life Sci* 81: 1130–1140.
- Wang P, Chen LL, Yan H, Li JC (2009) Trichosanthin suppresses HeLa cell proliferation through inhibition of the PKC/MAPK signaling pathway. *Cell Biol Toxicol* 25: 479–488.
- Wang P, Yan H, Li JC (2007) CREB-mediated Bcl-2 expression in trichosanthin-induced HeLa cell apoptosis. *Biochem Biophys Res Commun* 363: 101–105.
- Wang P, Xu J, Zhang C (2010) CREB, a possible upstream regulator of Bcl-2 in trichosanthin-induced HeLa cell apoptosis. *Mol Biol Rep* 37: 1891–1896.
- Wang P, Xu S, Zhao K, Xiao B, Guo J (2009) Increase in cytosolic calcium maintains plasma membrane integrity through the formation of microtubule ring structure in apoptotic cervical cancer cells induced by trichosanthin. *Cell Biol Int* 33: 1149–1154.
- Yamazoe S, Tanaka H, Sawada T, Amano R, Yamada N, et al. (2010) RNA interference suppression of mucin 5AC (MUC5AC) reduces the adhesive and invasive capacity of human pancreatic cancer cells. *J Exp Clin Cancer Res* 29: 53–61.
- Jiang Q, Bai T, Shen S, Li L, Ding H, et al. (2011) Increase of cytosolic calcium induced by trichosanthin suppresses cAMP/PKC levels through the inhibition of adenylyl cyclase activity in HeLa cells. *Mol Biol Rep* 38: 2863–2868.
- Park YG, Park S, Lim SO, Lee MS, Ryu CK, et al. (2001) Reduction in cyclin D1/Cdk4/retinoblastoma protein signaling by CRE-decoy oligonucleotide. *Biochem Biophys Res Commun* 281: 1213–1219.
- Yu Z, Shah DM (2007) Curcumin down-regulates Ets-1 and Bcl-2 expression in human endometrial carcinoma HEC-1-A cells. *Gynecol Oncol* 106: 541–548.
- Wang P, Li L, Zhang C, Lei Q, Fang W (2010) Effects of fractal surface on C6 glioma cell morphogenesis and differentiation in vitro. *Biomaterials* 31: 6201–6206.
- Peng B, Chang Q, Wang L, Hu Q, Wang Y, et al. (2008) Suppression of human ovarian SKOV-3 cancer cell growth by Duchesnea phenolic fraction is associated with cell cycle arrest and apoptosis. *Gynecol Oncol* 108: 173–181.
- Zhao K, Wang W, Guan C, Cai J, Wang P (2012) Inhibition of gap junction channel attenuates the migration of breast cancer cells. *Mol Biol Rep* 39: 2607–2613.
- Sherr CJ (2000) The Pezcoller lecture: cancer cell cycles revisited. *Cancer Res* 60: 3689–3695.
- Hanahan D, Weinberg RA (2000) The hallmarks of cancer. *Cell* 100: 57–70.
- Kahl CR, Means AR (2003) Regulation of cell cycle progression by calcium/calmodulin-dependent pathways. *Endocr Rev* 24: 719–736.
- Van den Heuvel S, Harlow E (1993) Distinct roles for cyclin-dependent kinases in cell cycle control. *Science* 262: 2050–2054.
- Gonzalez GA, Montminy MR (1989) Cyclic AMP stimulates somatostatin gene transcription by phosphorylation of CREB at serine 133. *Cell* 59: 675–680.
- Pugazhenthil S, Boras T, O'Connor D, Meintzer MK, Heidenreich KA, et al. (1999) Insulin-like growth factor I-mediated activation of the transcription factor cAMP response element-binding protein in PC12 cells. Involvement of p38 mitogen-activated protein kinase-mediated pathway. *J Biol Chem* 274: 2829–2837.
- Saeki K, Yuo A, Takaku F (1999) Cell-cycle-regulated phosphorylation of cAMP response element-binding protein: identification of novel phosphorylation sites. *Biochem J* 338: 49–54.
- Nakamura T, Okuyama S, Okamoto S, Nakajima T, Sekiya S, et al. (1995) Down-regulation of the cyclin A promoter in differentiating human embryonal

Conclusion

Our findings show, for the first time, that TCS induces specific cell cycle arrests in cancer cells by inhibiting the binding of CREB to CRE on genes related to cell proliferation.

Author Contributions

Conceived and designed the experiments: PW. Performed the experiments: SH FW YR XW JC. Analyzed the data: JC PW. Contributed reagents/materials/analysis tools: SH FW YR XW JC. Wrote the paper: PW MH.

- carcinoma cells is mediated by depletion of ATF-1 and ATF-2 in the complex at the ATF/CRE site. *Exp Cell Res* 216: 422–430.
- Shimizu M, Nomura Y, Suzuki H, Ichikawa E, Takeuchi A, et al. (1998) Activation of the rat cyclin A promoter by ATF2 and Jun family members and its suppression by ATF4. *Exp Cell Res* 239: 93–103.
- Watanabe G, Howe A, Lee RJ, Albanese C, Shu IW, et al. (1996) Induction of cyclin D1 by simian virus 40 small tumor antigen. *Proc Natl Acad Sci U S A* 93: 12861–12866.
- Beier F, Lee RJ, Taylor AC, Pestell RG, LuValle P (1999) Identification of the cyclin D1 gene as a target of activating transcription factor 2 in chondrocytes. *Proc Natl Acad Sci U S A* 96: 1433–1438.
- Freeland K, Boxer LM, Latchman DS (2001) The cyclic AMP response element in the Bcl-2 promoter confers inducibility by hypoxia in neuronal cells. *Mol Brain Res* 92: 98–106.
- Riccio A, Ahn S, Davenport CM, Blendy JA, Ginty DD (1999) Mediation by a CREB family transcription factor of NGF-dependent survival of sympathetic neurons. *Science* 286: 2358–2361.
- Wilson BE, Mochon E, Boxer LM (1996) Induction of bcl-2 expression by phosphorylated CREB proteins during B-cell activation and rescue from apoptosis. *Mol Cell Biol* 16: 5546–5556.
- Kang TH, Park DY, Kim W, Kim KT (2008) VRK1 phosphorylates CREB and mediates CCND1 expression. *J Cell Sci* 121: 3035–3041.
- Giampuzzi M, Oleggini R, Albanese C, Pestell R, Di Donato, A (2005) beta-catenin signaling and regulation of cyclin D1 promoter in NRK-49F cells transformed by down-regulation of the tumor suppressor lysyl oxidase. *Biochim Biophys Acta* 1745: 370–381.
- Kwon YJ, Sun Y, Kim NH, Huh SO (2009) Phosphorylation of CREB, a cyclic AMP responsive element binding protein, contributes partially to lysophosphatidic acid-induced fibroblast cell proliferation. *Biochem Biophys Res Commun* 380: 655–659.
- Catalano S, Giordano C, Rizza P, Gu G, Barone I, et al. (2009) Evidence that leptin through STAT and CREB signaling enhances cyclin D1 expression and promotes human endometrial cancer proliferation. *J Cell Physiol* 218: 490–500.
- Kim MJ, Kang JH, Park YG, Ryu GR, Ko SH, et al. (2006) Exendin-4 induction of cyclin D1 expression in INS-1 beta-cells: involvement of cAMP-responsive element. *J Endocrinol* 188: 623–633.
- Moriuchi A, Ido A, Nagata Y, Nagata K, Uto H, et al. (2003) A CRE and the region occupied by a protein induced by growth factors contribute to up-regulation of cyclin D1 expression in hepatocytes. *Biochem Biophys Res Commun* 300: 415–421.
- Schneider G, Oswald F, Wahl C, Greten FR, Adler G et al. (2002) Cyclosporine inhibits growth through the activating transcription factor/cAMP-responsive element-binding protein binding site in the cyclin D1 promoter. *J Biol Chem* 277: 43599–607.
- Pradeep A, Sharma C, Sathyanarayana P, Albanese C, Fleming JV, et al. (2004) Gastrin-mediated activation of cyclin D1 transcription involves beta-catenin and CREB pathways in gastric cancer cells. *Oncogene* 23: 3689–3699.
- Pines J, Hunter T (1990) Human cyclin A is adenovirus E1A-associated protein p60 and behaves differently from cyclin B. *Nature* 346: 760–763.
- Pines J, Hunter T (1989) Isolation of a human cyclin cDNA: evidence for cyclin mRNA and protein regulation in the cell cycle and for interaction with p34cdc2. *Cell* 58: 833–846.
- Kothapalli D, Stewart SA, Smyth EM, Azonobi I, Pure E, et al. (2003) Prostacyclin receptor activation inhibits proliferation of aortic smooth muscle cells by regulating cAMP response element-binding protein- and pocket protein-dependent cyclin a gene expression. *Mol Pharmacol* 64: 249–258.
- Desdouets C, Matesic G, Molina CA, Foulkes NS, Sassone-Corsi P, et al. (1995) Cell cycle regulation of cyclin A gene expression by the cyclic AMP-responsive transcription factors CREB and CREM. *Mol Cell Biol* 15: 3301–3309.