

# Cyclic Cubic Fields of Given Conductor and Given Index

Alan K. Silvester, Blair K. Spearman, and Kenneth S. Williams

*Abstract.* The number of cyclic cubic fields with a given conductor and a given index is determined.

## 1 Introduction

Let  $K$  be a cyclic cubic extension of  $\mathbb{Q}$  so that  $[K:\mathbb{Q}] = 3$  and  $\text{Gal}(K/\mathbb{Q}) \simeq \mathbb{Z}/3\mathbb{Z}$ . By the Kronecker–Weber theorem [10, p. 289] there exists a positive integer  $m$  such that the cyclotomic field  $\mathbb{Q}(e^{2\pi i/m}) \supseteq K$ . The smallest such  $m$  is called the conductor of  $K$  and is denoted by  $f(K)$ . The discriminant of  $K$  is given by  $d(K) = f(K)^2$  [8, p. 831]. The conductor  $f(K)$  of a cyclic cubic field is of the form

$$(1.1) \quad f = p_1 p_2 \cdots p_r,$$

where  $r \in \mathbb{N}$  and  $p_1, \dots, p_r$ , are distinct integers from the set

$$(1.2) \quad P = \{9\} \cup \{p \text{ (prime)} \equiv 1 \pmod{3}\} = \{7, 9, 13, 19, 31, 37, \dots\},$$

see [8, p. 831]. Moreover each positive integer  $f$  of the form (1.1) is the conductor of some cyclic cubic field; indeed it is the conductor of  $2^{r-1}$  cyclic cubic fields [8, p. 831]. For any cubic field  $K$  it is known that its field index  $i(K) = 1$  or  $2$  [5, p. 234]. For  $f$  of the form (1.1) and  $i \in \{1, 2\}$ , we define

$$(1.3) \quad N(f, i) = \text{number of cyclic cubic fields } K \text{ with } f(K) = f \text{ and } i(K) = i,$$

so that

$$(1.4) \quad N(f, 1) + N(f, 2) = 2^{r-1}.$$

In this paper we determine  $N(f, 1)$  and  $N(f, 2)$ .

It is well known that each prime  $p \equiv 1 \pmod{3}$  has a unique representation in the form

$$(1.5) \quad 4p = a^2 + 27b^2, \quad a, b \in \mathbb{N},$$

---

Received by the editors July 2, 2004; revised July 16, 2006.

The second and third authors were supported by grants from the Natural Sciences and Engineering Research Council of Canada.

AMS subject classification: Primary: 11R16; secondary: 11R29.

Keywords: Discriminant, conductor, index, cyclic cubic field.

©Canadian Mathematical Society 2006.

see [1, Theorem 3.1.3, p. 105; Lemma 3.0.1, p. 101]. Clearly for such a representation we have  $a \equiv b \pmod{2}$  and

$$(1.6) \quad \gcd(a, b) = 1 \text{ or } 2.$$

It is a classical result of Gauss that 2 is a cubic residue  $\pmod{p}$  if and only if  $\gcd(a, b) = 2$ , see [1, Theorem 7.1.1, p. 213]. We set

$$(1.7) \quad P_1 = \{9\} \cup \{p \text{ (prime)} \equiv 1 \pmod{3}, 4p = a^2 + 27b^2, \gcd(a, b) = 1\}$$

and

$$(1.8) \quad P_2 = \{p \text{ (prime)} \equiv 1 \pmod{3}, 4p = a^2 + 27b^2, \gcd(a, b) = 2\},$$

so that

$$(1.9) \quad P_1 \cup P_2 = P, \quad P_1 \cap P_2 = \phi.$$

Clearly

$$P_1 = \{7, 9, 13, 19, 37, \dots\}, \quad P_2 = \{31, 43, 109, 127, \dots\}.$$

If  $p$  is a prime in  $P_1$ , then  $a \equiv b \equiv 1 \pmod{2}$ . Replacing  $b$  by  $-b$ , if necessary, we may suppose that  $a \equiv b \pmod{4}$ . Set  $x = (a - b)/4 \in \mathbb{Z}$ ,  $y = b \in \mathbb{Z}$ . Then  $4x^2 + 2xy + 7y^2 = p$ . Conversely if  $p = 4x^2 + 2xy + 7y^2$  for some  $x, y \in \mathbb{Z}$  then  $y$  is odd,  $\gcd(x, y) = 1$  and  $4p = a^2 + 27b^2$  with  $a = |4x + y|$ ,  $b = |y|$  and  $\gcd(a, b) = \gcd(4x + y, y) = \gcd(4x, y) = \gcd(x, y) = 1$ . Thus the primes in  $P_1$  are precisely those which can be expressed in the form  $4x^2 + 2xy + 7y^2$  for some  $x, y \in \mathbb{Z}$ . The primes in  $P_2$  are precisely those which can be expressed in the form  $x^2 + 27y^2$  for some  $x, y \in \mathbb{Z}$ .

Now suppose that  $f$  is of the form (1.1) with

$$p_1, p_2, \dots, p_u \in P_1 \quad \text{and} \quad p_{u+1}, p_{u+2}, \dots, p_r \in P_2,$$

where  $u \in \{0, 1, \dots, r\}$ . In Section 5 we prove the following result.

**Theorem** *With the above notation, we have*

$$N(f, 1) = \frac{1}{3}(2^r - (-1)^u 2^{r-u}), \quad N(f, 2) = \frac{1}{3}(2^{r-1} + (-1)^u 2^{r-u}).$$

In Sections 2, 3, 4 we give some results on representations of integers by binary quadratic forms which will be needed in the proof of this theorem.

## 2 The Form Class Group $H(d)$

Let  $H(d)$  denote the set of classes of primitive, positive-definite, integral binary quadratic forms  $(a, b, c) = ax^2 + bxy + cy^2$  of discriminant  $d = b^2 - 4ac \equiv 0$  or  $1 \pmod{4}$  under the action of the modular group. As  $ax^2 + bxy + cy^2$  is positive-definite, we have  $a > 0$  and  $d < 0$ . The class of the form  $(a, b, c)$  is denoted by  $[a, b, c]$ . Multiplication of classes of  $H(d)$  is due to Gauss and is described, for example, in [2]. With respect to multiplication,  $H(d)$  is a finite abelian group called the form class group of discriminant  $d$ . The order of  $H(d)$  is called the form class number of discriminant  $d$  and is denoted by  $h(d)$ . The identity  $I$  of the group  $H(d)$  is the principal class

$$I = \begin{cases} [1, 0, -d/4] & \text{if } d \equiv 0 \pmod{4}, \\ [1, 1, (1-d)/4] & \text{if } d \equiv 1 \pmod{4}. \end{cases}$$

The inverse of the class  $K = [a, b, c] \in H(d)$  is the class  $K^{-1} = [a, -b, c] \in H(d)$ . Each class of  $H(d)$  contains one and only one form  $(a, b, c)$  with

$$(2.1) \quad -a < b \leq a \leq c, \quad b \geq 0 \text{ if } a = c, \quad b^2 - 4ac = d, \quad \gcd(a, b, c) = 1,$$

see [4, pp. 68-71]. Let  $n \in \mathbb{N}$ . If  $x$  and  $y$  are integers such that  $n = ax^2 + bxy + cy^2$ , then  $(x, y)$  is called a representation of the positive integer  $n$  by the form  $(a, b, c)$ . As  $(a, b, c)$  is a positive-definite form, the number  $R_{(a,b,c)}(n)$  of representations of  $n$  by the form  $(a, b, c)$  is finite. If in addition the representation  $(x, y)$  satisfies  $\gcd(x, y) = 1$ , then the representation is called primitive. The number of primitive representations of  $n$  by the form  $(a, b, c)$  is denoted by  $P_{(a,b,c)}(n)$ . Clearly,

$$(2.2) \quad R_{(a,b,c)}(n) = \sum_{e^2|n} P_{(a,b,c)}(n/e^2).$$

If  $(A, B, C)$  is a form equivalent to  $(a, b, c)$  it is well known that  $R_{(A,B,C)}(n) = R_{(a,b,c)}(n)$  and  $P_{(A,B,C)}(n) = P_{(a,b,c)}(n)$ . Hence we can define the number of representations of  $n \in \mathbb{N}$  by the class  $K \in H(d)$  by

$$(2.3) \quad R_K(n) = R_{(a,b,c)}(n) \text{ for any } (a, b, c) \in K$$

and the number of primitive representations of  $n \in \mathbb{N}$  by the class  $K \in H(d)$  by

$$(2.4) \quad P_K(n) = P_{(a,b,c)}(n) \text{ for any } (a, b, c) \in K.$$

From (2.2)–(2.4) we deduce that for  $n \in \mathbb{N}$  and  $K \in H(d)$

$$(2.5) \quad R_K(n) = \sum_{e^2|n} P_K(n/e^2).$$

In particular, if  $n \in \mathbb{N}$  is squarefree, we have

$$(2.6) \quad R_K(n) = P_K(n).$$

As each representation  $(x, y)$  of  $n$  by  $(a, b, c)$  gives a representation  $(x, -y)$  of  $n$  by  $(a, -b, c)$  and conversely, we have for  $n \in \mathbb{N}$  and  $K \in H(d)$

$$(2.7) \quad R_K(n) = R_{K^{-1}}(n), \quad P_K(n) = P_{K^{-1}}(n).$$

For  $n_1, n_2 \in \mathbb{N}$  with  $n_1$  squarefree,  $n_2$  squarefree and  $\gcd(n_1, n_2) = 1$ , it is known that

$$(2.8) \quad R_K(n_1 n_2) = \frac{1}{w(d)} \sum_{K_1 K_2 = K} R_{K_1}(n_1) R_{K_2}(n_2),$$

where  $K_1, K_2$  run through all the classes of  $H(d)$  whose product is  $K$ , and

$$(2.9) \quad w(d) = 6, 4 \text{ or } 2 \text{ according as } d = -3, d = -4 \text{ or } d < -4,$$

see [9, (29) and Lemma 5.5]. The largest positive integer  $f$  such that  $f^2 \mid d$  with  $\Delta = d/f^2 \equiv 0 \text{ or } 1 \pmod{4}$  is called the conductor of  $d$ . By a theorem of Dirichlet, see [6], we have for  $\gcd(n, f) = 1$

$$(2.10) \quad \sum_{K \in H(d)} R_K(n) = w(d) \sum_{e \mid n} \left(\frac{d}{e}\right) = w(d) \sum_{e \mid n} \left(\frac{\Delta}{e}\right),$$

where  $\left(\frac{d}{*}\right)$  is the Legendre–Jacobi–Kronecker symbol of discriminant  $d$ . If  $p$  is a prime such that  $\left(\frac{d}{p}\right) = 1$ , then there is at least one class  $C \in H(d)$  which represents  $p$ . If  $C = C^{-1}$ , then  $C$  is the only class of  $H(d)$  representing  $p$  and  $R_C(p) = 2w(d)$ . If  $C \neq C^{-1}$ , then  $C$  and  $C^{-1}$  are the only classes of  $H(d)$  representing  $p$  and  $R_C(p) = R_{C^{-1}}(p) = w(d)$ . See [9, Lemma 5.3].

### 3 Representations of Integers by [1,0,3]

From (2.1) with  $d = -12$  we find

$$H(-12) = \{I\}, \quad h(-12) = 1,$$

where

$$I = [1, 0, 3].$$

Here  $f = 2$  and  $\Delta = -3$ .

**Lemma 3.1** *Let  $p_1, \dots, p_t$  ( $t \geq 0$ ) be distinct primes  $\equiv 1 \pmod{3}$ . Then*

$$\begin{aligned} R_I(p_1 \cdots p_t) &= 2^{t+1}, & P_I(p_1 \cdots p_t) &= 2^{t+1}, \\ R_I(9p_1 \cdots p_t) &= 2^{t+1}, & P_I(9p_1 \cdots p_t) &= 0. \end{aligned}$$

**Proof** If  $n \in \mathbb{N}$  is such that  $\gcd(n, 2) = 1$ , by (2.9) and (2.10) with  $d = -12$ , we have

$$(3.1) \quad R_I(n) = 2 \sum_{e|n} \left( \frac{-3}{e} \right).$$

Taking  $n = p_1 \cdots p_t$ , as  $\left( \frac{-3}{p_i} \right) = 1$  ( $i = 1, \dots, t$ ), we obtain

$$R_I(p_1 \cdots p_t) = 2 \sum_{e|p_1 \cdots p_t} 1 = 2 \cdot 2^t = 2^{t+1}.$$

Then, appealing to (2.6), we obtain

$$P_I(p_1 \cdots p_t) = 2^{t+1}.$$

Taking  $n = 9p_1 \cdots p_t$  in (3.1), since  $\left( \frac{-3}{3} \right) = 0$  we obtain

$$R_I(9p_1 \cdots p_t) = 2 \sum_{e|9p_1 \cdots p_t} \left( \frac{-3}{e} \right) = 2 \sum_{e|p_1 \cdots p_t} \left( \frac{-3}{e} \right) = 2^{t+1}.$$

Finally, by (2.5), we have

$$R_I(9p_1 \cdots p_t) = P_I(9p_1 \cdots p_t) + P_I(p_1 \cdots p_t),$$

so that

$$P_I(9p_1 \cdots p_t) = 2^{t+1} - 2^{t+1} = 0.$$

This completes the proof of the lemma. ■

#### 4 Representations of Integers by [1,0,27] and [4,2,7]

From (2.1) with  $d = -108$  we find

$$H(-108) = \{I, A, A^2\} \simeq \mathbb{Z}/3\mathbb{Z}, \quad h(-108) = 3,$$

where

$$I = [1, 0, 27], \quad A = [4, 2, 7], \quad A^2 = [4, -2, 7], \quad A^3 = I.$$

Here  $f = 6$  and  $\Delta = -3$ .

Let  $p$  be a prime with  $p \equiv 1 \pmod{3}$ . Then

$$\left( \frac{d}{p} \right) = \left( \frac{-108}{p} \right) = \left( \frac{-2^2 \cdot 3^3}{p} \right) = \left( \frac{-3}{p} \right) = 1,$$

so that  $p$  is represented by some class in  $H(-108)$ . If  $p$  is represented by  $I$ , then (as  $I = I^{-1}$ )  $I$  is the only class representing  $p$ , and

$$(4.1) \quad R_I(p) = 4, \quad R_A(p) = R_{A^2}(p) = 0.$$

If  $p$  is represented by  $A$  or  $A^2$ , then (as  $A \neq A^{-1}$ ) the only classes of  $H(-108)$  representing  $p$  are  $A$  and  $A^2$ , and

$$(4.2) \quad R_I(p) = 0, \quad R_A(p) = R_{A^2}(p) = 2.$$

Now let  $m$  be a product of distinct primes  $\equiv 1 \pmod{3}$ . By (2.9) and (2.10) we have

$$R_I(m) + R_A(m) + R_{A^2}(m) = 2 \sum_{e|m} \left( \frac{-3}{e} \right) = 2^{\tau(m)+1},$$

where  $\tau(m)$  denotes the number of primes dividing  $m$ . As  $R_A(m) = R_{A^{-1}}(m) = R_{A^2}(m)$  by (2.7), we deduce that

$$(4.3) \quad R_A(m) = R_{A^2}(m) = 2^{\tau(m)} - \frac{1}{2}R_I(m).$$

By (2.8) we have for  $p \nmid m$

$$(4.4) \quad R_I(pm) = \frac{1}{2} (R_I(p)R_I(m) + R_A(p)R_{A^2}(m) + R_{A^2}(p)R_A(m)).$$

Appealing to (4.1)–(4.4), we obtain

$$(4.5) \quad R_I(pm) = \begin{cases} 2R_I(m) & \text{if } R_I(p) > 0, \\ 2^{\tau(m)+1} - R_I(m) & \text{if } R_A(p) > 0. \end{cases}$$

We now use (4.5) to prove the following result.

**Lemma 4.1** *Let  $p_1, \dots, p_l$  be  $l(\geq 0)$  distinct primes  $\equiv 1 \pmod{3}$ , which are represented by  $I = [1, 0, 27]$ , and let  $q_1, \dots, q_m$  be  $m(\geq 0)$  distinct primes  $\equiv 1 \pmod{3}$ , which are represented by  $A = [4, 2, 7]$ . Then*

$$R_I(p_1 \cdots p_l q_1 \cdots q_m) = \frac{1}{3} (2^{l+m+1} + (-1)^m 2^{l+2}),$$

$$R_A(p_1 \cdots p_l q_1 \cdots q_m) = R_{A^2}(p_1 \cdots p_l q_1 \cdots q_m) = \frac{1}{3} (2^{l+m+1} - (-1)^m 2^{l+1}).$$

**Proof** By (4.5) we obtain

$$\begin{aligned}
R_I(p_1 \cdots p_l q_1 \cdots q_m) &= 2R_I(p_1 \cdots p_{l-1} q_1 \cdots q_m) \\
&= 2^2 R_I(p_1 \cdots p_{l-2} q_1 \cdots q_m) \\
&= \cdots \\
&= 2^l R_I(q_1 \cdots q_m) \\
&= 2^l (2^m - R_I(q_1 \cdots q_{m-1})) \\
&= 2^l (2^m - 2^{m-1} + R_I(q_1 \cdots q_{m-2})) \\
&= \cdots \\
&= 2^l (2^m - 2^{m-1} + 2^{m-2} - \cdots + (-1)^{m-2} 2^2 + (-1)^{m-1} R_I(q_1)) \\
&= 2^l (2^m - 2^{m-1} + 2^{m-2} - \cdots + (-1)^{m-2} 2^2) \\
&= \frac{1}{3} (2^{l+m+1} + (-1)^m 2^{l+2}),
\end{aligned}$$

as required. Then, by (4.3), we obtain

$$\begin{aligned}
R_A(p_1 \cdots p_l q_1 \cdots q_m) &= 2^{l+m} - \frac{1}{2} \left( \frac{1}{3} (2^{l+m+1} + (-1)^m 2^{l+2}) \right) \\
&= \frac{1}{3} (2^{l+m+1} - (-1)^m 2^{l+1}),
\end{aligned}$$

as asserted. ■

## 5 Proof of Theorem

There is a one-to-one correspondence between cyclic cubic fields  $K$  and triples  $(a, b, f) \in \mathbb{N}^3$  with

$$\begin{aligned}
a^2 + 27b^2 &= 4f, \quad \gcd(a, b) = 1 \text{ or } 2, \quad f = p_1 \cdots p_r, \\
r \in \mathbb{N}, \quad p_1, \dots, p_r &\in P, \quad p_i \neq p_j \quad (1 \leq i < j \leq r),
\end{aligned}$$

see [3, Section 6.4.6, pp. 336–343]. The cyclic cubic field corresponding to the triple  $(a, b, f)$  is  $K = \mathbb{Q}(\theta)$ , where  $\theta^3 - 3f\theta + fa = 0$ . The conductor of  $K$  is  $f$ . The index of  $K$  is

$$i(K) = \begin{cases} 2 & \text{if } a \text{ is even,} \\ 1 & \text{if } a \text{ is odd,} \end{cases}$$

see [7, Theorem 4, p. 585]. If  $a$  is even, then  $b$  is even and  $(\frac{a}{2})^2 + 27(\frac{b}{2})^2 = f$  with  $\gcd(\frac{a}{2}, \frac{b}{2}) = 1$ . Thus

$$N(f, 2) = \frac{1}{4} P_{[1,0,27]}(f).$$

First suppose that  $9 \nmid f$ . We may suppose that  $p_1, \dots, p_u \in P_1$  (so they are represented by  $[4, 2, 7]$ ) and  $p_{u+1}, \dots, p_r \in P_2$  (so they are represented by  $[1, 0, 27]$ ) with  $u \in \{0, 1, 2, \dots, r\}$ . Then, by (2.6) and Lemma 4.1 (with  $l = r - u$  and  $m = u$ ), we have

$$P_{[1,0,27]}(f) = R_{[1,0,27]}(f) = \frac{1}{3}(2^{r+1} + (-1)^u 2^{r-u+2}),$$

so that

$$N(f, 2) = \frac{1}{3}(2^{r-1} + (-1)^u 2^{r-u}), \quad 9 \nmid f.$$

Now suppose that  $9 \mid f$ . We may suppose that  $p_1 = 9, p_2, \dots, p_u \in P_1$  (so they are represented by  $[4, 2, 7]$ ) and  $p_{u+1}, \dots, p_r \in P_2$  (so they are represented by  $[1, 0, 27]$ ). As  $9 \mid f$  we have

$$f = x^2 + 27y^2 \iff f/9 = (x/3)^2 + 3y^2$$

so that

$$R_{[1,0,27]}(f) = R_{[1,0,3]}(f/9).$$

As  $f/9$  is squarefree, we have

$$R_{[1,0,27]}(f/9) = P_{[1,0,27]}(f/9).$$

From (2.5) we deduce

$$R_{[1,0,27]}(f) = P_{[1,0,27]}(f) + P_{[1,0,27]}(f/9).$$

Thus

$$P_{[1,0,27]}(f) = R_{[1,0,3]}(f/9) - R_{[1,0,27]}(f/9).$$

Appealing to Lemma 3.1 (with  $t = r - 1$ ) and Lemma 4.1 (with  $l = r - u$  and  $m = u - 1$ ), we obtain

$$P_{[1,0,27]}(f) = 2^r - \frac{1}{3}(2^r + (-1)^{u-1} 2^{r-u+2}) = \frac{1}{3}(2^{r+1} + (-1)^u 2^{r-u+2})$$

so that

$$N(f, 2) = \frac{1}{3}(2^{r-1} + (-1)^u 2^{r-u}), \quad 9 \mid f.$$

Finally, from (1.4), we obtain in both cases

$$N(f, 1) = 2^{r-1} - N(f, 2) = \frac{1}{3}(2^r - (-1)^u 2^{r-u}).$$

This completes the proof of the theorem. ■



## References

- [1] B. C. Berndt, R. J. Evans and K. S. Williams, *Gauss and Jacobi Sums*. Wiley, New York, 1998.
- [2] D. A. Buell, *Binary Quadratic Forms. Classical Theory and Modern Computations*. Springer-Verlag, New York, 1989.
- [3] H. Cohen, *A Course in Computational Algebraic Number Theory*. Graduate Texts in Mathematics 138, Springer-Verlag, Berlin, 1993.
- [4] L. E. Dickson, *Introduction to the Theory of Numbers*. Dover, New York, 1957.
- [5] H. T. Engstrom, *On the common index divisors of an algebraic field*. Trans. Amer. Math. Soc. **32** (1930), no. 2, 223–237.
- [6] P. Kaplan and K. S. Williams, *On a formula of Dirichlet*. Far East J. Math. Sci. **5** (1997), no. 1, 153–157.
- [7] P. Llorente and E. Nart, *Effective determination of the decomposition of the rational primes in a cubic field*. Proc. Amer. Math. Soc. **87** (1983), no. 4, 579–585.
- [8] D. C. Mayer, *Multiplicities of dihedral discriminants*. Math. Comp. **58** (1992), 831–847.
- [9] H. Muzaffar and K. S. Williams, *Evaluation of Weber's functions at quadratic irrationalities*. JP J. Algebra Number Theory Appl. **4**(2004), no. 2, 209–259.
- [10] W. Narkiewicz, *Elementary and Analytic Theory of Algebraic Numbers*. Second edition. Springer-Verlag, Berlin, 1990.

*Department of Mathematics and Statistics*  
*Okanagan University College*  
*Kelowna, BC*  
*V1V 1V7*  
*e-mail: mascdman@canada.com*  
*bspearman@ouc.bc.ca*

*School of Mathematics and Statistics*  
*Carleton University*  
*Ottawa, ON*  
*K1S 5B6*  
*e-mail: williams@math.carleton.ca*