

# Cycloaspeptide A and Pseurotin A from the Endophytic Fungus *Penicillium janczewskii*

Guillermo Schmeda-Hirschmann<sup>a,\*</sup>, Emilio Hormazabal<sup>a</sup>, Jaime A. Rodríguez<sup>b</sup>, and Cristina Theodulóz<sup>b</sup>

<sup>a</sup> Laboratorio de Química de Productos Naturales, Instituto de Química de Recursos Naturales, Universidad de Talca, Casilla 747, Talca, Chile. Fax: +56 71-20 04 48. E-mail: schmeda@utalca.cl

<sup>b</sup> Departamento de Ciencias Básicas Biomédicas, Facultad de Ciencias de la Salud, Universidad de Talca, Casilla 747, Talca, Chile

\* Author for correspondence and reprint requests

Z. Naturforsch. **63c**, 383–388 (2008); received December 13, 2007

*Penicillium janczewskii* K. M. Zalessky was isolated as an endophytic fungus from the phloem of the Chilean gymnosperm *Prumnopitys andina*. When grown in liquid yeast extract-malt extract-glucose broth, the fungus produced two main secondary metabolites. The compounds were for the first time isolated from this species and identified by spectroscopic methods as pseurotin A and cycloaspeptide A. This is the first report on the production of cyclic peptides by endophytic fungi from Chilean gymnosperms. Pseurotin A and cycloaspeptide A presented low cytotoxicity towards human lung fibroblasts with  $IC_{50} \geq 1000 \mu M$ . Pseurotin A showed a moderate effect against the phytopathogenic bacteria *Erwinia carotovora* and *Pseudomonas syringae*, with  $IC_{50}$  values of 220 and  $112 \mu g ml^{-1}$ , respectively.

**Key words:** *Penicillium janczewskii*, Cycloaspeptide A, Pseurotin A, Endophyte

## Introduction

Little is known about the metabolites produced by endophytic fungi of Chilean native plants, including gymnosperms (Hormazabal *et al.*, 2005; Schmeda-Hirschmann *et al.*, 2005).

Endophytic fungi are widespread in plants and represent a relevant component of plant communities (Gunatillaka, 2006; Owen and Hundley, 2004). The relation of the plant and fungus can range from symbiosis to parasitism, depending on the host-invader balance (Kogel *et al.*, 2006). It has been shown that endophytes protect plants against pathogens and that host affinity is mediated by the leaf chemistry (Arnold *et al.*, 2003). Those fungi have been recognized as a valuable source of novel bioactive metabolites including taxol, cryptocin, cryptocandin, jesterone, oocydin, isopestacin, the pseudomycins and ambuic acid (Strobel, 2002; Li *et al.*, 1996) as well as podophyllotoxin (Puri *et al.*, 2006) and volatile organic compounds (Strobel, 2006).

The metabolite production of endophytic fungi has been shown to be related with the genotypic diversity. In a comparative study of *Cylindrocarpon destructans* and *Heliscus lugdunensis* populations isolated from the roots of a single tree,

Seymour *et al.* (2004) found a good correlation between genotype classification and the natural product patterns of the crude extracts obtained from the microbial cultures. The role of arbuscular mycorrhizal fungi and agricultural practices has been recently revised by Gosling *et al.* (2006).

Following our studies on bioactive products from endophytic fungi isolated from Chilean gymnosperms, we now report the secondary metabolites produced by *Penicillium janczewskii* obtained from the Podocarpaceae *Prumnopitys andina* (Poepp. ex Endl.) de Laub, known in Chile as “lleuque”.

## Materials and Methods

### Isolation and culture

The fungus was isolated from wood pieces collected on the western Andean slopes near Las Trancas, Chillán as described in a previous report (Schmeda-Hirschmann *et al.*, 2005). The microorganism was identified by Prof. Eduardo Piontelli, Universidad Católica de Valparaíso; it is kept at the microbial strain collection of Universidad de Talca. *Penicillium janczewskii* was cultured in yeast extract-malt extract-glucose broth (YMG broth, 4 g yeast extract, 10 g malt extract, 10 g glu-

cose in 1 l water, pH 5.5) in Erlenmeyer flasks (14.4 l) at 25 °C under constant shaking (150 rpm). Once glucose in the medium was consumed (13 d), mycelium and culture broth were separated by filtration and extracted separately with EtOAc (3 × 2 l). Some 640 mg of EtOAc-solubles were obtained from the culture filtrate and 2.15 g from the mycelium. The extract obtained from the culture medium was dissolved in MeOH and permeated in a Sephadex LH-20 column (length, 70 cm; internal diameter, 2 cm) with MeOH. Some 200 fractions of 4 ml each were collected and pooled together according to the TLC patterns. Fractions 56–59 (138 mg) were rechromatographed on silica gel with a petroleum ether (PE)/ethyl acetate (EtOAc)/EtOAc/acetone gradient. Some 190 fractions were collected and pooled together in 12 groups according to the TLC patterns. The pooled fractions 5 and 6 were purified by preparative TLC (silica gel, PE/EtOAc 3:7) affording 30 mg pseurotin A (**1**) and 10 mg cycloaspeptide A (**2**). The other fractions did not contained compounds of interest based on TLC analysis and NMR measurements and were not further investigated.

### Compounds

Melting points were determined on a Koffler hot stage apparatus (Electrothermal 9100, Essex, UK) and are uncorrected. Optical rotations were

obtained for solutions in CHCl<sub>3</sub> (concentrations expressed in g 100 ml<sup>-1</sup>) on a Jasco DIP 370 polarimeter. IR spectra were recorded on a Nicolet Nexus FT-IR instrument. <sup>1</sup>H NMR spectra were recorded at 400 MHz and <sup>13</sup>C NMR data were obtained at 100 MHz on a Bruker spectrometer ( $\delta$  scale). TLC spots were visualized by spraying the chromatograms with H<sub>2</sub>SO<sub>4</sub>/EtOH (10:90) and heating at 110 °C for 3 min. Column chromatography was performed over Merck Kieselgel 60, particle size 0.063–0.200 mm. Mass spectra are presented as *m/z* (% rel. int.).

*Pseurotin A (1)*: Colourless crystals, m.p. 162–163.5 °C. – MS (EI): *m/z* = 381 [M-50] (1), 365 (2), 347 (4), 315 (37), 242 (64), 155 (45), 105 (100). – C<sub>22</sub>H<sub>25</sub>NO<sub>8</sub> (calcd. for C<sub>22</sub>H<sub>25</sub>NO<sub>8</sub>: 431.441). – FT-IR:  $\nu$  = 3433, 3292, 2964, 2927, 1723, 1704, 1686, 1637, 1080 cm<sup>-1</sup>. –  $[\alpha]_D^{20}$  = -5° (c 0.5 methanol). – The spectroscopic data agree with those reported by Bloch and Tamm (1981) and Breitenstein *et al.* (1981). – <sup>1</sup>H and <sup>13</sup>C NMR: see Table I.

*Cycloaspeptide A (2)*: Colourless resin. – EI-MS: *m/z* = 641 (29), 497 (45) [M-144], 421 (10) (497–76), 378 (7), 352 (19), 306 (9), 266 (20), 232 (9), 150 (34), 134 (100), 120 (31). – C<sub>36</sub>H<sub>43</sub>N<sub>5</sub>O<sub>6</sub> (641). – FT-IR:  $\nu$  = 3316, 1670, 1656, 1632, 1593, 1516, 1447, 753, 702 cm<sup>-1</sup>. – <sup>1</sup>H and <sup>13</sup>C NMR: see Table II.

Table I. <sup>1</sup>H and <sup>13</sup>C NMR data of pseurotin A (**1**) (400 and 100 MHz, respectively, CDCl<sub>3</sub>).

H	$\delta_H$	$\delta_C$	HMBC
2	–	166.53 s	–
3	–	113.56 s	1.65
4	–	196.45 s	1.65
5	–	92.67 s	8.39
6	–	185.73 s	1.65
7 NH	8.39 s	–	92, 73
8	–	90.25 s	3.43
9	4.67 s	73.15 d	8.39, 3.43
10	4.57 d (4.4)	70.82 d	–
11	4.73 dd (9, 4.4)	70.63 d	5.56, 4.73
12	5.25 br dd (11, 9)	126.50 d	2.05, 5.25
13	5.56 ddd (11, 7.8, 7.8)	137.04 d	0.96, 2.09
14	2.09 dq (7.8, 7.3)	21.43 t	0.96, 5.25
15	0.96 t (7.3)	14.11 q	137, 21; 2.09
16	1.65 s	6.00 q	196, 185, 113
17	–	194.94 s	8.30
18	–	132.38 s	–
19, 23	8.30 d (8.30, 1.0)	130.70 d	194, 134; 7.63
20, 22	7.47 dd (7.83, 7.80, 1.0)	128.74 d	128, 132
21	7.63 ddd (8.30, 7.80, 1.0)	134.76 d	130; 8.30, 7.63
OMe	3.43 s (3 H)	51.76 q	90

Table II. <sup>1</sup>H and <sup>13</sup>C NMR data of cycloaspeptide A (2) (400 and 100 MHz, respectively, CDCl<sub>3</sub>).

Residue and position	$\delta_{\text{H}}$	$\delta_{\text{C}}$	HMBC
Alanine			
NH	6.96 d (6.9)	–	
C $\alpha$	4.41 dq (6.9, 6.9)	44.16 d	16.22, 173.72
C $\beta$	0.40 d (6.4)	16.22 q	44.16, 173.72
CO		173.93 s	0.40, 2.87
Phenylalanine			
N-CH <sub>3</sub>	2.87 s	30.10 q	63.37, 173.93
C $\alpha$	5.21 dd (11.7, 3.4)	63.37 d	30.10, 168.06
C $\beta$	2.96 dd (13, 12.2), 3.48 dd (14.2, 4)	34.07 t	129.40
C1	–	137.61 s	7.33
C2, C6	7.16 br d (8.2)	129.40 d (2 C)	34.07
C3, C5	7.33 m	129.18 d (2 C)	129.42, 137.61
C4	7.26 m	134.33 d	
CO	–	169.63 s	
Leucine			
NH	7.30 d (7.8)	–	48.89, 169.63
C $\alpha$	4.72 ddd (8, 8, 5)	48.89 d	24.88, 41.25, 169.73
C $\beta$	1.36 ddd (13.0, 8.0, 5.5), 1.86 ddd (13.0, 8.5, 5.5)	41.25 t	23.32, 48.89, 169.73
C $\gamma$	1.67 m	24.88 d	
CH <sub>3</sub>	0.98 d (6.4)	23.32 q	24.88,
CH <sub>3</sub>	0.99 d (6.4)	21.96 q	24.88, 41.25
CO	–	169.73 s	1.36
Tyrosine			
N-CH <sub>3</sub>	2.69 s	39.13 q	70.1, 169.7
C $\alpha$	3.80 dd (11.2, 4)	70.01 d	2.69
C $\beta$	3.35 dd (14.2, 11.3), 3.48 dd (14.2, 4)	32.12 t	70.1, 130.27
C1	–	130 s	
C2, C6	6.98 dbr (2 H) (8.6)	130.27 d (2 C)	32.12, 154.82
C3, C5	6.79 dbr (2 H) (8.6)	115.70 d (2 C)	130.27, 154.82
C4	–	154.82 s	6.79
CO	–	168.19 s	
<i>o</i> -Aminobenzoic acid			
NH	12.00 s	–	115.04, 122.27, 168.19
C1	–	115.04 s	
C2	–	141.52 s	
C3	7.44 dbr (7.8)	126.92 d	
C4	6.96 t (7.8)	122.27 d	115.17, 121.03
C5	7.46 t (8.6)	127.25 d	
C6	8.86 d (8.6)	120.90 d	115.17, 122.33
CO	–	168.06 s	

### Antibacterial and antifungal activity

Pseurotin A was assessed for antimicrobial activity by the agar diffusion method at a content of 50  $\mu\text{g disk}^{-1}$  towards the following microorganisms: Gram-positive bacteria: *Bacillus brevis* (local isolate), *Bacillus subtilis* (ATCC 6633) and *Micrococcus luteus* (ATCC 381); Gram-negative bacteria: *Enterobacter dissolvens* (LMG 2683); fungi: *Mucor miehei* (Cooney et Emerson Tü 284), *Paecilomyces variotti* (ETH 44646) and *Penicillium notatum* (Collection University of Kaiserslautern,

Germany). The activity of the extracts was estimated by the growth inhibition (in mm) as follows: < 8 mm, inactive; 8–12 mm, weak activity; 13–15 mm, moderate activity; > 15 mm, strong activity. As references, the commercial antifungal benlate [benomyl, methyl-1-(butylcarbamoyl)-2-benzimidazole carbamate, 50% active ingredient, 50  $\mu\text{g benomyl disk}^{-1}$ ] and chloramphenicol (30  $\mu\text{g disk}^{-1}$ ) were used.

The antifungal activity of pseurotin A against *Alternaria alternata* (Centro Micológico, Univer-

sidad de Rosario, Rosario, Argentina) and *Botrytis cinerea* (Instituto de Investigaciones Agropecuarias del Gobierno de Chile, INIA) was determined by the microdilution assay (Favre *et al.*, 2003) in 96-well microtiter plates. Stock solutions of pseurotin A in DMSO were diluted to give serial two-fold dilutions that were added to each medium resulting in concentrations ranging from 1.92 up to 250  $\mu\text{g ml}^{-1}$ . The final fraction of DMSO in the assay did not exceed 2%. The plates were incubated for 7 d at 25 °C. Minimal inhibitory concentration (MIC) was defined as the lowest extract concentration showing no visible fungal growth after the incubation time. As references, the common fungicides iprodione [3-(3,5-dichlorophenyl)-*N*-isopropyl-2,4-dioximidazolidine-1-carboxamide] (Systane) and myclobutanil [ $\alpha$ -butyl- $\alpha$ -(4-chlorophenyl)-1*H*-1,2,4-triazol-1-propanonitrile] (Rukon) were used.

The activity of pseurotin A against the phytopathogenic Gram-negative bacteria *Erwinia carotovora* (INIA) and *Pseudomonas syringae* (INIA) was assessed by the microdilution assay (Eloff, 1998) in 96-well microtiter plates. The bacterial inoculum was prepared to give approx.  $10^4$ – $10^5$  colony forming units (CFU) per well. Stock solutions of the compound in DMSO were diluted to give serial twofold dilutions that were added to each medium resulting in concentrations ranging from 1.92 up to 250  $\mu\text{g ml}^{-1}$ . The final fraction of DMSO in the assay did not exceed 2%. The plates were incubated overnight at 25 °C. To assess bacterial growth, the dye 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (MTT) was dissolved in water (0.5 mg  $\text{ml}^{-1}$ ), added to the microplate wells and incubated at 25 °C for 30 min. The absorbance was read in a universal microplate reader (Bio-Tek Instruments, INC, Winooski, Vermont, USA) at 515 nm. The results were transformed to percentage of controls and the  $\text{IC}_{50}$  values were graphically obtained from the dose-response curves. Penicillin G and streptomycin were used as standard antibacterials.

#### Cytotoxicity

The cytotoxic effect of the compounds, expressed as cell viability, was assessed on the permanent cell line derived from human lung fibroblasts (MRC-5) (ATCC Nr CCL-171). MRC-5 fibroblasts were grown as monolayers in minimum essential Eagle medium (MEM), with Earle's salts,

2 mM L-glutamine and 1.5 g  $\text{l}^{-1}$  sodium bicarbonate, supplemented with 10% heat-inactivated fetal bovine serum (FBS), 100 IU  $\text{ml}^{-1}$  penicillin and 100  $\mu\text{g ml}^{-1}$  streptomycin in a humidified incubator with 5%  $\text{CO}_2$  in air at 37 °C. Cells were seeded at a density of  $2.5 \cdot 10^3$  cells per well in 96-well plates. Confluent cultures were treated with medium containing the compounds at concentrations ranging from 3.8 up to 1000  $\mu\text{M}$ . The substance was firstly dissolved in DMSO and then in the medium supplemented with 2% FBS. The final content of DMSO in the test medium and controls was 1%. Cells were exposed for 24 h to the test medium with or without the compound (control). Each drug concentration was tested in quadruplicate, and repeated three times in separate experiments. At the end of the incubation, the neutral red uptake (NRU) assay was carried out as described by Rodríguez and Haun (1999). To calculate the  $\text{IC}_{50}$  values the results were transformed to percentage of controls, and the  $\text{IC}_{50}$  values were graphically obtained from the dose-response curves.

#### Results and Discussion

From the YMG broth culture of *Penicillium janczewskii* K. M. Zalesky, isolated from the phloem of the Chilean native gymnosperm *Prumnopitys andina*, two compounds were isolated and identified as pseurotin A (**1**) and cycloaspeptide A (**2**) by spectroscopic means (Fig. 1).

At a content of 50  $\mu\text{g disk}^{-1}$ , pseurotin A was inactive against *Enterobacter dissolvens*, *Micrococcus luteus* and *Penicillium notatum* but showed a weak effect on *Bacillus brevis*, *B. subtilis*, *Mucor miehei* and *Paecilomyces variotti* with growth inhibition of 8, 8, 7 and 9 mm, respectively. The antifungal effect of compound **1** against *Botrytis cinerea* and *Alternaria alternata* was low (Table III). In the antibacterial assay against the phytopathogenic bacteria *Erwinia carotovora* and *Pseudomonas syringae*, pseurotin A presented a moderate activity (Table III), with  $\text{IC}_{50}$  values of 220 and 112  $\mu\text{g ml}^{-1}$ , respectively. The effect of pseurotin A on *P. syringae* was close to the value observed for penicillin G, but much higher than that of streptomycin.

The cytotoxicity of pseurotin A (**1**) towards human lung fibroblasts was low, with an  $\text{IC}_{50}$  value of 1000  $\mu\text{M}$  while the  $\text{IC}_{50}$  value of cycloaspeptide A (**2**) was  $> 1000 \mu\text{M}$ .




Fig. 1. Structure of pseurotin A (1) and cycloaspeptide A (2) isolated from *Penicillium janczewskii*.

Pseurotin A has been previously reported from *Pseudeurotium ovalis* (Bloch and Tamm 1981; Breitenstein *et al.*, 1981) and *Aspergillus fumigatus* as well as from *Pochonia chlamydosporia* var. *catenulata* (Hellwig *et al.*, 2003). Cycloaspeptide A was first isolated from an *Aspergillus* species by Kobayashi *et al.* (1987) and was described from the psychrotolerant fungus *Penicillium ribeum* by Dalsgaard *et al.* (2004). The close related metabolites psychrophilin A and cycloaspeptide D were also reported from *Penicillium ribeum*. Cycloaspeptide A was also obtained from *P. algidum* cultures and exhibited moderate activity against *Plasmodium falciparum* with an  $IC_{50}$  value of  $3.5 \mu\text{g ml}^{-1}$  (Dalsgaard *et al.*, 2005).

Several natural products of different structure have been reported for *P. janczewskii*, including *cis*-fumagillin (Kwon *et al.*, 2000), quinolinone alkaloids (He *et al.*, 2005), peniprequinolone and gliovictin (Schmeda-Hirschmann *et al.*, 2005). A previous study on the secondary metabolites of endophytes obtained from *Prumnopitys andina*,

Table III. Antimicrobial activity ( $\mu\text{g ml}^{-1}$ ) of pseurotin A and cycloaspeptide A determined by the microdilution method.

Compound	<i>E. c.</i> $IC_{50}$	<i>P. s.</i> $IC_{50}$	<i>B. c.</i> MIC	<i>A. a.</i> MIC
Pseurotin A	220	112	>250	125
Cycloaspeptide A	–	>250	>250	–
Streptomycin	11.1	15.6	–	–
Penicillin G	15.6	122.7	–	–
Myclobutanil	–	–	31.3	3.9
Iprodione	–	–	15.6	31.3

*E. c.*, *Erwinia carotovora*; *P. s.*, *Pseudomonas syringae*; *B. c.*, *Botrytis cinerea*; *A. a.*, *Alternaria alternata*; –, not done.

including *P. janczewskii* grown on liquid potato-dextrose medium was reported (Schmeda-Hirschmann *et al.*, 2005). The chemical diversity of the compounds produced by the same species either isolated from different natural sources or grown in different media indicates the high versatility of the fungus in producing natural products and shows the potential of new research work on endophytic microorganisms when looking for bioactive products. The potential of this approach when looking for bioactive fungal metabolites has been outlined by Gunatilaka (2006) in a recent review. A related cyclic peptide was produced by a fungal strain obtained from the seed of *Avicennia marina*.

According to Madi and Katan (1998), when the filtrate of a liquid culture of *Penicillium janczewskii* was infiltrated into cotton and melon leaves, it induced systemic resistance protecting the lower stem portion of both plant species against the phytopathogenic fungus *Rhizoctonia solani*, reducing the incidence of damping-off up to 100%. The results suggested that the culture filtrate of *P. janczewskii* triggers the signal transduction cascade and activates defense genes. A similar effect can be expected in other plant species. Additional studies including plant pathogens of the endophytes host should be performed to disclose the relation of this fungus with the gymnosperm *Prumnopitys andina*.

#### Acknowledgements

E. H. thanks the Universidad de Talca for a Doctoral grant. We are grateful to the Centro de Investigación en Biotecnología Silvoagrícola de la Universidad de Talca and the Programa “Investigación y Desarrollo de Productos Bioactivos” for financial support.

- Arnold A.-E., Mejía L.-C., Kylo D., Rojas E.-I., Maynard Z., Robbins N., and Herre E.-A. (2003), Fungal endophytes limit pathogen damage in a tropical tree. *Proc. Natl. Acad. Sci. USA* **100**, 15649–15654.
- Bloch P. and Tamm C. (1981), Isolation and structure of pseurotin A, a microbial metabolite of *Pseudeurotium ovalis* Stolk with an unusual heterospirocyclic system. *Helv. Chim. Acta* **64**, 304–315.
- Breitenstein W., Chexal K.-K., Mohr P., and Tamm C. (1981), Pseurotin B, C, D and E. Further new metabolites of *Pseudeurotium ovalis* Stolk. *Helv. Chim. Acta* **64**, 379–388.
- Dalsgaard P.-W., Larsen T.-O., Frydenvang K., and Christophersen C. (2004), Psychrophilin A and cycloaspeptide D, novel cyclic peptides from the psychrotolerant fungus *Penicillium ribeum*. *J. Nat. Prod.* **67**, 878–881.
- Dalsgaard P.-W., Larsen T.-O., and Christophersen C. (2005), Bioactive cyclic peptides from the psychrotolerant fungus *Penicillium algidum*. *J. Antibiot.* **58**, 141–144.
- Eloff J. (1998), A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Med.* **64**, 711–713.
- Favre B., Hofbauer B., Hildering K.-S., and Ryder N. (2003), Comparison of *in vitro* activities of 17 antifungal drugs against a panel of 20 dermatophytes by using a microdilution assay. *J. Clin. Microbiol.* **41**, 4817–4819.
- Gosling P., Hodge A., Goodlass G., and Bending G.-D. (2006), Arbuscular mycorrhizal fungi and organic farming. *Agr. Ecosyst. Environ.* **113**, 17–35.
- Gunatilaka A. A. L. (2006), Natural products from plant-associated microorganisms: distribution, structural diversity, bioactivity, and implications of their occurrence. *J. Nat. Prod.* **69**, 509–526.
- He J., Lion U., Sattler I., Gollmick F.-A., Grabley S., Cai J., Meiners M., Schunke H., Schaumann K., Dechert U., and Krohn M. (2005), Diastereomeric quinolinone alkaloids from the marine-derived fungus *Penicillium janczewskii*. *J. Nat. Prod.* **68**, 1397–1399.
- Hellwig V., Mayer-Bartschmid A., Müller H., Greif G., Kleymann G., Zitzmann W., Tichy H., and Stadlert M. (2003), Pochonins A–F, new antiviral and antiparasitic resorcylic acid lactones from *Pochonia chlamydosporia* var. *catenulata*. *J. Nat. Prod.* **66**, 829–837.
- Hormazabal E., Schmeda-Hirschmann G., Astudillo L., Rodríguez J.-A., and Theoduloz C. (2005), Metabolites from *Microsphaeropsis olivacea* (Bonord.) Hohn, an endophytic fungus of *Pilgerodendron uviferum* (D. Don) Florin. *Z. Naturforsch.* **60c**, 11–21.
- Kobayashi R., Samejima Y., Nakajima S., Kawai K., and Udagawa S. (1987), Studies on fungal products XI. Isolation and structures of novel cyclic pentapeptides from *Aspergillus* sp. NE-45. *Chem. Pharm. Bull.* **35**, 1347–1352.
- Kogel K.-H., Franken P., and Hüchelhoven R. (2006), Endophyte or parasite – what decides? *Curr. Opin. Plant Biol.* **9**, 358–363.
- Kwon J.-Y., Jeong H.-W., Kim H.-K., Kang K.-H., Chang Y.-H., Bae K.-S., Choi J.-D., Lee U.-C., Son K.-H., and Kwon B.-M. (2000), *cis*-Fumagillin, a new methionine aminopeptidase (type 2) inhibitor produced by *Penicillium* sp. F2757. *J. Antibiot.* **53**, 799–806.
- Li J.-Y., Strobel G., Sidhu R., Hess W.-M., and Ford E.-J. (1996), Endophytic taxol-producing fungi from bald cypress, *Taxodium distichum*. *Microbiology* **142**, 2223–2226.
- Madi L. and Katan J. (1998), *Penicillium janczewskii* and its metabolites, applied to leaves, elicit systemic acquired resistance to stem rot caused by *Rhizoctonia solani*. *Physiol. Mol. Plant Pathol.* **53**, 163–175.
- Owen N. L. and Hundley N. (2004), Endophytes – the chemical synthesizers inside plants. *Sci. Prog.* **87**, 79–99.
- Puri S., Nazir A., Chawla R., Arora R., Riyaz-ul-Hasan S., Amna T., Ahmed B., Verma V., Singh S., Sagar R., Sharma A., Kumar R., Sharma R.-K., and Qazi G.-N. (2006), The endophytic fungus *Trametes hirsuta* as a novel alternative source of podophyllotoxin and related aryl tetralin lignans. *J. Biotechnol.* **122**, 494–510.
- Rodríguez J. and Haun M. (1999), Cytotoxicity of *trans*-dehydrocrotonin from *Croton cajucara* (Euphorbiaceae) on V79 cells and rat hepatocytes. *Planta Med.* **65**, 522–526.
- Schmeda-Hirschmann G., Hormazabal E., Astudillo L., Rodríguez J.-A., and Theoduloz C. (2005), Secondary metabolites from endophytic fungi isolated from the Chilean gymnosperm *Prumnopitys andina* (Lleuque). *World J. Microb. Biot.* **21**, 27–32.
- Seymour F.-A., Cresswell J.-E., Fisher P.-J., Lappin-Scott H.-M., Haag H., and Talbot N.-J. (2004), The influence of genotypic variation on metabolite diversity in populations of two endophytic fungal species. *Fungal Genet. Biol.* **41**, 721–734.
- Strobel G.-A. (2002), Rainforest endophytes and bioactive products. *Crit. Rev. Biotechnol.* **22**, 315–333.
- Strobel G. (2006), Harnessing endophytes for industrial microbiology. *Curr. Opin. Microbiol.* **9**, 240–244.