

Full Length Research Paper

Cytotoxicity and antibacterial studies of iridoids and phenolic compounds isolated from the latex of *Himatanthus sucuuba*

Jefferson Rocha A. Silva^{1*}, Claudia M. Rezende², Angelo C. Pinto² and Ana Claudia F. Amaral³

¹Departamento de Química, Universidade Federal do Amazonas, Av. Rodrigo Otávio, 3000 - Japiim - Manaus / AM, 69077-000, Brazil.

²Instituto de Química, UFRJ, Rio de Janeiro / RJ, 21945-970, R J, Brazil.

³Laboratório de Plantas Mediciniais e Derivados, Farmanguinhos/FIOCRUZ – R. Sizenando Nabuco, 100 - Manguinhos / RJ, 21041-250, RJ, Brazil.

Accepted 20 August, 2010

The latex of *Himatanthus sucuuba* (Spruce) Woodson, used popularly in the Amazon for the treatment of tumors, gastritis, inflammations and infections, was evaluated for cytotoxicity and antibacterial activities. The iridoid lactones, plumericin and isoplumericin were isolated from latex by bioassay fractionation and were found to be associated with DNA damage. Gallic acid exhibited the highest antimicrobial activity among the phenolic compounds isolated from the aqueous fraction. The compounds associated to cytotoxicity and antimicrobial activities could be responsible to the effects of this species used in traditional medicine.

Key words: *Himatanthus sucuuba*, iridoids, phenolics, cytotoxicity, antibacterial.

INTRODUCTION

In Brazil, there are about 90 genera and 850 species of the *Apocynaceae* family, divided into various formations (Souza and Lorenzi, 2005). Some species of this family such as *Himatanthus sucuuba* (Spruce) Woodson, are of therapeutic value and have a long history of use in folk medicine of Brazil. A recent review described the morphology, chemistry, pharmacology and ethnopharmacology of this species which is popularly known as *sucuuba*, *janaguba* and *janauba*, amongst other common

names (Amaral et al., 2007).

The latex and bark of this plant are mainly used in folk medicine for the treatment of ulcers, infections, inflammatory processes and tumors (Van Den Berg, 1982; Perdue and Blomster, 1978; Schultes, 1979; Bourdy et al., 2000; Villegas et al., 1997). Previous studies of *H. sucuuba* bark and latex have shown the antifungal and antiprotozoal activity of iridoid lactones (Silva et al., 1998; Castillo et al., 2007) and the anti-inflammatory activity of triterpenoids (Miranda et al., 2000). In this study, iridoids and phenolic compounds were isolated for the first time from the latex of this species. Bioassays-guided fractionations were used to identify the compounds responsible for the antitumor and antibacterial activities of the latex from this plant.

*Corresponding author. E mail: jrocha_01@yahoo.com.br or jrocha_01@ufam.edu.br. Tel: +55-092-36474031. Fax: +55-092-36474031.

Abbreviations: TLC, Thin layer chromatography; MIC, minimum inhibitory concentration; IC₁₂, concentration (µg/ml) is required to produce a zone of inhibition of 12 mm; MeOH, methanol; EtOAc, ethyl ethanoate; n-BuOH, n-butanol; GC/MS, gas chromatography coupled with mass spectrophotometer; NMR, nuclear magnetic resonance; MPLC, medium pressure liquid chromatography.

MATERIALS AND METHODS

Plant material

Collection

Plant samples were collected in Santarém city, Pará State and a

voucher specimen (number 5436) was deposited at the Herbarium of the Federal University of Amazonas, Manaus, AM, Brazil.

Obtaining Latex

The latex was removed from some cuts in the bark with the help of a damp sponge. The material was collected in a bottle and then stored at 4°C.

Extraction of latex and isolation of the compounds

Liquid-liquid extraction was performed on the latex of *H. sucuuba* as described previously (Barreto et al., 2007). The hexane fraction (12.0 g) was separated on a Sephadex LH-20 column, and eluted successively with hexane, dichloromethane (CH₂Cl₂) and methanol (MeOH). The hexane fraction was also subjected to preparative thin layer chromatography (TLC) (Hexane: EtOAc, 6:4) in order to isolate a mixture of plumericin and isoplumericin (8:2, 5.0 mg, 0.04% of the extract). Plumericin and isoplumericin were identified by spectroscopic methods (NMR and GC/MS) and by comparison with literature data (Trost et al., 1986; Abdel-Kader et al., 1997). The aqueous fraction was successively partitioned with CHCl₃ and after lyophilized. This powdered fraction (2.4 g) was submitted to reverse phase C-18 medium pressure liquid chromatography (MPLC) with water up to water/MeOH (1:1), flow 6 ml/min. 26 fractions of 40 ml were obtained and combined in two principal fractions: A (0.63 g) and B (1.8 g). The fraction A was chromatographed on Sephadex LH-20 column with water/MeOH (7:3) as eluent, for the isolation of catechol (39.5 mg). The fraction B was subjected to Sephadex LH-20 chromatography with MeOH as eluent and also on HPLC column (twice) RP-18 (Shimadzu, 45 x 250 mm, 10 µm particle size) and water/MeOH (6:4) as eluent, for the isolation of three compounds: gallic acid (16.6 mg), myricitrin (3.9 mg) and quercitrin (4.7 mg). The compounds were identified by spectroscopic methods and by comparison with data in the literature (Chanwitheesuk et al., 2007; Mabry et al., 1970). The flavonoids were also identified by UV spectroscopy using shift reagents (Mabry et al., 1970).

Antibacterial activity

Microorganisms

The microorganisms used in this study were *Staphylococcus aureus* (MRSA), *Staphylococcus epidermidis* (ATCC 12228), *Staphylococcus haemolyticus* (ATCC 2737), *Proteus mirabilis* (MRSA), *Shigella sonnei* (ATCC 25931), *Salmonella typhimurium* (ATCC 13311) and *Escherichia coli* (ATCC 25922). All MRSA bacteria were identified by traditional biochemical tests and according to National Committee for Clinical Laboratory Standards (NCCLS) (Machado et al., 2003).

Disc diffusion method

The antibacterial activity of the samples was determined using the disc diffusion method (Machado et al., 2003). Varying amounts of sample (25, 50, 100, 250, 500 and 1000 µg) were applied to sterile filter paper discs (6 mm in diameter). Antibacterial activity was determined by measuring the diameter (d) of the inhibition zone formed around the disk (considered 10 - 18 mm).

Minimum inhibitory concentration (MIC)

Culture conditions, media preparation and minimum inhibitory

concentration (MIC) assays were undertaken according to methodology adopted from National Council for Clinical Laboratory Standards (1993). Concentrations ranging from 15.5 to 500 µg/ml were used for each sample. The MIC is defined as the lowest concentration of the sample at which visible growth of the microorganism is completely inhibited.

Mechanism-based yeast bioassay for DNA damaging activity

The assay was evaluated using genetically engineered mutants of yeast of *Saccharomyces cerevisiae* as described previously (Gunatilaka et al., 1992). The IC₁₂ is defined as the concentration in µg/ml that is required to inhibit growth over a 12 mm diameter in a 100 µl well after 48 h incubation at 37°C. Camptothecin (RS188N and rad52) and streptonigrin (rad52.top1) were used as control drugs.


RESULTS AND DISCUSSION

Bioassay fractionation of the latex extract was used to identify substances that could be related to the ethnopharmacological use of *H. sucuuba*. The fractionation was performed as follows: *n*-BuOH was added to the crude latex of this species to promote polyisoprene precipitation. The organic (*n*-BuOH) and aqueous fractions were separated, evaporated and assayed against mutant strains of *S. cerevisiae*. This assay utilizes DNA repair or recombination deficient mutants of the yeast *S. cerevisiae* for the screening of compounds, which induce DNA damage. This mechanism-based yeast assay depends on the different responses of DNA repair-deficient and repair proficient yeast (*S. cerevisiae*) strains to the sample. The major DNA repair pathway is RAD52 pathway associated with the repair of double strand break and meiotic recombination (Gunatilaka et al., 1994). A mutant rad52 repair-deficient strain, rad52.top1, with the additional deletion of the DNA topoisomerase I gene is also available and can detect agents that produce DNA damage specifically by interacting with DNA topoisomerase (Gunatilaka et al., 1994). An extract is considered active if it shows selective activity against one or more repair-deficient yeasts and has an IC₁₂ less than 2000 µg/ml (Gunatilaka et al., 1994). The results are shown in Table 1. Only the *n*-BuOH fraction exhibited activity with an IC₁₂ = 1641 µg/ml (rad52.top1). The *n*-BuOH fraction was then partitioned with a solution of MeOH (80%) and hexane. Enhanced activity was exhibited by the hexane fraction, with an IC₁₂ = 542 µg/ml (rad52.top1). Column and preparative TLC were performed on the hexane fraction and the activity of isolated compounds was determined. The iridoid lactones (plumericin (1) and isoplumericin (2) showed relevant activity against rad52.top1 (IC₁₂ = 32.8 µg/ml). The activity of the iridoids against rad52 yeast strains was approximately four times lower (IC₁₂ = 112.3 µg/ml) and nine times less effective against repair-proficient strains (RS188N, IC₁₂ = 289.0 µg/ml) than against rad52.top1. These results indicated, for the first time, that this mixture

Table 1. Activity of the fractions and iridoids from *H. sucuuba* against mutant strains of *S. cerevisiae*.

Test samples	Zone of inhibition (mm)			IC ₁₂ (µg/ml)	
	RS188N	rad52	rad52.top1	rad52	rad52.top1
Aqueous	11.0	10.0	10.5	7160	4787
<i>n</i> -BuOH	11.0	12.5	13.0	1910	1641
MeOH (80%)	8.0	9.0	10.0	3104	2225
Hexane	10.0	12.0	15.5	1434	542
Plumericin and Isoplumericin	19.0	23.0	29.0	112.3	32.8

IC₁₂, the concentration (µg/ml) is required to produce a zone of inhibition of 12 mm.



1 Plumericin R = H; R¹ = CH₃

2 Isoplumericin R = CH₃; R¹ = H

3 Gallic acid

4 Myricetrin R₁=R₂=R₃=OH

5 Quercitrin R₁=R₂=OH, R₃=H

Figure 1. Chemical structures of phenolic and iridoids compounds isolated from *H. sucuuba*.

of iridoids could be associated with the DNA damaging activity of the latex extract and that it could be topoisomerase II inhibitor. The observed activity is important not only because it corroborates the ethnopharmacological use of the species, but also because it indicates a possible synergistic or additive action due to the two iridoids since plumericin alone results in less cytotoxicity activity (IC₁₂ = 70 µg/ml, rad52.top1) (Wood et al., 2001) than that observed in mixture (IC₁₂ = 32.8 µg/ml).

The antimicrobial activities of the hexane, chloroform and aqueous fractions of *H. sucuuba* latex were tested using the disc diffusion method with seven microorganisms and only the last fraction exhibited activity (d = 17 mm). The antimicrobial activity of the aqueous fraction was observed against five bacteria at a concentration of 500 µg/ml (*Proteus mirabilis* and *Escherichia coli*) and 350 µg/ml (*S. aureus*, *S. epidermidis* and *S. haemolyticus*). Among the phenolic compounds isolated from the aqueous fraction, gallic acid (3), a hydroxybenzoic acid, showed the most potent antimicrobial activity (MIC values: 31 µg/ml for *S. aureus* and *S. epidermidis*, 62 µg/ml for *P. mirabilis* and *S. haemolyticus* and 125 µg/ml for *E. coli*). Gallic acid has been previously reported to

possess anti-inflammatory, antifungal and antibacterial activities (Chanwitheesuk et al., 2007). The flavonoids, myricetrin (4) and quercitrin (5), were also isolated from the aqueous fraction, but they did not show antibacterial activity at any of the concentrations tested. The chemical structures of iridoids and phenolic compounds 1 - 5 are illustrated in Figure 1.

Conclusion

The results of this phytochemical work on the isolation and the identification of the active compounds of the latex of *H. sucuuba* provided scientific validation for the popular use of this species, and also confirmed a better activity of iridoids mixture on the DNA in a new plant material source, which is easier to obtain and less difficult to the plant than the bark removal.

ACKNOWLEDGEMENT

The authors thank professor Dr. Kátia R. N. -dos Santos for the use of bacteria to carry out this work.

REFERENCES

- Abdel-Kader MS, Wisse J, Evans R, Werff H, Kingston DGI (1997). Bioactive iridoids and a new lignan from *Allamanda cathartica* and *Himatanthus falax* from the Suriname rainforest. *J. Nat. Prod.* 60: 1294-1297.
- Amaral ACF, Ferreira JLP, Pinheiro MLB, Silva JRA (2007). Monograph of *Himatanthus sucuuba*, a plant of Amazonian folk medicine Phcog. *Rev.* 1: 77-85.
- Barreto AS, Silva JRA, Amaral ACF, Schripsema J, Rezende CM, Pinto AC (2007). Ácido 15-desmetilispolumierídeo, um novo iridóide isolado das cascas de *Plumeria rubra* e do látex de *Himatanthus sucuuba*. *Quim. Nova.* 30: 1133-1135.
- Bourdy G, DeWalt SJ, Michel LRC, Roca A, Deharo E, Munõz V, Balderrama L, Quenevo C, Gimenez A (2000). Medicinal plants uses of the Tacana, na Amazonian Bolivian ethnic group. *J. Ethnopharmacol.* 70: 87-109.
- Castillo D, Arevalo J, Herrera F, Ruiz C, Rojas R, Rengifo E, Vaisberg A, Lock O, Lemesre JL, Gornitzka H, Sauvain M (2007). Spirolactone iridoids might be responsible for the antileishmanial activity of a Peruvian traditional remedy made with *Himatanthus sucuuba* (*Apocynaceae*). *J. Ethnopharmacol.* 112: 410-414.
- Chanwitheesuk A, Teerawutgulrag A, Kilburn JD, Rakariyatham N (2007). Antimicrobial gallic acid from *Caesalpinia mimosoides* Lamk. *Food Chem.* 100: 1044-1048.
- Gunatilaka AAL, Kingston DGI, Johnson RK (1994). Mechanism-based isolation and structures of some anticancer active natural products. *Pure Appl. Chem.* 66: 2219-2222.
- Gunatilaka AAL, Samaranyake G, Kingston DGI, Hofmann G, Johnson RK. (1992). Bioactive ergost-5-ene-3 β , 7 α -diol derivatives from *Pseudobersama mossambicensis*. *J. Nat. Prod.* 55: 1648-1654.
- Mabry TJ, Markham KR, Thomas MB (1970). *The Systematic Identification of Flavonoids*. Springer Verlag, New York.
- Machado TB, Pinto AV, Pinto MCFR, Leal ICR, Silva MG, Amaral ACF, Kuster RM, Netto-dos Santos KR (2003). *In vitro* activity of Brazilian medicinal plants, naturally occurring naphthoquinones and their analogues, against methicillin-resistant *Staphylococcus aureus*. *Int. J. Antimicrob. Agents.* 21: 279-284.
- Miranda ALP, Silva JRA, Rezende CM, Pinto AC, Pinheiro MLB, Cordeiro MC, Tamborini E, Parrini JS (2000). Anti-inflammatory and analgesic activities of the latex containing triterpenes from *Himatanthus sucuuba*. *Planta Med.* 66: 284-286.
- National Council for Clinical Laboratory Standards (1993). *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: approved standard M7-A3*, third ed. Villanova, PA, USA: NCCLS.
- Perdue GP, Blomster RN (1978). South American Plants III: Isolation of fulvoplumierin from *Himatanthus sucuuba* (M. Arg.) Woodson (*Apocynaceae*). *J. Pharm. Sci.* 67: 1322-1323.
- Schultes RE (1979). De plantis toxicariie e mundo novo tropicale commentationes. XIX. Biodynamic apocynaceous plants of the northwest Amazon. *J. Ethnopharmacol.* 1: 165-192.
- Silva JRA, Rezende CM, Pinto AC, Pinheiro MLB, Cordeiro MC, Tamborini E, Young CM, Bolzani VS (1998). Ésteres triterpênicos de *Himatanthus sucuuba* (Spruce) Woodson. *Quim. Nova.* 21: 702-704.
- Souza VC, Lorenzi H. (2005). *Botânica Sistemática: Guia Ilustrado para a identificação das famílias de Angiospermas da Flora Brasileira*. Nova Odessa, Sp. Instituto Plantarum.
- Trost BM, Mao MKT, Balkovec JM, Buhlmayer P (1986). A total synthesis of plumericin, allamcin and allamandin. 1. Basic strategy. *J. Am. Chem. Soc.* 108: 4965-4973.
- Van Den Berg ME (1982). *Plantas medicinais na Amazônia: contribuição ao seu conhecimento sistemático*, CNPq/PRU/MPEG, Belém, Brasil.
- Villegas LF, Fernandez ID, Maldonado H, Torres R, Zavaleta A, Vaisberg AJ, Hammond GB (1997). Evaluation of the wound-healing activity of selected traditional medicinal plants from Peru. *J. Ethnopharmacol.* 55: 193-200.
- Wood CA, Lee K, Vaisberg AJ, Kingston DGI, Neto CC, Hammond GB (2001). A bioactive spirolactone iridoid and triterpenoids from *Himatanthus sucuuba*. *Chem. Pharm. Bull.* 49: 1477-1478.