

Published in final edited form as:

Parasitology. 2010 January ; 137(1): 159–171. doi:10.1017/S0031182009990321.

***daf-7-related* TGF- β homologues from Trichostrongyloid nematodes show contrasting life cycle expression patterns**

Henry J McSorley^{1,†}, John R Grainger¹, Yvonne Harcus¹, Janice Murray¹, Alasdair J Nisbet², David P Knox², and Rick M Maizels^{1,*}

¹Centre for Immunity, Infection and Evolution, and Institute for Immunology and Infection Research, Ashworth Laboratories, University of Edinburgh, West Mains Road, Edinburgh EH9 3JT

²Moredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik, Midlothian, EH26 0PZ

SUMMARY

OBJECTIVE—The transforming growth factor- β (TGF- β) gene family regulates critical processes in animal development, and plays a crucial role in regulating the mammalian immune response. We aimed to identify TGF- β homologues from two laboratory model nematodes (*Heligmosomoides polygyrus* and *Nippostrongylus brasiliensis*) and two major parasites of ruminant livestock (*Haemonchus contortus* and *Teladorsagia circumcincta*).

METHODS—Parasite cDNA was used as a template for gene-specific PCR and RACE.

RESULTS—Homologues of the TGH-2 subfamily were isolated, and found to differ in length (301, 152, 349 and 305 amino acids respectively), with variably truncated N-terminal pre-proteins. All contain conserved C-terminal active domains (>85% identical over 115 amino acids) containing 9 cysteine residues, as in *C. elegans* DAF-7, *Brugia malayi* TGH-2 and mammalian TGF- β . Surprisingly, only the *H. contortus* homologue retains a conventional signal sequence, absent from shorter proteins of other species. RT-PCR assays of transcription show that in *H. contortus* and *N. brasiliensis* expression is maximal in the infective larval stage, and very low in adult worms. In contrast, in *H. polygyrus* and *T. circumcincta*, *tgh-2* transcription is higher in adults than infective larvae.

CONCLUSION—The molecular evolution of this gene family in parasitic nematodes has diversified the pre-protein and life-cycle expression patterns of TGF- β homologues while conserving the structure of the active domain.

Keywords

development; immune modulator; stage-specific expression

*Corresponding author : Institute of Immunology and Infection Research, University of Edinburgh, Ashworth Laboratories, West Mains Road, Edinburgh EH9 3JT, r.maizels@ed.ac.uk, Fax +44 131 650 5450.

[†]Current address : Queensland Institute for Medical Research, Brisbane, Australia

INTRODUCTION

The TGF- β family is intimately involved in development and homeostasis of both vertebrate and invertebrate organisms (Newfeld *et al.*, 1999; Massagué *et al.*, 2000). In vertebrates, one branch of the family mediates immunological functions, in particular dampening potential immune pathology (Li *et al.*, 2006). In *Drosophila*, the TGF- β family member *decapentaplegic* (DPP) acts in morphogenesis (Affolter & Basler, 2007) while in *Caenorhabditis elegans*, *daf-7* regulates entry into the arrested Dauer larval stage (Ren *et al.*, 1996; Inoue & Thomas, 2000) and *dbl-1* controls body size (Suzuki *et al.*, 1999; Morita *et al.*, 2002). There is considerable interest in TGF- β homologues of parasitic helminths, and in *Schistosoma mansoni* one member of this family (SmInAct) is required for embryonic development (Freitas *et al.*, 2007). It has been suggested that other homologues may control parasite arrested development, in particular the hiatus of the infective third stage nematode larvae (L3) awaiting a mammalian host (Viney *et al.*, 2005). Furthermore, parasitic species may have even adapted TGF- β genes to interact with, and suppress, the host immune system (Maizels *et al.*, 2001).

To address these possibilities, nematode TGF- β homologues have been studied from *Brugia malayi* (Gomez-Escobar *et al.*, 1998; Gomez-Escobar *et al.*, 2000), *Ancylostoma caninum* (Brand *et al.*, 2005; Freitas & Arasu, 2005), two *Strongyloides* species and *Parastrongyloides trichosuri* (Crook *et al.*, 2005; Massey *et al.*, 2005). Only the *B. malayi* TGF- β homologue-2 (*Bm*-TGH-2) shows binding to mammalian TGF- β receptors (Gomez-Escobar *et al.*, 2000) with all other homologues appearing to fulfill a developmental function in the parasite, as is the case for *C. elegans daf-7* and *dbl-1*. However, the contrasting gene expression patterns suggests a reversal in developmental roles between *C. elegans* and parasitic species (Viney *et al.*, 2005). In the former, *daf-7* is expressed in early stages, but down-regulated when entering the arrested Dauer larval form (Ren *et al.*, 1996). In parasitic species expression is maximal in diapausal stages, including the arrested L3 larvae which may be developmentally analogous to Dauer (Hotez *et al.*, 1993). As larval arrest is constitutive in parasitic life cycles, but facultative in *C. elegans*, parasite TGF- β homologues may be required not to control entry into diapause, but rather to maintain arrest until the opportunity for infection arises (Viney *et al.*, 2005). This evolutionary switch is made more plausible by the recent discovery that the common molecular pathway controlling Dauer formation in *C. elegans* and parasite infective larval development is not *daf-7*-related, but signals through dafachronic acid ligation of the nuclear hormone receptor *daf-12* (Ogawa *et al.*, 2009).

The importance of TGF- β homologues in development and immune modulation may be dissected in model organisms such as the mouse intestinal helminth *Heligmosomoides polygyrus* and the rat parasite *Nippostrongylus brasiliensis*. Both are closely related to other Trichostrongyle nematodes of great economic significance as veterinary pathogens, including *Haemonchus contortus* and *Teladorsagia circumcincta*. These species offer fascinating and contrasting developmental and immunological features. *H. contortus*, *H. polygyrus* and *T. circumcincta* all share transmission via the faecal/oral route, with L3 larvae entering orally, developing to L4 larvae in the stomach of mice or the abomasum (4th stomach) of sheep, following which, they emerge into the lumen and deposit eggs which

egress in faeces. *H. contortus*, like many cattle nematodes (Armour & Duncan, 1987), may enter environmentally-induced arrest and *T. circumcincta* can undergo immune-induced developmental delay in the mammalian host. In *N. brasiliensis* infections, larvae penetrate the skin, and migrate through the lungs, trachea and oesophagus to reach the gastrointestinal tract. In the lumen, parasites develop to sexually-reproducing adults, depositing eggs which are voided in the faeces. *H. polygyrus* establishes chronic infections in mice, associated with immune suppression (Elliott *et al.*, 2004) and regulatory T cell activity (Wilson *et al.*, 2005; Finney *et al.*, 2007; Rausch *et al.*, 2008); however, *N. brasiliensis* is rapidly expelled from mice by a strong Th2-dependent immune response (Finkelman *et al.*, 2004; Anthony *et al.*, 2007). In both rodent and ovine hosts, a protective memory response can be generated following chemotherapy (Schallig, 2000; Finkelman *et al.*, 2004; Anthony *et al.*, 2006), which is also characterized by a dominant Th2 effector mechanisms. With the definition of a *daf-7* homologue from the dog hookworm *A. caninum* (Brand *et al.*, 2005; Freitas & Arasu, 2005) and with extensive genomic sequence data available for *H. contortus*, we initiated a project to identify new family members within this extremely important group of parasites, as we now report below.

MATERIALS AND METHODS

Parasites and Life Cycles

H. polygyrus bakeri was maintained in CBAx57BL/6 F1 mice by infection with 200 larvae through a gavage tube; adults were recovered at d14 post-infection. *N. brasiliensis* was maintained as described (Lawrence *et al.*, 1996), by infecting SD rats with 4000 L3 subcutaneously. L1, L2 and L3 of *H. polygyrus* and L3 of *N. brasiliensis* were collected by Baermannisation of faecal-charcoal cultures. *N. brasiliensis* L4 larvae were recovered from lungs 42 h post-infection, and adult parasites from the gut 6 days post-infection. Eggs were collected from adults cultured in vitro as described previously (Holland *et al.*, 2000). *H. contortus* and *T. circumcincta* lifecycles were maintained as described (Knox & Jones, 1990). L3 larvae from faecal cultures were exsheathed in 8% sodium hypochlorite solution and washed four times in saline at 38°C. Fourth-stage larvae and adult parasites were harvested 8 days and 21 days respectively after infecting worm-free sheep (7-12 months old) with 50,000 L3 of either *H. contortus* or *T. circumcincta*.

cDNA Preparations

H. polygyrus and *N. brasiliensis* harvested as above, were washed five times in PBS, and then stored in TRIzol reagent (Invitrogen) at -80°C, before use for RNA extraction following the manufacturer's methods. Briefly, around 100 µl packed volume of parasites (equivalent to around 50 adults or 1000 larvae) were first homogenised on ice, and after addition of 200 µl chloroform, centrifuged at 12,000 *g* for 15 min at 4°C. The aqueous layer was then isopropanol precipitated and centrifuged at 12,000 *g* for 10 min at 4°C. The RNA pellet was then washed in 500 µl 70 % ethanol, air-dried and resuspended in 20 µl DEPC water. The DNA-free kit (Ambion) was used to degrade DNA: 2 µl of DNase with 2.4 µl DNase buffer was added, and incubated at 37°C for, 30 min. 5 µl DNase inactivation reagent was then added, incubated at room temperature for 2 min, spun down at 12,000 *g* for

2 min, and supernatant removed. The concentration was determined by spectrophotometry at 260 nm.

For reverse transcription to cDNA, 500 ng of RNA was mixed with 2 µl 10× reverse transcriptase buffer, 2 µl 25 mM dNTP mix, 1 µl 50 U/µl MMLV reverse transcriptase (Stratagene), 0.5 µl 40 U/µl RNasin (Promega), 1 µl 0.4 µg/ml oligo dT primer (Promega) and DEPC-treated water to 20 µl. The reaction was incubated on a PCR block at 20°C for 10 min, 37°C for 60 min and 99°C for 5 min. cDNA from *H. contortus* and *T. circumcincta* was prepared as described previously (Redmond *et al.*, 2006).

PCR Amplifications and RACE

Polymerase Chain Reactions (PCRs) were run on an MJ Research DNA Engine with denaturing at 94°C for 1 min, annealing at 55-65°C for 1 min (depending on primers and product), and extended at 72°C for 0.5-2 min (depending on length of product). Reaction mixes comprised 0.2 µl 5 U/µl Taq (QIAGEN), 0.2 µl 25 mM dNTP mix, 2 µl 10 × PCR buffer (QIAGEN), 1 µl forward and reverse primers (10 µM each), 1 µl template (cDNA, colony pick or plasmid miniprep) and water to 20 µl. RACE-ready cDNA was made using the Invitrogen Generacer Core kit from adult parasites of all 4 species.

For 3' RACE reactions, reverse transcription was carried out with a modified oligo-dT primer containing a sequence insert for subsequent amplification (5'-GCTGTCAACGATACGCTACGTAACGGCATGACAG TG(T)₂₄-3'). The resulting cDNA was RNase-treated and then used in PCR with the gene-specific forward primers set out in Table 1. For 5' RACE, an oligonucleotide 5'-CGACUGGAGCACGAGGACACUGACAUGGACUGAAGGAGUAGAAA-3' was ligated to the 5' end of the RNA molecule prior to reverse transcription; 5 µg RNA was dephosphorylated with CIP (Calf Intestinal Phosphatase), phenol:chloroform precipitated, and the 5' cap structure then removed with TAP (Tobacco Acid Pyrophosphatase). Free phosphate groups created at the 5' end of mRNA were then ligated to the 5' RACE primer. The resulting cDNA was RNase-treated and used in PCR with the gene-specific reverse primers shown in Table 1.

Cloning

PCR products were excised under UV transillumination, purified using the QIAGEN gel extraction kit and ligated into pGEM-T (Promega) at a 3:1 molar ratio. Ligated plasmid preparations were then transformed into *E. coli* JM109 cells for overnight colony formation. Colonies positive by PCR screening were cultured overnight, and plasmid mini-prep DNA sequenced.

Real-time PCR

Real-time PCR was performed on 1 µl of a 1 in 5 dilution of cDNA, mixed with 5 µl Platinum SYBR green qPCR supermix (Invitrogen), 0.2 µl of forward and reverse primers (10 µM), and water to 10 µl. Triplicate reactions were set up, with standards of a mixture of all cDNA, with doubling dilutions from neat cDNA mix. The reactions were run on a Chromo4 real-time PCR machine (Alpha Innotech). Reactions were denatured for 20 sec at

94°C, annealed for 20 sec at 65°C and extended for 20 sec at 72°C (for SYBR green fluorescence) for 50 cycles. Primers used for the actin, tubulin and *tgh-2* genes of all 4 parasite species are given in Table 1. In all reactions a melting curve analysis was also carried out, in which the fluorescence profile was measured at 0.5°C increments from the annealing temperature until 94°C, to ensure a single peak, indicative of a single amplified product, was detected. Reaction products were also run on agarose gels to confirm that a single product had been amplified.

Bioinformatics

Homology searches of the partially sequenced *H. contortus* genome were performed using the Wellcome Trust Sanger Institute Blast server (www.sanger.ac.uk/Projects/H_contortus), using the datasets published on 27/01/2006, and more recently on 12/11/2007. Searches of nematode ESTs (including *T. circumcincta*) were carried out at <http://www.ebi.ac.uk/Tools/blast2/nucleotide.html>, selecting the “nucleic acid” and “EMBL EST Invertebrate” options. Signal peptide predictions were performed using the SignalP 3.0 server (<http://www.cbs.dtu.dk/services/SignalP/>). Alignments, translational predictions and sequence analysis were carried out using MacVector 9.5.2 (Symantec). The phylogenetic tree shown was constructed by best tree analysis, using a Poisson distribution.

Nomenclature

Naming of genes and proteins followed standard nomenclature for nematode species (Bird & Riddle, 1994), with a two letter species code followed by a three-letter gene name and a numeral if required. Gene names for DNA and mRNA are referred to in lower case italics, while proteins and amino acid sequences are Roman capitals. Names and numerals are conserved between orthologues from different species wherever possible. Hence, the orthologues of *B. malayi transforming growth factor homologue-2* (*Bm-tgh-2*) are each given the gene name *tgh-2*. In TGF- β superfamily members, proteolytic cleavage at an internal tetrabasic site is required to produce the receptor-binding C-terminal domain, variously termed mature, ligand or active domain; we use the latter term to denote this domain.

RESULTS

Identification of TGH-2 in *Haemonchus contortus*

Nematode homologues of TGF- β identified in recent years include *Ac-tgh-2* (also designated *Ac-daf-7*) from the dog hookworm *A. caninum* (Brand *et al.*, 2005; Freitas & Arasu, 2005). Because *Ancylostoma* is within the same taxonomic suborder as *H. contortus* (Strongyloida), the *Ac-TGH-2* protein sequence was selected to search for TGF- β homologues in *H. contortus*. A closely related sequence was found on a contig in the Wellcome Trust Sanger Institute dataset (http://www.sanger.ac.uk/Projects/H_contortus), covering 3 exons in the active domain, as shown in Fig. 1A. No ESTs from *H. contortus* (Parkinson *et al.*, 2004), *N. brasiliensis* (Harcus *et al.*, 2004), or *T. circumcincta* (Nisbet *et al.*, 2008) were found with similarity to either *Ac-tgh-2* or the *H. contortus* contig. When, as described below, 5' and 3' RACE were used to extend the cDNA sequence to the N- and C-termini of the protein, the additional sequences matched two additional contigs containing

five 5' and one 3' exons. The organisation of *Hc-tgh-2* into 10 exons closely resembles that of *Ac-tgh-2*, with all but the first exon being of identical length. There is considerable variation, however, in the intronic tracts, as far as genomic information permits comparison (Freitas & Arasu, 2005).

Isolation of TGH-2 homologues from other species

To isolate cDNAs coding for TGH-2 homologues from other gastrointestinal nematode parasites, an alignment of known human and nematode TGF- β amino acid sequences was made, including the newly-defined *H. contortus* homologue (Fig. 1B). Degenerate primers were designed for regions which were best conserved and offered the lowest degeneracy. Where sequences diverged between homologues, the *H. contortus*/*A. caninum* TGH-2 consensus was taken. These primer sequences are shown in Fig. 1C.

Combinations of these degenerate primers were used in PCRs with *H. polygyrus*, *N. brasiliensis* and *T. circumcineta* adult parasite cDNA, and products (Fig. 1D) excised from gels and sequenced. From both *H. polygyrus* and *N. brasiliensis*, bands of ~250 bp were amplified, which on sequencing showed similarity to *Ac-tgh-2*. Using a different forward primer, a 200-bp product was amplified from *T. circumcineta* cDNA, with homologous sequence. Non-degenerate primers (Table 1) were then designed to these newly-sequenced regions of each homologue, and were used in 5' and 3' Rapid Amplification of cDNA Ends (RACE) to identify flanking coding sequences.

Full sequences of TGF- β homologue genes

RACE was used to isolate the 5' and 3' ends of each gene, in separate reactions with products as shown in Fig. 2. The *H. contortus* and *T. circumcineta* 5' and 3' RACE products were amplified by single PCR reactions, while the *H. polygyrus* and *N. brasiliensis* 5' and 3' RACE products required amplification by nested PCR, using a second set of primers. Products were sequenced, and sequences of the newly-assigned *Hc*-TGH-2, *Hp*-TGH-2, *Nb*-TGH-2 and *Tc*-TGH-2 aligned with other TGF- β family members at the amino acid level (Fig. 3). The four trichostrongyloid parasite proteins were all identical at 100/117 positions (85.4%) of the C-terminal domain following , but only 50 amino acids (42.7%) were identical to *Bm*-TGH-2 over the same tract.

Particular scrutiny was given to the 5' termini of these genes, as in three cases they were truncated at the 5' end compared to canonical TGF- β s (Fig. 3). To verify the full-length sequence of *Hp-tgh-2* and *Nb-tgh-2*, 5' RACE was repeated several times and the 5' ends of each gene were also amplified from cDNA libraries prepared as described elsewhere (Harcus *et al.*, 2004); in no case did we detect a longer transcript. Furthermore, we were able to amplify *Hp-tgh-2* with an internal gene-specific reverse primer, and a forward primer for the nematode 22-nt spliced leader sequence, and this product contained the same start codon as present in all other PCR products (data not shown). Significantly, the *SmInAct* gene is also truncated at the 5' end (Freitas *et al.*, 2007), so this is not a feature confined to trichostrongyloid TGF- β homologues.

The truncated N-termini of *Hp*-TGH-2, *Nb*-TGH-2 and *Tc*-TGH-2 also result in the absence of a conventional signal peptide; interestingly, the N-terminus of *Nb*-TGH-2 fulfils the

criteria for a predicted signal peptide, although this sequence corresponds to an internal hydrophobic region in the longer homologues (Fig. 3). Notably, a potential N-glycosylation site is present in the active domain of the trichostrongyloid proteins, which is not observed in other family members (Fig. 3). N-glycosylation is known to be important for secretion of mammalian TGF- β (Brunner *et al.*, 1992) and it would be interesting to establish if this also pertains to the parasite products.

Phylogenetic analysis

Within the TGF- β superfamily, each of the homologues described here were found by phylogenetic analysis of the conserved C-terminus to group as a closely-spaced set on the same branch as *Bm-tgh-2* and *C. elegans daf-7* (Fig. 4). Morphological characteristics group *H. contortus* and *T. circumcincta* together in the Family Haemonchidae, while *H. polygyrus* and *N. brasiliensis* are placed in separate Families, Heligmonellidae and Heligmosomatidae respectively ((Hoberg & Lichtenfels, 1994) and <http://beta.uniprot.org/taxonomy/>). Our analysis lends support to this classification, but not at the highest level of confidence because of the limited number of variant positions available for comparison.

Real-time PCR analysis

To assess the expression profile of the novel TGF- β homologues in parasites at different developmental points, cDNA was prepared from eggs, larvae and adult worms of each organism. Larval stage availability varied between species, but it was possible to analyze L1, L2 and L3 larvae of *H. polygyrus*, L3 and lung L4 larvae of *N. brasiliensis*, L3 and exsheathed L3 (xL3) and abomasal L4 of *H. contortus*, and xL3 and abomasal L4 of *T. circumcincta*. Using real-time PCR, *tgh-2* gene transcription was quantified, and normalised by comparison to parasite β -actin gene expression. In *H. polygyrus*, transcription of *Hp-tgh-2* is lowest in the early larval stages, and increases through later stages to reach highest levels in the adults and eggs (Fig. 5 A). In contrast, *N. brasiliensis tgh-2* expression was maximal in L3 larvae and eggs, with lower expression in L4 larvae and minimal levels in adults (Fig. 5 B). RT-PCR analysis of cDNA from *H. contortus* and *T. circumcincta* also revealed a different developmental pattern of gene expression, the former species showing a larval maximum (Fig. 5 C), as has also been shown for *S. ratti* (Crook *et al.*, 2005), while in *T. circumcincta* adult stages contained high levels of *tgh-2* transcript (Fig. 5 D).

Discussion

Over recent years, it has become established that members of the TGF- β gene superfamily are required in helminth organisms to direct a suite of developmental processes, most clearly observed in the free-living nematode *C. elegans* (Patterson & Padgett, 2000). Strikingly, the complex signalling pathway, with two receptor subunits and intracellular kinases, is conserved from mammals to nematodes (Estevez *et al.*, 1993; Gomez-Escobar *et al.*, 1997; Newfeld *et al.*, 1999), trematodes (Beall & Pearce, 2001; Osman *et al.*, 2006; Freitas *et al.*, 2007) and cestodes (Zavala-Gongora *et al.*, 2006), indicating the potential for molecular cross-talk between evolutionarily distant organisms (Luckhart *et al.*, 2003). An important, but unproven, hypothesis is that parasites may have adapted genes first involved in endogenous body organisation to interact with the immune system of their host (Gomez-

Escobar *et al.*, 2000). TGF- β proteins may exemplify this process, if it is the case that helminth homologues can ligate host TGF- β receptors and thereby diminish immune responsiveness.

A number of TGF- β homologues have been characterised from parasitic nematodes, including the human filarial nematode *B. malayi* (Gomez-Escobar *et al.*, 1998; Gomez-Escobar *et al.*, 2000), *S. stercoralis* and *S. ratti*, and *P. trichosuri* (Crook *et al.*, 2005; Massey *et al.*, 2005), as well as the dog hookworm *A. caninum* (Brand *et al.*, 2005; Freitas & Arasu, 2005). Some important groups of parasites, however, have not been examined in this regard. For example, *H. polygyrus* and *N. brasiliensis* are species widely used for experimental infections with rodents and known to drive strong regulatory T cell (Wilson *et al.*, 2005; Finney *et al.*, 2007; Rausch *et al.*, 2008) and Th2 differentiation (Lawrence *et al.*, 1996; Voehringer *et al.*, 2004) respectively in laboratory mice. It is possible that parasite-derived TGF- β family members play a role in this very pronounced skewing of immune responsiveness. Two related nematodes, found worldwide in sheep and other ruminants and responsible for very considerable economic damage, are *H. contortus* and *T. circumcincta*. Identification of TGF- β homologues in these species could pave the way for new strategies to control infections which are proving recalcitrant to drug clearance (Wolstenholme *et al.*, 2004). These four species are closely related taxonomically within the Trichostrongyloidea superfamily of nematodes, and are each found in specialised habitats within the host gastrointestinal tract.

Our identification of new homologues was made possible, firstly, by the release of substantial genomic sequence from *H. contortus*, and secondly by the availability of sequences from closely related species, most particularly *A. caninum* (Brand *et al.*, 2005; Freitas & Arasu, 2005). While the four new homologues were found to be highly similar to each other in the conserved active domain, it was surprising to find that three of the four contained truncated pro-protein N-terminal domains, without conventional signal sequences. Although in *N. brasiliensis* it is possible that the truncated product employs an alternative hydrophobic tract as a signal sequence (see Fig. 3), no alternative is apparent for either *H. polygyrus* or *T. circumcincta*. However, it has recently been reported that the *S. mansoni* homologue SmInAct, which is a functional TGF- β like protein, is also truncated and lacks a conventional signal sequence (Freitas *et al.*, 2007). In the context of signal peptide-independent secretion, it should be noted that in a recent proteomic study of *B. malayi* secreted proteins, some 33% lacked signal peptides, including one of the most abundant secreted molecules (Hewitson *et al.*, 2008). Studies are now under way to ascertain if the trichostrongyloid parasites secrete TGH-2 proteins in a signal peptide-independent manner.

A notable level of diversity between the trichostrongyloid species was seen in the patterns of *tgh-2* expression throughout their developmental cycles. Real-time PCR showed that in *N. brasiliensis* and *H. contortus* expression is highest in the arrested L3 larvae, which is similar to the profile reported for *A. caninum*, *P. trichosuri*, *S. ratti* and *S. stercoralis* (Brand *et al.*, 2005; Crook *et al.*, 2005; Freitas & Arasu, 2005; Massey *et al.*, 2005). In contrast, the TGF- β homologue *daf-7* is down-regulated in *C. elegans* dauer larvae which have entered developmental arrest (Ren *et al.*, 1996). Thus, in parasitic species in which larval arrest is constitutive, TGF- β homologues may regulate later developmental steps which are

exquisitely dependent on the environmental or immunological cues accompanying infection (Freitas & Arasu, 2005; Viney *et al.*, 2005). Interestingly, among these cues may be host TGF- β (Arasu, 2001). In distinction to *N. brasiliensis* and *H. contortus*, two other species display continued *tgh-2* expression beyond the infective larval stage. Adult worms of *H. polygyrus* and *T. circumcincta* show high levels of gene transcription, as do the eggs. Because *H. contortus* eggs do not express *tgh-2*, it is difficult to attribute its function in either adult worms or eggs to any fundamental, conserved developmental pathway.

An alternative scenario is that these gene products exert an immunological effect in the host. Thus, expression by infective larvae allows them to release high levels of TGF- β homologue on infection of the host, thereby impairing the host immune response from the very outset. Where, as with *H. polygyrus*, the expression increases through the duration of mammalian infection, this could explain the dominance of regulatory T cell activity in mice harbouring adult worms of this parasite (Wilson *et al.*, 2005; Finney *et al.*, 2007; Rausch *et al.*, 2008). It is interesting to speculate whether the failure of *N. brasiliensis* adult worms to establish beyond 6-8 days in the murine gut is related to the absence of TGF- β homologue expression by this stage of the parasite.

These alternative hypotheses now require experimental investigation. While conventional genetic knockout techniques cannot be used with helminth parasites, RNAi offers a promising route to study individual gene function, and most recently knock-down of the *S. mansoni* TGF- β family member SmInAct confirmed its role in parasite development (Freitas *et al.*, 2007). Although progress with RNAi on trichostrongyloid nematodes has been limited (Hussein *et al.*, 2002; Geldhof *et al.*, 2006; Zawadzki *et al.*, 2006; Lendner *et al.*, 2008), in some instances this technique has been successful (Kotze & Bagnall, 2006) indicating that it may be possible to knock down TGF- β expression in a species such as *H. contortus*. Further, heterologous transfection of helminth genes into other organisms such as *Leishmania* has permitted the immunological function of individual genes to be elucidated (Gomez-Escobar *et al.*, 2005; Maizels *et al.*, 2008). We are pursuing these and other means of establishing the biological role of TGF- β family members in these important helminth parasite species.

ACKNOWLEDGEMENTS

HMcS thanks the Medical Research Council for Studentship support; JRG thanks the Wellcome Trust for a PhD studentship; YH, JM and RMM thank the Wellcome Trust for Programme Grant support. DPK and AJN gratefully acknowledge funding from the Scottish Government Rural and Environment Research and Analysis Directorate (RERAD).

REFERENCES

- Affolter M, Basler K. The Decapentaplegic morphogen gradient: from pattern formation to growth regulation. *Nature Reviews Genetics*. 2007; 8:663–674.
- Anthony RM, Urban JF Jr, Alem F, Hamed HA, Roza CT, Boucher JL, Van Rooijen N, Gause WC. Memory T_H2 cells induce alternatively activated macrophages to mediate protection against nematode parasites. *Nature Medicine*. 2006; 12:955–960.
- Anthony RM, Rutitzky LI, Urban JF Jr, Stadecker MJ, Gause WC. Protective immune mechanisms in helminth infection. *Nature Reviews Immunology*. 2007; 7:975–987.
- Arasu P. In vitro reactivation of *Ancylostoma caninum* tissue-arrested third-stage larvae by transforming growth factor-beta. *Journal of Parasitology*. 2001; 87:733–738. [PubMed: 11534634]

- Armour J, Duncan M. Arrested larval development in cattle nematodes. *Parasitol Today*. 1987; 3:171–176. [PubMed: 15462948]
- Beall MJ, Pearce EJ. Transforming growth factor-b and insulin-like signaling pathways in parasitic helminths. *International Journal of Parasitology*. 2001; 32:399–404. [PubMed: 11849636]
- Bird DM, Riddle DL. A genetic nomenclature for parasitic nematodes. *Journal of Nematology*. 1994; 26:138–143. [PubMed: 19279876]
- Brand AM, Varghese G, Majewski W, Hawdon JM. Identification of a DAF-7 ortholog from the hookworm *Ancylostoma caninum*. *International Journal for Parasitology*. 2005; 35:1489–1498. [PubMed: 16135366]
- Brunner AM, Lioubin MN, Marquardt H, Malacko AR, Wang WC, Shapiro RA, Neubauer M, Cook J, Madisen L, Purchio AF. Site-directed mutagenesis of glycosylation sites in the transforming growth factor-beta 1 (TGF beta 1) and TGF beta 2 (414) precursors and of cysteine residues within mature TGF beta 1: effects on secretion and bioactivity. *Molecular Endocrinology*. 1992; 6:1691–1700. [PubMed: 1448117]
- Crook M, Thompson FJ, Grant WN, Viney ME. *daf-7* and the development of *Strongyloides ratti* and *Parastongyloides trichosuri*. *Molecular and Biochemical Parasitology*. 2005; 139:213–223. [PubMed: 15664656]
- Elliott DE, Setiawan T, Metwali A, Blum A, Urban JF Jr, Weinstock JV. *Heligmosomoides polygyrus* inhibits established colitis in IL-10-deficient mice. *European Journal of Immunology*. 2004; 34:2690–2698. [PubMed: 15368285]
- Estevez M, Attisano L, Wrana JL, Albert PS, Massagué J, Riddle DL. The *daf-4* gene encodes a bone morphogenetic protein receptor controlling *C.elegans* dauer larva development. *Nature*. 1993; 365:644–649. [PubMed: 8413626]
- Finkelman FD, Shea-Donohue T, Morris SC, Gildea L, Strait R, Madden KB, Schopf L, Urban JF Jr. Interleukin-4- and interleukin-13-mediated host protection against intestinal nematode parasites. *Immunological Reviews*. 2004; 201:139–155. [PubMed: 15361238]
- Finney CA, Taylor MD, Wilson MS, Maizels RM. Expansion and activation of CD4⁺CD25⁺ regulatory T cells in *Heligmosomoides polygyrus* infection. *European Journal of Immunology*. 2007; 37:1874–1886. [PubMed: 17563918]
- Freitas T, Jung E, Pearce EJ. TGF-b signaling controls embryo development in the parasitic flatworm *Schistosoma mansoni*. *PLOS Pathogens*. 2007; 3:e52. [PubMed: 17411340]
- Freitas TC, Arasu P. Cloning and characterisation of genes encoding two transforming growth factor-b-like ligands from the hookworm, *Ancylostoma caninum*. *International Journal for Parasitology*. 2005; 35:1477–1487. [PubMed: 16140304]
- Geldhof P, Murray L, Couthier A, Gilleard JS, McLauchlan G, Knox DP, Britton C. Testing the efficacy of RNA interference in *Haemonchus contortus*. *International Journal for Parasitology*. 2006; 36:801–810. [PubMed: 16469321]
- Gomez-Escobar N, van den Biggelaar A, Maizels RM. A member of the TGF- β receptor gene family in the parasitic nematode *Brugia*. *Gene*. 1997; 199:101–109. [PubMed: 9358045]
- Gomez-Escobar N, Lewis E, Maizels RM. A novel member of the transforming growth factor- β (TGF- β) superfamily from the filarial nematodes *Brugia malayi* and *B.pahangi*. *Experimental Parasitology*. 1998; 88:200–209. [PubMed: 9562423]
- Gomez-Escobar N, Gregory WF, Maizels RM. Identification of *Bm-tgh-2*, a filarial nematode homolog of *C.elegans daf-7* and human TGF- β , expressed in microfilarial and adult stages of *Brugia malayi*. *Infection and Immunity*. 2000; 68:6402–6410. [PubMed: 11035752]
- Gomez-Escobar N, Bennett C, Prieto-Lafuente L, Aebischer T, Blackburn CC, Maizels RM. Heterologous expression of the filarial nematode *alt* gene products reveals their potential to inhibit immune function *BMC Biology*. 2005; 3:1. [PubMed: 15638945]
- Harcus YM, Parkinson J, Fernández C, Daub J, Selkirk ME, Blaxter ML, Maizels RM. Signal sequence analysis of expressed sequence tags from the nematode *Nippostrongylus brasiliensis* and the evolution of secreted proteins in parasites. *Genome Biology*. 2004; 5:R39. [PubMed: 15186490]
- Hewitson JP, Harcus YM, Curwen RS, Dowle AA, Atmadja AK, Ashton PD, Wilson RA, Maizels RM. The secretome of the filarial parasite, *Brugia malayi*: proteomic profile of adult excretory-

- secretory products. *Molecular and Biochemical Parasitology*. 2008; 160:8–21. [PubMed: 18439691]
- Hoberg EP, Lichtenfels JR. Phylogenetic systematic analysis of the Trichostrongylidae (Nematoda), with an initial assessment of coevolution and biogeography. *Journal of Parasitology*. 1994; 80:976–996. [PubMed: 7799171]
- Holland MJ, Harcus YM, Riches PL, Maizels RM. Proteins secreted by the parasitic nematode *Nippostrongylus brasiliensis* act as adjuvants for Th2 responses. *European Journal of Immunology*. 2000; 30:1977–1987. [PubMed: 10940887]
- Hotez P, Hawdon J, Schad GA. Hookworm larval infectivity, arrest and amphiparatensis: the *Caenorhabditis elegans* Daf-c paradigm. *Parasitology Today*. 1993; 9:23–26. [PubMed: 15463660]
- Hussein AS, Kichenin K, Selkirk ME. Suppression of secreted acetylcholinesterase expression in *Nippostrongylus brasiliensis* by RNA interference. *Molecular and Biochemical Parasitology*. 2002; 122:91–94. [PubMed: 12076773]
- Inoue T, Thomas JH. Targets of TGF- β signaling in *Caenorhabditis elegans* dauer formation. *Developmental Biology*. 2000; 217:192–204. [PubMed: 10625546]
- Knox DP, Jones DG. Studies on the presence and release of proteolytic enzymes (proteinases) in gastro-intestinal nematodes of ruminants. *International Journal of Parasitology*. 1990; 20:243–249. [PubMed: 2185170]
- Kotze AC, Bagnall NH. RNA interference in *Haemonchus contortus*: suppression of beta-tubulin gene expression in L3, L4 and adult worms in vitro. *Molecular and Biochemical Parasitology*. 2006; 145:101–110. [PubMed: 16253354]
- Lawrence RA, Gray C, Osborne J, Maizels RM. *Nippostrongylus brasiliensis*: Cytokine responses and nematode expulsion in normal and IL4-deficient mice. *Experimental Parasitology*. 1996; 84:65–73. [PubMed: 8888733]
- Lendner M, Doligalska M, Lucius R, Hartmann S. Attempts to establish RNA interference in the parasitic nematode *Heligmosomoides polygyrus*. *Molecular and Biochemical Parasitology*. 2008; 161:21–31. [PubMed: 18606194]
- Li MO, Wan YY, Sanjabi S, Robertson AK, Flavell RA. Transforming growth factor-beta regulation of immune responses. *Annual Review of Immunology*. 2006; 24:99–146.
- Luckhart S, Crampton AL, Zamora R, Lieber MJ, Dos Santos PC, Peterson TM, Emmith N, Lim J, Wink DA, Vodovotz Y. Mammalian transforming growth factor b1 activated after ingestion by *Anopheles stephensi* modulates mosquito immunity. *Infection and Immunity*. 2003; 71:3000–3009. [PubMed: 12761076]
- Maizels RM, Gomez-Escobar N, Gregory WF, Murray J, Zang X. Immune evasion genes from filarial nematodes. *International Journal of Parasitology*. 2001; 31:889–898. [PubMed: 11406138]
- Maizels RM, Gomez-Escobar N, Prieto-Lafuente L, Murray J, Aebischer T. Expression of helminth genes in *Leishmania*: an experimental transfection system to test immunological function. *Parasite Immunology*. 2008; 30:195–201. [PubMed: 18324922]
- Massagué J, Blain SW, Lo RS. TGFbeta signaling in growth control, cancer, and heritable disorders. *Cell*. 2000; 103:295–309. [PubMed: 11057902]
- Massey HC, Castelletto ML, Bhopale VM, Schad GA, Lok JB. Sst-tgh-1 from *Strongyloides stercoralis* encodes a proposed ortholog of *daf-7* in *Caenorhabditis elegans*. *Molecular and Biochemical Parasitology*. 2005; 142:116–120. [PubMed: 15907565]
- Morita K, Flemming AJ, Sugihara Y, Mochii M, Suzuki Y, Yoshida S, Wood WB, Kohara Y, Leroi AM, Ueno N. A *Caenorhabditis elegans* TGF-beta, DBL-1, controls the expression of LON-1, a PR-related protein, that regulates polyploidization and body length. *EMBO Journal*. 2002; 21:1063–1073. [PubMed: 11867534]
- Newfeld SJ, Wisotzkey RG, Kumar S. Molecular evolution of a developmental pathway: phylogenetic analyses of transforming growth factor-beta family ligands, receptors and Smad signal transducers. *Genetics*. 1999; 152:783–795. [PubMed: 10353918]
- Nisbet AJ, Redmond DL, Matthews JB, Watkins C, Yaga R, Jones JT, Nath M, Knox DP. Stage-specific gene expression in *Teladorsagia circumcincta* (Nematoda: Strongylida) infective larvae and early parasitic stages. *International Journal for Parasitology*. 2008; 38:829–838. [PubMed: 18062971]

- Ogawa A, Streit A, Antebi A, Sommer RJ. A conserved endocrine mechanism controls the formation of dauer and infective larvae in nematodes. *Current Biology*. 2009; 19:67–71. [PubMed: 19110431]
- Osman A, Niles EG, Verjovski-Almeida S, LoVerde PT. *Schistosoma mansoni* TGF- β receptor II: role in host ligand-induced regulation of a schistosome target gene. *PLOS Pathogens*. 2006; 2:e54. [PubMed: 16789838]
- Parkinson J, Mitreva M, Whitton C, Thomson M, Daub J, Martin J, Schmid R, Hall N, Barrell B, Waterston RH, McCarter JP, Blaxter ML. A transcriptomic analysis of the phylum Nematoda. *Nature Genetics*. 2004; 36:1259–1267. [PubMed: 15543149]
- Patterson GI, Padgett RW. TGF β -related pathways. Roles in *Caenorhabditis elegans* development. *Trends in Genetics*. 2000; 16:27–33. [PubMed: 10637628]
- Rausch S, Huehn J, Kirchhoff D, Rzepecka J, Schnoeller C, Pillai S, Loddenkemper C, Scheffold A, Hamann A, Lucius R, Hartmann S. Functional analysis of effector and regulatory T cells in a parasitic nematode infection. *Infection and Immunity*. 2008; 76:1908–1919. [PubMed: 18316386]
- Redmond DL, Smith SK, Halliday A, Smith WD, Jackson F, Knox DP, Matthews JB. An immunogenic cathepsin F secreted by the parasitic stages of *Teladorsagia circumcincta*. *International Journal for Parasitology*. 2006; 36:277–286. [PubMed: 16387304]
- Ren P, Lim C-S, Johnsen RJ, Albert PS, Pilgrim D, Riddle DL. Control of *C. elegans* larval development by neuronal expression of a TGF- β homolog. *Science*. 1996; 274:1389–1391. [PubMed: 8910282]
- Schallig HD. Immunological responses of sheep to *Haemonchus contortus*. *Parasitology*. 2000; 120(Suppl):S63–72. [PubMed: 10874710]
- Suzuki Y, Yandell MD, Roy PJ, Krishna S, Savage-Dunn C, Ross RM, Padgett RW, Wood WB. A BMP homolog acts as a dose-dependent regulator of body size and male tail patterning in *Caenorhabditis elegans*. *Development*. 1999; 126:241–250. [PubMed: 9847238]
- Viney ME, Thompson FJ, Crook M. TGF- β and the evolution of nematode parasitism. *International Journal for Parasitology*. 2005; 35:1473–1475. [PubMed: 16139836]
- Voehringer D, Shinkai K, Locksley RM. Type 2 immunity reflects orchestrated recruitment of cells committed to IL-4 production. *Immunity*. 2004; 20:267–277. [PubMed: 15030771]
- Wilson MS, Taylor M, Balic A, Finney CAM, Lamb JR, Maizels RM. Suppression of allergic airway inflammation by helminth-induced regulatory T cells. *Journal of Experimental Medicine*. 2005; 202:1199–1212. [PubMed: 16275759]
- Wolstenholme AJ, Fairweather I, Prichard R, von Samson-Himmelstjerna G, Sangster NC. Drug resistance in veterinary helminths. *Trends in Parasitology*. 2004; 20:469–476. [PubMed: 15363440]
- Zavala-Gongora R, Kroner A, Bernthaler P, Knaus P, Brehm K. A member of the transforming growth factor- β receptor family from *Echinococcus multilocularis* is activated by human bone morphogenetic protein 2. *Molecular and Biochemical Parasitology*. 2006; 146:265–271. [PubMed: 16434111]
- Zawadzki JL, Presidente PJ, Meeusen EN, De Veer MJ. RNAi in *Haemonchus contortus*: a potential method for target validation. *Trends in Parasitology*. 2006; 22:495–499. [PubMed: 16971180]




Figure 1. Primer design and amplification of *tgh-2* sequences

A. Schematic of *H. contortus tgh-2* genomic organisation, in comparison to the corresponding organisation of the *A. caninum* homologue, *Ac-tgh-2*, according to Freitas and Arasu (Freitas & Arasu, 2005). Based on cDNA sequence of *H. contortus tgh-2* (accession number FJ391183), exons were identified on contigs 0077796, 016772 and 0001624. Nucleotide lengths of exons (below or above box, bold) and introns (within box, italic) are given. Intron boundaries invariably conformed with the GT....AG consensus. Longer intronic tracts which contain gaps in existing sequence are indicated with broken

lines. Note that no genomic sequence yet corresponds with exon VI of *Hc-tgh-2*. The active domain is encoded within Exons VII-X (indicated in yellow). The positions of forward and reverse primers used for RT-PCR analysis (see Fig. 5 below) are indicated beneath the *H. contortus* diagram.

B. Alignment of the active domains (AD) of TGF- β family members from *C. elegans* (*Ce*-DAF-7, AAC47389), *B. malayi* (*Bm*-TGH-2, AF104016), *A. caninum* (*Ac*-TGH-2, AAY79430 and AAX36084, identical depositions), and the newly-identified *H. contortus* (*Hc*-TGH-2, accession number FJ391183). Numbering corresponds to the *Ce*-DAF-2 sequence. Identical amino acids are shaded, and the amino acid sequences coded by the degenerate primers shown in solid blocks.

C. Degenerate primer sequences used in this study (Y = C/T, R = A/G and N = A/C/G/T).

D. Ethidium bromide gels of degenerate PCR products amplified from *H. polygyrus*, *N. brasiliensis* and *T. circumcincta*. Sizes of marker polynucleotides are given in bp. *H. polygyrus* and *N. brasiliensis* products were amplified using F1 and R1 degenerate primers, while the *T. circumcincta* product was amplified using F2 and R1 degenerate primers as shown in **B**.




Figure 2. 5' and 3' RACE products of TGF- β homologues

Upper panel : 5' RACE of *H. contortus*, *H. polygyrus*, *N. brasiliensis* and *T. circumcincta* adult cDNA, amplified and visualised by ethidium bromide staining of agarose gels.

Following sequencing, the indicated bands were found to contain *tgh-2*-like sequences.

Lower panel : 3' RACE from the same cDNA samples as above.




Figure 3. Full amino acid sequences and alignment of TGF-β homologues
 Protein alignments of known TGF-β homologues from *Homo sapiens*, *B. malayi*, *A. caninum*, and novel TGF-β homologues from *H. contortus*, *H. polygyrus*, *N. brasiliensis* and *T. circumcincta*. Numbering corresponds to the human protein sequence. Predicted signal sequences are shown in red boxes, or where no signal sequence is predicted, start methionines are indicated in solid red boxes. Solid blue box indicates tetrabasic protease cleavage site, yellow boxes indicate conserved cysteines. Open turquoise boxes denote potential N-glycosylation motifs (N-X-S/T). Accession numbers for novel sequences are: *Hp-tgh-2* FJ410912, *Nb-tgh-2* FJ410912 and *Tc-tgh-2* FJ410912.




Figure 4. Phylogenetic tree of TGF- β superfamily members

Neighbour Joining phylogenetic tree of TGF- β superfamily members, generated using with Poisson correction on MacVector; numbers indicate bootstrap values (percentage of calculated trees branching at each point, based on 1000 repetitions). Where no numbers are shown, bootstrap values are below 50%.




Figure 5. Expression levels of *tgh-2* transcript in different life cycle stages

Levels of *tgh-2* and actin transcription were measured by real-time PCR in different developmental stages of *H. polygyrus* (A), *N. brasiliensis* (B), *H. contortus* (C) and *T. circumcincta* (D). The genomic position of the RT-PCR primers used for *H. contortus* are shown in Figure 1 (RT-F and RT-R). The relative transcription levels were calculated by dividing the relative levels of *tgh-2* transcription by relative levels of actin transcription. Standard errors shown are calculated from replicate determinations of *tgh-2* levels divided by the mean of replicate determinations of actin levels. Relative levels of *T. circumcincta tgh-2* were very similar when β -tubulin was used as the reference gene (data not shown). Data are representative of at least two independent experiments in each case.

TABLE 1

Gene-specific primers used in 3' and 5' RACE (A) and in RT-PCR (B). RACE reactions used a single gene-specific primer with a second primer to a sequence ligated onto all cDNA molecules; products and sizes are shown in Fig. 2. RT-PCR used the indicated primer pairs and yielded products of the sizes indicated. Numerals in brackets correspond to the nucleotide positions in the full-length cDNA sequences as deposited at GenBank.

A. RACE Primers		
Transcript	3' RACE (Forward primer)	5' RACE (Reverse primer)
<i>H. contortus tgh-2</i>	5'-ATGTTTCATGTCGGTTACGCTGG-3' (651-673)	5'-TTCAGCGGGTGACAGCACATTC-3' (954-932)
<i>H. polygyrus tgh-2</i>	5'-CCCGGATGTTGCCTCTACGAC-3' (589-609)	5'-CCCCTGATTCCAGTTGCGTC-3' (789-769)
<i>N. brasiliensis tgh-2</i>	5'-GGTTGCTGTCTCTACGATCTC-3' (148-168)	5'-CCCTTGTTACCCGTTGCGTC-3' (342-322)
<i>T. circumcincta tgh-2</i>	5'-GGGTGATTGTTCTGTGAATCAC-3' (699-720)	5'-GTGATTCACAGAACAATCACCC-3' (720-699)

B. Real-time PCR Primers			
Transcript	Forward primer	Reverse primer	Product size
Actin (all species)	5'-TGAGCACGGTATCGTCACCAAC-3' (216-237)	5'-TTGAAGGTCTCGAACATGATCTG-3' (386-364)	171 bp
<i>T. circumcincta</i> β -tubulin	5'-TTCCATFCCCTCGTCTTACAC-3' (773-792)	5'-AGCCATTTCAATCCACGAG-3' (1092-1073)	320 bp
<i>H. contortus tgh-2</i>	5'-ATGTTTCATGTCGGTTACGCTGG-3' (652-673)	5'-TTCAGCGGGTGACAGCACATTC-3' (954-932)	303 bp
<i>H. polygyrus tgh-2</i>	5'-CCCGGATGTTGCCTCTACGAC-3' (589-609)	5'-CCCCTGATTCCAGTTGCGTC-3' (789-769)	201 bp
<i>N. brasiliensis tgh-2</i>	5'-GGTTGCTGTCTCTACGATCTC-3' (148-168)	5'-CCCTTGTTACCCGTTGCGTC-3' (342-322)	195 bp
<i>T. circumcincta tgh-2</i>	5'-GGGTGATTGTTCTGTGAATCAC-3' (699-720)	5'-TCAAGAGCATGTACATTTACGG-3' (918-897)	220 bp