

**Dance improves motor, cognitive and social skills in children with developmental cerebellar anomalies**

Valentin Bégel<sup>a,1</sup>, Asaf Bachrach<sup>b</sup>, Simone Dalla Bella<sup>c,d</sup>, Julien Laroche<sup>b,2</sup>, Sylvain Clément<sup>a</sup>, Audrey Riquet<sup>e</sup> & Delphine Dellacherie<sup>a,e</sup>

<sup>a</sup>University of Lille, ULR 4072 – PSITEC – Psychologie : Interactions, Temps, Emotions, Cognition, F-59000 Lille, France

<sup>b</sup>UMR 7023 CNRS/Université Paris 8, 75017 Paris, Paris, France

<sup>c</sup>International Laboratory for Brain, Music and Sound Research (BRAMS), 1430 Boulevard du Mont-Royal, Montreal, QC H2V 2J2, Canada

<sup>d</sup>University of Montreal, Department of Psychology, CP 6128 Succursale Centre-Ville, Montreal, QC, H3C 3J7, Canada

<sup>e</sup>University Hospital of Lille, Department of Pediatric Neurology, 2 Av. Oscar Lambret, 59037, Lille, France

<sup>1</sup> Present address: McGill University, Department of Psychology, 1205 Dr Penfield Ave, Montreal, QC, H3A 1B1, Canada.

<sup>2</sup> Present address: Fondazione Istituto Italiano di Tecnologia (IIT), Center for Translational Neurophysiology of Speech and Communication (CTNSC), Ferrara, Italia

Manuscript accepted for publication in *The Cerebellum*

\* Corresponding author:

Delphine Dellacherie

University of Lille, ULR 4072 – PSITEC – Psychologie : Interactions, Temps, Emotions, Cognition, F-59000 Lille, France

Email: [delphine.dellacherie@univ-lille.fr](mailto:delphine.dellacherie@univ-lille.fr)

Phone: +33 320 41 6370

**Abstract:** In this multiple single-cases study, we used dance to train Sensorimotor Synchronization (SMS), motor and cognitive functions in children with Developmental Cerebellar Anomalies (DCA). DCA are rare dysfunctions of the cerebellum that affect motor and cognitive skills. The cerebellum plays an important role in temporal cognition including SMS which is, critical for motor and cognitive development. Dancing engages the SMS neuronal circuitry, composed of the cerebellum, the basal ganglia and the motor cortices. Thus, we hypothesized that dance has a beneficial effect on SMS skills and associated motor and cognitive functions in children with DCA. Seven children (aged 7-11) with DCA participated in a 2-month dance training protocol (3h/week). A test-retest design protocol with multiple baselines was used to assess children's SMS skills as well as motor, cognitive and social abilities. SMS skills were impaired in DCA before the training. The training led to improvements in SMS (reduced variability in paced tapping), balance and executive functioning (cognitive flexibility), as well as in social skills (social cognition). The beneficial effects of the dance training were visible in all participants. Notably, gains were maintained two months after the intervention. These effects are likely to be sustained by enhanced activity in SMS brain networks due to the dance training protocol.

**Keywords:** Cerebellar anomalies, sensorimotor synchronization, rhythm, rehabilitation, dance

## 1. Introduction

Developmental Cerebellar Anomalies (DCA) are rare diseases that affect cerebellar functions in less than one in 2000 (Musselman et al., 2014). The cerebellum is a pivotal brain region that contributes to motor and cognitive control as well as social and affective functions (Bareš et al., 2018; Hoche, Guell, Sherman, Vangel, & Schmahmann, 2016; Koziol et al., 2014; Manto et al., 2012; Schmahmann & Sherman, 1998; Therrien & Bastian, 2019). It achieves maturity a few years after birth, making it particularly vulnerable to developmental disorders (ten Donkelaar, Lammens, Wesseling, Thijssen, & Renier, 2003; Wang & Zoghbi, 2001). Unfortunately, very few specific remediation strategies to compensate for DCA currently exist (Brenna, 2014), possibly because of the rarity of this disease. The current research focuses on an innovative training protocol using dance to improve motor, cognitive and social skills in children with DCA.

DCA can have various causes (e.g., Joubert and Dandy-Walker syndromes, intrauterine trauma or intoxication). Cerebellar syndromes found in DCA are often explained by structural and volumetric anomalies of the cerebellum (Boltshauser, 2004; Poretti, Boltshauser, & Doherty, 2014). They can also be diagnosed on the basis of clinical observations in the absence of neuroimaging evidence (Ryan & Engle, 2003). Predominantly, DCA is associated with congenital forms of cerebellar ataxia (Fogel, 2012; Marsden, 2018). Ataxia refers to abnormal organization of movement and impaired coordination of balance, gait, extremity and eye movements, and dysarthria. DCA is also consistently associated with cognitive deficits (e.g., in executive functions, visual-spatial skills, linguistic abilities; Bodranghien et al., 2016; De Smet, Paquier, Verhoeven, & Mariën, 2013; Schmahmann, 2004; Schmahmann, Guell, Stoodley, & Halko, 2019; Schmahmann & Sherman, 1998) and emotional, social and communication deficits (Bolduc & Limperopoulos, 2009; Guell, Hoche, & Schmahmann, 2015; Hoche et al., 2016; Koziol et al., 2014; Schmahmann, 2004; Schmahmann & Sherman, 1998).

The cerebellum plays a crucial role in motor control (Manto et al., 2012); for example, it is involved in the control of oculomotor movements (Thier & Markanday, 2019), speech (Ackermann, 2008), grasping (Nowak, Topka, Timmann, Boecker, & Hermsdörfer, 2007), and voluntary limb movements (Ebner, Hewitt, & Popa, 2011). A specific contribution of the cerebellum to these tasks is the control of precise, online adaptive and predictive timing (Bastian, 2006; Manto et al., 2012; Therrien & Bastian, 2019). The cerebellum is involved in coordinating an action with temporally predictable, rhythmic events (Bareš et al., 2018; Braitenberg, Heck, & Sultan, 1997; Damm, Varoqui,

Cochen De Cock, Dalla Bella, & Bardy, 2019; Manto et al., 2012; Molinari et al., 2003; Molinari, Leggio, & Thaut, 2007; Paquette, Fujii, Li, & Schlaug, 2017), especially in auditory–motor interactions (Doyon, Penhune, & Ungerleider, 2003; Provasi et al., 2014; Schwartz, Keller, & Kotz, 2016; Zatorre, Chen, & Penhune, 2007). This is referred to as Sensorimotor Synchronization (SMS, Repp, 2005; Repp & Su, 2013). SMS is usually studied via finger tapping to predictable stimuli such as a metronome or music sequences (Dalla Bella et al., 2017). The cerebellum is involved in the perception of sequences (Molinari et al., 2008), a capacity that is required for SMS. In addition, cerebellar circuitry is necessary to produce and adapt the next motor response using error information (i.e., discrepancy between the tap and the sound) of the previous response (Shadmehr, Smith, & Krakauer, 2010).

More generally, the cerebellum is part of a distributed neural system, including the basal ganglia, motor and premotor cortices, that enables temporal processing (Breska & Ivry, 2016; Coull & Nobre, 2008; Coull, Cheng, & Meck, 2011; Grahn, 2009; Grahn & Brett, 2007; Kotz & Schwartz, 2011). The cerebellum is crucial for timing and rhythmic tasks as well as for other motor and cognitive capacities in which timing and SMS are involved, such as gait (Damm, Varoqui, Cochen De Cock, Dalla Bella & Bardy, 2019; Bengtsson, Ehrsson, Forssberg, & Ullén, 2005; Dalla Bella, Benoit, Farrugia, Schwartz, & Kotz, 2015; Lang et al., 2017), language (Murdoch, 2010; Schwartz & Kotz, 2013) and social interactions (Van Overwalle, Manto, Leggio, & Delgado-García, 2019). Therefore, it is not surprising that patients with DCA are usually impaired in a variety of the aforementioned motor, cognitive and social capacities associated with SMS (Bodranghien et al., 2016; Koziol et al., 2014; Schmähmann, 2004; Schmähmann & Sherman, 1998).

To date, there is no direct evidence that SMS skills are affected in DCA, as no research studied SMS skills such as tapping to a metronome or music in these patients. Nevertheless, studies have reported other timing impairments with other cerebellar acquired or degenerative pathologies such as spino cerebellar ataxia or infarction. They are visible in tasks requiring precise timing, such as eye movements, spatio-temporal prediction, predictive motor timing and finger movements in overarm throwing (Broersen et al., 2016; Ivry & Keele, 1989; Ivry, Spencer, Zelaznik, & Diedrichsen, 2006; Timmann, Watts, & Hore, 1999). Interestingly, timing disorders are not confined to motor tasks, as cerebellar anomalies are also associated with poorer performance in perceptual tasks such as duration discrimination (Ivry & Keele, 1989; Ivry, Spencer, Zelaznik, & Diedrichsen, 2002). Provasi et al. (2014) studied SMS and rhythm perception in children with acquired (not developmental) cerebellar tumors (medulloblastoma) and showed that patients displayed

more tapping variability in SMS task, despite normal rhythm perception. Therefore, it is likely that patients with DCA also have difficulties in SMS.

Because of the role played by the cerebellum in motor and sensorimotor skills, dance is of particular interest for persons with cerebellar dysfunctions. Dance is an engaging activity involving an elaborate form of full body synchronization to a musical beat in a nonverbal communicative, aesthetic context (Joufflineau, Vincent, & Bachrach, 2018; Laland, Wilkins, & Clayton, 2016). It engages neuronal networks including the premotor and parietal cortices (“action observation network”), the supplementary motor area (SMA), the motor cortex, the basal ganglia and the cerebellum (Bachrach, Jola, & Pallier, 2016; Brown, Martinez, & Parsons, 2006; Cross, Hamilton, & Grafton, 2006; Cross, Kraemer, Hamilton, Kelley, & Grafton, 2009; Giacosa, Karpati, Foster, Penhune, & Hyde, 2016; Karpati, Giacosa, Foster, Penhune, & Hyde, 2015). The anterior cerebellar vermis and cerebellar lobules V and VI appear as key structures for the entrainment of movement to external auditory timing cues in dance (Brown et al., 2006) and expert dancers reveal brain changes, structural and functional, in these regions. Expertise in dance is also associated with improved sensorimotor functions (Bläsing et al., 2012; Jin et al., 2019; Karpati, Giacosa, Foster, Penhune, & Hyde, 2016), and proprioceptive body representation (Jola, Davis, & Haggard, 2011), thus supporting the tight link between brain networks affected by dance training and the sensorimotor and proprioceptive functions.

During the last decade, promising results showed that dance training can be used successfully for improving motor and cognitive functions in a variety of conditions such as aging (Chuang, Hung, Huang, Chang, & Hung, 2015; Coubard, Duretz, Lefebvre, Lapalus, & Ferrufino, 2011), neurodegenerative diseases (de Natale et al., 2017), stroke (Sampaio, Subramaniam, Arena, & Bhatt, 2016) and neurodevelopmental diseases (Grönlund, Renck, & Weibull, 2005). For example, dance training showed beneficial effects on motor and cognitive skills in Parkinson’s disease (dos Santos Delabary, Komerowski, Monteiro, Costa, & Haas, 2018; Hackney et al., 2015; Hackney, Kantorovich, Levin, & Earhart, 2007; McKee & Hackney, 2013; Shanahan, Morris, Bhriain, Saunders, & Clifford, 2015), a condition associated with poorer rhythm and timing skills (Benoit et al., 2014; Grahn & Brett, 2009; Jones & Jahanshahi, 2014). Beneficial effects of a dance training protocol on motor functions were also observed in children with ADHD (Grönlund et al., 2005), a condition also affecting the cortico-cerebello-striatal pathways and rhythmic capacities (Bledsoe, Semrud-Clikeman, & Pliszka, 2011; Puyjarinet, Bégel, Lopez, Dellacherie, & Dalla Bella, 2017), and in children with cerebral palsy (López-Ortiz, Egan, & Gaebler-Spira, 2016; López-Ortiz et al., 2012). Moreover, consistent with the notion that

dance is a social and highly engaging activity (Chauvigné, Walton, Richardson, & Brown, 2019; Hackney & Gammon, 2011; Himberg, Laroche, Bigé, Buchkowski, & Bachrach, 2018; Joufflineau et al., 2018), positive effects of this activity on psychosocial behavior and emotional responses are reported, especially in children with developmental disorders (Grönlund et al., 2005; Jeong et al., 2005; Teixeira-Machado, Azevedo-Santos, & De Santana, 2017). Yet, it is not clear whether these motor and socio-affective improvements are mediated by SMS skills, because most studies did not include SMS testing. More generally, the effect of dance on cerebellar deficits has not been evaluated so far, except in one recent case study reporting the effects of an eight-week dance training protocol (based on an adapted form of tango) in an adult with a non-congenital cerebellar ataxia (Song, Ryu, Im, Lee, & Park, 2018). After the dance training, this patient displayed improved standing balance, gait characteristics, and functional mobility as well as self-declared improvements in emotion regulation and quality of life. This result is promising for the use of dance in cerebellar dysfunctions. However, so far, other physical therapy programs for patients with cerebellar dysfunctions, especially DCA, are scarce (Brenna, 2014), and, to the best of our knowledge, none included SMS or rhythmic components (Marsden & Harris, 2011; Martin, Tan, Bragge, & Bialocerkowski, 2009; Sartor-glittenberg & Brickner, 2014). We aimed to respond to this lack of rehabilitation methods in DCA by designing a sensorimotor training protocol based on dance.

The goal of this study was to test the effect of dance on a range of functions in children with DCA. This small-scale study included seven children with DCA who participated in a customized seven-week dance training program. Children with DCA were all diagnosed with cerebellar ataxia and displayed Coordination disorders. Nevertheless, participants displayed different cerebellar signs (Table 1). We implemented a concurrent multiple baseline design adapted to small sample studies. With this method, the number of baselines (i.e., evaluation sessions before the protocol begins) was set randomly for each participant (between 4 and 7 baselines). This allows an accurate and rigorous assessment of the patient's initial performance. Therefore, the effects of the intervention can be highlighted more reliably than with a single baseline or with the same number of baselines for all participants, reinforcing the causality between treatment and performance change (Himle, Miltenberger, Flessner, & Gatheridge, 2004; Morgan & Morgan, 2008). In addition, analyses methods adapted to single-case studies were used (Crawford & Garthwaite, 2007). Finally, a control group was tested in order to compare patients' skills with those of age-matched typically

developing children and to control for the test-retest effect and the normal developmental evolution of the main tested domains.

Children SMS's skills when tapping to an external stimulus were evaluated. A battery of fine motor, cognitive tasks and social questionnaires was also used. We expected improved SMS skills after the training, with possible associated effects on motor, cognitive and social skills.

## 2. Methods

### 2.1.1. Participants

#### Patients (CEREB)

Case histories: Seven children (2 females, all right-handed) with DCA, aged between 7.5 and 10.3 years at the inclusion stage, participated in the study (Table 1). Patients were recruited at the Centre de Référence Maladies Rares: Malformations et Maladies Congénitales du Cervelet (CRMR, University Hospital of Lille) where they each received a multidisciplinary assessment including a neurological standard examination and a neuropsychological evaluation. No specific scale to evaluate congenital ataxia currently exists. Cerebellar functionality was evaluated with the items of the Scale for the assessment and rating of ataxia (SARA) (Schmitz-Hübsch et al., 2006), although the quantitative items of this scale were not specifically developed for this type of congenital ataxia. The results at the SARA revealed the presence of cerebellar signs in all patients (see Table 1). All patients were diagnosed with cerebellar ataxia of unknown etiology (Patients 1 to 6) or, in one case (Patient 7), with cerebellar ataxia in the context of a Joubert's syndrome linked to mutations of the gene *CC2D2A* (see Table 2). Cerebellar ataxia was associated with malformations observable on structural brain imagery (MRI) in Patients 1 (cisterna magna and slightly enlarged fourth ventricle), 4 (vermian dysplasia), 5 (abnormal fissuring of the superior vermis) and 7 (molar tooth sign, cerebellar dysplasia of superior vermis, and malformation of the cerebellar peduncles) whereas Patients 2, 3 and 6 did not present any morphological malformation visible on a structural MRI. The exclusion criteria were as follows : (1) radiological picture or medical history consistent with a diagnosis of acquired lesions due to pre-natal or perinatal issues; (2) progressive cerebellar pathology; (3) metabolic diseases associated with congenital malformation of the cerebellum; (4) malformation affecting other cerebral structures besides the cerebellum; (5) epileptic seizures or febrile convulsions or even focal or generalized paroxysmal EEG abnormalities (spikes or spike-wave complexes), (6)

Intellectual disability ( $IQ < 70$ ) in the patients clinical history. Table 2 presents IQ indexes for each participant. All indexes are within the norms, although some patients' performances were in the low norms in verbal (VCI) and non-verbal (PRI) and visual-spatial (VSI) indexes. Most patients displayed borderline performances in processing speed (PSI). None of them had music or dance experience nor were engaged in extra-curricular musical, dance or sport activities during the study.

Control group (CONTROL):

A group of 19 children (CONTROL) aged between 7.00 and 10.58 years ( $M = 8.75$ ,  $SD = 1.27$ , 7 females) were tested twice with a 8-week interval on a subset of tasks in order to control the test-retest effect and the effect of normal development on the children performances. No children who displayed impairments in verbal comprehension and/or perceptual reasoning, with perceptual or motor impairments, neurological, psychiatric, or developmental disorders were included in the control group. Children with dance and/or musical training were not included.

The study was approved by the National Ethics Committee (CPP Sud Est III, France, EudraCT: 2017-A03300-53). All participants and their parents gave their informed consent before testing and intervention in accordance with the Declaration of Helsinki.

## 2.2. Procedure

Figure 1 shows the experimental design for the CEREB group. Children in the CEREB group took part in a seven-week dance training protocol at the Eurasport center (University of Lille, France). Before and after the training, a complete individual testing session was administered to each participant (see 2.4. Experimental measures, for a description of the tests). Testing included the evaluation of movement, sensorimotor and cognitive skills.

Questionnaires assessing social cognition, executive functioning, and anxiety were also administered. Two months after the training, a two-hour evaluation session was administered in order to evaluate the long-term effects of the training.

For a subset of tasks (Synchronization with a metronome and with music, Verbal inhibition and switching from the NESPY-II), a concurrent multiple baselines design was used. We administered randomized baseline periods (i.e., each participant received a different number of short pre-intervention evaluations). Children were randomly assigned from 4 to 7 multiple baselines. These short evaluation sessions lasted 15 minutes. Repeated measures with the same subset of



tasks were also administered on a weekly basis during treatment. Since the intervention starts for all patients at the same time after a different number of baselines between participants, if an improvement is observed after the start of the intervention regardless of the number of baselines, this improvement can be attributed with more certainty to the intervention. In other words, this multiple baseline methodological design reinforces the inference of causal links between the intervention and the observed improvement.

### 2.3. Training

CEREB children participated in a seven-week dance training protocol specifically developed for that purpose. A total of fourteen dance classes (one and a half hours each) were given regularly by a professional dance teacher. The dance training consisted in several playful exercises with different physical intensity. These exercises combined several components, with and without musical beat: 1) Whole body movement to rhythmic music or without music; 2) Sequence learning; 3) Coordination with one or more partners; 4) Expressivity and creativity through movement improvisation. The order of the exercises was changed across sessions to make them less repetitive and more interesting. Instrumental music suitable for children was used. All excerpts had a salient beat. The tempo was moderate (80-100BPM). Each exercise lasted about seven minutes. Each session started with warm-up and ended with relaxation and stretching exercises.

During the sessions, three main types of exercises were proposed:

Synchronization with rhythms: with these exercises, children learned to perceive, produce, synchronize with rhythms, and adapt to rhythmic changes. For example, they had to walk in the room synchronizing their steps with a tambourine sound. Either the dance teacher or one of the children played the tambourine and chose the tempo speed, trying to keep it steady. A variation of this exercise was to step in hoops laid out on the floor, synchronizing each step with the rhythm. Other exercises involved the production of rhythmic sequences in group, using hand clapping or drumming on small percussion instruments, sometimes accompanied with singing.

Body and space exploration: various exercises were used to foster children's consciousness of the body and space. These exercises were performed with or without rhythmic sounds, such as percussion sounds played by the dance teacher or music. They consisted in improvised movements or progressive movements (e.g., moving from one position

to another in a given number of beats). Other exercises involved moving a small plastic ball in the space with different instructions, such as touching other children with the ball or bringing it to different places in the room.

Choreographies: in each session, moments were dedicated to improvisation and creation. Choreographies were created using movements proposed by the children or the dance teacher. These choreographies were rehearsed at each session and new movements were added. These exercises were designed to encourage creativity and interactions between the children.

## 2.4. Experimental measures

### 2.4.1. Rhythm assessment

Perceptual rhythm assessment: the Beat Alignment Test (BAT, Iversen & Patel, 2008) was taken from The Battery for the Assessment of Auditory and Sensorimotor Timing Abilities (BAASTA, Dalla Bella et al., 2017). The task was to judge whether a metronome was aligned or not with the beat of music. Two 20-beat musical excerpts from Bach's « Badinerie » and two from Rossini's « William Tell Overture » were played at a tempo of 600ms of Inter-Beat Interval (IBI). After 7 beats, a metronome (isochronous tones with a triangle timbre) was superimposed onto the music. When unaligned, tones occurred earlier or later than the beat by 33% of the quarter note duration (phase shift, 12 stimuli), or the interval between the tones was increased or decreased by 10% of the quarter note duration (period shift, 12 stimuli). Stimuli were presented in pseudo-randomized order with PsychoPy 2 (<http://www.psychopy.org/>) installed on a Dell computer (XPS 13 9360 Model). Auditory stimuli were delivered via headphones (Sennheiser HD280) at a comfortable sound level (60 dB). Responses were provided verbally by the children, and entered by the Experimenter by pressing one of the two response keys (“yes” or “no”). “Yes” indicated the situation when the child considered that the metronome was aligned with the musical beat. The sensitivity index ( $d'$ ) was used as an unbiased measure of detection performance. It is obtained by calculating the standardized difference between the Hits (i.e., when a misaligned metronome was correctly detected) and False alarms (i.e., when a misalignment was erroneously reported):  $z(H) - z(FA)$ .

Motor and sensorimotor rhythmic tasks: motor and sensorimotor rhythmic skills were assessed using finger tapping tasks from the BAASTA (Dalla Bella et al., 2017). A self-paced finger tapping (referred to as Unpaced tapping) was used to assess participants' spontaneous motor tempo and variability. Sensorimotor rhythmic skills were assessed

using finger tapping to the beat of a metronome or music (referred to as Paced tapping). For tapping tasks, a tablet version of BAASTA was used (Acer Iconia Tab 10 model) (Bégel, Verga, Benoit, Kotz, & Dalla Bella, 2018; Puyjarinet et al., 2017). Note that the timing precision of the presentation of the stimuli on the tablet was 1% of the inter onset interval. Auditory stimuli were delivered via headphones (Sennheiser HD280). Stimuli were presented at a comfortable sound level (60 dB). Children had to tap with their index finger on the tablet. In Unpaced Tapping, participants had to produce regular finger taps with their dominant hand at a comfortable rate for 60 seconds. In Paced Tapping to metronome and music, children's capacity to track the beat was tested with finger tapping synchronization tasks. The metronome sequences included 60 piano tones (tone frequency = 1319 Hz) of 600 ms IBI. Two additional conditions with slow (750 ms) and fast (450 ms) IBIs were used at pre and post testing (but not in weekly evaluations). 64-beats musical excerpts taken from Bach's « Badinerie » and Rossini's « William Tell Overture » (Beat = quarter note) were used in Paced Tapping with music. The IBI was 600 ms, which is within the range for optimal rhythm perception (Drake & Botte, 1993; London, 2012; Repp, 2006). All stimuli were repeated twice. Artifacts and outliers were discarded as follows. Artifacts were taps leading to Inter-Taps Intervals (ITIs) smaller than 100 ms. In addition, taps were considered as outliers when the ITI between the actual tap and the preceding tap was smaller than  $Q1 - 3 * \text{Interquartile range (IQR)}$  or greater than  $Q3 + 3 * \text{IQR}$ , where  $Q1$  is the first quartile and  $Q3$  is the third quartile (Dalla Bella et al., 2017). The first ten taps of each trial were discarded. In Unpaced tapping, the mean ITI was computed. The coefficient of variation (CV) of the ITI (the ratio of the SD of the ITIs over the mean ITI) was calculated as a measure of motor variability. In Paced tapping, the CV of the ITI was used as a measure of motor variability during a synchronization task. In Paced tapping with a metronome, the mean performance at the two trials was calculated for each tempo. The four music trials were averaged for tapping with music.

#### 2.4.2. Balance and fine motor control

Subtests from the Movement Assessment Battery for Children (M-ABC-II, Henderson, Sugden, & Barnett, 2007) (Static and dynamic balance, Manual dexterity) and the NEPSY-II (Brooks, Sherman, & Strauss, 2010) (Hand imitation) were used to assess balance and motor skills. Three subtests from the M-ABC-II (Placing pegs in a peg board, Threading a lace and Drawing a line into a trail) were used to evaluate Manual dexterity. The percentile score at manual dexterity on the French version of the M-ABC-II (Soppelsa & Albaret, 2005) was calculated for each patient. In Static and dynamic balance (M-ABC-II), three subtests from the M-ABC-II (One-leg balance, Walking on a line

and Hopping on mats) were used. The final scores were the average time the child kept on the balance board for the One-leg balance task (Maximum = 30 seconds), the total number of steps for the Walking on a line task (Maximum = 15) and the number of hops averaged across the two legs for the Hopping on mats task (Maximum = 5). The percentile score at static and dynamic balance on the French version of the M-ABC-II (Soppelsa & Albaret, 2005) was also calculated for each patient. In Hand imitation (NEPSY-II), the experimenter presented up to 12 different positions with his hand. Children were asked to reproduce the position. The task was done first with the dominant hand and was repeated with the non-dominant hand. A total score (Maximum = 24) was calculated.

#### 2.4.3. Neuropsychological assessment

A battery of neuropsychological tests was submitted. Subtests were taken from the Test of Everyday Attention for Children (TEA-ch, Manly et al., 2001) (Motor inhibition, Cognitive flexibility) and the NEPSY-II (Brooks et al., 2010) (Verbal inhibition and switching, Verbal and Non-verbal fluencies). The Motor inhibition task (TEA-ch) is a go-no go task consisting of pointing at squares while listening to short sequences of tones and stopping to point when a different timbre occurs. The final score is the number of good responses. In Cognitive flexibility (TEA-ch), Children had to count the number of “creatures” (i.e., green little monsters) visible all along their burrow. Arrows are interspersed among the creatures and pointed either upwards or downwards. They were instructed to begin counting the creatures one by one and to change the direction of counting when the sense of the next arrow was downwards, until the last creature was presented. The final score was calculated based on the number of good responses (Quality index) and the time spent (Speed index) to complete the task. Verbal inhibition and switching (NEPSY-II) consisted in a series of black and white shapes (squares and circles or upwards or downwards arrows) that children had to denominate or tell the direction (denomination condition). In two other conditions, they were instructed to give the name of the other shape or of the other direction (inhibition), and to name the shape or give the direction when it was black and to give the name of the other shape or direction when it was white (change condition). The total number of errors and the time of execution were calculated for each condition. The total of errors across conditions was also calculated. In Verbal fluencies (NEPSY-II), participants were asked to generate as many words as they could within a specific semantic category (Animals and foods/drinks) or initial letter (S and F) category for 60 seconds. A score of semantic fluency and a score of initial letter category were calculated by cumulating the total number of correct words generated in each category. In Non-verbal fluencies (NEPSY-II), children had to connect structured or random dots presented in squares

taking care not to form the same pattern as in previous squares. The total number of correct squares filled was considered as the final score for each condition, and a composite score for non-verbal fluencies was calculated by adding the two conditions.

#### 2.4.4. Questionnaires

Questionnaires filled by the children and by the parents were used to assess executive functions in everyday life, social cognition, and anxiety. The Behavior Rating Inventory of Executive Function parent version (BRIEF, Gioia, Isquith, Guy, & Kenworthy, 2000), a validated questionnaire, was used to measure the child's executive function competence in the real-world setting. Six sub-scores (the higher the score the better) were provided: Inhibition, Cognitive flexibility, Emotional control, Initiation, Working Memory, Planification/Organisation, Organisation of materials and self-control. A global composite score was also computed. A Social cognition & theory of mind questionnaire created by our research team (Dellacherie et al, in preparation) was administered. It consisted in a series of 15 five-level Lickert scale questions evaluating children capacities in social cognition and theory of mind. The final score is the sum of the scores obtained for each question. Higher scores indicate higher estimated capacities. Finally, anxiety reported by the children was evaluated using the Revised Children's Manifest Anxiety Scale (R-CMAS, Reynolds & Richmond, 1979). A global score was calculated. Low scores indicated high anxiety.

#### 2.5. Statistical analysis

Exploratory group statistical analyses: we first conducted exploratory group statistical analyses. For all tasks, we compared CEREB children's performances before and after the training with Wilcoxon signed-rank test. When a difference was observed, we also compared the difference between the pre-test and the follow-up test and between the post-test and the follow-up test. For tasks that have been included in the multiple baselines (Synchronization with a metronome and with music, Verbal inhibition and switching from the NESPY-II), the mean of the last four multiple baselines was used as the pre-training measure. In addition, we calculated the difference between the post and the pre-training (Delta pre/post), the follow-up and pre-training (Delta pre/follow-up) in the CEREB group, and between the retest and the test (Delta test/retest) in CONTROL. For tasks in which a significant difference between the pre- and the post- training evaluations was observed, we used Wilcoxon-Mann-Whitney U test to compare the Delta of the CEREB and the CONTROL groups. It is important to note that given the small sample size of the CEREB group and the

multiple tests, statistical power of group analyses is very limited. We used them only to highlight overall tendencies for CEREB children to display improved performance after the training.

Single-cases statistical analyses: Statistical analyses adapted to the study of single cases (Crawford & Garthwaite, 2007) were used to evaluate patients' performance before and after the training. CEREB performance at the pre and post-training were compared to controls' using statistics adapted for the analysis of single cases (DissocBayes program, Crawford & Garthwaite, 2007). For each patient, a subgroup of six to eight age-matched (i.e., they were less than six months older or one year younger than the patient) control participants was constituted. The mean of the last four multiple baselines was used as the pre-training measure for multiple baselines tasks.

### 3. Results

#### 3.1. Exploratory group analyses

Table 3 shows tasks for which a significant difference between the pre and the post training results were found. Note that almost all patients displayed a significant impairment before the training at these tasks (see below, "Single-case analysis" section).

Perceptual, motor and sensorimotor rhythmic tasks: When patients synchronized with a metronome and with music, tapping variability (CV) was significantly reduced after the training. The change was significantly greater in CEREB than in CONTROL. The differences with baseline were maintained at follow-up (Metronome: follow-up,  $M = .18$ ,  $SD = .17$ ,  $Z = 2.20$ ,  $p < .05$ ,  $r = 0.83$ ; Music: follow-up,  $M = .30$ ,  $SD = .16$ ,  $Z = 2.36$ ,  $p < .01$ ,  $r = 0.89$ ). No overall significant change was observed in perceptual rhythmic skills (BAT) and unpaced tapping ( $p > .05$ ) in CEREB.

Balance and fine motor control: Hopping on mats was improved after the training in CEREB, and the difference was maintained at follow-up (follow-up,  $M = 3.79$ ,  $SD = 1.87$ ;  $Z = 2.38$ ,  $p < .05$ ,  $r = 0.90$ ). The change in hopping on mats was significantly larger in CEREB than in CONTROL. No differences were observed between the pre and the post tests in the other variables ( $p > .05$ ).

Neuropsychological assessment: in neuropsychological tests, cognitive flexibility was improved after the training, both in terms of number of correct answers and time; the difference was maintained at follow-up for good answers (follow-up,  $M = 5.66$ ,  $SD = 1.03$ ;  $Z = 2.13$ ,  $p < .01$ ,  $r = 0.87$ ). The change was significantly greater in CEREB than in

CONTROL for both correct answers and time. A difference was also found in the number of errors of the change condition in verbal inhibition and switching (before,  $M = 8.62$ ,  $SD = 5.64$ ; after,  $M = 5.71$ ,  $SD = 5.68$ ;  $Z = 2.37$ ,  $p < .01$ ,  $r = 0.89$ ) and in the total number of errors (before,  $M = 16.36$ ,  $SD = 13.29$ ; after,  $M = 10.86$ ,  $SD = 9.62$ ;  $Z = 1.95$ ,  $p < .05$ ,  $r = 0.74$ ). However, these differences were not confirmed at follow-up (number of errors of the change condition, follow-up,  $M = 8$ ,  $SD = 12.61$ ;  $Z = 1.10$ ,  $p < .05$ ,  $z = 0.42$ ; total number of errors, follow-up,  $M = 9.33$ ,  $SD = 7.03$ ;  $Z = 1.16$ ,  $p < .05$ ,  $r = 0.47$ ). In addition, the change was similar in CONTROL and in CEREB for both the number of errors in the change condition (CEREB,  $M = -2.9$ ,  $SD = 2.92$ ; CONTROL,  $M = -1.68$ ,  $SD = 2.89$ ;  $Z = 0.55$ ,  $p > .05$ ,  $r = 0.11$ ) and the total number of errors (CEREB,  $M = -5.5$ ,  $SD = -5.96$ ; CONTROL,  $M = -2.28$ ,  $SD = 5.52$ ;  $Z = 1.12$ ,  $p > .05$ ,  $r = 0.22$ ). No improvements were observed in other neuropsychological tests ( $p > .05$ ).

Questionnaires: no effect of the training was observed on the overall score of the executive function questionnaire, but there was an improvement in the cognitive flexibility sub-score (Cognitive flexibility at home). Scores reported on the social cognition questionnaire were also better after the training. In these two questionnaires, the differences were not maintained at follow-up (Cognitive flexibility at home, follow-up,  $M = 13.6$ ,  $SD = 3.78$ ;  $Z = 0.27$ ,  $p > .05$ ,  $r = 0.10$ ; Social cognition questionnaire, follow-up,  $M = 59.33$ ,  $SD = 10.38$ ;  $Z = 1.26$ ,  $p > .05$ ,  $r = 0.52$ ). The changes were significantly different in CEREB (Delta pre/post) as compared with CONTROL for both Cognitive flexibility at home and Social cognition. No effects were found at the R-CMAS ( $p > .05$ ).

Table 4 shows the results at the first and second sessions for controls. For Hopping on mats, the average performance of controls (4.97) before the training was very close to the maximum score (5). Consequently, an improvement of balance between the first and the second sessions would not be detected by this task. A significant difference at the speed index of the cognitive flexibility task was observed, meaning that control participants did the task more quickly in the second session. This difference is not as large as the one observed in the CEREB group though.

### 3.2. Single-case analysis

Motor rhythmic skills: statistics calculated with the DissocsBayes program (Crawford & Garthwaite, 2007) are presented in Table 5. In tapping with a metronome (CV), Patients 2, 3, 4, 6 and 7 were impaired at pre-training, but only patient 3 was still impaired at post training. Only patients 1 and 5 did not have a significant dissociation between

pre and post-testing. In tapping with music (CV), patients 2, 3, 5, 6 and 7 were significantly impaired at pre-training. Patients 5 and 6 did not show measurable impairment after the training.

Neuropsychological assessment: single-case statistics are presented in table 6. In Balance (Hopping on mats), all patients displayed an impairment before the training. Patients 1, 4 and 6 had comparable performances after the training. Note that most controls participants obtained the maximum score (5) at this task. In Cognitive flexibility (Speed), Patients 5, 6 and 7 had a performance significantly poorer than controls at pre-training. This impairment was still found at post-training. At the Quality index of Cognitive flexibility, Patients 2, 4, 6 and 7 were impaired before the training, and patient 1 and 5's performances were borderline. Patient 4 and 5 displayed a normal performance at post-training in comparison with the control group. Note that Patient 3 did not pass the Cognitive flexibility task. At the Cognitive flexibility at home index, Patients 1, 2, 3 and 7 were significantly impaired before the training. Among them, Patient 3 and 7 displayed a normal performance after the training. Finally, in Social cognition, Patients 2, 3 and 5 had significantly lower scores than controls at pre-training.

#### 4. Discussion

This interventional study explored the effect of a dance training protocol in children with DCA. Seven patients participated in fourteen dance classes over a period of seven weeks. Before and after the training, we tested children with DCA's movement, sensorimotor and cognitive skills, as well as social and affective functions. The results showed a positive change in rhythmic sensorimotor (Finger tapping with a metronome and music), motor (balance), cognitive (Cognitive flexibility) and social (Social cognition) skills in most patients. We compared these changes with the evolution of these skills in a group of age-matched control participants over the same period of time. This confirmed that the improvement was greater in the CEREB patients than in typically developing children, and not merely due to a test retest effect. Given the rarity of DCA and the complexity of the training protocol, only a small number of participants could be included in this study. We used single case analyses with a multiple baseline experimental design to compare the performance of each patient before and after the training.

Children with DCA's difficulties in SMS were manifest: six patients out of seven displayed a more variable performance than controls when tapping to the beat of a metronome or music. To the best of our knowledge, this is the



first time that rhythmic difficulties are reported in developmental anomalies affecting the cerebellum. This result highlights the important role played by the cerebellum in synchronization already suggested by studies showing rhythmic deficits in acquired cerebellar dysfunction (Provasi et al., 2014). Recent neuroimaging evidence also support the idea that the cerebellum is important for rhythm perception and neural tracking of beat and rhythm (Nozaradan, Schwartz, Obermeier, & Kotz, 2017; Molinari et al., 2003; Molinari, Leggio, & Thaut, 2007, Paquette et al., 2017; Schwartz et al., 2016).

After the dance training protocol, most patients had SMS performances comparable to those of controls. Their motor variability while synchronizing with a metronome or music was reduced (i.e., the performance was improved). Our study suggests that SMS skills can be improved in children with DCA. This confirms the hypothesis that dance, which enhances sensorimotor functions in other populations (Jin et al., 2019; Karpati et al., 2016), may boost these functions in cerebellar disorders. Dance training involves various sensorimotor tasks, and it is possible that similar effects would be obtained with other forms of sensorimotor training. This result could indicate that the cerebral network sustaining SMS can be trained to match an average functioning with specific remediation techniques. The cerebellum is belonging to this cerebral network that also involves the basal ganglia, the motor and premotor cortices (Breska & Ivry, 2016; Coull & Nobre, 2008; Coull, Cheng, & Meck, 2011; Grahn, 2009; Grahn & Brett, 2007; Kotz & Schwartz, 2011). Dance is a complex activity that recruits a vast array of brain regions, including the cerebellar-cortical and basal ganglia-cortical networks (i.e., the supplementary motor area, the motor cortex, the basal ganglia and the cerebellum; Bachrach et al., 2016; Brown et al., 2006; Cross et al., 2006, 2009; Giacosa et al., 2016; Karpati et al., 2015). It is reasonable to hypothesize that the recruitment of some of these dance associated regions would underlie the observed positive effects.

One question that remains unanswered is whether the training had a direct effect on the cerebellum. The observed improvement can also be underpinned by compensatory mechanisms in the brain involving the other brain structure sustaining timing, such as the basal ganglia (Schwartz & Kotz, 2013). Further studies involving neuroimaging could be used to confirm this hypothesis.

The cerebellar-cortical and basal ganglia-cortical networks that sustain SMS are also involved in various motor (Bengtsson, Ehrsson, Forssberg, & Ullén, 2005; Dalla Bella, Benoit, Farrugia, Schwartz, & Kotz, 2015; Lang et al.,

2017), cognitive (Murdoch, 2010; Kotz & Schwartz, 2010; Schwartz & Kotz, 2013) and social (Van Overwalle et al., 2019) skills. Therefore, it is not surprising that a dance protocol would affect other seemingly unrelated skills, confirming the idea that the sensorimotor training may exhibit a broader spectrum of remediation effects.

Balance involves some common structures with the SMS brain network such as the cerebellum and basal ganglia (Surgent, Dadalko, Pickett, & Travers, 2019). The effect of dance on balance associated brain network might be directly linked to the improvement of the SMS skills. Nevertheless, the changes are not necessarily limited to SMS brain networks. Dance involves balance training as well and hence possibly stimulates several other balance associated cerebral structures distinct from the SMS ones (e.g., frontal, temporal and parietal cortices, the thalamus, Brown et al., 2006; Karpati et al., 2015; Surgent et al., 2019).

Cognitive skills (Cognitive flexibility) were also improved by the dance training. This might seem surprising at first sight, as dance is a physical activity that mainly involves motor and SMS skills. It is important to note, however, that cognitive flexibility is a complex ability that involves different subdomains such as inhibition, working memory and switching (Dajani & Uddin, 2015). Recent studies showed that these capacities correlate with SMS skills in children (Puyjarinet et al., 2017; Tierney & Kraus, 2013; Bégel et al., under review). Therefore, we cannot rule out the possibility that the gain in cognitive flexibility is linked to the SMS training, even though some other factors such as the exercise of creativity through body improvisation might have acted in synergy. Indeed, cognitive flexibility is linked to creativity (De Dreu et al., 2011). Moreover, in a review on interventions to aid executive functions in children, Diamond & Lee (2011) concluded that addressing emotional, social and physical development was more efficient than focusing narrowly on executive functions only (Diamond & Lee, 2011). Therefore, with its social, physical, and creative components, dance appears as an ideal candidate to boost executive functions, and more specifically cognitive flexibility. Note that previous studies reported the effect of dance on cognitive flexibility in aging people (Coubard et al., 2011) and in pre-school children (Shen et al., 2020). This suggests that the effect of dance on cognitive flexibility is not limited to the current study's protocol and patients.

The improvement produced by the training extended to social skills. More specifically, parents reported enhancement of their children's social cognition and theory of mind, such as the capacities to express their own emotions, to adapt their behavior to others in a social context, and to infer on other's mental states. The collective training protocol

proposed in this study may have contributed to develop children's social skills by stimulating their capacity to anticipate others' movement. Motor synchronization between individuals leads to increased pro-social behaviour (Hove & Risen, 2009; Cirelli, Einarson & Trainor, 2014; Wiltermuth & Heath, 2009). In addition, interpersonal synchronization involved in dance increases the sense of connectedness with partners (Himberg, Laroche, Bigé, Buchkowski, & Bachrach, 2018). Dance also fosters the allocation of attention to others' cognitive and mental states (Joufflineau et al., 2018), and dance training is linked to enhanced empathic abilities (Gujing et al., 2019). Notably, cognitive flexibility is important for social behavior, especially in the ability to shape behavior to the social context (Turner, Oakes, Haslam, & McGarty, 1994). Cognitive flexibility is also correlated with measures of empathy (Grattan, & Eslinger, 1989; Milders, Ietswaart, Crawford, & Currie, 2008). In contrast, other cognitive skills, such as memory, processing speed, or attention, play a limited role in social behaviour (Milders, Ietswaart, Crawford, & Currie, 2008). Therefore, the concurrent improvement in synchronization, social skills and cognitive flexibility is not surprising. However, the causal relations between these competences are still to be explored. The effect of a dance training protocol on these capacities in typically developing children was not tested in this study and remains unexplored. Provasi et al. (2014) showed that a 30-minute practice of SMS reduced tapping variability in children with acquired lesions of the cerebellum but not in controls. Testing whether a dance training protocol also impacts differently children with DCA and controls will be important to further understand the mechanisms that sustain the effect of dancing.

The functions that were enhanced after the dance training are crucial for everyday life activities. This transfer effect may in turn significantly improve patients' quality of life. Notably, the effect found on a behavioral test of cognitive flexibility is corroborated by the parents' evaluation, meaning that the effect in everyday life was noticed by the parents and reflects a functional improvement. Cognitive flexibility is of great importance for the executive adaptation of cognitive and behavioral responses to the environment. It is of particular importance for different skills such as mathematics (Verschaffel, Torbeyns, De Smedt, Luwel, & Van Dooren, 2007), language (Deák, 2004) and planification (Snow, 1992). Enhancing cognitive flexibility abilities would improve patients' quality of life and boost their academic performance. In general, executive functions are a good predictor of school achievement (Becker, Miao, Duncan, & McClelland, 2014). In addition, social skills are also a good predictor of school achievement (Agostin & Bain, 1997; Bramlett, Scott, & Rowell, 2000). The fact that movement and balance skills were also

improved further indicates that patients' everyday life can be improved by the dance training protocol on several dimensions.

Because of their difficulties, children with DCA cannot take part in most extracurricular physical activities with children of their age. This in turn reduces their chances to progress. Thus, in addition to the lack of specific treatment or training available for DCA (Brenna, 2014), the range of activities for these children to participate in is limited. Some types of dance can be too challenging for them, but the results of this study confirm that dance can be proposed to DCA patients. The dance training protocol we proposed is promising because it is adapted to each child's skills. Children with various disorders and symptoms can participate in the same dance class. Besides, there is no competition and no judgment on the aesthetic production of the children, which helps to set trustful environment. Dance is also an activity that can be tailored to participants' special needs. Training protocols can therefore be adapted to different populations, such as adolescents or adults with cerebellar anomalies, using various types of dance.

In summary, very encouraging sensorimotor, motor, cognitive and social improvements were exhibited by children with DCA as a consequence of a dance training protocol. The cerebellum is a key structure of the central nervous system and its role in various motor (Bastian, 2006; Manto et al., 2012; Therrien & Bastian, 2019), cognitive (Murdoch, 2010; Schmahmann, 2004; Schmahmann, Guell, Stoodley, & Halko, 2019) as well as social (Van Overwalle et al., 2019) and affective functions is well known. Consequently, some or all of these domains are affected in patients with DCA. Finding ways to enhance patients with DCA's motor, cognitive and social skills is a challenge that we addressed using dance in this study. Dance is an activity that stimulates different aspects of behavior, especially sensorimotor functions. The beneficial effect of dance is likely to be partly due to increased activity in the sensorimotor cortico-cerebello-striatal pathways. This research paves the way for larger scale studies on dance as a tool for reeducation in cerebellar dysfunctions.

### **Acknowledgment**

We would like to thank the clinicians who conducted patients' evaluations: Dr. Marie-Pierre Lemaître, Camille Fallot and Yohan Bourez. We also thank Véronique Brunel, the dance teacher who created and conducted the dance program. Finally, we thank Séverine Fally for language editing.

### **Funding**

The research was supported by the “Fondation Maladies Rares” (“Temdance” grant).

### **Conflict of interest**

The authors have no relevant financial or non-financial interests to disclose

### **References**

- Ackermann, H. (2008). Cerebellar contributions to speech production and speech perception: psycholinguistic and neurobiological perspectives. *Trends in Neurosciences*, *31*(6), 265–272.  
<https://doi.org/10.1016/j.tins.2008.02.011>
- Agostin, T. M. K., & Bain, S. K. (1997). Predicting early school success with developmental and social skills screeners. *Psychology in the Schools*, *34*(3), 219–228. [https://doi.org/10.1002/\(SICI\)1520-6807\(199707\)34:3<219::AID-PITS4>3.0.CO;2-J](https://doi.org/10.1002/(SICI)1520-6807(199707)34:3<219::AID-PITS4>3.0.CO;2-J)
- Bachrach, A., Jola, C., & Pallier, C. (2016). Neuronal bases of structural coherence in contemporary dance observation. *NeuroImage*, *124*, 464–472. <https://doi.org/10.1016/j.neuroimage.2015.08.072>
- Bareš, M., Apps, R., Avanzino, L., Breska, A., D’Angelo, E., Filip, P., ... Petter, E. A. (2018). Consensus paper: Decoding the Contributions of the Cerebellum as a Time Machine. From Neurons to Clinical Applications. *Cerebellum*, *18*(2), 266–286. <https://doi.org/10.1007/s12311-018-0979-5>
- Bastian, A. J. (2006). Learning to predict the future: the cerebellum adapts feedforward movement control. *Current Opinion in Neurobiology*, *16*(6), 645–649. <https://doi.org/10.1016/j.conb.2006.08.016>
- Becker, D. R., Miao, A., Duncan, R., & McClelland, M. M. (2014). Behavioral self-regulation and executive function both predict visuomotor skills and early academic achievement. *Early Childhood Research Quarterly*.  
<https://doi.org/10.1016/j.ecresq.2014.04.014>
- Bégel, V., Verga, L., Benoit, C.-E., Kotz, S. A., & Dalla Bella, S. (2018). Test-retest reliability of the Battery for the Assessment of Auditory Sensorimotor and Timing Abilities (BAASTA). *Annals of Physical and Rehabilitation Medicine*, *61*(6), 395–400. <https://doi.org/10.1016/j.rehab.2018.04.001>
- Bengtsson, S. L., Ehrsson, H. H., Forssberg, H., & Ullén, F. (2005). Effector-independent voluntary timing:

behavioural and neuroimaging evidence. *The European Journal of Neuroscience*, 22(12), 3255–3265.

<https://doi.org/10.1111/j.1460-9568.2005.04517.x>

Benoit, C.-E., Dalla Bella, S., Farrugia, N., Obrig, H., Mainka, S., & Kotz, S. A. (2014). Musically cued gait-training improves both perceptual and motor timing in Parkinson's disease. *Frontiers in Human Neuroscience*, 8, 494.

<https://doi.org/10.3389/fnhum.2014.00494>

Bläsing, B., Calvo-Merino, B., Cross, E. S., Jola, C., Honisch, J., & Stevens, C. J. (2012). Neurocognitive control in dance perception and performance. *Acta Psychologica*, 139(2), 300–308.

<https://doi.org/10.1016/j.actpsy.2011.12.005>

Bledsoe, J. C., Semrud-Clikeman, M., & Pliszka, S. R. (2011). Neuroanatomical and Neuropsychological Correlates of the Cerebellum in Children with Attention-Deficit/Hyperactivity Disorder – Combined Type. *J Am Acad Child Adolesc Psychiatry*, 50(6), 593–601.

<https://doi.org/10.1016/j.jaac.2011.02.014>. Neuroanatomical

Bodranghien, F., Bastian, A., Casali, C., Hallett, M., Louis, E. D., Manto, M., ... van Dun, K. (2016). Consensus Paper: Revisiting the Symptoms and Signs of Cerebellar Syndrome. *Cerebellum*, 15(3), 369–391.

<https://doi.org/10.1007/s12311-015-0687-3>

Bolduc, M.-E., & Limperopoulos, C. (2009). Neurodevelopmental outcomes in children with cerebellar malformations : a systematic review. *Developmental Medicine & Child Neurology*, 51(4), 256–267.

<https://doi.org/10.1111/j.1469-8749.2008.03224.x>

Boltshauser, E. (2004). Cerebellum - Small Brain but Large Confusion: A Review of Selected Cerebellar Malformations and Disruptions. *American Journal of Medical Genetics*, 126(4), 376–385.

<https://doi.org/https://doi.org/10.1002/ajmg.a.20662>

Braitenberg, V., Heck, D., & Sultan, F. (1997). The detection and generation of sequences as a key to cerebellar function: Experiments and theory. *Behavioral and Brain Sciences*, 20(2), 229–277.

<https://doi.org/10.1017/S0140525X9700143X>

Bramlett, R. K., Scott, P., & Rowell, R. K. (2000). A Comparison of Temperament and Social Skills in Predicting Academic Performance in First Graders. *Special Services in the Schools*, 16(1–2), 147–158.

<https://doi.org/10.1300/J008v16n01>

Breska, A., & Ivry, R. B. (2016). Taxonomies of timing: Where does the cerebellum fit in? *Current Opinion in Behavioral Sciences*, 8, 282–288. <https://doi.org/10.1016/j.cobeha.2016.02.034>

Breska, A., & Ivry, R. B. (2018). Double dissociation of single-interval and rhythmic temporal prediction in cerebellar degeneration and Parkinson's disease. *Proceedings of the National Academy of Sciences*, 115(48), 12283–12288. <https://doi.org/10.1073/pnas.1810596115>

Brenna, V. (2014). Patients with cerebellar pathology: Rehabilitation of cognitive functions. In *Paediatric Neurological Disorders With Cerebellar Involvement—Diagnosis and Management* (John Libbey Eurotext, Vol. 27, pp. 221–234). D'arigo, S; Riva, D; Valente, EM.

Broersen, R., Onuki, Y., Abdelgabar, A. R., Owens, C. B., Picard, S., Willems, J., ... De Zeeuw, C. I. (2016). Impaired spatio-temporal predictive motor timing associated with spinocerebellar ataxia type 6. *PLoS ONE*, 11(8), 1–20. <https://doi.org/10.1371/journal.pone.0162042>

Brooks, B. L., Sherman, E. M. S., & Strauss, E. (2010). NEPSY-II: A Developmental neuropsychological assessment, second edition. *Child Neuropsychology*. <https://doi.org/10.1080/09297040903146966>

Brown, S., Martinez, M. J., & Parsons, L. M. (2006). The neural basis of human dance. *Cerebral Cortex*, 16(8), 1157–1167. <https://doi.org/10.1093/cercor/bhj057>

Chauvigné, L. A. S., Walton, A., Richardson, M. J., & Brown, S. (2019). Human Movement Science Multi-person and multisensory synchronization during group dancing. *Human Movement Science*, 63(May 2018), 199–208. <https://doi.org/10.1016/j.humov.2018.12.005>

Chuang, L. Y., Hung, H. Y., Huang, C. J., Chang, Y. K., & Hung, T. M. (2015). A 3-month intervention of Dance Revolution improves interference control in elderly females: a preliminary investigation. *Experimental Brain Research*, 233(4), 1181–1188. <https://doi.org/10.1007/s00221-015-4196-x>

Cochen De Cock, V., Dotov, D. G., Ihalainen, P., Bégel, V., Galtier, F., Lebrun, C., ... Dalla Bella, S. (2018). Rhythmic abilities and musical training in Parkinson's disease: do they help? *Npj Parkinson's Disease*, 4(1), 8.

<https://doi.org/10.1038/s41531-018-0043-7>

Coubard, O. A., Duretz, S., Lefebvre, V., Lapalus, P., & Ferrufino, L. (2011). Practice of contemporary dance improves cognitive flexibility in aging. *Frontiers in Aging Neuroscience*, *3*(SEP), 1–12.

<https://doi.org/10.3389/fnagi.2011.00013>

Coull, J., & Nobre, A. (2008). Dissociating explicit timing from temporal expectation with fMRI. *Current Opinion in Neurobiology*, *18*(2), 137–144. <https://doi.org/10.1016/j.conb.2008.07.011>

Coull, J. T., Cheng, R.-K., & Meck, W. H. (2011). Neuroanatomical and neurochemical substrates of timing. *Neuropsychopharmacology*, *36*(1), 3–25. <https://doi.org/10.1038/npp.2010.113>

Crawford, J. R., & Garthwaite, P. H. (2007). Comparison of a single case to a control or normative sample in neuropsychology: Development of a Bayesian approach. *Cognitive Neuropsychology*, *24*(4), 343–372.

<https://doi.org/10.1080/02643290701290146>

Cross, E. S., Hamilton, A. F. d. C., & Grafton, S. T. (2006). Building a motor simulation de novo: Observation of dance by dancers. *NeuroImage*, *31*(3), 1257–1267. <https://doi.org/10.1016/j.neuroimage.2006.01.033>

Cross, E. S., Kraemer, D. J. M., Hamilton, A. F. D. C., Kelley, W. M., & Grafton, S. T. (2009). Sensitivity of the action observation network to physical and observational learning. *Cerebral Cortex*, *19*(2), 315–326.

<https://doi.org/10.1093/cercor/bhn083>

Cirelli, L. K., Einarson, K. M., & Trainor, L. J. (2014). Interpersonal synchrony increases prosocial behavior in infants. *Developmental science*, *17*(6), 1003–1011. <https://doi.org/10.1111/desc.12193>

Dajani, D. R., & Uddin, L. Q. (2015). Demystifying cognitive flexibility: Implications for clinical and developmental neuroscience. *Trends in Neurosciences*, *38*(9), 571–578. <https://doi.org/10.1016/j.tins.2015.07.003>

Dalla Bella, S., Farrugia, N., Benoit, C. E., Begel, V., Verga, L., Harding, E., & Kotz, S. A. (2017). BAASTA: Battery for the Assessment of Auditory Sensorimotor and Timing Abilities. *Behavior Research Methods*, *49*(3), 1128–

1145. <https://doi.org/10.3758/s13428-016-0773-6>

Damm, L., Varoqui, D., Cochen De Cock, V., Dalla Bella, S., & Bardy, B. (2019). Why do we move to the beat? A



multi-scale approach, from physical principles to brain dynamics. *Neuroscience & Biobehavioral Reviews*, (December). <https://doi.org/10.1016/j.neubiorev.2019.12.024>

de Natale, E. R., Paulus, K. S., Aiello, E., Sanna, B., Manca, A., Sotgiu, G., ... Deriu, F. (2017). Dance therapy improves motor and cognitive functions in patients with Parkinson's disease. *NeuroRehabilitation*, *40*(1), 141–144. <https://doi.org/10.3233/NRE-161399>

Dreu, C. K. D., Nijstad, B. A., & Baas, M. (2011). Behavioral activation links to creativity because of increased cognitive flexibility. *Social Psychological and Personality Science*, *2*(1), 72-80. <https://doi.org/10.1177/1948550610381789>

De Smet, H. J., Paquier, P., Verhoeven, J., & Mariën, P. (2013). The cerebellum: Its role in language and related cognitive and affective functions. *Brain and Language*, *127*(3), 334–342. <https://doi.org/10.1016/j.bandl.2012.11.001>

Deák, G. O. (2004). The Development of Cognitive Flexibility and Language Abilities. *Advances in Child Development and Behavior*. [https://doi.org/10.1016/S0065-2407\(03\)31007-9](https://doi.org/10.1016/S0065-2407(03)31007-9)

Diamond, A., & Lee, K. (2011). Interventions shown to aid executive function development in children 4 to 12 years old. *Science*, *333*(6045), 959-964. <https://doi.org/10.1126/science.1204529>

dos Santos Delabary, M., Komerowski, I. G., Monteiro, E. P., Costa, R. R., & Haas, A. N. (2018). Effects of dance practice on functional mobility, motor symptoms and quality of life in people with Parkinson's disease: a systematic review with meta-analysis. *Aging Clinical and Experimental Research*, *30*(7), 727–735. <https://doi.org/10.1007/s40520-017-0836-2>

Doyon, J., Penhune, V., & Ungerleider, L. G. (2003). Distinct contribution of the cortico-striatal and cortico-cerebellar systems to motor skill learning. *Neuropsychologia*, *41*(3), 252–262. [https://doi.org/https://doi.org/10.1016/S0028-3932\(02\)00158-6](https://doi.org/https://doi.org/10.1016/S0028-3932(02)00158-6)

Drake, C., & Botte, M. C. (1993). Tempo sensitivity in auditory sequences: Evidence for a multiple-look model. *Perception & Psychophysics*, *54*(3), 277–286. <https://doi.org/10.3758/BF03205262>

- Ebner, T. J., Hewitt, A. L., & Popa, L. S. (2011). What features of limb movements are encoded in the discharge of cerebellar neurons? *Cerebellum*, *10*(4), 683–693. <https://doi.org/10.1007/s12311-010-0243-0>
- Fogel, B. L. (2012). Childhood Cerebellar Ataxia. *Journal of Child Neurology*, *27*(9), 1138–1145. <https://doi.org/https://doi.org/10.1177/0883073812448231>
- Giacosa, C., Karpati, F. J., Foster, N. E. V., Penhune, V. B., & Hyde, K. L. (2016). Dance and music training have different effects on white matter diffusivity in sensorimotor pathways. *NeuroImage*, *135*, 273–286. <https://doi.org/10.1016/j.neuroimage.2016.04.048>
- Gioia, G. a, Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). Behavior rating inventory of executive function. *Child Neuropsychology*, *6*(3), 235–238. <https://doi.org/10.1076/chin.6.3.235.3152>
- Grahn, J. A. (2009). The role of the basal ganglia in beat perception: Neuroimaging and neuropsychological investigations. *Annals of the New York Academy of Sciences*, *1169*, 35–45. <https://doi.org/10.1111/j.1749-6632.2009.04553.x>
- Grahn, J. A., & Brett, M. (2007). Rhythm and beat perception in motor areas of the brain. *Journal of Cognitive Neuroscience*, *19*(5), 893–906. <https://doi.org/10.1162/jocn.2007.19.5.893>
- Grahn, J. A., & Brett, M. (2009). Impairment of beat-based rhythm discrimination in Parkinson’s disease. *Cortex*, *45*(1), 54–61. <https://doi.org/10.1016/j.cortex.2008.01.005>
- Grattan, L. M., & Eslinger, P. J. (1989). Higher cognition and social behavior: Changes in cognitive flexibility and empathy after cerebral lesions. *Neuropsychology*, *3*(3), 175. <https://doi.org/10.1037/h0091764>
- Grönlund, E., Renck, B., & Weibull, J. (2005). Dance/movement therapy as an alternative treatment for young boys diagnosed as ADHD: A pilot study. *American Journal of Dance Therapy*, *27*(2), 63–85. <https://doi.org/10.1007/s10465-005-9000-1>
- Grube, M., Cooper, F. E., Chinnery, P. F., & Griffiths, T. D. (2010). Dissociation of duration-based and beat-based auditory timing in cerebellar degeneration. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(25), 11597–11601. <https://doi.org/10.1073/pnas.0910473107>

- Guell, X., Hoche, F., & Schmahmann, J. D. (2015). Metalinguistic Deficits in Patients with Cerebellar Dysfunction: Empirical Support for the Dysmetria of Thought Theory. *Cerebellum*, *14*(1), 50–58.  
<https://doi.org/10.1007/s12311-014-0630-z>
- Hackney, M. E., Byers, C., Butler, G., Sweeney, M., Rossbach, L., & Bozzorg, A. (2015). Adapted tango improves mobility, motor-cognitive function, and gait but not cognition in older adults in independent living. *Journal of the American Geriatrics Society*, *63*(10), 2105–2113. <https://doi.org/10.1111/jgs.13650>
- Hackney, M. E., & Gammon, E. M. (2011). Comparison of Partnered and Non-Partnered Dance Movement. *Neurorehabilitation and Neural Repair*, *24*(4), 384–392. <https://doi.org/10.1177/1545968309353329>.Effects
- Hackney, M. E., Kantorovich, S., Levin, R., & Earhart, G. M. (2007). Effects of tango on functional mobility in Parkinson's disease: a preliminary study. *Journal of Neurologic Physical Therapy*, *31*(4), 173–179.  
<https://doi.org/10.1097/NPT.0b013e31815ce78b>
- Himberg, T., Laroche, J., Bigé, R., Buchkowski, M., & Bachrach, A. (2018). Coordinated Interpersonal Behaviour in Collective Dance Improvisation: The Aesthetics of Kinaesthetic Togetherness. *Behavioral Sciences*, *8*(2), 23.  
<https://doi.org/10.3390/bs8020023>
- Himle, M. B., Miltenberger, R. G., Flessner, C., & Gatheridge, B. (2004). Teaching Safety Skills To Children To Prevent Gun Play. *Journal of Applied Behavior Analysis*, *37*(1), 1–9. <https://doi.org/10.1901/jaba.2004.37-1>
- Hoche, F., Guell, X., Sherman, J. C., Vangel, M. G., & Schmahmann, J. D. (2016). The cerebellar contribution to social cognition. *The Cerebellum*, *15*(6), S23. <https://doi.org/10.1002/ana.24247>
- Hove, M. J., & Risen, J. L. (2009). It's all in the timing: Interpersonal synchrony increases affiliation. *Social cognition*, *27*(6), 949-960. <https://doi.org/10.1521/soco.2009.27.6.949>
- Iversen, J. R., & Patel, A. D. (2008). The Beat Alignment Test (BAT): Surveying beat processing abilities in the general population. In K. Miyazaki, Y. Hiraga, M. Adachi, Y. Nakajima, & M. Tsuzaki (Eds.), *Proceedings of the 10th International Conference on Music Perception & Cognition (ICMPC10)* (pp. 465–468). Adelaide: Causal Productions.

- Ivry, R B, & Keele, S. W. (1989). Timing functions of the cerebellum. *Journal of Cognitive Neuroscience*, 1(2), 136–152. <https://doi.org/10.1162/jocn.1989.1.2.136>
- Ivry, Richard B, Spencer, R. M. C., Zelaznik, H. N., & Diedrichsen, J. (2002). The cerebellum and event timing. *Annals of the New York Academy of Sciences*, 978(1), 302–317. <https://doi.org/https://doi.org/10.1111/j.1749-6632.2002.tb07576.x>
- Jeong, Y. J., Hong, S. C., Myeong, S. L., Park, M. C., Kim, Y. K., & Suh, C. M. (2005). Dance movement therapy improves emotional responses and modulates neurohormones in adolescents with mild depression. *International Journal of Neuroscience*, 115(12), 1711–1720. <https://doi.org/10.1080/00207450590958574>
- Jin, X., Wang, B., Lv, Y., Lu, Y., Chen, J., & Zhou, C. (2019). Does dance training influence beat sensorimotor synchronization? Differences in finger-tapping sensorimotor synchronization between competitive ballroom dancers and nondancers. *Experimental Brain Research*, 237(3), 743–753. <https://doi.org/10.1007/s00221-018-5410-4>
- Jola, C., Davis, A., & Haggard, P. (2011). Proprioceptive integration and body representation: Insights into dancers' expertise. *Experimental Brain Research*, 213(2–3), 257–265. <https://doi.org/10.1007/s00221-011-2743-7>
- Jones, C. R. G., & Jahanshahi, M. (2014). Motor and perceptual timing in Parkinson's disease. *Advances in Experimental Medicine and Biology*, 829, 265–290. [https://doi.org/10.1007/978-1-4939-1782-2\\_14](https://doi.org/10.1007/978-1-4939-1782-2_14)
- Joufflineau, C., Vincent, C., & Bachrach, A. (2018). Synchronization, Attention and Transformation: Multidimensional Exploration of the Aesthetic Experience of Contemporary Dance Spectators. *Behavioral Sciences*, 8(2), 24. <https://doi.org/10.3390/BS8020024>
- Karpati, F. J., Giacosa, C., Foster, N. E. V, Penhune, V. B., & Hyde, K. L. (2015). Dance and the brain: A review. *Annals of the New York Academy of Sciences*, 1337(1), 140–146. <https://doi.org/10.1111/nyas.12632>
- Karpati, F. J., Giacosa, C., Foster, N. E. V, Penhune, V. B., & Hyde, K. L. (2016). Sensorimotor integration is enhanced in dancers and musicians. *Experimental Brain Research*, 234(3), 893–903. <https://doi.org/10.1007/s00221-015-4524-1>

- Kotz, S. A., & Schwartz, M. (2011). Differential input of the supplementary motor area to a dedicated temporal processing network: functional and clinical implications. *Frontiers in Integrative Neuroscience*, *5*(86), 86. <https://doi.org/10.3389/fnint.2011.00086>
- Koziol, L. F., Budding, D., Andreasen, N., D'Arrigo, S., Bulgheroni, S., Imamizu, H., ... Yamazaki, T. (2014). Consensus paper: The cerebellum's role in movement and cognition. *Cerebellum*, *13*(1), 151–177. <https://doi.org/10.1007/s12311-013-0511-x>
- Laland, K., Wilkins, C., & Clayton, N. (2016). The evolution of dance. *Current Biology*, *26*(1), R5–R9. <https://doi.org/10.1016/j.cub.2015.11.031>
- Lang, E. J., Apps, R., Bengtsson, F., Cerminara, N. L., De Zeeuw, C. I., Ebner, T. J., ... Xiao, J. (2017). The Roles of the Olivocerebellar Pathway in Motor Learning and Motor Control. A Consensus Paper. *Cerebellum*, *16*(1), 230–252. <https://doi.org/10.1007/s12311-016-0787-8>
- London, J. (2012). *Hearing in time: psychological aspects of musical meter* (Oxford Uni). <https://doi.org/10.1093/acprof:oso/9780199744374.001.0001>
- López-Ortiz, C., Egan, T., & Gaebler-Spira, D. J. (2016). Pilot study of a targeted dance class for physical rehabilitation in children with cerebral palsy. *SAGE Open Medicine*, *4*, 205031211667092. <https://doi.org/10.1177/2050312116670926>
- López-Ortiz, C., Gladden, K., Deon, L., Schmidt, J., Girolami, G., & Gaebler-Spira, D. (2012). Dance program for physical rehabilitation and participation in children with cerebral palsy. *Arts and Health*, *4*(1), 39–54. <https://doi.org/10.1080/17533015.2011.564193>
- Manly, T., Anderson, V., Nimmo-Smith, I., Turner, A., Watson, P., & Robertson, I. H. (2001). The Differential Assessment of Children's Attention: The Test of Everyday Attention for Children (TEA-Ch), Normative Sample and ADHD Performance. *Journal of Child Psychology and Psychiatry*, *42*(8), 1065–1081. <https://doi.org/10.1111/1469-7610.00806>
- Manto, M., Bower, J. M., Conforto, A. B., Delgado-García, J. M., Da Guarda, S. N. F., Gerwig, M., ... Timmann, D. (2012). Consensus paper: Roles of the cerebellum in motor control-the diversity of ideas on cerebellar

involvement in movement. *Cerebellum*, 11(2), 457–487. <https://doi.org/10.1007/s12311-011-0331-9>

Marsden, J. F. (2018). Cerebellar ataxia. In B. L. Day & S. R. Lord (Eds.), *Handbook of Clinical Neurology* (pp. 261–281). <https://doi.org/10.1016/B978-0-444-63916-5.00017-3>

Marsden, J. F., & Harris, C. (2011). Cerebellar ataxia: Pathophysiology and rehabilitation. *Clinical Rehabilitation*, 25(3), 195–216. <https://doi.org/10.1177/0269215510382495>

Martin, C. L., Tan, D., Bragge, P., & Bialocerkowski, A. (2009). Systematic review of the effectiveness of physiotherapy for cerebellar dysfunction. *Clinical Rehabilitation*, 23(8), 15–26. <https://doi.org/10.1177/0269215509337272>

McKee, K. E., & Hackney, M. E. (2013). The effects of adapted tango on spatial cognition and disease severity in Parkinson's disease. *Journal of Motor Behavior*, 45(6), 519–529. <https://doi.org/10.1080/00222895.2013.834288>

Mogadam, A., Keller, A. E., Taylor, M. J., Lerch, J. P., Anagnostou, E., & Pang, E. W. (2018). Mental flexibility: An MEG investigation in typically developing children. *Brain and Cognition*, 120, 58–66. <https://doi.org/10.1016/j.bandc.2017.10.001>

Molinari, M., Chiricozzi, F. R., Clausi, S., Tedesco, A. M., Lisa, M. De, & Leggio, M. G. (2008). Cerebellum and Detection of Sequences, from Perception to Cognition. *Cerebellum*, 7, 611–615. <https://doi.org/10.1007/s12311-008-0060-x>

Molinari, M., Leggio, M. G., De Martin, M., Cerasa, A., & Thaut, M. (2003). Neurobiology of rhythmic motor entrainment. *Annals of the New York Academy of Sciences*, 999(1), 313–321. <https://doi.org/10.1196/annals.1284.042>

Molinari, M., Leggio, M. G., & Thaut, M. H. (2007). The cerebellum and neural networks for rhythmic sensorimotor synchronization in the human brain. *The Cerebellum*, 6(1), 18–23. doi: 10.1080/14734220601142886.

Morgan, D. L., & Morgan, R. K. (2008). *Single-case research methods for the behavioral and health sciences*. (SAGE publi). <https://doi.org/http://dx.doi.org/10.4135/9781483329697>

Milders, M., Ietswaart, M., Crawford, J. R., & Currie, D. (2008). Social behavior following traumatic brain injury and

its association with emotion recognition, understanding of intentions, and cognitive flexibility. *Journal of the International Neuropsychological Society*, 14(2), 318-326. <https://doi.org/10.1017/S1355617708080351>

Murdoch, B. E. (2010). The cerebellum and language: historical perspective and review. *Cortex*, 46(7), 858-868. <https://doi.org/10.1016/j.cortex.2009.07.018>

Musselman, K. E., Stoyanov, C. T., Marasigan, R., Jenkins, M. E., Konczak, J., Morton, S. M., & Bastian, A. J. (2014). Prevalence of ataxia in children: A systematic review. *Neurology*, 82(1), 80–89. <https://doi.org/10.1212/01.wnl.0000438224.25600.6c>

Nowak, D. A., Topka, H., Timmann, D., Boecker, H., & Hermsdörfer, J. (2007). The role of the cerebellum for predictive control of grasping. *Cerebellum*, 6(1), 7–17. <https://doi.org/10.1080/14734220600776379>

Nozaradan, S., Schwartz, M., Obermeier, C., & Kotz, S. A. (2017). Specific contributions of basal ganglia and cerebellum to the neural tracking of rhythm. *Cortex*, 95, 156–168. <https://doi.org/10.1016/j.cortex.2017.08.015>

Paquette, S., Fujii, S., Li, H. C., & Schlaug, G. (2017). The cerebellum's contribution to beat interval discrimination. *NeuroImage*, 163, 177–182. <https://doi.org/10.1016/j.neuroimage.2017.09.017>

Poretti, A., Boltshauser, E., & Doherty, D. (2014). Cerebellar hypoplasia: Differential diagnosis and diagnostic approach. *American Journal of Medical Genetics, Part C: Seminars in Medical Genetics*, 166(2), 211–226. <https://doi.org/10.1002/ajmg.c.31398>

Provasi, J., Doyère, V., Zélanti, P. S., Kieffer, V., Perdry, H., El Massioui, N., ... Droit-Volet, S. (2014). Disrupted sensorimotor synchronization, but intact rhythm discrimination, in children treated for a cerebellar medulloblastoma. *Research in Developmental Disabilities*, 35(9), 2053–2068. <https://doi.org/10.1016/j.ridd.2014.04.024>

Puyjarinet, F., Bégel, V., Lopez, R., Dellacherie, D., & Dalla Bella, S. (2017). Children and adults with Attention-Deficit/Hyperactivity Disorder cannot move to the beat. *Scientific Reports*, 7(1), 11550. <https://doi.org/10.1038/s41598-017-11295-w>

Repp, B. H. (2005). Sensorimotor synchronization: a review of the tapping literature. *Psychonomic Bulletin & Review*,

12(6), 969–992. <https://doi.org/https://doi.org/10.3758/BF03206433>

Repp, B. H. (2006). Rate Limits of Sensorimotor Synchronization. *Advances in Cognitive Psychology*, 2(2), 163–181.

<https://doi.org/10.2478/v10053-008-0053-9>

Repp, B. H., & Su, Y.-H. (2013). Sensorimotor synchronization: a review of recent research (2006-2012).

*Psychonomic Bulletin & Review*, 20(3), 403–452. <https://doi.org/10.3758/s13423-012-0371-2>

Ryan, M. M., & Engle, E. C. (2003). Acute Ataxia in Childhood. *Journal of Child Neurology*, 18, 309–316.

<https://doi.org/https://doi.org/10.1177/08830738030180050901>

Sampaio, L. M. M., Subramaniam, S., Arena, R., & Bhatt, T. (2016). Does virtual reality-based kinect dance training

paradigm improve autonomic nervous system modulation in individuals with chronic stroke? *Journal of*

*Vascular and Interventional Neurology*, 9(2), 21–29.

Sartor-glittenberg, C., & Brickner, L. (2014). A multidimensional physical therapy program for individuals with

cerebellar ataxia secondary to traumatic brain injury : a case series. *Physiotherapy Theory and Practice*, 30(2),

138–148. <https://doi.org/10.3109/09593985.2013.819952>

Schmahmann, J. D. (2004). Disorders of the Cerebellum. *The Journal of Neuropsychiatry and Clinical Neurosciences*,

16, 367–378. <https://doi.org/10.1212/wnl.32.7.790>

Schmahmann, J. D., Guell, X., Stoodley, C. J., & Halko, M. A. (2019). The theory and neuroscience of cerebellar

cognition. *Annual Review Of Neuroscience*, 42, 337–364. <https://doi.org/10.1146/annurev-neuro-070918-050258>

Schmahmann, J. D., & Sherman, J. C. (1998). The cerebellar cognitive affective syndrome. *Brain*, 121(4), 561–579.

<https://doi.org/https://doi.org/10.1093/brain/121.4.561>

Schmitz-Hübsch, T., Du Montcel, S. T., Baliko, L., Berciano, J., Boesch, S., Depondt, C., ... Fancellu, R. (2006).

Scale for the assessment and rating of ataxia: Development of a new clinical scale. *Neurology*, 66(11), 1717–

1720. <https://doi.org/10.1212/01.wnl.0000219042.60538.92>

Schwartz, M., Keller, P. E., & Kotz, S. A. (2016). Spontaneous, synchronized, and corrective timing behavior in

cerebellar lesion patients. *Behavioural Brain Research*, 312, 285–293. <https://doi.org/10.1016/j.bbr.2016.06.040>



- Schwartz, M., & Kotz, S. A. (2013). A dual-pathway neural architecture for specific temporal prediction. *Neuroscience & Biobehavioral Reviews*, *37*(10), 2587-2596.  
<https://doi.org/10.1016/j.neubiorev.2013.08.005>
- Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error Correction , Sensory Prediction , and Adaptation in Motor Control. *Annual Review of Neuroscience*, *33*, 89–108. <https://doi.org/10.1146/annurev-neuro-060909-153135>
- Shanahan, J., Morris, M. E., Bhriain, O. N., Saunders, J., & Clifford, A. M. (2015). Dance for people with Parkinson disease: What is the evidence telling us? *Archives of Physical Medicine and Rehabilitation*, *96*(1), 141–153.  
<https://doi.org/10.1016/j.apmr.2014.08.017>
- Shen, Y., Zhao, Q., Huang, Y., Liu, G., & Fang, L. (2020). Promotion of Street-Dance Training on the Executive Function in Preschool Children. *Frontiers in Psychology*, *11*. <https://doi.org/10.3389/fpsyg.2020.585598>
- Snow, J. H. (1992). Mental flexibility and planning skills in children and adolescents with learning disabilities. *Journal of Learning Disabilities*, *25*(4), 265–270. <https://doi.org/10.1177/002221949202500408>
- Song, Y. G., Ryu, Y. U., Im, S. J., Lee, Y. S., & Park, J. H. (2018). Effects of dance-based movement therapy on balance, gait, and psychological functions in severe cerebellar ataxia: A case study. *Physiotherapy Theory and Practice*, *35*(8), 756–763. <https://doi.org/10.1080/09593985.2018.1457119>
- Soppelsa, R., & Albaret, J. M. (2005). La batterie d'évaluation du mouvement chez l'enfant (M-ABC) : étalonnage sur une population d'enfants de 4 à 12 ans. *Psymotricité*, 195–200.
- Surgent, O. J., Dadalko, O. I., Pickett, K. A., & Travers, B. G. (2019). Balance and the brain: A review of structural brain correlates of postural balance and balance training in humans. *Gait and Posture*, *71*, 245–252.  
<https://doi.org/10.1016/j.gaitpost.2019.05.011>
- Teixeira-Machado, L., Azevedo-Santos, I., & De Santana, J. M. (2017). Dance improves functionality and psychosocial adjustment in cerebral palsy: A randomized controlled clinical trial. *American Journal of Physical Medicine and Rehabilitation*, *96*(6), 424–429. <https://doi.org/10.1097/PHM.0000000000000646>

- ten Donkelaar, H. J., Lammens, M., Wesseling, P., Thijssen, H. O. M., & Renier, W. O. (2003). Development and developmental disorders of the human cerebellum. *Journal of Neurology*, *250*(09), 1025–1036.  
<https://doi.org/10.1007/s00415-003-0199-9>
- Therrien, A. S., & Bastian, A. J. (2019). The cerebellum as a movement sensor. *Neuroscience Letters*, *688*, 37–40.  
<https://doi.org/10.1016/j.neulet.2018.06.055>
- Thier, P., & Markanday, A. (2019). Role of the Vermal Cerebellum in Visually Guided Eye Movements and Visual Motion Perception. *Annual Review of Vision Science*, *5*(1), 247–268. <https://doi.org/10.1146/annurev-vision-091718-015000>
- Tierney, A. T., & Kraus, N. (2013). The ability to tap to a beat relates to cognitive, linguistic, and perceptual skills. *Brain and Language*, *124*(3), 225–231. <https://doi.org/10.1016/j.bandl.2012.12.014>
- Timmann, D., Watts, S., & Hore, J. (1999). Failure of cerebellar patients to time finger opening precisely causes ball high-low inaccuracy in overarm throws. *Journal of Neurophysiology*, *82*(1), 103–114.  
<https://doi.org/10.1152/jn.1999.82.1.103>
- Turner, J. C., Oakes, P. J., Haslam, S. A., & McGarty, C. (1994). Self and collective: Cognition and social context. *Personality and social psychology bulletin*, *20*(5), 454-463. <https://doi.org/10.1177/0146167294205002>
- Van Overwalle, F., Baetens, K., Mariën, P., & Vandekerckhove, M. (2014). Social cognition and the cerebellum: a meta-analysis of over 350 fMRI studies. *Neuroimage*, *86*, 554-572.  
<https://doi.org/10.1016/j.neuroimage.2013.09.033>
- Van Overwalle, F., Manto, M., Leggio, M., & Delgado-García, J. M. (2019). The sequencing process generated by the cerebellum crucially contributes to social interactions. *Medical Hypotheses*, *128*, 33–42.  
<https://doi.org/10.1016/j.mehy.2019.05.014>
- Verschaffel, L., Torbeyns, J., De Smedt, B., Luwel, K., & Van Dooren, W. (2007). Strategy flexibility in children with low achievement in mathematics. *Educational and Child Psychology*, *24*(2), 16–27.
- Wang, V. Y., & Zoghbi, H. Y. (2001). Genetic regulation of cerebellar development. *Nature Reviews Neuroscience*,

2(7), 484–491. <https://doi.org/10.1038/35081558>

Wiltermuth, S. S., & Heath, C. (2009). Synchrony and cooperation. *Psychological science*, 20(1), 1-5.

<https://doi.org/10.1111/j.1467-9280.2008.02253.x>

Zatorre, R. J., Chen, J. L., & Penhune, V. B. (2007). When the brain plays music: auditory-motor interactions in music perception and production. *Nature Reviews. Neuroscience*, 8(7), 547–558. <https://doi.org/10.1038/nrn2152>

## Tables

**Table 1.** Demographics and Cerebellar signs at the latest neurological evaluation

Patient	Age	Genre	A	HT	CD	DM	DA	T	S	F	N	OA	OM	OBF
1	10.3	M	+	-	+	-	-	-	+	+	-	-	+	-
2	7.8	F	+	-	+	-	+	-	-	-	-	-	+	+
3	7.5	F	+	+	+	-	+	-	-	-	-	-	-	+
4	8.2	M	+	+	+	-	-	-	-	+	-	-	-	-
5	9.0	M	+	+	+	+	+	+	+	+	-	-	-	+
6	7.11	M	+	+	+	-	+	-	+	-	-	-	+	-
7	8.3	M	+	+	+	+	+	-	+	+	+	+	+	+

A = Ataxia, HT = Hypotonia, CD = Coordination disorder/Dysdiadochokinesis, DM = Dysmetria, DA = Dysarthria/Speech disorder (articulation), T = Intentional tremors, S = Slowness, F = Fatigability, N = Nystagmus, OA = Oculomotor apraxia, OM = Other Oculomotor disorders, OBF = Oro-bucco-Facial apraxia

**Table 2.** IQ scores of children with DCA. IQ indexes were calculated with the WISC-IV for patients 1, 4,5 and 6, and with the WPPSI-IV for patients 2, 3 and 7 in accordance with the age of the patient at last intellectual evaluation.

Patient	VCI	PRI	VSI	FRI	WMI	PSI
1	98	94	NT	NT	97	112
2	84	NT	106	118	106	74
3	95	NT	77	95	97	77
4	138	114	NT	NT	100	96
5	82	79	NT	NT	NT	76
6	89	NT	102	91	88	77
7	100	NT	85	95	NT	71

**VCI** = Verbal Comprehension Index, **PRI** = Perceptual Reasoning Index, **VSI** = Visual Spatial Index, **FRI** = Fluid Reasoning Index, **WMI** = Working Memory Index, **PSI** = Processing Speed Index; note that in the WISC IV, the main indices are VCI, PRI, WMI and PSI. In the WPPSI IV, the main indices are VCI, VSI, FRI, WMI and PSI. NT = not tested (the index does not exist in the battery used).

**Table 3. Differences between the pre and the post training and between CEREB and CONTROLS group. Tasks for which a significant difference were observed are presented**

Variable	CEREB Pre-Post Comparison				Comparison CEREB and CONTROLS					
	Mean Pre (SD)	Mean Post (SD)	Z	p	Effect Size (r)	Delta Patients (SD)	Delta Controls (SD)	Z	p	Effect Size (r)
Tapping with a metronome (CV)	0.35 (0.05)	0.19 (0.11)	2.37	< 0.01**	0.89	-0.16 (0.08)	0.03 (0.15)	3.32	< 0.001***	0.65
Tapping with music (CV)	0.41 (0.12)	0.22 (0.09)	2.37	< 0.01**	0.89	-0.19 (0.11)	0.00 (0.03)	2.86	< 0.005***	0.56
Hopping on mats	1.86 (1.38)	3.57 (2.09)	2.30	< 0.05*	0.87	1.71 (1.19)	-0.05 (0.28)	3.88	< 0.001***	0.76
Cognitive flexibility Speed	7.24 (2.87)	12.8 (10.3)	2.20	< 0.05*	0.90	-5.19 (8.47)	-1.08 (1.22)	1.84	< 0.05*	0.36
Cognitive flexibility Quality	4.67 (1.86)	3 (1.91)	2.27	< 0.05*	0.93	1.33 (0.52)	0.32 (0.75)	2.71	< 0.05*	0.46
Cognitive flexibility at home	16.57 (3.6)	13.86 (3.39)	1.87	< 0.05*	0.71	2.71 (3.30)	0.18 (1.13)	2.11	< 0.05*	0.47
Social cognition	59.43 (8.28)	55.14 (7.78)	1.86	< 0.05*	0.70	4.29 (4.89)	-0.47 (4.89)	2.71	< 0.01**	0.53

**Table 4. Differences between the first and second testing sessions CONTROLS group.**

Variable	Mean first session ( <i>SD</i> )	Mean second session ( <i>SD</i> )	Z	<i>p</i>
Tapping with a metronome (CV)	0.09 (0.08)	0.12 (0.12)	1.25	= 0.22
Tapping with music (CV)	0.14 (0.11)	0.13 (0.08)	0.20	= 0.86
Hopping on mats	4.95 (0.16)	4.89 (0.31)	-0.61	= 0.75
Cognitive flexibility Speed	5.50 (1.37)	4.42 (1.05)	-3.06	< 0.05*
Cognitive flexibility Quality	6.26 (0.73)	6.58 (0.61)	1.73	= 0.15
Cognitive flexibility at home	20.00 (2.83)	20.06 (3.07)	0.45	= 0.70
Social cognition	67.74 (5.90)	66.41 (4.20)	-1.48	= 0.22

**Table 5. Individual scores in Perceptual and motor rhythmic skills for each DCA patient in comparison with an age-matched control group. Statistics are calculated with Bayesian Single-Case Methods (Crawford & Garthwaite, 2007). Columns Pre show the comparison between the control group and patients' performance before the testing and columns Post show the comparison between the control**

**group and patients' performance after the testing.**

<i>Patient 1</i>						
Pre			Post			
Variable	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Tapping with a metronome (CV)	0.29	0.12 (0.14)	= 0.15	0.10	0.12 (0.13)	= 0.38
Tapping with music (CV)	0.42	0.15 (0.15)	= 0.07	0.13	0.14 (0.12)	= 0.47
<i>Patient 2</i>						
Pre			Post			
Variable	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Tapping with a metronome (CV)	0.33	0.01 (0.01)	< 0.005***	0.25	0.11 (0.07)	= 0.06
Tapping with music (CV)	0.43	0.12 (0.05)	< 0.005***	0.26	0.12 (0.04)	< 0.05*
<i>Patient 3</i>						
Pre			Post			
Variable	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Tapping with a metronome (CV)	0.42	0.07 (0.01)	< 0.005***	0.39	0.11 (0.07)	< 0.01**
Tapping with music (CV)	0.55	0.12 (0.05)	< 0.005***	0.34	0.12 (0.04)	< 0.005***
<i>Patient 4</i>						
Pre			Post			
Variable	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Tapping with a metronome (CV)	0.31	0.08 (0.02)	< 0.001***	0.09	0.13 (0.15)	= 0.41

Tapping with music (CV)	0.15	0.12 (0.07)	= .35	0.13	0.14 (0.06)	0.44
<i>Patient 5</i>						
	Pre			Post		
Variable	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Tapping with a metronome (CV)	0.31	0.07 (0.03)	< 0.001***	0.11	0.06 (0.01)	< 0.005***
Tapping with music (CV)	0.46	0.14 (0.09)	< 0.005***	0.13	0.09 (0.04)	0.19
<i>Patient 6</i>						
	Pre			Post		
Variable	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Tapping with a metronome (CV)	0.39	0.08 (0.02)	< 0.001***	0.26	0.13 (0.15)	= 0.23
Tapping with music (CV)	0.39	0.12 (0.07)	< 0.01*	0.25	0.14 (0.06)	= 0.07
<i>Patient 7</i>						
	Pre			Post		
Variable	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Tapping with a metronome (CV)	0.39	0.08 (0.02)	< 0.001***	0.13	0.13 (0.15)	0.5
Tapping with music (CV)	0.42	0.12 (0.07)	< 0.01**	0.31	0.14 (0.06)	0.023*

**Table 6. Individual scores in Motor and cognitive skills for each DCA patient in comparison with an age-matched control group. Statistics are calculated with Bayesian Single-Case Methods with the DissocsBayes program (Crawford & Garthwaite, 2007). Columns Pre show the comparison between the control group and patients' performance before the testing and columns Post show the comparison between the control group and patients' performance after the testing.**



*Patient 1*

Variable	Pre			Post		
	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Hopping on mats	4.5	5.00 (0.00)	< 0.001***	5	5.00 (0.00)	0.5
Cognitive flexibility Speed	5.19	5.31 (0.76)	= 0.44	4.5	3.96 (0.46)	0.15
Cognitive flexibility Quality	4	6.00 (1.00)	= 0.06	4	6.43 (0.79)	0.07
Cognitive flexibility at home	14	19.33 (2.25)	< 0.05*	13	20 (2.83)	< 0.05*
Social cognition	61	68.14 (4.77)	= 0.11	69	65.67 (3.27)	= 0.19

*Patient 2*

Variable	Pre			Post		
	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Hopping on mats	2	4.92 (0.20)	< 0.000***	3.5	4.67 (0.52)	< 0.05*
Cognitive flexibility Speed	9.43	6.24 (1.91)	= 0.09	6.78	5.31 (1.24)	= 0.16
Cognitive flexibility Quality	4	6.33 (0.42)	< 0.005***	5	6.83 (0.41)	< 0.005***
Cognitive flexibility at home	10	20.83 (2.4)	< 0.005***	13	20.67 (2.73)	< 0.05*
Social cognition	39	67.17 (8.93)	< 0.05*	48	67.50 (5.86)	< 0.05*

*Patient 3*

Variable	Pre			Post		
	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Hopping on mats	0	4.92 (0.20)	< 0.005***	0.5	4.67 (0.62)	< 0.005***

Cognitive flexibility Speed	NA	NA	NA	NA		
Cognitive flexibility Quality	NA	NA	NA	NA		
Cognitive flexibility at home	12	20.83 (2.40)	< 0.005***	21	20.67 (2.73)	= 0.46
Social cognition	61	67.17 (8.93)	< 0.05*	62	67.50 (5.86)	= 0.212

*Patient 4*

Variable	Pre			Post		
	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Hopping on mats	2	4.83 (0.26)	< 0.001***	5	4.66 (0.52)	= 0.29
Cognitive flexibility Speed	6.05	6.2 (2.01)	= 0.47	5.35	4.45 (1.09)	= 0.24
Cognitive flexibility Quality	5	6.33 (0.52)	< 0.05*	7	6.83 (0.41)	= 0.36
Cognitive flexibility at home	14	19.83 (3.54)	= 0.09	15	19.4 (4.16)	= 0.19
Social cognition	58	64.83 (8.25)	= 0.24	68	64 (4.30)	= 0.21

*Patient 5*

Variable	Pre			Post		
	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Hopping on mats	2	4.94 (0.18)	< 0.001***	4	5 (0.00)	< 0.001***
Cognitive flexibility Speed	8.60	4.76 (0.80)	< 0.001***	6.43	4.02 (0.69)	< 0.01**
Cognitive flexibility Quality	5	6.37 (0.74)	= 0.06	6	6.37 (0.52)	= 0.26
Cognitive flexibility at home	20	18.87 (3.52)	= 0.38	20	19.28 (0.59)	= 0.43
Social cognition	59	66.37 (2.67)	< 0.05*	60	66.12 (2.70)	< 0.05*

*Patient 6*

Variable	Pre			Post		
	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Hopping on mats	2	4.83 (0.26)	< 0.001***	5	4.66 (0.52)	= 0.29
Cognitive flexibility Speed	10.33	6.2 (2.01)	= 0.057	7.77	4.45 (1.09)	< 0.05*
Cognitive flexibility Quality	1	6.33 (0.52)	< 0.001***	3	6.83 (0.41)	< 0.001***
Cognitive flexibility at home	16	19.83 (3.54)	= 0.18	20	19.4 (4.16)	= 0.45
Social cognition	52	64.83 (8.25)	= 0.1	49	64 (4.30)	< 0.01**

*Patient 7*

Variable	Pre			Post		
	Patient's score	Control Group Mean (SD)	<i>p</i>	Patient's score	Control Group Mean (SD)	<i>p</i>
Hopping on mats	1	4.83 (0.26)	< 0.001***	1	4.66 (0.52)	< 0.001***
Cognitive flexibility Speed	35	6.2 (2.01)	< 0.001***	12.62	4.45 (1.09)	< 0.001***
Cognitive flexibility Quality	1	6.33 (0.52)	0.000***	2	6.83 (0.41)	0.000***
Cognitive flexibility at home	11	19.83 (3.54)	< 0.05*	14	19.4 (4.16)	= 0.14
Social cognition	56	64.83 (8.25)	= 0.18	60	64 (4.30)	= 0.21

**Figures**




Figure 1. Experimental design for the CEREB children.