

Abstract Submitted  
for the DNP07 Meeting of  
The American Physical Society

**Decoupling via the Similarity Renormalization Group for Nucleon-Nucleon Forces**<sup>1</sup> E.D. JURGENSON, R.J. FURNSTAHL, Ohio State Univ., S.K. BOGNER, Michigan State Univ. — The Similarity Renormalization Group (SRG) provides a compelling new method for decoupling low-energy nuclear physics from high-energy details. While observables are unchanged by the SRG's unitary transformations, the dependence of matrix elements on high-momentum contributions is modified by the running transformation. The SRG has the effect of partially diagonalizing the potential to a width of order the evolution parameter  $\lambda$ . Because of this diagonalization, one expects a simple decoupling of the low-energy observables for these high-energy degrees of freedom. In a previous work,<sup>2</sup> evidence for decoupling in phase shifts and the deuteron was shown for the Argonne V18 potential. Here we extend the demonstration of decoupling to other NN potentials and up to  $A=6$  nuclei to verify its universal nature and to show quantitatively that the residual coupling is perturbative above the energy corresponding to the SRG evolution parameter.

<sup>1</sup>Supported in part by the NSF under grants Nos. PHY-0354916 and PHY-0653312.

<sup>2</sup>S.K. Bogner, R.J. Furnstahl, R.J. Perry and A. Schwenk, Phys. Lett. B 649 (2007) 488.

Richard Furnstahl  
Ohio State University

Date submitted: 02 Jul 2007

Electronic form version 1.4