

Decrease Properties of Truncated Correlation Functions and Analyticity Properties for Classical Lattices and Continuous Systems

M. Duneau

Centre de Physique Théorique, Ecole Polytechnique, Paris

D. Iagolnitzer

Service de Physique Théorique, Centre d'Etudes Nucléaires de Saclay, Gif-sur-Yvette,
France

B. Souillard

Centre de Physique Théorique, Ecole Polytechnique, Paris, France

Received January 2, 1973

Abstract. We present and discuss some physical hypotheses on the decrease of truncated correlation functions and we show that they imply the analyticity of the thermodynamic limits of the pressure and of all correlation functions with respect to the reciprocal temperature β and the magnetic field h (or the chemical potential μ) at all (real) points (β_0, h_0) (or (β_0, μ_0)) where they are supposed to hold. A decrease close to our hypotheses is derived in certain particular situations at the end.

Introduction

In a recent work [1], Lebowitz considers a ferromagnetic Ising spin lattice and presents a method which, starting from decrease properties of the correlations, proved in certain cases, allows him to derive regularity properties of the infinite-volume free energy $\psi(\beta, h)$ and correlations $\langle \sigma_A \rangle(\beta, h)$, at real points (β_0, h_0) where other methods have given so far no information (e.g. at points close to β_c). However, even for finite range potentials, only infinite differentiability, rather than analyticity has been obtained. The purpose of the present paper is to present and discuss certain “physically reasonable” hypotheses, not proved in general so far, on the decrease of the truncated functions¹ and to show that they do yield analyticity.

¹ Also called “cluster functions” or “Ursell functions” of the correlations in statistical mechanics.

The hypotheses are described in Part I. In Section A, we recall the general type of decrease, called below “weak decrease”, obtained and used in [1] and we see that a stronger decrease might be expected, following the idea that the *truncated* functions correspond to processes or correlations “involving all the particles jointly”. We outline the mathematical reasons why the “weak decrease properties” do not yield analyticity in Section B. In Section C, we then present our hypotheses, called “strong decrease properties”; we indicate in Section D how they will yield analyticity, in contrast to the “weak properties” and we see that they are completely consistent with all known results, (in particular those by Ruelle [2] on the integrals of truncated functions at low activity).

In Section E, we finally describe an extension, whose usefulness will appear in Parts II, III of the hypotheses to functions which are truncated only with respect to various clusters of points.

In Parts II, III, we respectively consider classical lattices and continuous systems. In Section II.A (resp. III.A) we present general formulae expressing the successive derivatives of the pressure and of the correlation functions with respect to β and h , (resp. β and μ), in terms of (generalized) truncated functions, and in II.B, (resp. III.B), we prove that our hypotheses do imply analyticity. In the continuous case, the truncation does not refer in general to the correlation functions $\rho(X)$ themselves, but to related quantities which generalize the functions $\hat{\rho}(X)$ of [3] and which are introduced at the beginning of Part III.

In Part IV, we consider lattice systems with finite range interactions and we show that, whenever there is a gap in the spectrum of the transfer matrix, the n -point truncated correlations have a decrease of the type $e^{-\chi D}$ where D is the diameter of the configuration $X = (x_1, \dots, x_n)$ ($D = \text{Max}_{i,j=1,\dots,n} |x_i - x_j|$), and χ is independent of n (and of the configuration). This decrease is better than the “weak decrease”, which, for instance in the case of equidistant points on a line, provides only a factor $e^{-\chi D/n}$. It coincides with the “strong decrease” whenever the points are lined up, but is weaker in the other situations. Concerning the problem of analyticity, the replacement of $e^{-\chi D/n}$ by $e^{-\chi D}$ is already an improvement (see footnote 2 in Part I.B). However, this is not yet sufficient to derive analyticity.

I. Physical Hypotheses on the Truncated Functions

A. Decrease Properties of the Truncated Functions

We consider factorization properties of the following type for the (non-truncated) correlation functions $f(X)$ ($X = (x_1 \dots x_n)$), when the

distances between several clusters $X_1 \dots X_M$ of points tend to infinity:

$$\begin{cases} f(X_1, X_2) - f(X_1)f(X_2) \simeq \mathcal{O}[u(d(X_1, X_2))], & (1) \\ f(X_1, \dots, X_M) - \prod_{j=1}^M f(X_j) \simeq \mathcal{O}\left[u\left(\inf_{\substack{i,j=1, \dots, M \\ i \neq j}} d(X_i, X_j)\right)\right] & (2) \end{cases}$$

where u is a monotonous decreasing function vanishing at infinity, and $d(X_1, X_2) = \text{Inf}\{d(x_i, x_j)\} (x_i \in X_1, x_j \in X_2)$. We remark that (2) is a consequence of (1) and that we cannot expect physically better than the infimum of the distances in (2).

The truncated functions, denoted below by T.F., are defined recurrently by:

$$\begin{cases} f^T(x) = f(x) \\ f^T(x_1, \dots, x_n) = f(x_1, \dots, x_n) - \sum_{\pi_1 \dots \pi_k} \prod_{j=1}^k f^T(x(\pi_j)) \end{cases} \quad (3)$$

where the sum $\sum_{\pi_1 \dots \pi_k}$ runs over all (non trivial, i.e. $k > 1$) partitions of x_1, \dots, x_n .

When factorization properties are satisfied, it is well known that the T.F. vanish at infinity as soon as certain clusters are separated from each other. In the present framework, one shows more precisely that (1), (2) are equivalent to the following decrease property, that we shall call "weak decrease":

$$f^T(x_1, \dots, x_n) \simeq \mathcal{O}(u(\text{Max}_{X_1, X_2} d(X_1, X_2))) \quad (4)$$

where X_1, X_2 is any partition of x_1, \dots, x_n into two subsets.

Starting from (1) (2), it is not possible to get better than (4). But better decrease properties might be physically expected in various situations, namely a decrease taking into account the separation of all particles with respect to each other, and not only the maximum distance $\text{Max}d(X_1, X_2)$. Truncated functions do verify such ideas in particle physics [4]. We note that we do not want conversely to get better, unexpected, asymptotic factorization properties than type (1), (2). The "strong decrease properties" of the T.F. introduced in Section C satisfy this requirement.

B. Weak Decrease Properties and the Problem of Analyticity

For the case of a ferromagnetic spin system, Lebowitz does obtain in [1] factorization properties of type (1), (2) in the form:

$$|\langle \sigma_{X_1 X_2} \rangle_A - \langle \sigma_{X_1} \rangle_A \langle \sigma_{X_2} \rangle_A| < |X_1| |X_2| u(d(X_1, X_2)) \quad (5)$$

where A is any finite region and where u is independent of A , (and depends only on β and h). The corresponding weak decrease of the T.F., of type (4) is:

$$|\langle \sigma_{i_1, \dots, i_n} \rangle_A^T| < C^{n-1} (n-1)! u(\text{Max}_{X_1, X_2} d(X_1, X_2)). \tag{6}$$

The relationship between the derivatives of the free energy ψ_A and of the correlations $\langle \sigma_A \rangle_A$ with respect to β, h , and the T.F., then allows to obtain the infinite differentiability of the thermodynamic limits ψ and $\langle \sigma_A \rangle$, if u has a rapid decrease at infinity, (i.e. faster than any inverse power of the distance). The decrease of u has actually been proved in certain cases to be exponential, for finite-range potentials and in appropriate domains in β and h : see [1, 5].

For instance, let us consider the behaviour of ψ with respect to h . It is proved in [1] (see Part II.A) that

$$\frac{\partial^n \psi_A}{\partial \tilde{h}^n}(\beta, \tilde{h}) = \frac{1}{|A|} \sum_{i_1, \dots, i_n \in A} \langle \sigma_{i_1}, \dots, \sigma_{i_n} \rangle_A^T(\beta, \tilde{h}), \quad (\tilde{h} = \beta h) \tag{7}$$

Then (6) yields bounds on $\frac{\partial^n \psi_A}{\partial \tilde{h}^n}$ which are uniform with respect to A as soon as $u(d)$ decreases at least like $d^{-nv-\epsilon}$ at infinity, where v is the space dimension. It is shown in [1] that those bounds entail the existence of the thermodynamic limit ψ and of its derivatives.

The n^{th} derivative of ψ still possesses the same uniform bound as $\frac{\partial^n \psi_A}{\partial \tilde{h}^n}$.

But the trouble which prevents from getting analyticity even when u is rapidly or exponentially decreasing, is that the bounds, thus derived from (6), increase too fast with n : in particular, the use of the weak decrease factor $u(\text{Max}(d(X_1, X_2)))$ induces terms such as n^{nv} and $(nv)!^2$.

In our approach, we shall only use integrability properties of u but we will require stronger decrease properties of the T.F. than (6).

C. Strong Decrease Properties of the Truncated Functions

Asymptotic Physics of the T.F.

We now want to express in a stronger way the idea that the T.F. correspond to correlations “involving all the particles jointly”.

Consider for simplicity a three-point function $f^T(x, y, z)$ and the situation in which x, y, z are separated from each other in a way such that $d(x, z) \gg d(x, y), d(y, z)$, see Fig. 1:

² The decrease $e^{-\epsilon d}$ proved in Part IV allows one to remove the terms n^{nv} (but not $(nv)!$).



Fig. 1

We may then expect a first decrease of the order of $u(d(x, y))$ due to separation of x with respect to y (inside the “cluster” x, y) and a second factor of the order of $u(d(y, z))$ due to the separation of z with respect to y . Effects due to the separation of z with respect to x are masked asymptotically since $u(d(z, x)) \ll u(d(x, y)), u(d(y, z))$. Therefore we are led to a decrease of the order of $u(d(x, y))u(d(y, z))$ whereas (4) provides a decrease of the order of $u(d(y, z))$ alone.

For general n -point functions $f^T(x_1, \dots, x_n)$, the same physical ideas lead to consider all trees \mathcal{T} (i.e. connected graphs without closed loops) of $n-1$ lines joining x_1, \dots, x_n and to assume a decrease of the type:

$$|f^T(x_1, \dots, x_n)| \simeq \mathcal{O} \left(\prod_{l \in \mathcal{T}_{\min}} u(d_l) \right) \quad (8)$$

where \mathcal{T}_{\min} is the tree whose length L is minimal and d_l is the length of line l (The length $\mathcal{L}(\mathcal{T})$ of a tree \mathcal{T} is the sum of the lengths of its lines).

An alternative type of decrease will be presented in Section D for finite-range interactions, but will be shown to be equivalent.

The above tree \mathcal{T}_{\min} can be obtained as the n -th term of a sequence of (connected) trees \mathcal{T}_p joining p points: \mathcal{T}_1 may be anyone of the points x_1, \dots, x_n and \mathcal{T}_{p+1} is obtained from \mathcal{T}_p by adding the line of smallest length joining a point outside \mathcal{T}_p to a point of \mathcal{T}_p . Alternatively, \mathcal{T}_{\min} can be constructed by considering the set of all lines joining two points, and by successively choosing the line of smallest length, avoiding however closed loops. (These two constructions correspond to slightly different physical considerations).

Bounds on the T.F

In the following we need not only asymptotic properties, but general bounds which would extend the bound on the 2-point-function $\langle \sigma_{ij} \rangle^T (= \langle \sigma_{ij} \rangle - \langle \sigma_i \rangle \langle \sigma_j \rangle)$ which follows from (5) or (6).

We shall consider the assumption:

$$|f^T(x_1, \dots, x_n)| < \sum_{\mathcal{T}} \prod_{l=1}^{n-1} u(d_l) \quad (9)$$

where the sum Σ runs over all trees \mathcal{T} (of $n-1$ lines) joining x_1, \dots, x_n .

D. Links with Analyticity and Consistency with Known Results

For simplicity, we here consider again the free energy ψ of a spin lattice and its behaviour with respect to h . The function $f^T(x_1, \dots, x_n)$ is here anyone of the $\langle \sigma_{i_1, \dots, i_n} \rangle_A^T$ obtained for finite A ; (9) will be supposed to hold in appropriate domains in β, h , and for any allowed point β, h the corresponding function u is assumed to be independent of A and integrable ($\sum_{i \in Z^v} u(|i|) < \infty$): see Part II.

The contribution to the sum $\frac{1}{|A|} \sum_{i_1, \dots, i_n}^A$ of (7) corresponding to a given tree is clearly bounded by $\left[\sum_{i \in Z^v} u(|i|) \right]^{n-1}$. Since the number of trees is itself bounded, as is easily checked, by $D^{n-1}(n-1)!$ (where D is independent of n), we get:

$$\left| \frac{\partial^n \psi_A}{\partial \tilde{h}^n} \right| < \frac{1}{|A|} \sum_{i_1, \dots, i_n \in A} |\langle \sigma_{i_1, \dots, i_n} \rangle_A^T| < \left[D \left(\sum_{i \in Z^v} u(|i|) \right) \right]^{n-1} (n-1)! \quad (10)$$

It is the use of such bounds (generalized in Parts II, III to derivatives of ψ_A and $\langle \sigma_A \rangle_A$ with respect to β and h and to the continuous case) which will ensure the convergence of the appropriate Taylor series and will thus lead to analyticity (see Parts II and III).

Consistency with Known Results

a) *Results Obtained from the Explicit Form of the T.F.* [6]. 1) We below consider the example of a classical gas. The truncated correlation is then given explicitly [2] by:

$$\varrho^T(x_1, \dots, x_n) = \sum_{N \geq 0} \frac{z^{N+n}}{N!} \int \varphi(x_1, \dots, x_n, y_1, \dots, y_N) dy_1, \dots, dy_N$$

where

$$\varphi(x_1, \dots, y_N) = \sum_{\Gamma^c} \prod_{l \in \Gamma^c} (e^{-\beta \Phi(d_l)} - 1).$$

The sum \sum_{Γ^c} runs over all *connected* graphs joining $x_1, \dots, x_n, y_1, \dots, y_N$ and d_l is the length of line l .

In the case of a finite-range interaction, $\Phi(d) = 0$ when $d > d_0$, the product $\prod_{l \in \Gamma^c}$ vanishes whenever $d(l_0) > d_0$ for at least one line l_0 and is otherwise bounded by $e^{-\bar{\chi} N(\Gamma^c)}$ where $\bar{\chi}$ is positive if $|e^{-\beta \Phi} - 1| < 1^3$, and $N(\Gamma^c)$ is the number of lines in Γ^c . For given points x_1, \dots, x_n , we remark that $e^{-\bar{\chi} N(\Gamma^c)}$ is always less than $e^{-\chi L'}$ where L' is the minimal length of all (connected) trees \mathcal{T}' joining x_1, \dots, x_n and possibly other vertices.

³ Or possibly equal to 1 at $d = 0$.

Although no such result has been proved so far, a detailed analysis of the above form of $q^T(x_1, \dots, x_n)$ then suggests that it might have a decrease of the type $e^{-\kappa L'}$, under appropriate conditions.

We remark that this decrease is in actual fact *equivalent* to (8) if we admit that the function $u(d)$ of (8) is of the form $e^{-\kappa d}$ for finite-range interactions: the product $\prod_{l \in \mathcal{T}_{\min}} u(d_l)$ is then equal to $e^{-\kappa L}$ where L is the length of \mathcal{T}_{\min} , and the equivalence follows from the inequalities $\frac{1}{2}L \leq L' \leq L$.

The inequality $L' \leq L$ is obvious, since the trees \mathcal{T} of Section C may not include other vertices than x_1, \dots, x_n . To obtain the inequality $\frac{1}{2}L \leq L'$ [7], we note that a tree \mathcal{T}_0 , whose length $\mathcal{L}(\mathcal{T}_0)$ satisfies $(L \leq) \mathcal{L}(\mathcal{T}_0) \leq 2L'$, can always be constructed from the minimal tree \mathcal{T}_{\min} by joining directly the points x which appear successively when "turning around \mathcal{T}'_{\min} ".

2) Ruelle has proved [2], at low activity and under general assumptions of stability and integrability of the potential, that the infinite-volume T.F. $q_n^T(x_1, \dots, x_n)$ are absolutely integrable with respect to the differences $x_i - x_j$. More precisely, he obtains a bound of the following type:

$$\int |q_n^T(x_1, \dots, x_n)| dx_2, \dots, dx_n < C^{n-1}(n-1)! \quad (11)$$

(where C depends on β and μ).

The "strong decrease properties" do lead to such bounds [see for instance (10)] in contrast to the "weak decrease properties". The latter yield the infinite volume integrability of the T.F. for all n if u is rapidly decreasing, but the corresponding bound is even then much worse than (11) (see Section B).

A final remark: a bound better than (11) is not expected [8]. The hypotheses (9) do not indeed lead to a better bound since the number of trees is larger than $(n-1)!$.

b) *Results Obtained from the Transfer Matrix Formalism.* As already mentioned, (8) has been proved in certain cases for the 2-point function $\langle \sigma_{ij} \rangle^T$, with u being exponentially decreasing for finite-range potentials [1, 5]. Those results will be generalized in Part IV to the n -point function when the points are lined up.

E. Generalized Truncated Functions

We now introduce functions $f^{T(X_1, \dots, X_M)}(X)$, called G.T.F., which are truncated only with respect to certain subsets X_1, \dots, X_M of a partition of x_1, \dots, x_n , i.e. are defined by the following formulae where

X_1, \dots, X_M replace the individual variables x_1, \dots, x_n of (2):

$$f^{T(X)}(X) = f(X) - \sum_{(\pi_1, \dots, \pi_k)}^{(X_1, \dots, X_M)} \prod_{j=1}^k f^{T(X(\pi_j))}(X(\pi_j)) \quad (12)$$

where $\sum_{\pi_1, \dots, \pi_k}^{X_1, \dots, X_M}$ is the sum over all (non trivial) partitions of (X_1, \dots, X_M) ; $\{X(\pi_j)\}$ denotes the subsets X_i belonging to π_j and $X(\pi_j)$ is the set of all x_i belonging to the various subsets of $\{X(\pi_j)\}$. We note that $f^{T(X_1, \dots, X_M)}$ reduces to $f(X)$ when $M=1$, to the usual T.F. $f^T(X)$ when each $X_t (t=1, \dots, M)$ is one point ($M=n$), and that it can be generally expressed in terms of usual T.F. through the following formulae, which can be proved by a recurrence procedure:

$$f^{T(X_1, \dots, X_M)}(X) = f^T(X) + \sum_{(\pi_1, \dots, \pi_k)^{\neq} / X_1, \dots, X_M}^{(x_1, \dots, x_n)} \prod_{j=1}^k f^T(X(\pi_j)) \quad (13)$$

where the sum Σ in the right-hand side runs over all (non trivial) partitions of (x_1, \dots, x_n) which are “connected” with respect to X_1, \dots, X_M : i.e. such that a *connected* diagramm of M vertices is obtained when each X_t is identified with one vertex ($t=1, \dots, M$) and when X_{t_1}, X_{t_2} are joined by a line whenever one (or more) variable x_{i_1} in X_{t_1} and one (or more) variable x_{i_2} in X_{t_2} belong to a common $\pi_j (j=1, \dots, k)$.

Strong Decrease Hypotheses for the G.T.F.

From the same physical ideas as before, we shall now assume the following “strong decrease properties” for the G.T.F, analogous to those described for the usual T.F.:

$$|f^{T(X_1, \dots, X_M)}(X)| < \left(\prod_{i=1}^M \chi(|X_i|) \right) \sum_{\mathcal{T}}^{(X_1, \dots, X_M)} \prod_{l=1}^{M-1} d(X'_l, X''_l) \quad (14)$$

where $\sum_{\mathcal{T}}^{X_1, \dots, X_M}$ is a sum over all trees of $M-1$ lines joining the subsets X_1, \dots, X_M , X'_l and X''_l are the subsets joined by line l and $d(X'_l, X''_l)$ is their distance.

A factor $\prod_{i=1}^M \chi(|X_i|)$ has been introduced [we shall assume that $\chi(1)=1$]. We shall not discuss it in detail from physical considerations, but we remark that a bound of type (14) is actually obtained by Lebowitz in [1] in the case of two subsets X_1, X_2 : see (5) [Note that $f^{T(X_1, X_2)}(X) = f(X_1, X_2) - f(X_1) f(X_2)$]. As a matter of fact, (5) in Lebowitz’s proof

is a consequence of the more refined bound:

$$|\langle \sigma \rangle_A^{T(X_1, X_2)}| < \sum_{\substack{x \in X_1 \\ y \in X_2}} u(d(x, y)). \quad (15)$$

A direct generalization of (15) when $M > 2$, from which (14) would follow with $\chi(|X|)$ being an appropriate power of $|X|$, would be:

$$|f^{T(X_1, \dots, X_M)}(X)| < \sum_{\mathcal{F}}^{(X_1, \dots, X_M)} \sum_{\substack{\{x'_i \in X_i(\mathcal{F}) \\ \{x''_i \in X'_i(\mathcal{F})\}}} \prod_{l=1}^{M-1} u(d(x'_l, x''_l)). \quad (16)$$

A Final Remark

We note that (9) is a particular case of (14). Conversely by using (13) the following bound may be derived from (9) for the G.T.F.:

$$|f^{T(X_1, \dots, X_M)}(X)| < \sum_{\Gamma^c / X_1, \dots, X_M}^{x_1, \dots, x_n} \prod_{l \in \Gamma^c} u(d(x'_l, x''_l)) \quad (17)$$

where the sum Σ runs over all graphs Γ^c without closed loops, which have n vertices (x_1, \dots, x_n) and which are connected with respect to X_1, \dots, X_M ⁴.

The bound (17) does provide the expected asymptotic behaviour of $f^{T(X_1, \dots, X_M)}(X)$: each term in the sum Σ of (17) is in fact bounded by (at least) one of the terms involved in the sums Σ of (14) (or (15)) multiplied by an appropriate power of $u(0)$. However the number of terms involved in (17) is too large for the proofs of parts II, III referring to the derivatives with respect to β . We shall then use the formulation (14) (or (15)) of our physical assumptions. Concerning the derivatives with respect to h (or μ), we shall see that (17) and thus (9) are sufficient.

II. Ferromagnetic Ising Spin Systems

In this part, we consider classical spin lattices.

For simplicity, we shall restrict ourselves to Ising spin systems with ferromagnetic pair interactions, but the essential parts of what follows apply equally well to more general lattices. The space dimension ν will not be specified. In the following, we shall be interested only in cases where there is no phase transition. Therefore, the thermodynamic limits $A \rightarrow \infty$ of the correlation functions are known, from the results of [9], to be independent of the boundary conditions. For simplicity,

⁴ The definition is here completely analogous to that given above for the connected partitions.

we thus consider below free boundary conditions, but formulae analogous to those below can be written more generally.

We recall the following definitions:

$$H_A = -1/2 \sum_{\substack{i,j \in A \\ i \neq j}} J_{ij} \sigma_i \sigma_j - h \sum_{i \in A} \sigma_i, \quad (18)$$

$$Z_A = \sum_{\sigma} \exp(-\beta H_A(\sigma, h)), \quad (19)$$

$$\psi_A = |A|^{-1} \log Z_A, \quad (20)$$

$$\langle \sigma_A \rangle_A = (Z_A)^{-1} \sum_{\sigma} \sigma_A \exp(-\beta H_A(\sigma, h)),$$

where $\sigma_A = \prod_{i \in A} \sigma_i$. In the following, we consider β and $\tilde{h} = \beta h$ as independent variables.

A. Derivatives of ψ_A and $\langle \sigma_A \rangle_A$

The following formulae [generalizing (7)] hold⁵:

$$\frac{\partial^{q+r} \psi_A}{\partial \tilde{h}^q \partial \beta^r}(\beta, \tilde{h}) = |A|^{-1} \sum_{\substack{i_1, \dots, i_q \in A \\ \gamma_1, \dots, \gamma_r \in A^2}} \left(\prod_{j=1}^r J_{\gamma_j} \right) \langle \sigma_{i_1, \dots, i_q, \gamma_1, \dots, \gamma_r} \rangle_A^{T_{q,r}}(\beta, \tilde{h}), \quad (21)$$

$$\frac{\partial^{q+r} \langle \sigma_A \rangle_A}{\partial \tilde{h}^q \partial \beta^r}(\beta, \tilde{h}) = \sum_{\substack{i_1, \dots, i_q \in A \\ \gamma_1, \dots, \gamma_r \in A^2}} \left(\prod_{j=1}^r J_{\gamma_j} \right) \langle \sigma_{A, i_1, \dots, i_q, \gamma_1, \dots, \gamma_r} \rangle_A^{T_{q,r}(A)}(\beta, \tilde{h}) \quad (22)$$

where $T_{q,r}$ and $T_{q,r}(A)$ mean the truncation with respect to the q subsets i_1, \dots, i_q of one point, the r subsets $\gamma_1, \dots, \gamma_r$ of two points and possibly the subset A .

When $r=0$ and A is one point, (21) (22) involve only usual T.F. and are given in [1]. On the other hand, they can be extended without difficulty for k -body potentials ($k > 2$), the functions involved in the right-hand sides being then truncated with respect to subsets of k points and less. They are readily derived, once the following formulae are proved:

$$\frac{\partial \psi_A}{\partial \tilde{h}} = |A|^{-1} \frac{1}{Z_A} \frac{\partial Z_A}{\partial \tilde{h}} = |A|^{-1} \sum_{i \in A} \langle \sigma_i \rangle_A, \quad (23)$$

$$\frac{\partial \psi_A}{\partial \beta} = |A|^{-1} \frac{1}{Z_A} \frac{\partial Z_A}{\partial \beta} = |A|^{-1} \sum_{\gamma \in A^2} J_{\gamma} \langle \sigma_{\gamma} \rangle_A, \quad (24)$$

$$\frac{\partial}{\partial \tilde{h}} \langle \sigma_X \rangle_A^{T(X_1, \dots, X_M)} = \sum_{i \in A} \langle \sigma_{X,i} \rangle_A^{T(X_1, \dots, X_M, i)}, \quad (25)$$

$$\frac{\partial}{\partial \beta} \langle \sigma_X \rangle_A^{T(X_1, \dots, X_M)} = \sum_{\gamma \in A^2} J_{\gamma} \langle \sigma_{X,\gamma} \rangle_A^{T(X_1, \dots, X_M, \gamma)}. \quad (26)$$

⁵ In the following $J_{\gamma} = 0$ when $\gamma = (i, i)$.

(23) (24) are easily obtained by explicit computation as well as (25) (26) when $M=1$. The general proof of (25) (26) follows by a recurrence procedure.

B. Analyticity Properties of ψ and $\langle \sigma_A \rangle$

The existence and uniqueness of the thermodynamic limits of ψ_A and $\langle \sigma_A \rangle_A$ are known at all points (β, h) outside the phase transition region ($h=0, \beta > \beta_c$): see for instance [9].

We then state:

Theorem. Let β_0, \tilde{h}_0 be a (real) point at which the following bound holds uniformly for any region A :

$$|\langle \sigma_{A_1, \dots, A_M} \rangle_A^{T(A_1, \dots, A_M)}(\beta_0, \tilde{h}_0)| < \prod_{i=1}^M \chi(|A_i|) \times \sum_{\mathcal{F}}^{(A_1, \dots, A_M)} \prod_{l=1}^{M-1} u(d(A'_l, A''_l)) \quad (27)$$

where A_1, \dots, A_M are finite subsets of Z^v , u is integrable ($\sum_{i \in Z^v} u(|i|) < \infty$) and the meaning of the right-hand side is the same as in (14).

Then the thermodynamic limits $\psi(\beta, \tilde{h})$ and $\langle \sigma_A \rangle(\beta, \tilde{h})$ are analytic at (β_0, \tilde{h}_0) with respect to both variables.

Before giving the proof, we make some remarks:

1) The uniqueness of the thermodynamic limit $\langle \sigma_A \rangle$ which is already known from the results of [9] is also implied by a factorisation criterion on the probabilities of configurations which is due to Dobrushin [10] and which is akin to the decrease property (27) in the case of two subsets A_1, A_2 .

2) It seems physically reasonable to assume that the bounds (27) are valid at all points (β_0, \tilde{h}_0) outside the phase transition region. We recall that they have actually been proved by Lebowitz [1] in the case of two subsets A_1, A_2 , under various conditions.

3) Analyticity is already known when $\tilde{h}_0 \neq 0$ or $\tilde{h}_0 = 0$ and $\beta < \beta'$ where $\beta' < \beta_c$. (For references see for instance [11]). Thus the above theorem will give further information at all points $\tilde{h}_0 = 0$ and $\beta' \leq \beta_0 < \beta_c$ where (27) is valid, and the assumptions of the theorem can be reduced to those points.

4) As mentioned in Part I, the analyticity of ψ and $\langle \sigma_A \rangle$ with respect to \tilde{h} alone can be derived from assumptions involving only usual T.F.

(This follows from the remarks at the end of part I and from the fact that the number of graphs T^c is bounded by $C^{|A|+q}(|A|+q)!$ where C depends only on $|A|$, in the case of q subsets of one point besides the

subset A . Since $|A|$ is fixed, this factor will not modify the arguments given below).

Proof. For simplicity, we first consider $\psi(\beta, \tilde{h})$. For any finite A , ψ_A is analytic at (β_0, \tilde{h}_0) . Thus we have in a complex neighborhood of (β_0, \tilde{h}_0) :

$$\psi_A(\beta, \tilde{h}) = \sum_{\substack{q \geq 0 \\ r \geq 0}} \frac{(\tilde{h} - \tilde{h}_0)^q (\beta - \beta_0)^r}{q! r!} \frac{\partial^{q+r} \psi_A}{\partial \tilde{h}^q \partial \beta^r}(\beta_0, \tilde{h}_0). \quad (28)$$

The successive derivatives of ψ_A are given by (21) and the following bounds, uniform with respect to A , are then readily obtained from (27) in a way similar to the derivation of (10) in Part I⁶

$$\left| \frac{\partial^{q+r} \psi_A}{\partial \tilde{h}^q \partial \beta^r}(\beta_0, \tilde{h}_0) \right| \leq (q+r)! \left[C \sum_{i \in \mathbb{Z}^v} u(|i|) \right]^{q+r-1} \left[\chi(2) \sum_{i \in \mathbb{Z}^v} J_{0i} \right]^r. \quad (29)$$

From (29), it follows that the series defining $\psi_A(\beta, \tilde{h})$ are convergent in a complex disk D independent of A , in which a complex neighborhood of (β_0, \tilde{h}_0) of the form $D_1(\beta_0) \times D_2(\tilde{h}_0)$ can always be included, and that the ψ_A are moreover bounded in D uniformly with respect to A .

For any given real β_1 in $D_1(\beta_0)$, [resp. any real h_1 in $D_2(\tilde{h}_0)$], it follows readily from Vitali's theorem [12] and from the convergence at real points that ψ_A converges to a function ψ which is analytic with respect to \tilde{h} , in $D_2(\tilde{h}_0)$ [resp. with respect to β , in $D_1(\beta_0)$]. From that convergence, it follows from a second application of Vitali's theorem that for any fixed complex \tilde{h} in $D_2(\tilde{h}_0)$ [resp. complex β in $D_1(\beta_0)$] $\psi = \lim \psi_A$ exists and is analytic with respect to β in $D_1(\beta_0)$ [resp. \tilde{h} in $D_2(\tilde{h}_0)$]. Then Hartog's theorem [13] entails that ψ is analytic with respect to both variables β and \tilde{h} in $D_1(\beta_0) \times D_2(\tilde{h}_0)$.

The analyticity of $\langle \sigma_A \rangle(\beta, \tilde{h})$ is derived similarly.

III. Classical Continuous Systems

We again restrict ourselves in this part to systems of classical identical particles interacting through a two-body potential Φ . As in Part II, similar results may be obtained in more general cases. We shall assume that Φ is real symmetric, stable and absolutely integrable $\left(\int_{\mathbb{R}^v} |\Phi(x)| dx < \infty \right)$.

(Some results might however also be obtained by assuming only integrability at infinity.) For any finite region A of \mathbb{R}^v , we consider the

⁶ Use $u(d(A'_i, A''_i)) \leq \sum_{\substack{a'_i \in A'_i \\ a''_i \in A''_i}} u(d(a'_i, a''_i))$, and the fact that the number of products $\prod_l u(d(a'_i, a''_i))$ in $\prod_l \sum_{\substack{a'_i \in A'_i \\ a''_i \in A''_i}} u(d(a'_i, a''_i))$ is bounded by 2^{q+2r-1} when there are q subsets of one point and r subsets of two points.

grand canonical system for the reciprocal temperature β and the chemical potential μ . The partition function is:

$$Z_A(\beta, \tilde{\mu}) = \sum_{N \geq 0} \frac{e^{N\tilde{\mu}}}{N!} \int_{A^N} e^{-\beta U(x_1, \dots, x_N)} dx_1, \dots, x_N \tag{30}$$

when $\tilde{\mu} = \text{Log } z$ and z is the activity.

The pressure and the correlation functions are defined (for finite A) by:

$$\beta p_A(\beta, \tilde{\mu}) = |A|^{-1} \text{Log } Z_A(\beta, \tilde{\mu}), \tag{31}$$

$$\varrho_A(X) = (Z_A(\beta, \tilde{\mu}))^{-1} \sum_{N \geq 0} \frac{e^{(N+n)\tilde{\mu}}}{N!} \int_{A^N} e^{-\beta U(X, Y^N)} dY^N. \tag{32}$$

The function $\varrho_A(X)$ is the ‘‘probability density’’ for finding n different particles at points x_1, \dots, x_n . In the following, quantities which appear naturally in the continuous case are functions $\hat{\varrho}^{(X_1, \dots, X_M)}(X)$ where X_1, \dots, X_M is a partition of (x_1, \dots, x_n) , defined by:

$$\hat{\varrho}^{(X_1, \dots, X_M)}(X) = \varrho(X) + \sum_L^{(X_1, \dots, X_M)} \delta_L(X) \varrho(X_L) \tag{33}$$

where the sum $\sum_L^{(X_1, \dots, X_M)}$ runs over all ‘‘liaisons’’ L between X_1, \dots, X_M : a liaison L is composed of a certain number of ‘‘contractions’’, each contraction being a set of variables x_i with at most one variable in each $X_t (t=1, \dots, M)$. (A variable x_i may not belong to more than one contraction of L .) With a contraction of p variables is associated a product of $(p-1)$ δ -functions linking together the variables of the contraction, and $\delta_L(X)$ is the product of all δ -functions associated with all contractions of L . Finally, the set X_L is obtained from X by identifying all the variables of each contraction⁷.

The function $\hat{\varrho}^{(X_1, \dots, X_M)}(X)$ represents the ‘‘probability density’’ for finding $n_t = |X_t|$ different particles at the points of $X_t (t=1, \dots, M)$, the particles belonging to different X_t being no longer necessarily different. When each subset X_t is reduced to one point x_t , $\hat{\varrho}^{(X_1, \dots, X_M)}(X)$ reduces to the function $\hat{\varrho}(X)$ introduced by Ruelle [3].

Formula (33) can be shown to be equivalent to:

$$\int_{(A^N)} \hat{\varrho}_A^{(X_1, \dots, X_M)}(X) \varphi(X) dX = Z_A^{-1} \sum_{N \geq 0} \frac{e^{N\tilde{\mu}}}{N!} \int e^{-\beta U(X^N)} dX^N \tag{34}$$

$$\times \sum_{J_1}^{(x_1, \dots, x_N)} \dots \sum_{J_M}^{(x_1, \dots, x_N)} \varphi(x(J_1), \dots, x(J_M))$$

⁷ $|X_L| = |X| - \sum_{\text{contractions of } L} (p-1)$.

where φ is a test function (with support in A), and each sum $\sum_{(x_1, \dots, x_N)}^{(t=1, \dots, M)}$ runs over all the ordered subsets of $|J_t| = n_t (= |X_{t_i}|)$ variables taken from $X^N = (x_1, \dots, x_N)$; $x(J_t)$ denotes the set of x_j in J_t . This formula is a generalization of Ruelle's formula (7.1) of [3] in which each X_t is a point ($M = n$).

From the functions $\hat{\varrho}^{(X_1, \dots, X_M)}(X)$, the functions $(\hat{\varrho}^{(X_1, \dots, X_M)})^{T(X_1, \dots, X_M)}$, which are truncated with respect to the same subsets X_1, \dots, X_M , and are denoted below $\varrho^{\hat{T}(X_1, \dots, X_M)}(X)$, are defined by the recurrence formulae:

$$\varrho^{\hat{T}(X_1, \dots, X_M)}(X) = \hat{\varrho}^{(X_1, \dots, X_M)}(X) - \sum_{(\pi_1, \dots, \pi_k)}^{(X_1, \dots, X_M)} \prod_{j=1}^k \varrho^{\hat{T}(\{X(\pi_j)\})}(X(\pi_j)) \quad (35)$$

where the notations are the same as in (17). The functions $\varrho^{\hat{T}(X_1, \dots, X_M)}(X)$ thus defined can be shown to possess a property analogous to (33):

$$\varrho^{\hat{T}(X_1, \dots, X_M)}(X) = \varrho^{T(X_1, \dots, X_M)}(X) + \sum_L^{(X_1, \dots, X_M)} \delta_L(X) \varrho^{T(X_1, \dots, X_M; L)}(X_L) \quad (35')$$

where $\{X_1, \dots, X_M; L\}$ is the partition of X_L obtained by gathering all variables x_i of any two subsets $X_{t_1}, X_{t_2}(t_1, t_2 = 1, \dots, M)$ as soon as there is a contraction of L joining X_{t_1} and X_{t_2} .

A. Derivatives of βp_A and $\varrho_A(A)$

Formulae analogous to (21) (22) hold:

$$\frac{\partial^{q+r}(\beta p_A)}{\partial \tilde{\mu}^q \partial \beta^r}(\beta, \tilde{\mu}) = \frac{(-1)^r}{A} \int_{(A^q \times A^{2r})} dx_1, \dots, dx_q dY_1, \dots, dY_r \prod_{j=1}^r \Phi(Y_j) \cdot e^{\hat{T}^{q,r}(x_1, \dots, x_q, Y_1, \dots, Y_r)}, \quad (36)$$

$$\frac{\partial^{q+r} \varrho_A(A)}{\partial \tilde{\mu}^q \partial \beta^r}(\beta, \tilde{\mu}) = (-1)^{r+1} \int_{(A^q \times A^{2r})} dx_1, \dots, dx_q dY_1, \dots, dY_r \prod_{j=1}^r \Phi(Y_j) \cdot \varrho_A^{\hat{T}^{q,r}(A)}(A, x_1, \dots, x_q, Y_1, \dots, Y_r) \quad (37)$$

where $dY_j = du_j dv_j$ and $\Phi(Y_j) = \Phi(u_j - v_j)$ ⁸.

As in Part II, those formulae are readily extended for k -body potentials, $k > 2$, and are derived from:

$$\frac{\partial(\beta p_A)}{\partial \tilde{\mu}}(\beta, \tilde{\mu}) = \frac{1}{|A|} \frac{1}{Z_A} \frac{\partial Z_A}{\partial \tilde{\mu}} = \frac{1}{|A|} \int_A \varrho_A(x) dx, \quad (38)$$

$$\frac{\partial(\beta p_A)}{\partial \beta}(\beta, \tilde{\mu}) = \frac{1}{|A|} \frac{1}{Z_A} \frac{\partial Z_A}{\partial \beta} = -\frac{1}{|A|} \int_{A^2} \varrho_A(Y) \Phi(Y) dY, \quad (39)$$

⁸ $\hat{T}_{q,r}$ and $\hat{T}_{q,r}(A)$ have the same meaning as in Part II.A.

$$\frac{\partial}{\partial \tilde{\mu}} \varrho_A^{\hat{T}(X_1, \dots, X_M)}(X) = \int_A \varrho_A^{\hat{T}(X_1, \dots, X_M, x)}(X, x) dx, \tag{40}$$

$$\frac{\partial}{\partial \beta} \varrho_A^{\hat{T}(X_1, \dots, X_M)}(X) = - \int_{A^2} \varrho_A^{\hat{T}(X_1, \dots, X_M, Y)} \Phi(Y) dY. \tag{41}$$

(38) (39) are obtained by explicit computation. In contrast to Part II, (40) (41) are not quite trivial at $M = 1$; (41) for instance is then a particular case of the following formula which is verified by using (34):

$$\begin{aligned} & \frac{\partial}{\partial \beta} \varrho_A^{(X_1, \dots, X_M)}(X) \\ &= - \int_{A^2} \varrho_A^{(X_1, \dots, X_M, Y)} \Phi(Y) dY + \varrho_A^{(X_1, \dots, X_M)}(X) \int_{A^2} \varrho_A(Y) \Phi(Y) dY. \end{aligned} \tag{42}$$

The general proofs of (40) (41) follow by a recurrence procedure as in Part II.

We finally note that the derivatives with respect to $\tilde{\mu}$ alone can also be written [14]:

$$\frac{\partial^q}{\partial z^q} (\beta p_A) = \frac{1}{|A|} \frac{1}{z^q} \int_{A^q} \varrho_A^T(x_1, \dots, x_q) dx_1, \dots, dx_q, \tag{43}$$

$$\frac{\partial^q}{\partial z^q} \left(\frac{\varrho_A(A)}{z^{|A|}} \right) = \frac{1}{z^{|A|+q}} \int_{A^q} \varrho_A^{T_q(A)}(A, x_1, \dots, x_q) dx_1, \dots, dx_q \tag{44}$$

where z is the activity ($z = e^{\tilde{\mu}}$).

B. Analyticity Properties

We now state:

Theorem. *Let $(\beta_0, \tilde{\mu}_0)$ be a (real) point at which:*

- 1) *the thermodynamic limits $p(\beta, \tilde{\mu})$ and $\varrho(A)(\beta, \tilde{\mu})$ of the pressure and the correlation functions exist and are uniquely defined;*
- 2) *the following bounds hold uniformly with respect to Λ for any test function $\varphi(y_1, \dots, y_M)$ with sufficiently small support around the origin⁹:*

$$\begin{aligned} & \left| \int_{\mathbb{R}^{M \times d}} dy_1, \dots, dy_M \varrho_A^{\hat{T}(X_1, \dots, X_M)}(X_1 + y_1, \dots, X_M + y_M) \varphi(y_1, \dots, y_M) \right| \\ & \leq C(\varphi) \prod_{j=1}^M \chi(|X_j|) \times \sum_{\mathcal{J}}^{X_1, \dots, X_M} \prod_{l=1}^{M-1} \int_{I_l} u(d(X'_l, X''_l)) \end{aligned} \tag{45}$$

where $X_j + y_j$ is the set X_j translated by the common vector y_j , u is an integrable function $\left(\int_{\mathbb{R}^d} u(|x|) dx < \infty \right)$ and $C(\varphi)$ depends only upon φ .

⁹ $\varrho^{\hat{T}(X_1, \dots, X_M)} \equiv \varrho^{\hat{T}(X_1 + y_1, \dots, X_M + y_M)}$.

Then the thermodynamic limits p and $q(A)$ are analytic at $(\beta_0, \tilde{\mu}_0)$ with respect to both variables.

The proof is completely analogous to that of Part II¹⁰. We finally make some remarks:

1) The uniqueness of the infinite Gibbs state is probably linked with fall-off properties of the correlation functions, as for the case of lattices (see remark about Dobrushin's criterion in Part II).

2) In contrast to Part II, the $q^{\hat{T}(X_1, \dots, X_M)}(X)$ are now distributions and thus the bounds (45) refer to integrals with some test function $\varphi(y_1, \dots, y_M)$. Bounds of the type (45) may seem plausible (provided the support of φ is sufficiently small) outside the phase transition region and the solid phases.

3) We do not know from physical considerations whether the bounds should apply to $q^{T(X_1, \dots, X_M)}$ or $q^{\hat{T}(X_1, \dots, X_M)}$. Thus it may be useful to remark that analyticity with respect to $\tilde{\mu}$ can also be derived from assumptions analogous to (45) but referring to $q^{T(X_1, \dots, X_M)}$ (or alternatively can be derived directly from analogous assumptions referring to the usual T.F. as in Part II). To see this, it is sufficient to use (43) (44), in place of (36) (37), in the proofs.

IV. Some Results on the Decrease of the Truncated Correlations in the Ising Model

For simplicity we consider a ν -dimensional Ising lattice with nearest neighbour interaction, at least in one direction \hat{l} , and the splitting into corresponding layers.

With any given finite width of the layers is associated the set of configurations σ of each layer, which generates a vector space \mathcal{H} , and the symmetric transfer matrix T defined by:

$$T(\sigma, \sigma') = \exp \left\{ -\beta \left[\frac{H(\sigma)}{2} + \frac{H(\sigma')}{2} + I(\sigma, \sigma') \right] \right\} \tag{46}$$

where $H(\sigma)$ is the energy of the configuration σ and $I(\sigma, \sigma')$ is the interaction between σ and σ' on two successive layers.

We shall denote by $\lambda_0, \lambda_1, \dots, \lambda_j, \dots, (\lambda_0 > \lambda_1 \geq \dots \geq \lambda_j \dots)$ and $|0\rangle, |1\rangle, \dots, |j\rangle, \dots$ the eigenvalues and eigenvectors of T .

Let A_0, A_1, \dots, A_n be arbitrary operators defined on the layers of coordinates $l_0, l_1, \dots, l_n (l_0 < l_1 < \dots < l_n)$ along \hat{l} .

¹⁰ The bounds on the integrals of (36) (37) are readily obtained from (45) by choosing φ such that $\int \varphi(y_1, \dots, y_M) dy_1, \dots, dy_M \neq 0$ and by remarking that the function $\prod_{j=1}^r \Phi(Y_j)$ is invariant by translations y_j of each set Y_j .

Theorem. *In the limit of an infinite number of layers, the truncated correlation function $\langle A_0 A_1 \dots A_n \rangle^T$ decreases like*

$$\left(\frac{\lambda_1}{\lambda_0}\right)^{(l_n - l_0)} \equiv \left(\frac{\lambda_1}{\lambda_0}\right)^{(l_1 - l_0)} \times \left(\frac{\lambda_1}{\lambda_0}\right)^{(l_2 - l_1)} \times \dots \times \left(\frac{\lambda_1}{\lambda_0}\right)^{(l_n - l_{n-1})},$$

when the distances $l_1 - l_0, l_2 - l_1, \dots, l_n - l_{n-1}$ tend to infinity.

Proof. It is easily shown ([15]) that:

$$\langle A_0 A_1 \dots A_n \rangle = \langle 0 | \tilde{A}_0 \tilde{A}_1 \dots \tilde{A}_n | 0 \rangle \quad (47)$$

where

$$\tilde{A}_i = \left(\frac{T}{\lambda_0}\right)^{l_i} A_i \left(\frac{T}{\lambda_0}\right)^{-l_i}$$

and correspondingly:

$$\langle A_0 A_1 \dots A_n \rangle^T = \langle 0 | \tilde{A}_0 \tilde{A}_1 \dots \tilde{A}_n | 0 \rangle^T. \quad (48)$$

The following lemma is readily checked at $n=1$ and is then proved by recurrence.

Lemma. *Let B_0, B_1, \dots, B_n be arbitrary operators defined on \mathcal{H} ; let $\mathbf{Q} = \mathbf{1} - |0\rangle\langle 0|$. Then:*

$$\langle 0 | B_0 \mathbf{Q} B_1 \dots \mathbf{Q} B_n | 0 \rangle = \sum_{\{0, \dots, n\}_r}^{(0, \dots, n)_r} \prod_{i=1}^k \langle 0 | B(\pi_i) | 0 \rangle^T \quad (49)$$

where $\sum_{\{\pi_1, \dots, \pi_k\}}^{(0, \dots, n)_r}$ runs over all partitions of $(0, \dots, n)$ (including the trivial one) which "cover" the interval $(0, n)$, i.e. are such that the set of open intervals (α_i, β_i) , where $\alpha_i = \text{Inf } \pi_i, \beta_i = \text{Sup } \pi_i$, is a covering of $(0, n)$.

(For the recurrence, use the operator $B'_{n-1} = B_{n-1} \cdot B_n$ and formula (13).)

By applying this lemma to $\tilde{A}_0, \dots, \tilde{A}_n$, we get:

$$\langle 0 | \tilde{A}_0 \tilde{A}_1 \dots \tilde{A}_n | 0 \rangle^T = \langle 0 | \tilde{A}_0 \mathbf{Q} \tilde{A}_1 \dots \mathbf{Q} \tilde{A}_n | 0 \rangle^T - \sum_{\substack{\{\pi_1, \dots, \pi_k\} \\ (\text{non trivial})}}^{(0, \dots, n)_r} \prod_{i=1}^k \langle 0 | \tilde{A}(\pi_i) | 0 \rangle^T \quad (50)$$

and the announced decrease of $\langle A_0 A_1 \dots A_n \rangle^T$ is then easily obtained by recurrence, using the fact that

$$\left\| \left(\frac{T}{\lambda_0}\right)^l \mathbf{Q} \right\| = \left(\frac{\lambda_1}{\lambda_0}\right)^l.$$

Theorem. *The same decrease is obtained in the limit of infinite width of the layers, whenever $\frac{\lambda_1}{\lambda_0}$ is strictly less than one in this limit.*

This result, which generalizes the decrease obtained for the two point function ([15]), can be extended for any finite range potential and non symmetric transfer matrices.

When A_0, A_1, \dots, A_n have the same localization on their respective layers l_0, l_1, \dots, l_n , the asymptotic decrease obtained coincides with the

strong decrease of Part I.C. [However we do not obtain the precise bound (9).]

For a general localization x_0, x_1, \dots, x_n of A_0, A_1, \dots, A_n the asymptotic decrease is, at least, of the type $e^{-\chi^D}$ where D is the diameter of the configuration (x_0, x_1, \dots, x_n) , $(D = \text{Max}_{i,j=0,\dots,n} |x_i - x_j|)$, and is thus better than the “weak decrease” of part I A, which provides essentially $e^{-\chi \frac{D}{n}}$ ([1]). However it is not yet sufficient to derive analyticity.

Acknowledgments. We are very grateful to Drs. J. Ginibre, J. Lascoux, J. L. Lebowitz, E. H. Lieb and D. Ruelle for most interesting discussions. (However none of them should be held responsible for our “mystical” belief in the properties of the truncated functions.)

Notes added in proof: 1. In various cases, a factor of the type $N_1! \dots N_p!$ is to be included in our bounds (for physical reasons) when the points x_1, \dots, x_n are not all different from each other. This factor does not change the derivations of analyticity.

2. The “strong decrease properties” assumed in this paper have in actual fact been proved in some situations, as described in a following paper by the same authors.

References

1. Lebowitz, J. L.: Commun. math. Phys. **28** (4), 313—321 (1972).
2. Ruelle, D.: Statistical mechanics, rigorous results, pp. 90—92. New York: Benjamin 1969.
3. Ruelle, D.: Statistical mechanics, rigorous results, p. 100. New York: Benjamin 1969.
4. Iagolnitzer, D., Stapp, H. P.: Commun. math. Phys. **14**, 15—55 (1969);
Iagolnitzer, D.: Introduction to S-matrix theory (to be published).
5. Fisher, M. E.: Physical Rev. **162**, 475 (1967).
Camp, W. J., Fisher, M. E.: Phys. Rev. B, August 1, (1972).
Marinero, M., Sewell, G. L.: Commun. math. Phys. **24**, 310 (1972).
Minlos, R. A., Sinai, J. G.: Investigations of the spectra of some stochastic operators arising in the lattice gas models. Teor. i Matem. Fizika **2** (2), 230 (1970) (russian).
6. We owe the ideas of this subsection to D. Ruelle.
7. We owe this elegant proof to C. Billionnet and C. DeCalan.
8. Arguments about this point have been given to us by D. Ruelle.
9. Lebowitz, J. L., Martin-Löf, A.: Commun. math. Phys. **25**, 276 (1972).
10. Dobrushin, R. I.: Gibbsian random fields for lattice systems with pairwise interactions. Funktsional'. Analiz i Ego Prilozhen, **2** (4), 31 (1968) (russian).
11. Ruelle, D.: Statistical mechanics, rigorous results. New York: Benjamin 1969.
12. For Vitali's theorem see for instance: Goluzin, G. M.: Geometrical theory of functions of a complex variable, p. 18. AMS, Providence, Rhode Island 1969.
13. For Hartog's theorem see for instance: Bochner, S., Martin, W. T.: Several complex variables. Princeton 1948, VII, 4, Th. 4.
14. Lebowitz, J. L., Penrose, O.: Commun. math. Phys. **11**, 99 (1968).
15. See for instance Camp, W. J., Fisher, M. E.: Phys. Rev. B, **6** (2), 946 (1972) where references to the transfer matrix formalism will be found.

M. Duneau
B. Souillard
Centre de Physique Théorique
Ecole Polytechnique
17 rue Descartes
F-75005 Paris, France

D. Iagolnitzer
Service de Physique Théorique
Centre d'Études Nucléaires de Saclay
B.P. n° 2
F-91190 Gif-sur-Yvette, France