

THE OFFICIAL MAGAZINE OF THE OCEANOGRAPHY SOCIETY

# Oceanography

#### CITATION

Abbriano, R.M., M.M. Carranza, S.L. Hogle, R.A. Levin, A.N. Netburn, K.L. Seto, S.M. Snyder, SIO280, and P.J.S. Franks. 2011. Deepwater Horizon oil spill: A review of the planktonic response. *Oceanography* 24(3):294–301, <http://dx.doi.org/10.5670/oceanog.2011.80>.

#### COPYRIGHT

This article has been published in *Oceanography*, Volume 24, Number 3, a quarterly journal of The Oceanography Society. Copyright 2011 by The Oceanography Society. All rights reserved.

#### USAGE

Permission is granted to copy this article for use in teaching and research. Republication, systematic reproduction, or collective redistribution of any portion of this article by photocopy machine, reposting, or other means is permitted only with the approval of The Oceanography Society. Send all correspondence to: [info@tos.org](mailto:info@tos.org) or The Oceanography Society, PO Box 1931, Rockville, MD 20849-1931, USA.



# DEEPWATER HORIZON OIL SPILL

A Review  
of the Planktonic Response

BY RAFFAELA M. ABBRIANO, MAGDALENA M. CARRANZA,  
SHANE L. HOGLE, RACHEL A. LEVIN, AMANDA N. NETBURN,  
KATHERINE L. SETO, STEPHANIE M. SNYDER, SIO280,  
AND PETER J.S. FRANKS

NASA image by Jeff Schmaltz,  
MODIS Rapid Response Team

## INTRODUCTION

On April 20, 2010, the explosion of the Deepwater Horizon (DWH) oil rig resulted in the loss of 11 lives and the largest oil spill in US history (Graham et al., 2010) and perhaps the second largest in the world, after the first Gulf War Oil Spill from Kuwait. Over the 84 days following the explosion, an estimated  $6.7 \times 10^5$  mT of Louisiana Sweet Crude oil (United States Government, 2011) and up to 500,000 mT of methane and gases (Joye et al., 2011) were released from 1,480 m below the ocean's surface into the Gulf of Mexico (GoM). As oil continued to escape from the seafloor throughout the summer of 2010, images of oiled wildlife pervaded the news. These pictures, though troubling, only hinted at the fate of the plankton that form the foundation of the GoM ecosystem. This review discusses the potential effects of the DWH oil spill on the overlooked, but extremely important, members of the GoM ecosystem—the plankton. Our assessment is based on data collected in the aftermath of the DWH spill and supplemented with studies from past oil spills when information on the GoM spill was limited or unavailable. The time line we develop traces the spill from a “planktonic perspective,” emphasizing the population dynamics of marine bacteria, phytoplankton, zooplankton, and fish larvae.

## ECONOMICS AND ECOSYSTEM OF THE GULF OF MEXICO

With its white sandy beaches and productive waters, the GoM sustains a number of lucrative industries, including

shrimp, oysters, menhaden, and bluefin tuna. The GoM shrimp fishery was valued at \$367 million dollars in 2008 (NOAA, 2010). The larval stages of shrimp and other organisms are particularly vulnerable in oil-contaminated

waters. Revenue losses from fisheries in Louisiana alone are estimated to be between \$100–200 million (IEM, 2010). Tourism brings in an estimated \$20 billion to the Gulf region (EPA, 2011), and huge losses are expected in the foreseeable future due to avoidance of areas impacted by the Deepwater Horizon disaster.

From April through June, the GoM serves as the sole breeding ground for western Atlantic bluefin tuna, a commercially important and endangered species. Atlantic bluefin tuna exhibit broadcast spawning in the oligotrophic waters of the GOM lower continental slope, including the area of surface oil dispersal (Figure 1; Teo et al., 2007; Muhling et al., 2010). They are the highest valued Atlantic tuna species in the Asian sushi and sashimi markets (\$400 a pound at Tokyo's Tsukiji fish market), and they are overfished.

The GoM is also one of the United States' main sources of oil, with ~ 4,000 oil platforms that collectively produce almost a quarter of America's petroleum (Considine et al., 2004; Trevors and Saier, 2010). Up to 350 natu-

“WHILE STUDIES OF THE DWH OIL SPILL WILL CONTINUE TO PROVIDE NEW INFORMATION IN THE COMING MONTHS AND YEARS, EARLY RESEARCH SHOWS THAT THE PLANKTONIC COMMUNITY EXHIBITS AN ENCOURAGING LEVEL OF RESILIENCE.”

rally occurring seeps distributed throughout the Gulf (Kvenvolden and Cooper, 2003) support microbial assemblages with a well-developed ability to oxidize hydrocarbons (Hazen et al., 2010). Respiration by these oil-degrading bacteria can create thin oxygen-minimum layers close to the seafloor (Bender et al., 2005), potentially affecting the communities that can live in those environments. However, GoM waters are well ventilated, and benthic oil-derived hypoxia tends to be local and transient (Bender et al., 2005).

### APRIL 20, 2010

*The DWH oil platform explodes and sinks 66 km off the coast of Louisiana, allowing oil to escape uncontrolled from the Macondo wellhead.*

Immediately following the explosion of the DWH oil rig, surface slicks of Louisiana Sweet Crude oil




Figure 1. Map of the Gulf of Mexico showing the oil spill area as of June 29, 2010 (shading). Black circles show the probability of collecting at least one Atlantic bluefin tuna larva at sampling stations between 1982 and 2006. Black dashed line shows approximate location of the Loop Current. Adapted from Muhling et al. (2010)

began appearing, covering as much as 75,000 km<sup>2</sup> within one month (Cleveland et al., 2011). This crude oil is a structurally complex, heterogeneous mixture comprised of simple aliphatic hydrocarbons (SAHs), aromatic hydrocarbons (including polycyclic aromatic hydrocarbons, PAHs), resins, and asphaltenes. While dissolution—the chemical stabilization of crude oil components in seawater—is a minor pathway for oil loss compared with evaporation and emulsification, the dissolved fraction of crude oil is most toxic to aquatic species. This water-soluble fraction (WSF) tends to be comprised

of low-molecular-weight aliphatic compounds, aromatic hydrocarbons, and PAHs. All oil types and grades contain WSFs of varying concentrations, but degraded crude varieties tend to have higher concentrations (National Research Council, 2003; Head et al., 2006). WSF components were detected at the DWH blowout with greatest signal intensities near 1,000 and 400 m depth. Although absolute concentrations could not be measured in the water column, it is likely that some portion of the total WSF migrated into the euphotic zone after the DWH spill (Camilli et al., 2010).

The chemical diversity of crude oil

creates niches for bacterial substrate specialists during the course of biodegradation. Through the activities of various specialist bacteria, the composition and concentration of the chemical components of raw crude oil change over time (Dutta and Harayama, 2000; Head et al., 2006). A boom-and-bust cycle of bacterial succession is common, though not certain, after oiling events. A general community progression beginning with consumers of SAHs, followed by the consumers of PAHs and methane (Figure 2), frequently arises after marine oil spills, although this hierarchy can be altered or arrested depending on environmental conditions such as temperature, nutrient concentrations, salinity, and pressure (National Research Council, 2003; Valentine et al., 2010).

While the oil may have been a boon for certain bacterial types, both the oil and oil biodegradation can cause problems for phytoplankton in the immediate vicinity of a spill. Thick, buoyant oil slicks inhibit air-sea gas exchange and light penetration, both essential to photosynthesis and phytoplankton growth (González et al., 2009). The PAHs in the oil also affect phytoplankton growth, with responses ranging from stimulation at low concentrations of oil (1 mg L<sup>-1</sup>) to inhibition at higher concentrations (100 mg L<sup>-1</sup>; Harrison et al., 1986).

Like the phytoplankton, many zooplankton species are sensitive to the chemicals found in the oil. Copepods in direct contact with the spill were likely to experience increased mortality and decreased feeding and reproduction (Suchanek, 1993), potentially allowing blooms of phytoplankton (Figure 2). Tolerance to oil varies by species, and a study of GoM zooplankton communities

---

Raffaela M. Abbriano, Magdalena M. Carranza, Shane L. Hogle, Rachel A. Levin, Amanda N. Netburn, Katherine L. Seto, and Stephanie M. Snyder are students in the biological oceanography class SIO280, and Peter J.S. Franks (pfranks@ucsd.edu) is Professor, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA, USA.

found that mortality tended to be more dependent upon exposure time than concentration of oil (Figure 3), though the highest oil concentrations led to the highest mortalities (50% after 50 hours; Lee and Nicol, 1977). Copepods may also be able to sense and avoid oiled areas (Seuront, 2010), thus reducing their contact and potential mortality (Figure 4).

The same may not be true of Atlantic bluefin tuna larvae. PAHs are known to be highly toxic to larval herring, topsmelt, minnow, and salmon (Petersen and Kristensen, 1998; Heintz et al., 1999; Couillard et al., 2005). Toxic effects include hemorrhages, spinal deformities, growth retardation, and death (Billiard et al., 1999; Carls et al., 1999). While larvae of different fish species exhibit variable sensitivities to PAHs, deformities and/or death are probable for larvae that come into direct contact with the oil.

**MAY 15–JULY 12, 2010**  
*Of the estimated 2.1 million gallons of chemical dispersant (Corexit 9500 and Corexit 9527) released, more than one-third was injected at the wellhead site at depths greater than 1.5 km (Kujawinski et al., 2011).*

The use of subsurface dispersants combined with high pressure and turbulence at the wellhead likely contributed to the formation of the subsurface oil plumes that were unique to the DWH (Kujawinski et al., 2011). Oil dispersants contain both surfactants and solvents, which lower interfacial tension between oil and water boundaries and reduce the tendency of oil to aggregate (Singer et al., 1994). The concentration of dissolved hydrocarbons increases by up to five times when Corexit is applied to crude oil (Fucik et al., 1994). Use of dispersant can expedite the removal of oil from the water column by increasing the rate of

biodegradation by bacteria (Venosa and Holder, 2007). However, the literature is inconclusive as to the toxicological effects of dispersant on bacteria, with studies showing both growth enhancement and retardation (Mulkins-Philips and Stewart, 1974; Bruheim et al., 1999; Garcia et al., 2001). Still, key investigations have cautiously concluded that chemical dispersants are appropriate for enhancing microbial degradation (National Research Council, 2005). By late May, SAH-degrading bacteria, particularly the order *Oceanospirillales* (Hazen et al., 2010) had increased to the point that they represented over 90% of the bacteria inside the plume compared to only 5% in samples acquired outside the plume (Figure 2). Following the bloom of *Oceanospirillales*, another  $\gamma$ -proteobacteria from the genus *Cycloclasticus* known to degrade SAHs and PAHs bloomed in late June





Figure 2. (left panel) Hypothesized (based on previous studies) time lines of oil-spill-driven changes in oil constituents (three lower time lines) and plankton. Vertical arrows indicate times discussed in the text. (right panel) Schematic food web indicating the oil constituents and their pathways into the planktonic food web. Colors correspond to the left panel.

(Kasai et al., 2002; Valentine et al., 2010). Respiration rates due to the degradation of SAHs were measured to be  $\sim 1 \mu\text{M O}_2 \text{ d}^{-1}$  (Camilli et al. 2010), accounting for  $\sim 70\%$  of the low-oxygen anomaly associated with the main plume (Valentine et al., 2010). Models of oxygen depletion due to bacterial respiration predicted hypoxia would occur within a few hundred kilometer radius of the wellhead at depths greater than 1,000 m (Adcroft et al., 2010). Though measurements showed oxygen draw-down, there was no evidence of hypoxia (Kessler et al., 2011).

Biodegradation by bacteria can also incorporate chemically dispersed oil into the food web. Graham et al. (2010) showed a decrease in the  $\delta^{13}\text{C}$  of the small suspended particles (including phytoplankton) and mesozooplankton after the spill toward the  $\delta^{13}\text{C}$  value of the local crude oil. This decrease implies uptake and transfer of components of the oil through the planktonic food web. Bioaccumulation of hydrocarbons at

the base of the food web could increase exposure of higher-trophic-level organisms, with potentially delayed negative effects (Wolfe et al., 1998).

The mixture of crude oil and Corexit has been found to be more toxic to phytoplankton and fish larvae than oil alone (Hsiao et al., 1978: a mixture of four different oils; Middaugh and Whiting, 1995: WSF of No. 2 Fuel Oil). Acute toxicity tests on the larval stages of several invertebrates indigenous to the GoM, such as shrimp, oysters, and crabs, showed that mixtures of the GoM oil from different wells and Corexit were about as toxic as the WSF of the oil alone (Fucik et al., 1994). Most of the toxicity of the oil-Corexit mixture occurred in the first 24 hours, probably due to the increased volatilization of certain components of the oil (Fucik et al., 1994). In some experiments, increases in toxicity could be attributed to the chemical properties of dispersants, which can affect the cellular membranes of planktonic organisms by

increasing permeability to toxic chemicals, disrupting respiration, and causing membrane lysis (Singer et al., 1991).

**JULY 15–SEPTEMBER 19, 2010**  
*After three months, the DWH well was finally capped. Relief wells were drilled, and the wellhead was declared effectively dead.*

At the time of the well capping, studies were beginning to reveal a resilient GoM planktonic ecosystem, reflecting a local adaptation to natural oil seeps. Changes in the structure of bacterial communities were seen mainly in the relatively short-term shifts in dominance from *Oceanospirillales* (SAH degraders) to *Cycloclasticus* (PAH degraders) (Figure 2). Subsequently, almost all of the released methane from the spill was eliminated by a bloom of bacterial methanotrophs in late August that oxidized  $\text{CH}_4$  to  $\text{CO}_2$  (Kessler et al., 2011). Although this rapid oxidation of methane raised concern about the potential for hypoxia, measurements by Kessler et al. (2011) showed that dissolved  $\text{O}_2$  concentrations—while low—were about three times higher than the level at which fisheries would be threatened (Camilli et al., 2010).

Meanwhile, we speculate that phytoplankton community structure changed and biomass increased (Teal and Howarth, 1984) due to a combination of the detrimental effects of oil contamination and the beneficial effects of decreased predation. Similarly, a predicted rapid recovery of the zooplankton would be due to their short generation times and high fecundity, their ability to avoid oily patches (Seuront, 2010), and their recruitment from unaffected areas. In most historical




Figure 3. Mortality of a mixture of coastal zooplankton exposed to the water-soluble fraction (WSF) of No. 2 fuel oil at various concentrations. Note that mortality depends primarily upon exposure time rather than on concentration, though the highest concentrations lead to the highest mortality after 50 h. From data in Lee and Nicol (1977)

oil spills, zooplankton abundance and community structure did not appear to be affected beyond several weeks following the incident (Davenport, 1982; Johansson et al., 1980; Varela et al., 2006). Therefore, we hypothesize that any initial phytoplankton increase or zooplankton population decline would be transient (Figure 2).

Hydrocarbon-degrading microbes provided a link between the plumes of oil generated at depth and the rest of the oceanic food web. Graham et al.'s (2010) study demonstrated that both the mesozooplankton and small particulate matter had incorporated oil-derived (low  $\delta^{13}\text{C}$ ) carbon; the overall impact of this transfer to zooplankton is yet to be determined. However, sampling in the area showed that recovery to pre-spill  $\delta^{13}\text{C}$  values only took about two to four weeks (Graham et al., 2010).

As the oil makes its way up the food chain, its effects on Atlantic bluefin tuna larvae remain unclear. While contact with the oil probably resulted in larval mutation or death, the boom-and-bust cycles in the microbial loop may have led to increases in the main prey sources of these larvae, namely heterotrophic microplankton (Llopiz et al., 2010; Nakagawa et al., 2007). Therefore, for the Atlantic bluefin tuna, being in the right place at the right time may prove to be the deciding factor for the population.

## CONCLUSION

While studies of the DWH oil spill will continue to provide new information in the coming months and years, early research shows that the planktonic community exhibits an encouraging level of resilience. Its pre-existing acclimation to the presence of hydrocarbons, as well

as its diversity and specialization, may have predisposed the community to respond to and even exploit a seemingly catastrophic event such as the DWH oil spill. Specialists within the diverse bacterial communities exhibited rapid boom-and-bust cycles, showing signs of returning to background levels as early as 60 days after the blowout. Although individual phytoplankton species may have experienced relative mortality or enhanced growth, the direct negative effects of oil were probably largely offset by a decrease in predation. Dispersion and degradation of oil in surface seawater, high rates of reproduction of marine planktonic organisms, and circulation and mixing in the ocean may also have contributed to rapid recovery of phytoplankton populations within weeks to months. Zooplankton, due to rapid reproduction and ability to avoid direct contact with oil, may have been minimally affected in the long run. Lastly, Atlantic bluefin tuna would likely suffer significant mortality with direct oil contact, but secondary effects such as an increase in food supply are still to be determined. Therefore, while delayed impacts of bioaccumulation in higher trophic levels may still prove significant, the direct impacts of DWH oil on the planktonic community—marine bacteria, phytoplankton, zooplankton, and fish larvae—provide some hope for a resilient and thriving GoM.

## ACKNOWLEDGEMENTS

The topic of this paper was assigned as a term project to SIO280, the graduate-level introductory biological oceanography course at Scripps Institution of Oceanography taught by Peter Franks. The 37 students,




Figure 4. (A) Movement of a copepod around an uncontaminated sample (open circle). (B) Movement of a copepod around a patch containing the water-soluble fraction of diesel oil, showing patch-avoidance response. (C) Distribution of copepods around a patch of the water-soluble fraction of diesel oil, showing that they avoid entering the patch. Adapted from Seuront (2010)

representing all the disciplines at Scripps, submitted eight group papers that were subsequently synthesized into this paper by the named authors. The remaining 30 authors were: N. Arakawa, C. Archer, A. Beaubien, N. Ben-Aderet,

S. Billheimer, J. Campana, S. Cusatti, X. Fan, C. Feehery, L. Gleason, D. Gonzalez-Paredes, C. Jew, T. Joyce, D. Krause, B. Ludka, J. McQuaid, N. Millan Aguinaga, L. Miller, R. Morrison, C. Perretti, M. Pham, N. Pierce, A. Redman, E. Sanchez Alvarez, E. Sawicka, J. Stiller, B. Thayre, N. Vahora, S. Walkup, and S. Walstead. The authors would like to thank an anonymous reviewer and John Farrington, whose comments led to a much improved manuscript. Particular thanks to Ellen Kappel, who encouraged this venture from the outset. ☒

## REFERENCES

- Adcroft, A., R. Hallberg, J.P. Dunne, B.L. Samuels, J.A. Galt, C.H. Barker, and D. Payton. 2010. Simulations of underwater plumes of dissolved oil in the Gulf of Mexico. *Geophysical Research Letters* 37, L18605, <http://dx.doi.org/10.1029/2010GL044689>.
- Bender, L.C., S.F. DiMarco, M.K. Howard, A.E. Jochens, M.C. Kennicutt II, J.W. Morse, and W.D. Nowlin. 2005. *Understanding the Processes That Maintain the Oxygen Levels in the Deep Gulf of Mexico: Synthesis Report*. Minerals Management Service access number 85080, 6 pp.
- Billiard, S.M., K. Querbach, and P.V. Hodson. 1999. Toxicity of retene to early life stages of two freshwater fish species. *Environmental Toxicology and Chemistry* 18:2,070–2,077, <http://dx.doi.org/10.1002/etc.5620180927>.
- Bruheim, P., H. Bredholt, and K. Eimhjellen. 1999. Effects of surfactant mixtures, including Corexit 9527, on bacterial oxidation of acetate and alkanes in crude oil. *Applied and Environmental Microbiology* 65:1,658–1,661.
- Camilli, R., C.M. Reddy, D. Yoerger, B.A.S. Van Mooy, M.V. Jakuba, J.C. Kinsey, C.P. McIntyre, S.P. Sylva and J.V. Maloney. 2010. Tracking hydrocarbon plume transport and biodegradation at Deepwater Horizon. *Science* 330:201–204, <http://dx.doi.org/10.1126/science.1195223>.
- Carls, M.G., S.D. Rice, and J.E. Hose. 1999. Sensitivity of fish embryos to weathered crude oil. Part I. Low-level exposure during incubation causes malformations, genetic damage, and mortality in larval Pacific herring (*Clupea pallasii*). *Environmental Toxicology and Chemistry* 18:481–493, <http://dx.doi.org/10.1002/etc.5620180317>.
- Cleveland, C.J., M. Hogan, and P. Saundry. 2011. Deepwater Horizon oil spill. In *Encyclopedia of Earth*. C. Cleveland, ed., Environmental Information Coalition, National Council for Science and the Environment, Washington, DC. Available online at: [http://www.eoearth.org/article/Deepwater\\_Horizon\\_oil\\_spill](http://www.eoearth.org/article/Deepwater_Horizon_oil_spill) (accessed July 25, 2011).
- Considine, T., C. Jablonowski, B. Posner, and C. Bishop. 2004. Value of hurricane forecasts to oil and gas producers in the Gulf of Mexico. *Journal of Applied Meteorology* 43:1,270–1,281, [http://dx.doi.org/10.1175/1520-0450\(2004\)043<1270:TVOHFT>2.0.CO;2](http://dx.doi.org/10.1175/1520-0450(2004)043<1270:TVOHFT>2.0.CO;2).
- Couillard, C.M., K. Lee, B. Legare, and T.L. King. 2005. Effect of dispersant on the composition of the water-accommodated fraction of crude oil and its toxicity to larval marine fish. *Environmental Toxicology and Chemistry* 24:1,496–1,504, <http://dx.doi.org/10.1897/04-267R.1>.
- Davenport, J. 1982. Oil and planktonic ecosystems. *Philosophical Transactions of the Royal Society of London, Series B, Biological Sciences* 297:369–384, <http://dx.doi.org/10.1098/rstb.1982.0048>.
- Dutta, T., and S. Harayama. 2000. Fate of crude oil by the combination of photooxidation and biodegradation. *Environmental Science and Technology* 34:1,500–1,505, <http://dx.doi.org/10.1021/es991063o>.
- EPA (Environmental Protection Agency). 2011. *General Facts about the Gulf of Mexico*. Available online at: <http://www.epa.gov/gmpo/about/facts.html> (accessed July 22, 2011).
- Fucik, K.W., K.A. Carr, and B.J. Balcom. 1994. *Dispersed Oil Toxicity Tests with Biological Species Indigenous to the Gulf of Mexico*. OCS Study MMS 94-0021, 97 pp.
- Garcia, J.M., L.Y. Wick, and H. Harms. 2001. Influence of the nonionic surfactant Brij 35 on the bioavailability of solid and sorbed dibenzofuran. *Environmental Science and Technology* 35:2,033–2,039, <http://dx.doi.org/10.1021/es001552k>.
- Gonzalez, J., F.G. Figueiras, M. Aranguren-Gassis, B.G. Crespo, E. Fernandez, X.A.G. Moran, and M. Nieto-Cid. 2009. Effect of a simulated oil spill on natural assemblages of marine phytoplankton enclosed in microcosms. *Estuarine, Coastal, and Shelf Science* 83:265–276, <http://dx.doi.org/10.1016/j.ecss.2009.04.001>.
- Graham, W.M., R.H. Condon, R.H. Carmichael, I. D'Ambra, H.K. Patterson, L.J. Linn, and F.J. Hernandez Jr. 2010. Oil carbon entered the coastal planktonic food web during the Deepwater Horizon oil spill. *Environmental Research Letters* 5, 045301, <http://dx.doi.org/10.1088/1748-9326/5/4/045301>.
- Harrison, P.J., W.P. Cochlan, J.C. Acreman, T.R. Parsons, P.A. Thompson, and H.M. Dovey. 1986. The effects of crude oil and Corexit 9527 on marine phytoplankton in an experimental enclosure. *Marine Environmental Research* 18:93–109, [http://dx.doi.org/10.1016/0141-1136\(86\)90002-4](http://dx.doi.org/10.1016/0141-1136(86)90002-4).
- Hazen, T.C., E.A. Dubinsky, T.Z. DeSantis, G.L. Andersen, Y.M. Piceno, N. Singh, J.K. Jansson, A. Probst, S.E. Borglin, J.L. Fortney, and others. 2010. Sea oil plume enriches indigenous oil-degrading bacteria. *Science* 330:204–207, <http://dx.doi.org/10.1126/science.1195979>.
- Head, I.M., D. Jones, and W.M. Roling. 2006. Marine microorganisms make a meal of oil. *Nature Reviews Microbiology* 4:173–182, <http://dx.doi.org/10.1038/nrmicro1348>.
- Heintz, R.A., J.W. Short, and S.D. Rice. 1999. Sensitivity of fish embryos to weathered crude oil. Part II. Increased mortality of pink salmon (*Oncorhynchus gorbuscha*) embryos incubating downstream from weathered Exxon Valdez crude oil. *Environmental Toxicology and Chemistry* 18:494–503, <http://dx.doi.org/10.1002/etc.5620180318>.
- Hsiao, S.I.C., D.W. Kittle, and M.G. Foy. 1978. Effects of crude oils and the oil dispersant Corexit on primary production of arctic marine phytoplankton and seaweed. *Environmental Pollution* 15:209–221, [http://dx.doi.org/10.1016/0013-9327\(78\)90066-6](http://dx.doi.org/10.1016/0013-9327(78)90066-6).
- IEM (Innovative Emergency Management). 2010. *A Study of the Economic Impact of the Deepwater Horizon Oil Spill*. Available online at: <http://gnoinc.org/news-events/gno-inc-unveils-oil-spill-economic-impact-study> (accessed July 22, 2011).
- Johansson, S., U. Larsson, and P. Boehm. 1980. The Tsesis oil spill: Impact on the pelagic ecosystem. *Marine Pollution Bulletin* 11:284–293, [http://dx.doi.org/10.1016/0025-326X\(80\)90166-6](http://dx.doi.org/10.1016/0025-326X(80)90166-6).
- Joye, S.B., I.R. MacDonald, I. Leifer, and V. Asper. 2011. Magnitude and oxidation potential of hydrocarbon gases released from the BP oil well blowout. *Nature Geoscience*, <http://dx.doi.org/10.1038/NGEO1067>.
- Kasai, Y., H. Kishira, and S. Harayama. 2002. Bacteria belonging to the genus *Cycloclasticus* play a primary role in the degradation of aromatic hydrocarbons released in a marine environment. *Applied Environmental Microbiology* 68:5,625–5,633, <http://dx.doi.org/10.1128/AEM.68.11.5625-5633.2002>.
- Kessler, J.D., D.L. Valentine, M.C. Redmond, M. Du, E.W. Chan, S.D. Mendes, E.W. Quiroz, C.J. Villanueva, S.S. Shusta, L.M. Werra, and others. 2011. A persistent oxygen anomaly reveals the fate of spilled methane in the deep Gulf of Mexico. *Science* 331:312–315, <http://dx.doi.org/10.1126/science.1199697>.
- Kujawinski, E.B., M.C.K. Soule, D.L. Valentine, A.K. Boysen, K. Longnecker, and M.C. Redmond. 2011. Fate of dispersants associated with the Deepwater Horizon



- oil spill. *Environmental Science and Technology* 45:1,298–1,306, <http://dx.doi.org/10.1021/es103838p>.
- Kvenvolden, K.A., and C.K. Cooper. 2003. Natural seepage of crude oil into the marine environment. *Geo-Marine Letters* 23:140–146, <http://dx.doi.org/10.1007/s00367-003-0135-0>.
- Lee, W.Y., and J.A.C. Nicol. 1977. The effects of the water-soluble fractions of No. 2 fuel oil on the survival and behaviour of coastal and oceanic zooplankton. *Environmental Pollution* 12:279–292, [http://dx.doi.org/10.1016/0013-9327\(77\)90022-2](http://dx.doi.org/10.1016/0013-9327(77)90022-2).
- Llopiz, J.K., D.E. Richardson, A. Shiroza, S.L. Smith, and R.K. Cowen. 2010. Distinctions in the diets and distributions of larval tunas and the important role of appendicularians. *Limnology and Oceanography* 55:983–996, <http://dx.doi.org/10.4319/lo.2010.55.3.0983>.
- Middaugh, D.P., and D.D. Whiting. 1995. Responses of embryonic and larval inland silversides, *Menidia beryllina*, to No. 2 fuel oil and oil dispersants in seawater. *Archives of Environmental Contamination and Toxicology* 29:535–539, <http://dx.doi.org/10.1007/BF00208385>.
- Muhling, B.A., J.T. Lamkin, and M.A. Roffer. 2010. Predicting the occurrence of Atlantic bluefin tuna (*Thunnus thynnus*) larvae in the northern Gulf of Mexico: Building a classification model from archival data. *Fisheries Oceanography* 19:526–539, <http://dx.doi.org/10.1111/j.1365-2419.2010.00562.x>.
- Mulkins-Phillips, G.J., and J.E. Stewart. 1974. Effect of four dispersants on biodegradation and growth of bacteria on crude oil. *Applied and Environmental Microbiology* 28:547–552.
- Nakagawa, Y., M. Eguchi, and S. Miyashita. 2007. Pacific bluefin tuna, *Thunnus orientalis*, larvae utilize energy and nutrients of microbial loop. *Aquaculture* 267:83–93, <http://dx.doi.org/10.1016/j.aquaculture.2007.02.024>.
- NOAA (National Oceanic and Atmospheric Administration). 2010. *Fish Stocks in the Gulf of Mexico Fact Sheet*. Available online at: [http://sero.nmfs.noaa.gov/sf/deepwater\\_horizon/Fish\\_economics\\_FACT\\_SHEET.pdf](http://sero.nmfs.noaa.gov/sf/deepwater_horizon/Fish_economics_FACT_SHEET.pdf) (accessed July 22, 2011).
- National Research Council. 2003. *Oil in the Sea III: Inputs, Fates, and Effects*. Committee on Oil in the Sea: Inputs Fates and Effects. The National Academies Press, 280 pp. Available online at: <http://www.nap.edu/openbook.php?isbn=0309084385> (accessed July 22, 2011).
- National Research Council. 2005. *Oil Spill Dispersants: Efficacy and Effects*. Committee on Understanding Oil Spill Dispersants: Efficacy and Effects. 2005. The National Academies Press. Available online at: [http://www.nap.edu/openbook.php?record\\_id=11283](http://www.nap.edu/openbook.php?record_id=11283) (accessed July 22, 2011).
- Petersen, G.I., and P. Kristensen. 1998. Bioaccumulation of lipophilic substances in fish early life stages. *Environmental Toxicology and Chemistry* 17:1,385–1,395, [http://dx.doi.org/10.1897/1551-5028\(1998\)017<1385:BOLSIF>2.3.CO;2](http://dx.doi.org/10.1897/1551-5028(1998)017<1385:BOLSIF>2.3.CO;2).
- Seuront, L. 2010. Zooplankton avoidance behaviour as a response to point sources of hydrocarbon-contaminated water. *Marine and Freshwater Research* 61:263–270, <http://dx.doi.org/10.1071/MF09055>.
- Singer, M.M., S. George, S. Jacobson, I. Lee, R.S. Tjeerdema, and M.L. Sowby. 1994. Comparative effects of oil dispersants to the early life stages of topsmelt (*Atherinops affinis*) and kelp (*Macrocystis pyrifera*). *Environmental Toxicology and Chemistry* 13:649–655, <http://dx.doi.org/10.1002/etc.5620130415>.
- Singer, M.M., D.L. Smalheer, and R.S. Tjeerdema. 1991. Effects of spiked exposure to an oil dispersant on the early life stages of four marine species. *Environmental Toxicology and Chemistry* 10:1,367–1,374, [http://dx.doi.org/10.1897/1552-8618\(1991\)10\[1367:EOSETA\]2.0.CO;2](http://dx.doi.org/10.1897/1552-8618(1991)10[1367:EOSETA]2.0.CO;2).
- Suchanek, T.H. 1993. Oil impacts on marine invertebrate populations and communities. *American Zoologist* 33:510–523, <http://dx.doi.org/10.1093/icb/33.6.510>.
- Teal, J.M., and R.W. Howarth. 1984. Oil spill studies: A review of ecological effects. *Environmental Management* 8:27–44, <http://dx.doi.org/10.1007/BF01867871>.
- Teo, S.L.H., A.M. Boustany, and B.A. Block. 2007. Oceanographic preferences of Atlantic bluefin tuna, *Thunnus thynnus*, on their Gulf of Mexico breeding grounds. *Marine Biology* 152:1,105–1,119, <http://dx.doi.org/10.1007/s00227-007-0758-1>.
- Trevors, J.T., and M.H. Saier. 2010. The legacy of oil spills. *Water, Air, & Soil Pollution* 211:1–3, <http://dx.doi.org/10.1007/s11270-010-0527-5>.
- United States Government. 2011. *One Year Later Press Pack*. Available online at: <http://www.restorethegulf.gov/release/2011/04/10/one-year-later-press-pack> (accessed July 25, 2011).
- Valentine, D.L., J.D. Kessler, M.C. Redmond, S.D. Mendes, M.B. Heintz, C. Farwell, L. Hu, F.S. Kinnaman, S. Yvon-Lewis, M. Du, and others. 2010. Propane respiration jump-starts microbial response to a deep oil spill. *Science* 330:208–211, <http://dx.doi.org/10.1126/science.1196830>.
- Varela, M., A. Bode, J. Lorenzo, M.T. Álvarez-Ossorio, A. Miranda, T. Patrocinio, R. Anadón, L. Viesca, N. Rodríguez, L. Valdéz, and others. 2006. The effect of the “Prestige” oil spill on the plankton of the N-NW Spanish coast. *Marine Pollution Bulletin* 53:272–286, <http://dx.doi.org/10.1016/j.marpolbul.2005.10.005>.
- Venosa, A.D., and E.L. Holder. 2007. Biodegradability of dispersed crude oil at two different temperatures. *Marine Pollution Bulletin* 54:545–553, <http://dx.doi.org/10.1016/j.marpolbul.2006.12.013>.
- Wolfe, M.F., J.A. Schlosser, G.J.B. Schwartz, S. Singaram, E.E. Mielbrecht, R.S. Tjeerdema, and M.L. Sowby. 1998. Influence of dispersants on the bioavailability and trophic transfer of petroleum hydrocarbons to primary levels of a marine food chain. *Aquatic Toxicology* 42:211–227, [http://dx.doi.org/10.1016/S0166-445X\(97\)00096-9](http://dx.doi.org/10.1016/S0166-445X(97)00096-9).