

UC Davis

UC Davis Previously Published Works

Title

Demazure crystals, Kirillov-Reshetikhin crystals, and the energy function

Permalink

<https://escholarship.org/uc/item/35d851r4>

Journal

The Electronic Journal of Combinatorics, 19(2)

Authors

Schilling, Anne
Tingley, Peter

Publication Date

2011-04-12

Peer reviewed

DEMAZURE CRYSTALS, KIRILLOV–RESHETIKHIN CRYSTALS, AND THE ENERGY FUNCTION

ANNE SCHILLING AND PETER TINGLEY

ABSTRACT. It has previously been shown that, at least for non-exceptional Kac–Moody Lie algebras, there is a close connection between Demazure crystals and tensor products of Kirillov–Reshetikhin crystals. In particular, certain Demazure crystals are isomorphic as classical crystals to tensor products of Kirillov–Reshetikhin crystals via a canonically chosen isomorphism. Here we show that this isomorphism intertwines the natural affine grading on Demazure crystals with a combinatorially defined energy function. As a consequence, we obtain a formula of the Demazure character in terms of the energy function, which has applications to Macdonald polynomials and q -deformed Whittaker functions.

CONTENTS

1. Introduction	2
1.1. Results	3
1.2. Applications	4
1.3. Exceptional types	4
1.4. Outline	4
Acknowledgments	4
2. Kac–Moody algebras and Crystals	5
2.1. General setup	5
2.2. $U_q(\mathfrak{g})$ crystals	5
2.3. $U'_q(\mathfrak{g})$ crystals	5
2.4. Tensor products of crystals	6
2.5. Abstract crystals	6
2.6. Perfect crystals	6
2.7. Kirillov–Reshetikhin modules and their crystals	7
2.8. Extended affine Weyl group	8
2.9. Demazure modules and crystals	10
3. Kashiwara–Nakashima tableaux and branching rules	11
3.1. Kashiwara–Nakashima tableaux	11
3.2. Branching rules and \pm diagrams	12
3.3. Nested \pm diagrams	14
4. Realizations of KR crystals	15
4.1. Classical decompositions of KR crystals	16
4.2. Classically irreducible KR crystals (case (IRR))	16
4.3. KR crystals constructed via Dynkin automorphisms (case (AUT))	16
4.4. $B^{r,s}$ of type $C_n^{(1)}$ for $r < n$ (in case (VIR))	17

1991 *Mathematics Subject Classification.* Primary 81R50, 81R10; Secondary: 05E99.

4.5.	$B^{n,s}$ in type $B_n^{(1)}$ (in case (VIR))	19
4.6.	The KR crystals of type $A_{2n}^{(2)}$ and $D_{n+1}^{(2)}$ in case (VIR)	19
5.	Energy functions	20
5.1.	The function E^{int}	20
5.2.	The D function	21
6.	Perfect KR crystals and Demazure crystals	22
7.	The affine grading via the energy function	25
8.	Generalizations to other tensor products of KR crystals	27
9.	Applications	29
9.1.	Demazure characters	29
9.2.	Nonsymmetric Macdonald polynomials	29
9.3.	q -deformed Whittaker functions	32
	References	33

1. INTRODUCTION

Kashiwara’s theory of crystal bases [28] provides a remarkable combinatorial tool for studying highest weight representations of symmetrizable Kac–Moody algebras and their quantizations. Here we consider finite-dimensional representations of the quantized universal enveloping algebra $U'_q(\mathfrak{g})$ corresponding to the derived algebra \mathfrak{g}' of an affine Kac–Moody algebra. These representations do not extend to representations of $U_q(\mathfrak{g})$, but one can nonetheless define the notion of a crystal basis. In this setting crystal bases do not always exist, but there is an important class of finite-dimensional modules for $U'_q(\mathfrak{g})$ that are known to admit crystal bases: tensor products of the Kirillov–Reshetikhin modules $W^{r,s}$ from [34] (denoted $W(s\omega_r)$ in that paper), where r is a node in the classical Dynkin diagram and s is a positive integer.

The modules $W^{r,s}$ were first conjectured to admit crystal bases $B^{r,s}$ in [17, Conjecture 2.1], and moreover it was conjectured that these crystals are perfect whenever s is a multiple of a particular constant c_r (perfectness is a technical condition which allows one to use the finite crystal to construct highest weight crystals, see [25]). This conjecture has now been proven in all non-exceptional cases (see [49, 50] for a proof that the crystals exist, and [11, Theorem 1.2] for a proof that they are perfect). We call $B^{r,s}$ a Kirillov–Reshetikhin (KR) crystal.

The perfectness of KR crystals ensures that they are related to highest weight affine crystals via the construction in [25]. In [31], Kashiwara proposed that this relationship is connected to the theory of Demazure crystals [27, 42], by conjecturing that perfect KR crystals are isomorphic as classical crystals to certain Demazure crystals (which are subcrystals of affine highest weight crystals). This was proven in most cases in [8, 9]. More relations between Demazure crystals and tensor products of perfect KR crystals were investigated in [36, 37, 38, 12, 47].

There is a natural grading deg on a highest weight affine crystal $B(\Lambda)$, where $\text{deg}(b)$ records the number of f_0 in a string of f_i ’s that act on the highest weight element to give b (this is well-defined by weight considerations). Due to the ideas discussed above, it seems natural that this grading should transfer to a grading on a tensor product of KR crystals.

Gradings on tensor products of KR crystals have in fact been studied, and are usually referred to as “energy functions.” They are vast generalizations of the major index statistics on words, which can be viewed as elements in $(B^{1,1})^{\otimes L}$. The idea dates to the earliest works on perfect crystals [25, 26], and was expanded in [51] following conjectural definitions in [16]. A function D , which we will refer to as the D -function, is defined as a sum involving local energy functions for each pair of factors in the tensor product and an ‘intrinsic energy’ of each factor. It has been suggested that there is a simple global characterization of intrinsic energy related to the affine grading on a corresponding highest weight crystal (see [52, Section 2.5], [16, Proof of Proposition 3.9]). However, as far as we know, it has previously never been shown that the explicit definition of intrinsic energy actually satisfies this condition.

1.1. Results. In the present work, we restrict to non-exceptional type (i.e. all affine Kac–Moody algebras except $A_2^{(2)}$, $G_2^{(1)}$, $F_4^{(1)}$, $E_6^{(1)}$, $E_7^{(1)}$, $E_8^{(1)}$, $E_6^{(2)}$ and $D_4^{(3)}$), where KR crystals are known to exist. The relationship between KR crystals and Demazure crystals discussed above can be enhanced: There is in fact a unique isomorphism of classical crystals between any tensor product $B = B_1 \otimes \cdots \otimes B_k$ of perfect level ℓ KR crystals and a certain Demazure crystal such that all 0 arrows on the Demazure side correspond to 0 arrows on the KR sides (although there are more 0 arrows in the KR crystal). See Theorem 6.1 for the precise statement. This was proven by Fourier, Shimozono and the first author in [12, Theorem 4.4] under certain assumptions since at the time KR crystals were not yet known to exist. Here we point out that in most cases the assumptions from [12] follow from [49, 50, 10, 11]. In the remaining cases of type $A_{2n}^{(2)}$ and the spin nodes for type $D_n^{(1)}$ we prove the statement separately, thereby firmly establishing this relationship between KR crystals and Demazure crystals in all non-exceptional types.

We then establish the correspondence between the D function and the affine grading discussed earlier in this introduction. That is, we show that the unique isomorphism discussed above intertwines the basic grading on the Demazure crystal with the D -function on the KR crystal, up to a shift (i.e. addition of a global constant). This also allows for a new characterization of the D function. Define the intrinsic energy function E^{int} on B by letting $E^{\text{int}}(b)$ record the minimal number of f_0 in a path from a certain fixed $u \in B$ to b . We show that E^{int} agrees with the D -function up to a shift (i.e. addition of a global constant). The isomorphism between KR crystals and Demazure crystals actually intertwines the basic grading with E^{int} exactly. In particular this shows that for any Demazure crystal arrow (see Remark 7.7) the energy changes by 1 on the corresponding classical crystal components.

We also consider the more general setting when B is a tensor product of KR crystals which are not assumed to be perfect or of the same level. The D function is still well-defined, and we give a precise relationship between D and the affine grading on a related direct sum of highest weight modules in Corollaries 8.3 and 8.4. In this case we no longer give an interpretation in terms of Demazure modules, although for tensor products of perfect crystals of various levels it was subsequently shown by Naoi [47] that the result is the disjoint union of Demazure crystals; see also Remark 8.5.

1.2. Applications. Our results express the characters of certain Demazure modules in terms of the intrinsic energy on a related tensor product of KR crystals (see Corollary 9.1). This has potential applications whenever those Demazure characters appear.

For untwisted simply-laced root systems, Ion [21], generalizing results of Sanderson [53] in type A , showed that the specializations $E_\lambda(q, 0)$ of nonsymmetric Macdonald polynomials at $t = 0$ coincide with specializations of Demazure characters of level one affine integrable modules. If λ is anti-dominant, then $E_\lambda(q, 0)$ is actually a symmetric Macdonald polynomial $P_\lambda(q, 0)$. In this case, the relevant Demazure module is associated to a tensor product B of level one KR crystals as above, so our results imply that $P_\lambda(q, 0)$ is the character of B , where the powers of q are given by $-D$, see Corollary 9.5. It follows that the coefficients in the expansion of $P_\lambda(q, 0)$ in terms of the irreducible characters are the one-dimensional configuration sums defined in terms of the intrinsic energy in [16].

There is also a relation between Demazure characters and q -deformed Whittaker functions for \mathfrak{gl}_n [13, Theorem 3.2]. Hence our results allow one to study Whittaker functions via KR crystals.

1.3. Exceptional types. KR crystals are still expected to exist in exceptional types, although in most cases this has not yet been established (see for example [22, 30, 32, 56] for some cases where it has). Furthermore, it is expected that the relationship between KR crystals and Demazure crystals holds in general (see [31] for the conjecture, and [8, 9] for a proof that it holds in some exceptional cases). Our expectation is that Theorem 6.1 and Corollary 7.6 would also continue to hold in all cases.

1.4. Outline. In Section 2 we briefly review the theory of crystals. In Section 3 we review combinatorial models for non-exceptional finite type crystals and their branching rules. Section 4 is devoted to combinatorial models for KR crystals. In Section 5, we introduce the two energy functions E^{int} and D . Section 6 discusses the isomorphism between Demazure crystals and tensor products of perfect KR crystals. In Section 7 we relate the basic grading on a Demazure crystal with the energy function on a tensor product of KR crystals and show that the two energy functions agree up to a shift (Theorems 7.4 and 7.5). Some of these results are generalized to tensor products of (not necessarily perfect) KR crystals of different level in Section 8. In Section 9 we discuss applications to Demazure characters, nonsymmetric Macdonald polynomials, and Whittaker functions.

Acknowledgments. We would like to thank Dan Bump, Ghislain Fourier, Stavros Koussidis, Cristian Lenart, Sergey Oblezin, and Masato Okado for enlightening discussions, and Nicolas Thiéry for his help with Sage. Most of the KR crystals have been implemented in the open-source mathematics system Sage (sagemath.org) by the first author. A thematic tutorial on crystals and affine crystals written by Dan Bump and the first author is available in Sage. A copy can be found at <http://www.math.ucdavis.edu/~anne/sage/lie/crystals.html> and http://www.math.ucdavis.edu/~anne/sage/lie/affine_crystals.html.

We would like to thank the Fields Institute in Toronto and the Hausdorff Institut in Bonn for their hospitality, where part of this work was done. AS was in part supported by NSF grants DMS-0652641, DMS-0652652, and DMS-1001256. PT was partially supported by NSF grant DMS-0902649.

2. KAC–MOODY ALGEBRAS AND CRYSTALS

2.1. General setup. Let \mathfrak{g} be a Kac–Moody algebra. Let $\Gamma = (I, E)$ be its Dynkin diagram, where I is the set of vertices and E the set of edges. Let Δ be the corresponding root system and $\{\alpha_i \mid i \in I\}$ the set of simple roots. Let P, Q, P^\vee , and Q^\vee denote the weight lattice, root lattice, coweight lattice, and coroot lattice respectively.

Let $U_q(\mathfrak{g})$ be the corresponding quantum enveloping algebra over $\mathbb{Q}(q)$. Let $\{E_i, F_i\}_{i \in I}$ be the standard elements in $U_q(\mathfrak{g})$ corresponding to the Chevalley generators of the derived algebra \mathfrak{g}' . We recall the triangular decomposition

$$U_q(\mathfrak{g}) \cong U_q(\mathfrak{g})^{<0} \otimes U_q(\mathfrak{g})^0 \otimes U_q(\mathfrak{g})^{>0},$$

where $U_q(\mathfrak{g})^{<0}$ is the subalgebra generated by the F_i , $U_q(\mathfrak{g})^{>0}$ is the subalgebra generated by the E_i , $U_q(\mathfrak{g})^0$ is the abelian group algebra generated by the usual elements K_w for $w \in P^\vee$, and the isomorphism is as vector spaces. Let $U'_q(\mathfrak{g})$ be the subalgebra generated by E_i, F_i and $K_i := K_{\alpha_i^\vee}$ for $i \in I$.

We are particularly interested in the case when \mathfrak{g} is of affine type. In that situation, we use the following conventions: Λ_i denotes the fundamental weight corresponding to $i \in I$. For $i \in I \setminus \{0\}$, ω_i denotes the fundamental weight corresponding to that node in the related finite type Lie algebra. The corresponding finite type weight lattice and Weyl group are denoted \overline{P} and \overline{W} , respectively.

2.2. $U_q(\mathfrak{g})$ crystals. Here we give a very quick review, and refer the reader to [18] more details. For us, a crystal is a nonempty set B along with operators $e_i : B \rightarrow B \cup \{0\}$ and $f_i : B \rightarrow B \cup \{0\}$ for $i \in I$, which satisfy some conditions. The set B records certain combinatorial data associated to a representation V of $U_q(\mathfrak{g})$, and the operators e_i and f_i correspond to the Chevalley generators E_i and F_i . The relationship between the crystal B and the module V can be made precise using the notion of a crystal basis for V .

Often the definition of a crystal includes three functions $\text{wt}, \varphi, \varepsilon : B \rightarrow P$, where P is the weight lattice. In the case of crystals of integrable modules, these functions can be recovered (up to a global shift in a null direction in cases where the Cartan matrix is not invertible) from the knowledge of e_i and f_i , as we discuss in a slightly different context in Section 2.3.

An important theorem of Kashiwara shows that every integrable $U_q(\mathfrak{g})$ highest weight module $V(\lambda)$ has a crystal basis and hence a corresponding crystal $B(\lambda)$.

2.3. $U'_q(\mathfrak{g})$ crystals. In the case when the Cartan matrix is not invertible, one can define an extended notion of $U'_q(\mathfrak{g})$ crystal bases and crystals that includes some cases which do not lift to $U_q(\mathfrak{g})$ crystals. Note however that not all integrable $U'_q(\mathfrak{g})$ representations have corresponding crystals. See e.g. [30]. We consider only integrable crystals B , i.e. crystals where each e_i, f_i acts locally nilpotently. Define a weight function on such a B as follows: First set

$$\begin{aligned} \varepsilon_i(b) &:= \max\{m \mid e_i^m(b) \neq 0\}, \\ \varphi_i(b) &:= \max\{m \mid f_i^m(b) \neq 0\}. \end{aligned}$$

Let Λ_i be the fundamental weight associated to $i \in I$. For each $b \in B$, define

$$(i) \quad \varphi(b) := \sum_{i \in I} \varphi_i(b) \Lambda_i,$$

$$(ii) \quad \varepsilon(b) := \sum_{i \in I} \varepsilon_i(b) \Lambda_i,$$

$$(iii) \quad \text{wt}(b) := \varphi(b) - \varepsilon(b).$$

Then $\text{wt}(b)$ corresponds to the classical weight grading of the $U_q(\mathfrak{g})$ module associated to B , and is referred to as the weight function. Notice that $\text{wt}(b)$ is always in

$$(2.1) \quad P' := \text{span}\{\Lambda_i : i \in I\}.$$

If the Cartan matrix of \mathfrak{g} is not invertible, P' is a proper sublattice of P .

Remark 2.1. One would expect that f_i should have weight $-\alpha_i$. This is true for $U_q(\mathfrak{g})$ crystals, but for $U'_q(\mathfrak{g})$ crystals the weight of f_i is actually the projection of $-\alpha_i$ onto P' under the projection that sends null roots to 0.

2.4. Tensor products of crystals. The tensor product rule for $U'_q(\mathfrak{g})$ or $U_q(\mathfrak{g})$ modules leads to a tensor product rule for the corresponding crystals. Following [12], we use the opposite conventions from Kashiwara [28]. If A and B are two crystals, the tensor product $A \otimes B$ is the crystal whose underlying set is the Cartesian set $\{a \otimes b \mid a \in A, b \in B\}$ with operators e_i and f_i defined by:

$$(2.2) \quad \begin{aligned} e_i(a \otimes b) &= \begin{cases} e_i(a) \otimes b & \text{if } \varepsilon_i(a) > \varphi_i(b), \\ a \otimes e_i(b) & \text{otherwise,} \end{cases} \\ f_i(a \otimes b) &= \begin{cases} f_i(a) \otimes b & \text{if } \varepsilon_i(a) \geq \varphi_i(b), \\ a \otimes f_i(b) & \text{otherwise.} \end{cases} \end{aligned}$$

2.5. Abstract crystals.

Definition 2.2. Let \mathfrak{g} be a Kac–Moody algebra with Dynkin diagram $\Gamma = (I, E)$. Let B be a nonempty set with operators $e_i, f_i : B \rightarrow B \sqcup \{0\}$. We say B is a *regular abstract crystal* of type \mathfrak{g} if, for each pair $i \neq j \in I$, B along with the operators e_i, f_i, e_j, f_j is a union of (possibly infinitely many) integrable highest weight $U_q(\mathfrak{g}_{i,j})$ crystals, where $\mathfrak{g}_{i,j}$ is the Lie algebra with Dynkin diagram containing i and j , and all edges between them.

Remark 2.3. If \mathfrak{g} is an affine algebra other than $\widehat{\mathfrak{sl}}_2$, then any $U'_q(\mathfrak{g})$ crystal is a regular abstract crystal of type \mathfrak{g} .

2.6. Perfect crystals. In this section, \mathfrak{g} is of affine type. Let $c = \sum_{i \in I} a_i^\vee \alpha_i^\vee$ be the canonical central element associated to \mathfrak{g} and P^+ the set of dominant weights, that is, $P^+ = \{\Lambda \in P \mid \Lambda(\alpha_i^\vee) \in \mathbb{Z}_{\geq 0}\}$. We define the *level* of $\Lambda \in P^+$ by $\text{lev}(\Lambda) := \Lambda(c)$. For each $\ell \in \mathbb{Z}$, we consider the sets

$$P_\ell = \{\Lambda \in P \mid \text{lev}(\Lambda) = \ell\} \quad \text{and} \quad P_\ell^+ = \{\Lambda \in P^+ \mid \text{lev}(\Lambda) = \ell\},$$

the sets of level- ℓ weights and level- ℓ dominant weights respectively. Note that $P_\ell^+ = \emptyset$ if $\ell < 0$. The following important notion was introduced in [25].

Definition 2.4. For a positive integer $\ell > 0$, a $U'_q(\mathfrak{g})$ -crystal B is called a *perfect crystal of level ℓ* if the following conditions are satisfied:

- (i) B is isomorphic to the crystal graph of a finite-dimensional $U'_q(\mathfrak{g})$ -module.
- (ii) $B \otimes B$ is connected.
- (iii) There exists a $\lambda \in \overline{P}$, such that $\text{wt}(B) \subset \lambda + \sum_{i \in I \setminus \{0\}} \mathbb{Z}_{\leq 0} \alpha_i$ and there is a unique element in B of classical weight λ .




FIGURE 1. Examples of perfect and non-perfect $U'_q(\mathfrak{g})$ crystals. One can verify that the crystal on the right is perfect of level 1, and the crystal on the left is not perfect of any level. These crystals are both KR crystals, as indicated, and the vertices are indexed by KN tableaux as in Section 3.1. In fact, all known finite $U'_q(\mathfrak{g})$ crystals are KR crystals or tensor products of KR crystals.

- (iv) $\forall b \in B$, $\text{lev}(\varepsilon(b)) \geq \ell$.
- (v) $\forall \Lambda \in P_\ell^+$, there exist unique elements $b_\Lambda, b^\Lambda \in B$, such that

$$\varepsilon(b_\Lambda) = \Lambda = \varphi(b^\Lambda).$$

Examples of a perfect and nonperfect crystal is given in Figure 1.

2.7. Kirillov–Reshetikhin modules and their crystals. In this section \mathfrak{g} is of affine type. The Kirillov–Reshetikhin modules were first introduced for the Yangian of \mathfrak{g}' in [34]. One can characterize the KR module $W^{r,s}$ for $U'_q(\mathfrak{g})$ [5, 6], where $r \in I \setminus \{0\}$ and $s \geq 1$, as the irreducible representations of $U'_q(\mathfrak{g})$ whose Drinfeld polynomials are given by

$$(2.3) \quad P_i(u) = \begin{cases} (1 - q_i^{1-s}u)(1 - q_i^{3-s}u) \cdots (1 - q_i^{s-1}u) & \text{if } i = r, \\ 1 & \text{otherwise.} \end{cases}$$

Here $q_i = q^{(\alpha_i|\alpha_i)/2}$.

It was shown in [26] for type $A_n^{(1)}$ and many special cases, and in [49, 50] for general non-exceptional types, that the modules $W^{r,s}$ have crystal bases. We denote the resulting crystals by $B^{r,s}$, and refer to them as *KR crystals*.

Theorem 2.5. [50, 11] *In all non-exceptional types, $W^{r,s}$ has a crystal base $B^{r,s}$. Furthermore, if s is a multiple of c_r (see Figure 4) the resulting crystals are perfect. \square*

The following is a slight strengthening of results from [10].

Lemma 2.6. *Let \mathfrak{g} be a non-exceptional affine Kac–Moody algebra with index set $I = \{0, 1, \dots, n\}$. Fix $r \in I \setminus \{0\}$ and $s > 0$. Then any regular abstract crystal B (see Definition 2.2) of type \mathfrak{g} which is isomorphic to $B^{r,s}$ as a $\{1, 2, \dots, n\}$ -crystal is also isomorphic to $B^{r,s}$ as an I -crystal.*

Proof. First, recall that any finite type crystal C is uniquely determined by its character

$$\text{ch}(C) := \sum_{c \in C} e^{\text{wt}^{\text{fin}}(c)},$$

where wt^{fin} denotes the weight as a $\{1, 2, \dots, n\}$ -crystal. Since B is isomorphic to $B^{r,s}$ as a $\{1, 2, \dots, n\}$ -crystal, it is finite and hence of level 0. Thus $\text{wt}(b)$ can be recovered from $\text{wt}^{\text{fin}}(b)$. In particular, we can recover the character of B as an $I \setminus \{j\}$ crystal for all $j \in I$, and from there recover the isomorphism class of B as an $I \setminus \{j\}$ -crystal.

The lemma now follows by the following uniqueness results from [50, 10]:

- By [50, Proposition 6.1] (for most nodes) and [10] (for exceptional nodes), in types $D_n^{(1)}$, $B_n^{(1)}$ and $A_{2n-1}^{(2)}$, any regular abstract crystal B which is isomorphic to $B^{r,s}$ as both a $\{1, 2, \dots, n\}$ -crystal and a $\{0, 2, \dots, n\}$ -crystal is isomorphic to $B^{r,s}$ as an affine crystal.
- As in [10, Sections 5.2 and 6.1], in types $C_n^{(1)}$, $D_{n+1}^{(2)}$ and $A_{2n}^{(2)}$, any regular abstract crystal B which is isomorphic to $B^{r,s}$ as both a $\{1, 2, \dots, n\}$ -crystal and a $\{0, 1, \dots, n-1\}$ -crystal is isomorphic to $B^{r,s}$ as an affine crystal. \square

By [39, Proposition 3.8], a tensor product $B = B^{r_1, s_1} \otimes \dots \otimes B^{r_N, s_N}$ of KR crystals is connected. We refer to such a B as a *composite KR crystal*. As in [25], if the factors are all perfect KR crystals of the same level, then $B = B^{r_1, \ell c_{r_1}} \otimes \dots \otimes B^{r_N, \ell c_{r_N}}$ is also perfect of level ℓ . We refer to such a perfect crystal as a *composite KR crystal of level ℓ* .

Explicit combinatorial models for KR crystals of non-exceptional type were constructed in [10], and will be reviewed in Section 4. Two examples are given in Figure 1.

2.8. Extended affine Weyl group. Write the null root as $\delta = \sum_{i \in I} a_i \alpha_i$. Let $\theta = \delta - a_0 \alpha_0$. As in [23, Section 6.4], θ is a root in the finite type root system corresponding to $I \setminus \{0\}$. Following [16], for each $i \in I \setminus \{0\}$, define $c_i = \max(1, a_i/a_i^\vee)$. It turns out that $c_i = 1$ except for $c_i = 2$ for: $\mathfrak{g} = B_n^{(1)}$ and $i = n$, $\mathfrak{g} = C_n^{(1)}$ and $1 \leq i \leq n-1$, $\mathfrak{g} = F_4^{(1)}$ and $i = 3, 4$, and $c_2 = 3$ for $\mathfrak{g} = G_2^{(1)}$. Here we use Kac's

indexing of affine Dynkin diagrams from [23, Table Fin, Aff1 and Aff2]. Consider the sublattices of \overline{P} given by

$$M = \bigoplus_{i \in I \setminus \{0\}} \mathbb{Z}c_i\alpha_i = \mathbb{Z}\overline{W} \cdot \theta/a_0,$$

$$\widetilde{M} = \bigoplus_{i \in I \setminus \{0\}} \mathbb{Z}c_i\omega_i.$$

Let \overline{W} be the finite type Weyl group for the Dynkin diagram $I \setminus \{0\}$, which acts on \overline{P} by linearizing the rules $s_i\lambda = \lambda - \langle \alpha_i^\vee, \lambda \rangle \alpha_i$. Clearly $M \subset \widetilde{M}$ and the action of \overline{W} on \overline{P} restricts to actions on M and \widetilde{M} . Let $T(\widetilde{M})$ (resp. $T(M)$) be the subgroup of $T(\overline{P})$ generated by the translations t_λ by $\lambda \in \widetilde{M}$ (resp. $\lambda \in M$).

There is an isomorphism [23, Prop. 6.5]

$$(2.4) \quad W \cong \overline{W} \ltimes T(M)$$

as subgroups of $\text{Aut}(\overline{P})$, where W is the affine Weyl group. Under this isomorphism we have

$$(2.5) \quad s_0 = t_{\theta/a_0} s_\theta,$$

where s_θ is the reflection corresponding to the root θ . Define $\theta^\vee \in \mathfrak{h}^*$ so that $s_\theta(\lambda) = \lambda - \langle \theta^\vee, \lambda \rangle \theta$.

Define the extended affine Weyl group to be the subgroup of $\text{Aut}(P)$ given by

$$(2.6) \quad \widetilde{W} = \overline{W} \ltimes T(\widetilde{M}).$$

Let $C \subset \overline{P} \otimes_{\mathbb{Z}} \mathbb{R}$ be the fundamental chamber, the set of elements λ such that $\langle \alpha_i^\vee, \lambda \rangle \geq 0$ for all $i \in I \setminus \{0\}$, and $\langle \theta^\vee, \lambda \rangle \leq 1/a_0$.

Let $\Sigma \subset \widetilde{W}$ be the subgroup of \widetilde{W} consisting of those elements that send C into itself. Then $\widetilde{W} = W\Sigma$, and in particular every element $x \in \widetilde{W}$ can be written uniquely as

$$(2.7) \quad x = w\tau$$

for some $w \in W$ and $\tau \in \Sigma$.

The usual affine Weyl group W is a normal subgroup of \widetilde{W} , so Σ acts on W by conjugation. Each $\tau \in \Sigma$ induces an automorphism (also denoted τ) of the affine Dynkin diagram Γ , which is characterized as the unique automorphism so that:

$$(2.8) \quad \tau s_i \tau^{-1} = s_{\tau(i)} \quad \text{for each } i \in I.$$

Remark 2.7. When \mathfrak{g} is of untwisted type, $M \cong Q^\vee$, $\widetilde{M} \cong P^\vee$, with the isomorphism ν given by $c_i\omega_i = \nu(\omega_i^\vee)$, and $c_i\alpha_i = \nu(\alpha_i^\vee)$ for $i \in I \setminus \{0\}$.

Lemma 2.8. Fix \mathfrak{g} with affine Dynkin diagram Γ , and let $\tau \in \text{Aut}(\Gamma)$. Then $\tau \in \Sigma$ if and only if there exists $w_\tau \in \overline{W}$ such that the following diagram commutes

$$\begin{array}{ccc} I & \xrightarrow{\iota} & \Delta \\ \downarrow \tau & & \downarrow w_\tau \\ I & \xrightarrow{\iota} & \Delta, \end{array}$$

where Δ is the underlying finite type root system, and ι is the map that takes i to α_i for all $i \neq 0$ and takes 0 to $-\theta$.




FIGURE 2. Standard crystals. The boxes represent the vertices of the crystal, and each arrow labeled i shows the action of f_i .

Remark 2.10. We mainly consider the case when \mathfrak{g} is affine, and the following Demazure modules: Fix an anti-dominant weight $\mu \in \overline{P}$, and write $t_\mu \in \widetilde{W}$ as $t_\mu = w\tau$, where $w \in W$ and $\tau \in \Sigma$. For a dominant $\Lambda \in P$, we consider $V_w(\tau(\Lambda))$ and its crystal $B_w(\tau(\Lambda))$. In [12] these were denoted by $V_\mu(\Lambda)$ and $B_\mu(\Lambda)$, respectively.

Remark 2.11. It is well-known [7, 35, 45] that the Demazure character can be expressed in terms of the Demazure operator $D_i : \mathbb{Z}[P] \rightarrow \mathbb{Z}[P]$

$$D_i(e^\mu) = \frac{e^\mu - e^{\mu - (1 + \langle \alpha_i^\vee, \mu \rangle) \alpha_i}}{1 - e^{-\alpha_i}},$$

where α_i is a simple root and α_i^\vee the corresponding coroot. Then for $w = s_{i_N} \cdots s_{i_1} \in W$ a reduced expression

$$\text{ch}V_w(\lambda) = D_{i_N} \cdots D_{i_1}(e^\lambda).$$

3. KASHIWARA-NAKASHIMA TABLEAUX AND BRANCHING RULES

3.1. Kashiwara-Nakashima tableaux. We briefly review a method, due to Kashiwara and Nakashima [33], of realizing all highest weight crystals of non-exceptional finite type \mathfrak{g} . The first observation is that many crystals occur inside high enough tensor powers of the “standard” crystals shown in Figure 2. Many here means all in type A and C , but not those involving spin weights in types B and D .

We call the set of symbols that show up in the boxes of the standard crystal of type $X_n = A_n, B_n, C_n, D_n$ the type X_n alphabet. Impose a partial order \prec on this alphabet by saying $x \prec y$ iff x is to the left of y in the presentation of the standard crystals in Figure 2 (in type D_n , the symbols n and \bar{n} are incomparable).

Definition 3.1. Fix \mathfrak{g} of type X_n , for $X = A, B, C, D$. Fix a dominant integral weight γ for $\mathfrak{g} = X_n$. Write $\gamma = m_1\omega_1 + m_2\omega_2 + \cdots + m_{n-1}\omega_{n-1} + m_n\omega_n$. Define a generalized partition $\Lambda(\gamma)$ associated to γ , which is defined case by case as follows:

- If $X = A, C$, $\Lambda(\gamma)$ has m_i columns of height i for each $1 \leq i \leq n$;
- If $X = B$, $\Lambda(\gamma)$ has m_i columns of height i for each $1 \leq i \leq n-1$, and $m_n/2$ columns of height n ;




FIGURE 3. An element of $B(\omega_3 + 2\omega_2)$ of type C_4 as realized inside the tensor product $B(\omega_1)^{\otimes 7}$ of seven copies of the standard crystal, and the corresponding tableau. For explicit conditions on which tableaux appear in this correspondence, see [33].

- If $X = D$, $\Lambda(\gamma)$ has m_i columns of each height i for each $1 \leq i \leq n - 2$, $\min(m_{n-1}, m_n)$ columns of height $n - 1$, and $|m_n - m_{n-1}|/2$ columns of height n . Color columns of height n using color 1 if $m_n > m_{n-1}$ and color 2 if $m_n < m_{n-1}$.

We use French notation for partitions here, where we adjust the columns at the bottom. In cases where the above formulas involve a fractional number x of columns at some height, we denote this by putting $\lfloor x \rfloor$ columns in addition to a single column of half width. Notice that this can only happen for columns of height n , and at worst we get a single column of width $1/2$.

Definition 3.2. Fix a dominant integral weight γ . We say γ is a *non-spin weight* if

- In types A_n and C_n , no conditions.
- In type B_n , $\Lambda(\gamma)$ does not contain any column of width $1/2$ (or equivalently, m_n is even).
- In type D_n , $\Lambda(\gamma)$ has no columns of height n (or equivalently, $m_{n-1} = m_n$).

In the case when $\Lambda = \Lambda(\gamma)$ does not contain a column of width $1/2$, the highest weight crystal $B(\gamma)$ embeds in the M -th tensor power of $B(\omega_1)$, where M is the number of boxes in the partition Λ . Furthermore, the image of $B(\gamma)$ is contained in the set of $x_M \otimes \cdots \otimes x_1$ such, when entered into the Young diagram Λ in order moving up columns and right to left (see Figure 3), the result is weakly increasing along rows and, if you ignore the symbols 0 in type B and n, \bar{n} in type D , also strictly increasing up columns. We refer to fillings which occur in the image of $B(\gamma)$ as Kashiwara–Nakashima (KN) tableaux. Precise conditions describing KN tableaux are given in [33]. The crystal structure on the set of KN tableaux is inherited from the crystal structure on $B(\omega_1)^{\otimes M}$. Modifications of this construction dealing with spin weights are given in [33].

Remark 3.3. The above construction of $B(\lambda)$ in type D_n goes through without modification in the case $n = 3$, when $D_3 \cong A_3$. Thus we have two realizations of the crystal in this case.

3.2. Branching rules and \pm diagrams. We now describe branching rules for certain representations of X_n , where $X = B, C$ or D .

Definition 3.4. Fix a non-spin weight γ (see Definition 3.2), and set $\Lambda = \Lambda(\gamma)$. A \pm *diagram* P with outer shape Λ is a sequence of partitions $\lambda \subseteq \mu \subseteq \Lambda$ such that Λ/μ and μ/λ are horizontal strips (i.e. every column contains at most one box). We depict a \pm diagram by filling μ/λ with the symbol $+$ and those of Λ/μ with the symbol $-$. We make the additional type dependent modifications and restrictions:

- (i) In type C_n , λ has no columns of height n .
- (ii) In type B_n , λ has no columns of height n . Additionally, we allow the \pm diagram to contain an extra symbol 0. There can be at most one 0, this must occur at height n , must be to the right of all $+$ at height n , and to the left of all 0 at height n , and must be the only symbol in its column.
- (iii) In type D_n , columns of λ of height $n - 1$ are either all colored 1 or all colored 2.

We denote by $\text{outer}(P) = \Lambda$ the outer shape of P and by $\text{inner}(P) = \lambda$ the inner shape of P .

Embed X_{n-1} into X_n by removing node 1 from the Dynkin diagram of type X_n . In the special cases of B_2, C_2 and D_3 , removing the node 1 gives a Dynkin diagram of a different type (A_1, A_1 , and $A_1 \times A_1$, respectively). By abuse of notation, we use the symbol X_{n-1} to mean this new diagram in these special cases. Although these special cases are not explicitly mentioned in [10], the proof of Theorem 3.5 goes through without change.

Theorem 3.5. (see [10, Section 3.2]) *Fix a non-spin dominant integral weight γ (see Definition 3.2) for $\mathfrak{g} = X_n$, and let $\Lambda = \Lambda(\gamma)$ (see Definition 3.1). Then there is a bijection between \pm diagrams P with outer shape Λ and X_{n-1} highest weight elements in $B(\Lambda)$. This can be realized in terms of KN tableaux by the following algorithm.*

- (i) For each $+$ at height n , fill that column with $12 \dots n$.
- (ii) Replace each $-$ with a $\bar{1}$ and, if there is a 0 in the \pm diagram, place a 0 in that position of the tableaux.
- (iii) Fill the remainder of all columns by strings of the form $23 \dots k$.
- (iv) Let S be the multi-set containing the heights of all the $+$ in P of height less than n . Move through the columns of Λ from top to bottom, left to right, ignoring the columns $12 \dots n$ of step (i), modifying the tableaux as follows. Each time you encounter a $\bar{1}$, replace it with $\overline{h+1}$, where h is the largest element of S , and delete h from S . Each time you encounter a 2 which is at the bottom of a column, replace the string $23 \dots k$ in that column by $12 \dots h h + 2 \dots k$, where h is the largest element of S , and remove h from S . Once S is empty, stop.
- (v) In type D_n , if the \pm diagram has empty columns of height $n - 1$ colored 2, change all occurrences of n at height $n - 1$ to \bar{n} .

Furthermore, two \pm diagrams P and P' correspond to X_{n-1} highest weight vectors of the same X_{n-1} weight if and only if $\text{inner}(P) = \text{inner}(P')$. \square

Remark 3.6. Definition 3.4 part (iii) differs from the statement in [10], since in that paper there were no colorings for columns of height $n - 1$. These colorings are needed, as can be seen by considering the branching rules for $B(\omega_{n-1} + \omega_n)$ for D_4 :

$$\begin{array}{|c|} \hline 3 \\ \hline 2 \\ \hline 1 \\ \hline \end{array} \leftrightarrow \begin{array}{|c|} \hline + \\ \hline \\ \hline \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bar{1} \\ \hline 3 \\ \hline 2 \\ \hline \end{array} \leftrightarrow \begin{array}{|c|} \hline - \\ \hline \\ \hline \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bar{3} \\ \hline 3 \\ \hline 2 \\ \hline \end{array} \leftrightarrow \begin{array}{|c|} \hline - \\ \hline + \\ \hline \\ \hline \end{array} \quad \begin{array}{|c|} \hline 4 \\ \hline 3 \\ \hline 2 \\ \hline \end{array} \leftrightarrow \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bar{4} \\ \hline 3 \\ \hline 2 \\ \hline \end{array} \leftrightarrow \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \end{array}$$

where the fourth \pm diagram is considered of color 1 and the fifth of color 2.

Definition 3.7. Let γ be an integral highest weight of type X_n and write $\gamma = m_1\omega_1 + m_2\omega_2 + \dots + m_{n-1}\omega_{n-1} + m_n\omega_n$. We call γ a *case (AUT) weight* if it satisfies:

- If $X = C$, no conditions;
- If $X = B$, assume $m_n = 0$;
- If $X = D$, $m_n = m_{n-1} = 0$.

Note that in all these cases, $\Lambda(\gamma)$ is an ordinary partition.

Remark 3.8. The term “case (AUT)” in Definition 3.7 is motivated by the fact that these are exactly the classical highest weights that appear in case (AUT) KR crystals, as defined in Section 4 below.

Remark 3.9. All case (AUT) weights are non-spin, as in Definition 3.2. Furthermore, if one starts with a case (AUT) weight γ , then the X_{n-1} weights which appear in the decomposition from Theorem 3.5 will all be non-spin weights, and the X_{n-1} highest weight γ' corresponding to the \pm diagram P will satisfy $\Lambda(\gamma') = \text{inner}(P)$.

3.3. Nested \pm diagrams. If γ is a case (AUT) weight, $n \geq 3$ in type B_n, C_n , and $n \geq 4$ in type D_n , it follows immediately from Section 3.2 that the branching rule from X_n to X_{n-2} can be described using pairs of \pm diagrams P and p with $\text{outer}(P) = \Lambda(\gamma)$ and $\text{inner}(P) = \text{outer}(p)$. We call such pairs of \pm diagrams *nested*. The following result from [55] gives an explicit description of the action of e_1 on an X_{n-2} highest weight vector in terms of pairs of \pm diagrams. Since e_1 commutes with all e_j for $j \geq 3$, this completely describes e_1 .

Pair off the symbols in (P, p) according to the rules

- Successively run through all $+$ in p from left to right and, if possible, pair it with the leftmost yet unpaired $+$ in P weakly to the left of it.
- Successively run through all $-$ in p from left to right and, if possible, pair it with the rightmost yet unpaired $-$ in P weakly to the left.
- Successively run through all yet unpaired $+$ in p from left to right and, if possible, pair it with the leftmost yet unpaired $-$ in p .

Notice that a \pm diagram is uniquely determined by its outer (or inner) shape along with the number of each symbol $+$, $-$, and 0 on each row. Similarly, a pair (P, p) of nested \pm diagrams is uniquely determined by its intermediate shape $\text{int}(P, p) = \text{outer}(p) = \text{inner}(P)$, along with the data of how many of each symbol is in each row of P and of p . Since e_1 commutes with e_j for all $j \geq 3$, the following gives a complete description of the action of e_1 on the crystal:

Lemma 3.10. [55, Lemma 5.1] *Let γ be a case (AUT) weight and fix a pair (P, p) of nested \pm diagrams such that $\text{outer}(P) = \Lambda(\gamma)$. Let b be the corresponding X_{n-2} highest weight vector in $B(\gamma)$. If there are no unpaired $+$ in p and no unpaired $-$ in P , then $e_1(b) = 0$. Otherwise, $e_1(b)$ is the X_{n-2} highest weight element corresponding to the pair of \pm diagrams (P', p') described as follows.*

If there is an unpaired $+$ in p , let k be the height the rightmost unpaired $+$ in p .

- *If there is a $-$ directly above the chosen $+$ in p , then $\text{int}(P, p)/\text{int}(P', p')$ is a single box at height $k + 1$. All rows of P and p have the same number of each symbol except: There is one more $+$ in P at height $k + 1$; there is one less $-$ in p at height $k + 1$; there is one less $+$ and one more $-$ in p at height k .*
- *Otherwise, $\text{int}(P, p)/\text{int}(P', p')$ is a single box at height k . All rows of P and p have the same number of each symbol except: There is one more $+$ in P at height k ; there is one less $+$ in p at height k .*

Otherwise let k be the height of the leftmost unpaired $-$ in P .

- If there is a $+$ directly below the chosen $-$ in P , then $\text{int}(P', p')/\text{int}(P, p)$ is a single box at height $k - 1$. All rows of P and p have the same number of each symbol except: There is one less $-$ and one more $+$ in P at height k ; there is one less $+$ in P at height $k - 1$; there is one more $-$ in p at height $k - 1$.
- Otherwise, $\text{int}(P', p')/\text{int}(P, p)$ is a single box at height k . All rows of P and p have the same number of each symbol except: There is one less $-$ in P at height k ; there is one more $-$ in p at height k .

In type B , in every case, the number of 0s in p remains unchanged (P cannot contain 0 since we are assuming that γ is in case (AUT)).

Remark 3.11. In principle, one could build a new combinatorial model for type B_n, C_n, D_n crystals using nested \pm diagrams, where the crystal operators would be given by Lemma 3.10. That is, the tableaux would consist of $n - 1$ nested \pm diagrams in type B_n, C_n and $n - 2$ nested \pm diagrams in type D_n , along with some extra data recording an element in a representation of A_1 (or $A_1 \times A_1$ in type D_n). This would have some advantages over KN tableaux, in that branching rules would be more readily visible. Such tableaux may also be more convenient for working with KR crystals.

4. REALIZATIONS OF KR CRYSTALS

Explicit combinatorial models for KR crystals $B^{r,s}$ for the non-exceptional types were constructed in [10]. In this section we recall and prove some properties that we need for the definition of the energy function in Section 5 and Lemma 7.2.

The results are presented for three different cases:

(IRR) The classically irreducible KR crystals $B^{r,s}$:

$$\begin{aligned} A_n^{(1)} & 1 \leq r \leq n \\ D_n^{(1)} & r = n - 1, n \\ D_{n+1}^{(2)} & r = n \\ C_n^{(1)} & r = n \end{aligned}$$

(AUT) KR crystals $B^{r,s}$ constructed via Dynkin automorphisms:

$$\begin{aligned} D_n^{(1)} & 1 \leq r \leq n - 2 \\ B_n^{(1)} & 1 \leq r \leq n - 1 \\ A_{2n-1}^{(2)} & 1 \leq r \leq n \end{aligned}$$

(VIR) The remaining virtually constructed KR crystals:

$$\begin{aligned} B_n^{(1)} & B^{n,s} \\ C_n^{(1)} & B^{r,s} \text{ for } 1 \leq r < n \\ D_{n+1}^{(2)} & B^{r,s} \text{ for } 1 \leq r < n \\ A_{2n}^{(2)} & B^{r,s} \text{ for } 1 \leq r \leq n. \end{aligned}$$

In Section 4.1 we present the classical decomposition of the KR crystals in the various cases. Case (IRR) is discussed in Section 4.2. This case is well-understood, and we simply state the facts we need. Case (AUT) is handled in Section 4.3. This time we discuss the realizations in more detail, and prove a technical lemma. Case (VIR) is handled in Sections 4.4, 4.5 and 4.6. Following [10], we realize these KR crystals as virtual crystals, using a combination of the similarity method of

Kashiwara [29] and the technique of virtual crystals developed in [51, 52, 10]. We refer to crystals obtained using both of these methods as virtual crystals.

4.1. Classical decompositions of KR crystals. Set

$$(4.1) \quad \diamond = \begin{cases} \emptyset & \text{for type } A_n^{(1)} \text{ and } 1 \leq r \leq n \\ & \text{for types } C_n^{(1)}, D_{n+1}^{(2)} \text{ and } r = n \\ & \text{for type } D_n^{(1)} \text{ and } r = n - 1, n \\ \text{vertical domino} & \text{for type } D_n^{(1)} \text{ and } 1 \leq r \leq n - 2 \\ & \text{for types } B_n^{(1)}, A_{2n-1}^{(2)} \text{ and } 1 \leq r \leq n \\ \text{horizontal domino} & \text{for types } C_n^{(1)}, D_{n+1}^{(2)} \text{ and } 1 \leq r < n \\ \text{box} & \text{for type } A_{2n}^{(2)} \text{ and } 1 \leq r \leq n. \end{cases}$$

Then every $B^{r,s}$ decomposes as a classical crystal as

$$(4.2) \quad B^{r,s} \cong \bigoplus_{\lambda} B(\lambda),$$

where the sum is over those λ such that $\Lambda(\lambda)$ (see Definition 3.1) can be obtained from the rectangle $\Lambda(s\omega_r)$ by removing some number of \diamond , each occurring with multiplicity 1. In the untwisted case this was obtained by Chari [3]. In the twisted case, it was conjectured in [16, Appendix A] and proven by Hernandez (see [19, Section 5]).

4.2. Classically irreducible KR crystals (case (IRR)). As a $\{1, 2, \dots, n\}$ -crystal, $B^{r,s} \cong B(r\omega_s)$. For example, the realization of the KR crystal $B^{r,s}$ of type $A_n^{(1)}$ is well-known in terms of rectangular Young tableaux of shape (s^r) . We refer the reader to e.g. [10, Section 4.1] for details. The construction of the other irreducible KR crystals can be found in [10, Section 6]. We only need the following fact, which follows immediately from the explicit models.

Lemma 4.1. *Let $B^{r,s}$ be a KR crystal of type \mathfrak{g} for one of the cases in case (IRR). Then for all $b \in B^{r,s}$, $\varepsilon_0(b) \leq s$.*

4.3. KR crystals constructed via Dynkin automorphisms (case (AUT)).

Let \mathfrak{g} be of type $D_n^{(1)}$, $B_n^{(1)}$ or $A_{2n-1}^{(2)}$, with the underlying finite type Lie algebra of type $X_n = D_n$, B_n or C_n , respectively. Fix $s > 0$ and r so that $B^{r,s}$ is in case (AUT). Consider the classical crystal

$$C^{r,s} := \bigoplus B(\lambda),$$

where the sum is over all λ such that $\Lambda(\lambda)$ (see Definition 3.1) can be obtained from $\Lambda(s\omega_r)$ by removing vertical dominos. As in Section 3.2, the X_{n-1} highest weight elements in $C^{r,s}$ (i.e. the highest weight element for the algebra with Dynkin diagram $I \setminus \{0, 1\}$) are indexed by \pm diagrams whose outer shape can be obtained from $\Lambda(s\omega_r)$ by removing vertical dominos.

Definition 4.2. Define the involution ζ on the X_{n-1} highest weight vectors of $C^{r,s}$, as indexed by \pm diagrams, as follows. Let P be a \pm diagram with $\text{outer}(P) = \Lambda$ and $\text{inner}(P) = \lambda$. For each $1 \leq i \leq r - 1$:

- (i) If $i \equiv r - 1 \pmod{2}$ then above each column of λ of height i , there must be a $+$ or a $-$. Interchange the number of such $+$ and $-$.

- (ii) If $i \equiv r \pmod{2}$ then above each column of λ of height i , either there are no signs or a \mp pair. Interchange the number of columns of each type.

By Theorem 3.5, this can be extended in a unique way to an involution ς on $C^{r,s}$.

Theorem 4.3. [55, 10] *Fix \mathfrak{g} and a KR crystal $B^{r,s}$ in case (AUT). Define operators e_0 and f_0 on $C^{r,s}$ by $e_0 := \varsigma \circ e_1 \circ \varsigma$ and $f_0 := \varsigma \circ f_1 \circ \varsigma$. Then $C^{r,s}$ along with these new operators is isomorphic to $B^{r,s}$. \square*

Lemma 4.4. *Fix \mathfrak{g} and a KR crystal $B^{r,s}$ as in case (AUT). Consider the realization of $B^{r,s}$ given in Theorem 4.3. Fix $b \in B^{r,s}$, and assume b lies in the component $B(\gamma)$ and $e_0(b)$ lies in the classical component $B(\gamma')$. Then*

- (i) $\Lambda(\gamma')$ is either equal to $\Lambda(\gamma)$, or else is obtained from $\Lambda(\gamma)$ by adding or removing a single vertical domino.
- (ii) If $\varepsilon_0(b) > s$, then $\Lambda(\gamma')$ is obtained from $\Lambda(\gamma)$ by removing a vertical domino.

Proof. Recall that for $b \in B^{r,s}$ by definition $e_0(b) = \varsigma \circ e_1 \circ \varsigma(b)$. Since e_0 and e_1 commute with e_i for $i = 3, 4, \dots, n$, they are defined on X_{n-2} components, which are described by pairs of \pm diagrams (P, p) . Consider the 1-string $\{b_0, b_1, \dots, b_k\}$ where $b_i = e_1^i \circ \varsigma(b)$ with corresponding \pm diagrams (P_i, p_i) . By Lemma 3.10 there exists a $0 \leq j \leq k$ such that:

- (a) For $0 \leq i < j$, P_{i+1} is obtained from P_i by the addition of one box containing $+$; the $+$ are added from right to left with increasing i .
- (b) For $j \leq i < k$, P_{i+1} is obtained from P_i by the removal of one box containing $-$; the $-$ are removed from left to right with increasing i .

Recall that ς interchanges $+$ and $-$ in columns of height congruent to 1 (mod r) and \mp -pairs and empty columns of height congruent to 0 (mod r). Empty columns of height r are left unchanged. Since the width of the diagrams is at most s , we can only have $\varepsilon_0(b) > s$ if we are in case (a) above and the $+$ is added at height strictly smaller than r . Then there are two cases:

- The $+$ is added below a $-$ in P_0 .
- The $+$ is added in an empty column in P_0 .

In both cases it is easy to check from the rules of ς that a vertical domino is removed from the outer shape.

In all other cases, applying e_1 to $\varsigma(b)$ only adds or removes a single symbol to/from the corresponding \pm diagram P , so from the definition of ς and the fact that $e_0 = \varsigma \circ e_1 \circ \varsigma$, it is clear that $\Lambda(\gamma')$ differs from $\Lambda(\gamma)$ by at most one vertical domino, and the lemma holds. \square

4.4. $B^{r,s}$ of type $C_n^{(1)}$ for $r < n$ (in case (VIR)). The KR crystals of type $C_n^{(1)}$ are constructed inside an ambient crystal of type $A_{2n+1}^{(2)}$.

Theorem 4.5. [10, Section 4.3, Theorem 5.7] *Fix $n \geq 2$ and $r < n$. Let $\hat{B}^{r,s}$ be the KR crystal corresponding to type $A_{2n+1}^{(2)}$, with crystal operators $\hat{e}_0, \dots, \hat{e}_{n+1}$. Let V be the subset of all $b \in \hat{B}^{r,s}$ which are invariant under the involution ς from Definition 4.2. Define the virtual crystal operators $e_i := \hat{e}_{i+1}$ for $1 \leq i \leq n$ and $e_0 := \hat{e}_0 \hat{e}_1$. Then V along with the operators e_0, \dots, e_n is isomorphic to $B^{r,s}$ for type $C_n^{(1)}$.*

Here the operators f_i and \hat{f}_i are defined by the condition $f_i(b) = b'$ if and only if $e_i(b') = b$. \square

Recall that as a classical crystal, $B^{r,s}$ of type $C_n^{(1)}$ decomposes as in (4.2).

Lemma 4.6. *Let $B^{r,s}$ with $r < n$ be the KR crystal of type $C_n^{(1)}$. Fix $b \in B^{r,s}$, and assume b lies in the component $B(\gamma)$ and $e_0(b)$ lies in the classical component $B(\gamma')$. Then*

- (i) $\Lambda(\gamma')$ is either equal to $\Lambda(\gamma)$, or else is obtained from $\Lambda(\gamma)$ by adding or removing a single horizontal domino.
- (ii) If $\varepsilon_0(b) > \lceil s/2 \rceil$, then $\Lambda(\gamma')$ is obtained from $\Lambda(\gamma)$ by removing a horizontal domino.

Proof. Realize the crystal $B^{r,s}$ of type $C_n^{(1)}$ inside the ambient crystal $\hat{B}^{r,s}$ of type $A_{2n+1}^{(2)}$ using Theorem 4.5. Denote the embedding by $S : B^{r,s} \hookrightarrow \hat{B}^{r,s}$, and set $\hat{b} = S(b)$. Let (P, p) be the pair of \pm diagrams associated to the $\{3, 4, \dots, n+1\}$ highest weight corresponding to \hat{b} , and let γ be the highest weight of b as a $C_n \subset C_n^{(1)}$ crystal. Since $e_0 = \hat{e}_1 \hat{e}_0$ and $e_i = \hat{e}_{i+1}$, $\Lambda(\gamma) = \text{inner}(P)$.

Recall from [10, Section 4.3] that $S(b)$ is invariant under ς , hence in particular P is invariant under ς . Also, as operators on $\hat{B}^{r,s}$,

$$(4.3) \quad S(e_0) = \hat{e}_1 \hat{e}_0 = \hat{e}_1 \circ \varsigma \circ \hat{e}_1 \circ \varsigma = \hat{e}_1 \circ \varsigma \circ \hat{e}_1.$$

By Lemma 3.10, \hat{e}_1 either moves a $+$ from p to P , or moves a $-$ from P to p . That is, $\text{outer}(P)$ is unchanged, $\text{inner}(P)$ only changes by a single box, and, except for the new $+$ or missing $-$, the number of $+$ and $-$ on each row of P is unchanged. Also, in all situations ς preserves $\text{inner}(P)$. Thus (4.3) implies that $\text{inner}(P)$ can change by at most 2 boxes. This along with the description of the classical decomposition of $B^{r,s}$ implies part (i).

Now assume that $\varepsilon_0(b) > \lceil s/2 \rceil$. Since P is ς -invariant, it is clear that there can be at most $\lfloor s/2 \rfloor$ symbols $-$ in P . Since $\varepsilon_0(b) = \hat{\varepsilon}_1(\hat{b}) > s/2$, by the description of \hat{e}_1 from Lemma 3.10 we see that \hat{e}_1 must move a $+$ from p to P . There are three cases:

- (a) The $+$ is added below a $-$ in P .
- (b) The $+$ is added in an empty column in P at height less than r .
- (c) The $+$ is added in an empty column in P at height r .

It is not hard to check that in case (a), in $\varsigma \circ \hat{e}_1(P)$, there is one less $+$ in row $k+1$ and one more empty column of height $k-1$ compared to P , and everything else is unchanged. Since both P and $\hat{e}_1 \circ \varsigma \circ \hat{e}_1(P)$ are invariant under ς , the only possibility is that the final \hat{e}_1 must add another $+$ to row k . Thus two symbols have been added to P in row k , which has the effect of removing a horizontal domino from $\text{inner}(P)$.

In case (b), in $\varsigma \circ \hat{e}_1(P)$, there is one less \mp -pair in row $k+1$ and one more $-$ in row k and everything else is unchanged. Again since $\hat{e}_1 \circ \varsigma \circ \hat{e}_1(b)$ has to be invariant under ς , the only possibility is that \hat{e}_1 adds another $+$ to row k of $\varsigma \circ \hat{e}_1(P)$. Thus $S(e_0)$ removes a horizontal domino from $\text{inner}(P)$.

In case (c), P has at least one column with no symbols, and so P contains at most $\lfloor (s-1)/2 \rfloor$ symbols $-$. Since $\hat{\varepsilon}_1(\hat{b}) = \varepsilon_0(b) > \lceil s/2 \rceil$, there are at least 2 uncanceled $+$ in p . Consider the double \pm diagram (P', p') for $\varsigma \circ \hat{e}_1(\hat{b})$. This differs from (P, p) by: P' has an extra $-$ at height r , and p' has one less $+$. Looking at the cancellation rules before Lemma 3.10, there is still an uncanceled $+$ in p' (corresponding to the second uncanceled $+$ in p). Thus the second \hat{e}_1 moves another $+$ into P' . So in

total $S(e_0) = \hat{e}_1 \circ \varsigma \circ \hat{e}_1$ decreases the inner shape of P by 2 boxes, which must remove a horizontal domino. \square

4.5. $B^{n,s}$ in type $B_n^{(1)}$ (in case (VIR)).

Theorem 4.7. (see [10, Lemma 4.2]) *Let $\hat{B}^{n,s}$ be the KR crystal of type $A_{2n-1}^{(2)}$ with crystal operators $\hat{e}_0, \dots, \hat{e}_n$. Let V be the subset of $\hat{B}^{n,s}$ which can be reached from the classical highest weight element in $\hat{B}^{n,s}$ of weight $s\omega_n$ using the virtual crystal operators $e_i = \hat{e}_i^2$ for $0 \leq i \leq n-1$, $e_n = \hat{e}_n$. Then V along with the operators e_0, \dots, e_n is isomorphic to $B^{n,s}$ for $B_n^{(1)}$.*

Furthermore, the type B_n highest weight vectors in V are exactly the type C_n highest weight vectors in $\hat{B}^{n,s}$ that can be obtained from $\Lambda(s\omega_r)$ by removing 2×2 blocks.

Proof. The virtual crystal realization is proven in [10, Lemma 4.2]. The classical decomposition of $B^{n,s}$ as a type B_n crystal is

$$\bigoplus_{\mathbf{k}} B(k_\iota \omega_\iota + k_{\iota+2} \omega_{\iota+2} + \dots + k_{n-2} \omega_{n-2} + k_n \omega_n),$$

where $\iota = 0, 1$ so that $\iota \equiv n \pmod{2}$, $\omega_0 = 0$, and the sum is over all nonnegative integer vectors \mathbf{k} such that $2k_\iota + \dots + 2k_{n-2} + k_n < s$. As discussed in [10, Lemma 4.2], the highest weight vectors for each of these components must be classical type C_n highest weight vector in the ambient crystal $\hat{B}^{n,s}$ of weight

$$2k_\iota \omega_\iota + 2k_{\iota+2} \omega_{\iota+2} + \dots + 2k_{n-2} \omega_{n-2} + k_n \omega_n.$$

The weight of these highest weight vectors in the ambient crystal are exactly those γ such that $\Lambda(\gamma)$ is obtained from $\Lambda(s\omega_r)$ by removing 2×2 blocks. The result follows. \square

Lemma 4.8. *Let $B^{n,s}$ be the KR crystal of type $B_n^{(1)}$. Fix $b \in B^{n,s}$, and assume b lies in the component $B(\gamma)$ and $e_0(b)$ lies in the classical component $B(\gamma')$. Then*

- (i) $\Lambda(\gamma')$ is either equal to $\Lambda(\gamma)$, or else is obtained from $\Lambda(\gamma)$ by adding or removing a single vertical domino.
- (ii) If $\varepsilon_0(b) > \lceil s/2 \rceil$, then $\Lambda(\gamma')$ is obtained from $\Lambda(\gamma)$ by removing a vertical domino.

Proof. For part (i), use the fact that by the virtual realization from Theorem 4.7, $e_0 = \hat{e}_0^2$. So, it follows from Lemma 4.4 that e_0 adds or subtracts at most 2 vertical dominoes. By Theorem 4.7, the only possibilities are that e_0 either leaves the shape unchanged, or adds or subtracts a single 2×2 block.

Part (ii) follows since, if $\varepsilon_0(b) > \lceil s/2 \rceil$, then $\hat{\varepsilon}_0(b) \geq s + 2$. Hence applying $e_0 = \hat{e}_0^2$ subtracts 2 vertical dominoes by Lemma 4.4, which as above must fit together as a 2×2 block. \square

4.6. The KR crystals of type $A_{2n}^{(2)}$ and $D_{n+1}^{(2)}$ in case (VIR). We now present the virtual crystal construction of the KR crystals $B^{r,s}$ of types $A_{2n}^{(2)}$ and $D_{n+1}^{(2)}$ of case (VIR), and then prove the analogue of Lemma 4.4 in this setting.

Theorem 4.9. [10, Section 4.4, Theorem 5.7] *Fix $n \geq 2$. Let $\hat{B}^{r,2s}$ be the KR crystal corresponding to type $C_n^{(1)}$, with crystal operators $\hat{e}_0, \dots, \hat{e}_n$. In each case below, let V be the subset of $\hat{B}^{r,2s}$ which can be reached from the classical highest weight element in $\hat{B}^{r,2s}$ of weight $2s\omega_r$ using the listed virtual crystal operators:*

- (i) Let $1 \leq r < n$. Define $e_i := \hat{e}_i^2$ for $1 \leq i \leq n-1$, $e_0 := \hat{e}_0$, and $e_n := \hat{e}_n$. Then V , along with the operators e_0, \dots, e_n , is isomorphic to the KR crystal $B^{r,s}$ for type $D_{n+1}^{(2)}$.
- (ii) Let $1 \leq r \leq n$. Define $e_i := \hat{e}_i^2$ for $1 \leq i \leq n$ and $e_0 := \hat{e}_0$. Then V along with the operators e_0, \dots, e_n is $B^{r,s}$ for type $A_{2n}^{(2)}$.

Here the operators f_i and \hat{f}_i are defined by the condition $f_i(b) = b'$ if and only if $e_i(b') = b$.

Lemma 4.10. Consider $B^{r,s}$ of type $D_{n+1}^{(2)}$ with $r < n$, or of type $A_{2n}^{(2)}$ with $1 \leq r \leq n$. Fix $b \in B^{r,s}$, and assume b lies in the classical (type B_n or C_n , respectively) component $B(\gamma)$, and $e_0(b)$ lies in the classical component $B(\gamma')$. Then

- (i) $\Lambda(\gamma')$ is either equal to $\Lambda(\gamma)$, or else is obtained from $\Lambda(\gamma)$ by adding or removing a single box.
- (ii) If $\varepsilon_0(b) > s$, then $\Lambda(\gamma')$ is obtained from $\Lambda(\gamma)$ by removing a box.

Proof. Let $\hat{B}^{r,2s}$ be the ambient KR crystal of type $C_n^{(1)}$ with crystal operators \hat{e}_i as in Theorem 4.9. Denote the virtualization map $S : B^{r,s} \rightarrow V \subseteq \hat{B}^{r,2s}$. Recall that V is the subset of $\hat{B}^{r,2s}$ generated by the element of weight $2s\omega_r$ by applying the virtual crystal operators \hat{e}_i :

- (i) In type $D_{n+1}^{(2)}$, $e_0 = \hat{e}_0$, $e_n = \hat{e}_n$, and $e_i = \hat{e}_i^2$ for $1 \leq i \leq n-1$.
- (ii) In type $A_{2n}^{(2)}$, $e_0 = \hat{e}_0$, and $e_i = \hat{e}_i^2$ for $1 \leq i \leq n$.

By Lemma 4.6, \hat{e}_0 changes the classical component of the underlying KR crystal of type $C_n^{(1)}$ by adding or subtracting at most one horizontal domino, and furthermore if $\hat{e}_0(S(b)) > s$, then \hat{e}_0 always removes a horizontal domino. The result now follows immediately from the description of the classical components of these virtual crystals as given in [10, Lemma 4.9].

Note that in the case of $A_{2n}^{(2)}$ with $r = n$, $\hat{B}^{n,2s}$ is not a KR crystal of type $C_n^{(1)}$. However, Theorem 4.5 still gives a type $C_n^{(1)}$ combinatorial crystal in this case and Lemma 4.6 still holds. So the proof still goes though in this case. \square

5. ENERGY FUNCTIONS

We define two energy functions on tensor products of KR crystals. The function E^{int} is given by a fairly natural ‘‘global’’ condition on tensor products of level- ℓ KR crystals. The function D is defined by summing up combinatorially defined ‘‘local’’ contributions, and makes sense for general tensor products of KR crystals. It was suggested (but not proven) in [52, Section 2.5] that, when E^{int} is defined, these two functions agree up to a shift. This will be proven in Theorem 7.5 below.

5.1. The function E^{int} .

Definition 5.1. For each node $r \in I \setminus \{0\}$ and each $\ell \in \mathbb{Z}_{>0}$, let $u_{r,\ell c_r}$ be the unique element of $B^{r,\ell c_r}$ such that $\varepsilon(u_{r,\ell c_r}) = \ell\Lambda_0$ (which exists as $B^{r,\ell c_r}$ is perfect).

The following is essentially the definition of a ground state path from [25].

Definition 5.2. Let $B = B^{r_N, \ell c_{r_N}} \otimes \dots \otimes B^{r_1, \ell c_{r_1}}$ be a composite level- ℓ KR crystal. Define $u_B = u_B^N \otimes \dots \otimes u_B^1$ to be the unique element of B such that

- (i) $u_B^1 = u_{r_1, \ell c_{r_1}}$ and
- (ii) for each $1 \leq j < N$, $\varepsilon(u_B^{j+1}) = \varphi(u_B^j)$.

This u_B is well-defined by condition (v) in Definition 2.4 of a perfect crystal. The element u_B is called the *ground state path* of B .

Definition 5.3. Let B be a composite KR crystal of level ℓ and consider u_B as in Definition 5.2. The *intrinsic energy function* E^{int} on B is defined by setting $E^{\text{int}}(b)$ to be the minimal number of f_0 in a string $f_{i_N} \cdots f_{i_1}$ such that $f_{i_N} \cdots f_{i_1}(u_B) = b$.

5.2. The D function.

Definition 5.4. The D -function on $B^{r,s}$ is the function defined as follows:

- (i) $D_{B^{r,s}} : B^{r,s} \rightarrow \mathbb{Z}$ is constant on all classical components.
- (ii) On the component $B(\lambda)$, $D_{B^{r,s}}$ records the maximum number of \diamond that can be removed from $\Lambda(\lambda)$ such that the result is still a (generalized) partition, where \diamond is as in (4.1).

In those cases when $\diamond = \emptyset$, this is interpreted as saying that $D_{B^{r,s}}$ is the constant function 0.

Let B_1, B_2 be two affine crystals with generators v_1 and v_2 , respectively, such that $B_1 \otimes B_2$ is connected and $v_1 \otimes v_2$ lies in a one-dimensional weight space. By [39, Proposition 3.8], this holds for any two KR crystals. The generator v for the KR crystal $B^{r,s}$ is chosen to be the unique element of classical weight $s\omega_r$.

The *combinatorial R -matrix* [25, Section 4] is the unique crystal isomorphism

$$\sigma : B_2 \otimes B_1 \rightarrow B_1 \otimes B_2.$$

By weight considerations, this must satisfy $\sigma(v_2 \otimes v_1) = v_1 \otimes v_2$.

As in [25] and [51, Theorem 2.4], there is a function $H = H_{B_2, B_1} : B_2 \otimes B_1 \rightarrow \mathbb{Z}$, unique up to global additive constant, such that, for all $b_2 \in B_2$ and $b_1 \in B_1$,

$$(5.1) \quad H(e_i(b_2 \otimes b_1)) = H(b_2 \otimes b_1) + \begin{cases} -1 & \text{if } i = 0 \text{ and LL,} \\ 1 & \text{if } i = 0 \text{ and RR,} \\ 0 & \text{otherwise.} \end{cases}$$

Here LL (resp. RR) indicates that e_0 acts on the left (resp. right) tensor factor in both $b_2 \otimes b_1$ and $\sigma(b_2 \otimes b_1)$. When B_1 and B_2 are KR crystals, we normalize H_{B_2, B_1} by requiring $H_{B_2, B_1}(v_2 \otimes v_1) = 0$, where v_1 and v_2 are the generators defined above.

Definition 5.5. For $B = B^{r_N, s_N} \otimes \cdots \otimes B^{r_1, s_1}$, $1 \leq i \leq N$ and $i < j \leq N$, set

$$D_i := D_{B^{r_i, s_i}} \sigma_1 \sigma_2 \cdots \sigma_{i-1} \quad \text{and} \quad H_{j,i} := H_i \sigma_{i+1} \sigma_{i+2} \cdots \sigma_{j-1},$$

where σ_i and H_i act on the i -th and $(i+1)$ -st tensor factors and $D_{B^{r_i, s_i}}$ is the D -function on the rightmost tensor factor B^{r_i, s_i} as given in Definition 5.4. The D -function $D_B : B \rightarrow \mathbb{Z}$ is defined as

$$(5.2) \quad D_B := \sum_{N \geq j > i \geq 1} H_{j,i} + \sum_{i=1}^N D_i.$$

Where it does not cause confusion, we shorten D_B to simply D .

6. PERFECT KR CRYSTALS AND DEMAZURE CRYSTALS

We now state a precise relationship between KR crystals and Demazure crystals (see Theorem 6.1). This was proven by Fourier, Schilling, and Shimozono [12], under a few additional assumptions on the KR crystals since at the time the existence and combinatorial models for KR crystals did not yet exist. Here we point out that most of the assumptions in [12] follow from the later results of [49, 50, 10, 11] showing that the relevant KR crystals exist, have certain symmetries related to Dynkin diagram automorphisms, and are perfect. In the special cases of type $A_{2n}^{(2)}$ and the exceptional nodes in type $D_n^{(1)}$, we give separate proofs as the assumptions from [12] do not follow directly from the above papers or need to be slightly modified.

Theorem 6.1. *Let $B = B^{r_N, \ell c_{r_N}} \otimes \cdots \otimes B^{r_1, \ell c_{r_1}}$ be a level- ℓ composite KR crystal of non-exceptional type. Define $\lambda = -(c_{r_1} \omega_{r_1^*} + \cdots + c_{r_N} \omega_{r_N^*})$, where r^* is defined by $\omega_{r^*} = -w_0(\omega_r)$ with w_0 the longest element of \overline{W} , and write $t_\lambda \in T(\widetilde{M}) \subset \widetilde{W}$ as $t_\lambda = v\tau$. Then there is a unique isomorphism of affine crystals*

$$j : B(\ell\Lambda_{\tau(0)}) \rightarrow B \otimes B(\ell\Lambda_0).$$

This satisfies

$$j(u_{\ell\Lambda_{\tau(0)}}) = u_B \otimes u_{\ell\Lambda_0},$$

where u_B is the distinguished element from Definition 5.2, and

$$(6.1) \quad j(B_v(\ell\Lambda_{\tau(0)})) = B \otimes u_{\ell\Lambda_0},$$

where $B_v(\ell\Lambda_{\tau(0)})$ is the Demazure crystal corresponding to the translation t_λ as defined in Section 2.9.

We delay the proof of Theorem 6.1 until the end of this section.

Remark 6.2. For non-exceptional types, we have $r^* = r$ except for type $A_n^{(1)}$ where $r^* = n + 1 - r$, and $D_n^{(1)}$ for n odd where $n^* = n - 1$ and $(n - 1)^* = n$.

Lemma 6.3. *Assume \mathfrak{g} is of non-exceptional type, and let $B^{r, \ell c_r}$ be a level- ℓ KR crystal. Then:*

- (i) *There is a unique element $u \in B^{r, \ell c_r}$ such that*

$$\varepsilon(u) = \ell\Lambda_0 \quad \text{and} \quad \varphi(u) = \ell\Lambda_{\tau(0)},$$

where $t_{-c_r \omega_{r^*}} = v\tau$ with $v \in W$ and $\tau \in \Sigma$.

- (ii) *Let ς be the Dynkin diagram automorphism defined by*

- *For type $A_n^{(1)}$, $i \mapsto i + 1 \pmod{n + 1}$.*
- *For types $B_n^{(1)}$, $D_n^{(1)}$, $A_{2n-1}^{(2)}$, exchange nodes 0 and 1.*
- *For types $C_n^{(1)}$ and $D_{n+1}^{(2)}$, $i \mapsto n - i$ for all $i \in I$.*

In all cases other than type $A_{2n}^{(2)}$ and $B^{n-1, s}$ and $B^{n, s}$ for type $D_n^{(1)}$, there is a unique involution of $B^{r, \ell c_r}$, also denoted ς , such that for all $b \in B^{r, \ell c_r}$ and $i \in I$,

$$f_i(b) = \varsigma^{-1} \circ f_{\varsigma(i)} \circ \varsigma(b).$$

Remark 6.4. Lemma 6.3 was stated as [12, Assumption 1] (with the misprint $t_{-c_r \omega_r}$ instead of $t_{-c_r \omega_{r^*}}$ in (i)). This Assumption also included the requirement that $B^{r, \ell c_r}$ is regular. Since we now know that $B^{r, \ell c_r}$ is the crystal of a KR module by [49, 50, 10], this is immediate, and so we do not include it in Lemma 6.3. The

Type	$u \in B^{r, \ell c_r}$	r	c_r	τ
$A_n^{(1)}$	$u(\ell\Lambda_r)$	$1 \leq r \leq n$	1	pr^r
$B_n^{(1)}$	$u(\emptyset)$	$1 \leq r < n, r \text{ even}$	1	id
	$u(\ell\Lambda_1)$	$1 \leq r < n, r \text{ odd}$	1	$\varsigma_{0,1}$
	$u(\emptyset)$	$r = n \text{ even}$	2	id
	$u(\ell\Lambda_1)$	$r = n \text{ odd}$	2	$\varsigma_{0,1}$
$C_n^{(1)}$	$u(\emptyset)$	$1 \leq r < n$	2	id
	$u(\ell\Lambda_n)$	$r = n$	1	$\varsigma_{\leftrightarrow}$
$D_n^{(1)}$	$u(\emptyset)$	$1 \leq r \leq n-2, r \text{ even}$	1	id
	$u(\ell\Lambda_1)$	$1 \leq r \leq n-2, r \text{ odd}$	1	$\varsigma_{0,1}\varsigma_{n-1,n}$
	$u(\ell\Lambda_{n-1})$	$r = n-1$	1	$\varsigma_{\leftrightarrow}\varsigma_{0,1}\varsigma_{n-1,n}^{n+1}$
	$u(\ell\Lambda_n)$	$r = n$	1	$\varsigma_{\leftrightarrow}\varsigma_{n-1,n}^n$
$A_{2n-1}^{(2)}$	$u(\emptyset)$	$1 \leq r \leq n, r \text{ even}$	1	id
	$u(\ell\Lambda_1)$	$1 \leq r \leq n, r \text{ odd}$	1	$\varsigma_{0,1}$
$D_{n+1}^{(2)}$	$u(\emptyset)$	$1 \leq r < n$	1	id
	$u(\ell\Lambda_n)$	$r = n$	1	$\varsigma_{\leftrightarrow}$
$A_{2n}^{(2)}$	$u(\emptyset)$	$1 \leq r \leq n$	1	id

FIGURE 4. The elements u and related data. Here $u(\lambda)$ is the highest weight vector in the classical component $B(\lambda)$, pr is the map $i \mapsto i+1 \pmod{n+1}$, and $\varsigma_{0,1}$ (resp. $\varsigma_{n-1,n}$) is the map that interchanges 0 and 1 (resp. $n-1$ and n) and fixes all other i . The map $\varsigma_{\leftrightarrow}$ is $i \mapsto n-i$. In the expression for τ the maps act on the left, so that e.g. $\varsigma_{\leftrightarrow}\varsigma_{0,1}(1) = \varsigma_{\leftrightarrow}(0) = n$.

statement in [12] also included an additional assumption for type $A_{2n}^{(2)}$, which we omit as we deal with type $A_{2n}^{(2)}$ separately.

Proof of Lemma 6.3. By perfectness of $B^{r, \ell c_r}$ for non-exceptional types [11], there exists a unique element $u \in B^{r, \ell c_r}$ such that $\varepsilon(u) = \ell\Lambda_0$. These elements are listed for all non-exceptional types in Figure 4. It is easy to check explicitly that these also satisfy $\varphi(u) = \ell\Lambda_{\tau(0)}$, where τ is as listed. Furthermore, one can easily check that all these τ satisfy the remaining conditions of (i).

The combinatorial model for $B^{r,s}$ of type $A_n^{(1)}$ uses the promotion operator which corresponds to the Dynkin diagram automorphism mapping $i \mapsto i+1 \pmod{n+1}$. Similarly, the KR crystals in case (AUT) are constructed using the automorphism ς (see Section 4.3).

By Theorem 4.7 the KR crystal $B^{n,2\ell}$ of type $B_n^{(1)}$ can be constructed as a virtual crystal inside $\hat{B}^{n,2\ell}$ of type $A_{2n-1}^{(2)}$ with $e_0 = \hat{e}_0^2$ and $e_1 = \hat{e}_1^2$. The virtual crystal is constructed using the analogue of the Dynkin automorphism $\hat{\varsigma}$ exchanging 0 and 1, so that $e_0 = \hat{e}_0^2 = (\hat{\varsigma}\hat{e}_1\hat{\varsigma})(\hat{\varsigma}\hat{e}_1\hat{\varsigma}) = \hat{\varsigma}\hat{e}_1^2\hat{\varsigma} = \hat{\varsigma}e_1\hat{\varsigma}$. Furthermore, by [10, Lemma 3.5] the image of $B^{n,2\ell}$ in the ambient crystals $\hat{B}^{n,2\ell}$ is closed under $\hat{\varsigma}$. Hence on $B^{n,2\ell}$ of type $B_n^{(1)}$ there also exists an isomorphism interchanging nodes 0 and 1, induced by $\hat{\varsigma}$.

It was shown in [10, Theorem 7.1] that $B^{r,s}$ of types $C_n^{(1)}$ and $D_{n+1}^{(2)}$ admit a twisted isomorphism ς corresponding to the Dynkin automorphism mapping $i \mapsto n - i$ for $i \in I$. This shows (ii). \square

For B a crystal of affine type and $\tau \in \text{Aut}(\Gamma)$, where Γ is the affine Dynkin diagram, let B^τ be the crystal with the same underlying set as B , but where $e_i^\tau = \tau \circ e_i \circ \tau^{-1}$ and $f_i^\tau = \tau \circ f_i \circ \tau^{-1}$.

Lemma 6.5. *Let Γ be a non-exceptional affine Dynkin diagram and $\tau \in \Sigma$. Then $(B^{r,s})^\tau \cong B^{r,s}$ for all $r \in I \setminus \{0\}$ and $s \geq 1$.*

Proof. Fix $\tau \in \Sigma$. By Lemma 2.8, there exists $w_\tau \in \overline{W}$ such that

$$(6.2) \quad \begin{array}{ccc} I & \xrightarrow{\iota} & \Delta \\ \downarrow \tau & & \downarrow w_\tau \\ I & \xrightarrow{\iota} & \Delta \end{array}$$

commutes, where ι is the map that takes i to α_i for all $i \neq 0$, and 0 to $-\theta$ (where $\theta = \delta - \alpha_0$). Thus the character of $(B^{r,s})^\tau$ as a $\{1, 2, \dots, n\}$ -crystal is in the Weyl group orbit of the character of $B^{r,s}$ as a $\{1, 2, \dots, n\}$ -crystal, and hence by Weyl group invariance these are equal. By the classification of classical crystals, it follows that $(B^{r,s})^\tau \cong B^{r,s}$ as a $\{1, 2, \dots, n\}$ -crystal. Thus the lemma follows by Lemma 2.6. \square

Lemma 6.6. *Theorem 6.1 holds in type $D_n^{(1)}$.*

Proof. For $B = B^{r,\ell}$ with $1 \leq r \leq n - 2$, Lemma 6.3 shows that [12, Assumption 1] holds, and so Theorem 6.1 holds by [12, Theorem 4.4].

Now consider $B^{n,\ell}$, noting that $c_n = 1$. We need the following notation:

- As in Theorem 6.1, n^* is defined by $\omega_{n^*} = -w_0(\omega_n)$, so that $(w_0^{n^*})^{-1} = w_0^n$.
- $w_0^n \in \overline{W}$ is of minimal length such that $w_0^n(\omega_n) = w_0(\omega_n)$,
- $w_0^{n^*}$ is of minimal length such that $w_0^{n^*}(\omega_{n^*}) = w_0(\omega_{n^*})$,
- τ_n and τ_{n^*} are the τ from Table 4 for $r = n, n^*$ respectively.

As in [12, Equation (2.10)] (but noting that due to differing conventions their τ_{n^*} is our τ_n),

$$(6.3) \quad t_{-\omega_{n^*}} = (w_0^{n^*})^{-1} \tau_n = w_0^n \tau_n.$$

Since $w_0^n(\ell\Lambda_n) = w_0(\ell\Lambda_n)$, we see that

$$(6.4) \quad B_{w_0^n}(\ell\Lambda_n) = B_{w_0}(\ell\Lambda_n) = \{f_{i_N}^{m_N} \cdots f_{i_1}^{m_1} u_{\ell\Lambda_n} \mid m_1, \dots, m_N \geq 0\},$$

where $s_{i_1} \cdots s_{i_N}$ is a reduced word for w_0 and the last equality follows by [27, Proposition 3.2.3] (or see Section 2.9 above).

As in Section 4.1, $B^{n,\ell}$ is isomorphic to $B(\ell\Lambda_n)$ as a classical crystal. The element u from Figure 4 is the highest weight element $u(\ell\Lambda_n)$ of weight $\ell\Lambda_n$. It satisfies $\varepsilon(u(\ell\Lambda_n)) = \ell\Lambda_0$ and $\varphi(u(\ell\Lambda_n)) = \ell\Lambda_n$. Furthermore $B^{n,\ell}$ is perfect of level ℓ so, as in [25], there is a unique isomorphism

$$(6.5) \quad \begin{aligned} j : B(\ell\Lambda_n) &\rightarrow B^{n,\ell} \otimes B(\ell\Lambda_0) \\ u_{\ell\Lambda_n} &\mapsto u(\ell\Lambda_n) \otimes u_{\ell\Lambda_0}. \end{aligned}$$

Since $\varepsilon_i(u_{\ell\Lambda_0}) = \varphi_i(u_{\ell\Lambda_0}) = 0$ for all $i \neq 0$, (6.4) and (6.5) imply

$$(6.6) \quad j(B_{w_0^n}(\ell\Lambda_n)) = B^{n,\ell} \otimes u_{\ell\Lambda_0}.$$

Since $\tau_n(0) = n$, we have proven Theorem 6.1 for $B = B^{n,s}$. A similar argument shows that Theorem 6.1 holds for $B = B^{n-1,s}$.

By Lemma 6.5, each automorphism $\tau \in \Sigma$ induces a bijection of $B^{r,\ell}$ to itself that sends i arrows to $\tau(i)$ arrows. Thus, as in [12, Theorem 4.7], a straightforward induction argument shows that Theorem 6.1 holds for all composite level- ℓ KR crystals. \square

Lemma 6.7. *Theorem 6.1 holds in type $A_{2n}^{(2)}$.*

Proof. In [12] a proof of Theorem 6.1 was given, but not with the concrete combinatorial model given in Theorem 4.9. The only place, where the concrete combinatorial model for type $A_{2n}^{(2)}$ is used in [12] is in the proof of [12, Lemma 4.3]. Thus we must verify that this lemma holds with the explicit realization of KR crystals for this type from [10] or equivalently Theorem 4.9. Let $S : B^{r,s} \rightarrow \hat{B}_{C_n^{(1)}}^{r,2s}$ be the unique injective map from the combinatorial KR crystal of type $A_{2n}^{(2)}$ to the KR crystal of type $C_n^{(1)}$ such that

$$S(e_i b) = \hat{e}_i^{m_i} S(b) \quad \text{and} \quad S(f_i b) = \hat{f}_i^{m_i} S(b) \quad \text{for all } i \in I \text{ and } b \in B^{r,s},$$

where $m = (m_0, \dots, m_n) = (1, 2, \dots, 2)$. Using the explicit realization of $\hat{B}_{C_n^{(1)}}^{r,2s}$ it is not too hard to check that for the classically highest weight element $u(s\omega_k) \in B(s\omega_k) \subset B^{r,s}$ for $k < r$ and $y := S_1 \cdots S_k(u(s\omega_k))$ (where S_i is f_i raised to the maximal power), we have $S(f_0^s y) = S(u(s\omega_{k+1}))$. Thus [12, Lemma 4.3] holds. Furthermore, in this case τ is always the identity. Thus Theorem 6.1 holds by [12, Theorem 4.7]. \square

Proof of Theorem 6.1. If $X = D_n^{(1)}$ or $A_{2n}^{(2)}$, the Theorem holds by Lemmas 6.6 and 6.7. In all other cases, Lemma 6.3 shows that [12, Assumption 1] holds. Furthermore, in all these cases Σ is generated by the diagram automorphism ς from Lemma 6.3. Thus the theorem holds by [12, Theorem 4.7]. \square

7. THE AFFINE GRADING VIA THE ENERGY FUNCTION

In this section, we show that for $B = B^{r_N, \ell c_{r_N}} \otimes \cdots \otimes B^{r_1, \ell c_{r_1}}$ a composite level- ℓ KR crystal the map j from Theorem 6.1 intertwines the D function from Section 5.2 with the affine degree deg given in Definition 7.1 below up to a shift. This allows us to show that j intertwines E^{int} with deg exactly, and in particular E^{int} agrees with D up to a shift.

Definition 7.1. For any Demazure crystal $B_w(\Lambda)$, where $w \in W$ and Λ is a dominant integral weight, let

$$\text{deg} : B_w(\Lambda) \rightarrow \mathbb{Z}_{\geq 0}$$

be the affine degree map defined by $\text{deg}(u_\Lambda) = 0$ and each f_i has degree $\delta_{i,0}$.

We begin by two preliminary lemmas, first for a single KR crystal and then tensor products of KR crystals.

Lemma 7.2. *Fix a KR crystal $B^{r,s}$ of type \mathfrak{g} . Then $D(e_0(b)) \geq D(b) - 1$ for all $b \in B^{r,s}$. Furthermore, if $\varepsilon_0(b) > \lceil s/c_r \rceil$, then we have $D(e_0(b)) = D(b) - 1$.*

Proof.

- **Case (IRR):** In this case $\diamond = \emptyset$, so that by Definition 5.4 the D function is the constant function 0 which satisfies $D(e_0(b)) \geq D(b) - 1$ for all $b \in B^{r,s}$. By Lemma 4.1 we always have $\varepsilon_0(b) \leq s/c_r$ since $c_r = 1$.
- **Case (AUT):** In this case \diamond is a vertical domino. The statements follow immediately from Lemma 4.4 since again $c_r = 1$ in all cases.
- For $B^{r,s}$ of type $C_n^{(1)}$ with $r < n$, we have $c_r = 2$ and \diamond is a horizontal domino. The result follows from Lemma 4.6.
- For $B^{n,s}$ of type $B_n^{(1)}$, we have $c_r = 2$ and \diamond is a vertical domino. The result follows from Lemma 4.8.
- For $B^{r,s}$ of types $A_{2n}^{(2)}$ with $r \leq n$ and $D_{n+1}^{(2)}$ with $r \leq n-1$, we have $c_r = 1$ and \diamond is a single box. The result follows from Lemma 4.10.

□

Lemma 7.3. *Let $B = B^{r_N, s_N} \otimes \cdots \otimes B^{r_1, s_1}$ be a tensor product of KR crystals and fix an integer ℓ such that $\ell \geq \lceil s_k/c_k \rceil$ for all $1 \leq k \leq N$. If $e_0(b) \neq 0$ then $D(e_0(b)) \geq D(b) - 1$, and if $\varepsilon_0(b) > \ell$ then this is an equality.*

Proof. Write $b = b_N \otimes \cdots \otimes b_1$. For some $N \geq k \geq 1$,

$$e_0(b) = b_N \otimes \cdots \otimes e_0(b_k) \otimes \cdots \otimes b_1.$$

Let

$$b'_N \otimes \cdots \otimes b'_1 = \sigma_1 \cdots \sigma_{k-1}(b).$$

Since each σ_j commutes with the action of e_0 ,

$$e_0(b'_N \otimes \cdots \otimes b'_1) = b'_N \otimes \cdots \otimes e_0(b'_c) \otimes \cdots \otimes b'_1$$

for some $k \geq c \geq 1$. For each $1 \leq x \leq N$, let

$$(7.1) \quad D^{[x]} = \sum_{1 \leq y < x} H_{x,y} + D_x,$$

where $H_{x,y}$ and D_x are as in Definition 5.5, so that $D = \sum_{1 \leq x \leq N} D^{[x]}$. Let $\Delta(b) = D(b) - D(e_0(b))$ and $\Delta^{[x]}(b) = D^{[x]}(b) - D^{[x]}(e_0(b))$. We find each $\Delta^{[x]}(b)$, considering three cases.

Case 1, $k > x$: Here $\Delta^{[x]}(b) = 0$, as all terms in $D^{[x]}(b)$ and $D^{[x]}(e_0(b))$ agree.

Case 2, $x = k$: If $c \neq 1$, then, by Definitions 5.4 and 5.5, $D_x(b) = D_x(e_0(b))$ and by (5.1), $H_{x,c-1}(e_0(b)) = H_{x,c-1}(b) - 1$. For all other $y < x$ it is clear that $H_{x,y}(b) = H_{x,y}(e_0(b))$. Hence $\Delta^{[x]}(b) = 1$.

Otherwise $c = 1$ and

$$\Delta^{[x]}(b) = D^{[x]}(b) - D^{[x]}(e_0(b)) = D_{B^{r_1, s_1}}(b'_1) - D_{B^{r_1, s_1}}(e_0(b'_1)).$$

In this case $\varepsilon(b'_1) = \varepsilon(b)$, so by Lemma 7.2, $\Delta^{[x]}(b) \leq 1$, with equality if $\varepsilon_0(b) > \ell \geq s_{r_1}/c_{r_1}$.

Case 3, $x > k$: It is clear that $H_{x,y}(b) = H_{x,y}(e_0(b))$ for $x > y > k$. Let

$$d = d_N \otimes \cdots \otimes d_1 := \sigma_k \cdots \sigma_{x-1}(b).$$

Since each σ_j commutes with the action of e_0 , one of the following must hold:

- $e_0(d_N \otimes \cdots \otimes d_1) = d_N \otimes \cdots \otimes e_0(d_{k+1}) \otimes d_k \otimes \cdots \otimes d_1$,
- $e_0(d_N \otimes \cdots \otimes d_1) = d_N \otimes \cdots \otimes d_{k+1} \otimes e_0(d_k) \otimes \cdots \otimes d_1$.

If (a) holds, then for $y < k$ we have $H_{x,y}(b) = H_{x,y}(e_0(b))$, since $H_{x,y}$ is calculated on exactly the same tensor product on each side. By (5.1) we have $H_{x,k}(b) = H_{x,k}(e_0(b))$ and by Definition 5.4 $D_x(b) = D_x(e_0(b))$. Hence $\Delta^{[x]}(b) = 0$.

If (b) holds, then, by (5.1), $H_{x,k}(e_0(b)) = H_{x,k}(b) + 1$, and for all $k < y < x$, $H_{x,y}(e_0(b)) = H_{x,y}(b)$. Furthermore, by Definition 5.5 $D_x(b) = D_k(d)$ and $D_x(e_0b) = D_k(e_0d)$. Comparing terms and noticing that $H_{x,r}(b) = H_{k,r}(d)$ and $H_{x,r}(e_0b) = H_{k,r}(e_0d)$ for all $r < x$ shows

$$\Delta^{[x]}(b) = \Delta^{[k]}(d) - 1,$$

so it follows by the argument in Case 2 that $\Delta^{[x]}(b) \leq 0$, with equality if $\varepsilon_0(b) = \varepsilon_0(d) > \ell$.

The result now follows by adding all the $\Delta^{[x]}(b)$. \square

Theorem 7.4. *With the same assumptions and notation as in Theorem 6.1, let $\tilde{j} : B_v(\ell\Lambda_{\tau(0)}) \rightarrow B$ be the restriction of the map j to $B_v(\ell\Lambda_{\tau(0)})$, where $B \otimes u_{\ell\Lambda_0}$ is identified with just B . Then for all $b \in B_v(\ell\Lambda_{\tau(0)})$ we have $\deg(b) = D(\tilde{j}(b)) - D(u_B)$.*

Proof. Since $B_v(\ell\Lambda_{\tau(0)})$ is connected with the highest weight element $u_{\ell\Lambda_{\tau(0)}}$, $\tilde{j}(u_B) = u_{\ell\Lambda_{\tau(0)}}$, and $\deg(u_{\ell\Lambda_{\tau(0)}}) = 0$, it suffices to show that $D(\tilde{j}(b)) = D(\tilde{j}(e_0(b))) - 1$ for each $b \in B_v(\ell\Lambda_{\tau(0)})$ such that $e_0(b) \neq 0$. So, choose such a $b \in B_v(\ell\Lambda_{\tau(0)})$. Since e_0 acts non-trivially on $\tilde{j}(b) \otimes u_{\ell\Lambda_0}$, we have $\varepsilon_0(\tilde{j}(b)) > \ell$, and the result follows by Lemma 7.3. \square

Theorem 7.5. *For any composite level- ℓ KR crystal $B = B^{r_N, \ell c_{r_N}} \otimes \dots \otimes B^{r_1, \ell c_{r_1}}$ we have $E^{\text{int}} = D - D(u_B)$.*

Proof. By the definition of E^{int} (see Section 5.1), for any b there exists a string of f_i involving exactly $E^{\text{int}}(b)$ factors f_0 taking u_B to b . By Lemma 7.3, applying f_0 increases the value of D by at most 1, and so it follows that $E^{\text{int}}(b) \geq D(b) - D(u_B)$. It follows from Theorem 7.4 and the fact that the Demazure crystal $B_v(\ell\Lambda_{\tau(0)})$ is connected that $E^{\text{int}}(b) \leq D(b) - D(u_B)$. \square

Corollary 7.6. *The map j intertwines E^{int} and the basic grading \deg .*

Proof. This is immediate from Theorems 7.4 and 7.5. \square

Remark 7.7. Let B be a composite level- ℓ KR crystal. If $\varepsilon_0(b) > \ell$, call the corresponding 0 arrow from $b' = e_0(b)$ to b in B a *Demazure arrow*. We have shown that for such an arrow $D(b) = D(b') + 1$. Furthermore, we have shown that u_B is the unique source in the subgraph of B consisting only of classical arrows (i.e. arrows colored i for some $i \neq 0$) and Demazure arrows. This could be used to give an algorithm for calculating the energy function: given b , move backwards along arrows until one reach u_B , only using Demazure 0-arrows. Then $D(b) - D(u_B)$ is the number of 0 arrows in this path.

8. GENERALIZATIONS TO OTHER TENSOR PRODUCTS OF KR CRYSTALS

Notice that Lemma 7.3 holds for more general tensor products of KR crystals, not just tensor products of level- ℓ perfect KR crystals. We now show how Theorem 7.4 and Corollary 7.6 can be generalized as well.

For this section, fix \mathfrak{g} of non-exceptional affine type, $\ell > 0$, and a tensor product $B = B^{r_N, s_N} \otimes \cdots \otimes B^{r_1, s_1}$ of KR crystals, such that $\ell \geq \lceil s_k/c_k \rceil$ for all $1 \leq k \leq N$. We call such a crystal a *composite KR crystal of level bounded by ℓ* .

The proof of the following proposition is similar to [25, Proof of Theorem 4.4.1].

Proposition 8.1. *For B a composite KR crystal of level bounded by ℓ ,*

$$B \otimes B(\ell\Lambda_0) \cong \bigoplus_{\Lambda'} B(\Lambda'),$$

where the sum is over a finite collection of (not necessarily distinct) $\Lambda' \in P_\ell^+$.

Proof. Fix $b \otimes c \in B \otimes B(\ell\Lambda_0)$. If $c \neq u_{\ell\Lambda_0}$, then for some $i \in I$ we have $e_i(c) \neq 0$. This implies that, for some $k \geq 1$,

$$(8.1) \quad e_i^k(b \otimes c) = e_i^{k-1}(b) \otimes e_i(c) \neq 0.$$

In this way, one can apply raising operators to $b \otimes c$ to obtain an element $b' \otimes u_{\ell\Lambda_0}$.

Set $M = \dim(B)$, and assume that, for some choice of e_{i_1}, \dots, e_{i_M} , and some $b \in B$, $e_{i_M} \cdots e_{i_1}(b \otimes u_{\ell\Lambda_0}) \neq 0$. Then some element $b' \otimes u_{\ell\Lambda_0}$ must appear twice. Thus for some $1 \leq r \leq s \leq M$, we have $e_{i_s} \cdots e_{i_r}(b' \otimes u_{\ell\Lambda_0}) = b' \otimes u_{\ell\Lambda_0}$. By weight considerations, at least one e_0 must appear in this sequence. Furthermore, whenever $e_{i_k} = e_0$, we must have $\varepsilon_0(e_{i_{k-1}} \cdots e_{i_r}(b')) > \ell$. Thus Lemma 7.3 implies that $D(b') < D(b')$, which is a contradiction. Hence the left hand side does decompose as a finite direct sum of $B(\Lambda')$ for $\Lambda' \in P^+$. It follows by weight considerations that all Λ' are in P_ℓ^+ . \square

Definition 8.2. For each $b \in B$, let $u_b^{\ell\Lambda_0}$ be the unique element of B such that $u_b^{\ell\Lambda_0} \otimes u_{\ell\Lambda_0}$ is the highest weight in the component from Proposition 8.1 containing $b \otimes u_{\ell\Lambda_0}$.

Define the function \deg on a direct sum of highest weight crystals to be the basic grading on each component, with all highest weight elements placed in degree 0.

Corollary 8.3. *Choose an isomorphism $m : B \otimes B(\ell\Lambda_0) \cong \bigoplus_{\Lambda'} B(\Lambda')$. Then for all $b \in B$, we have $D(b) - D(u_b^{\ell\Lambda_0}) = \deg(m(b \otimes u_{\ell\Lambda_0}))$.*

Proof. This follows from Lemma 7.3, just as in the proof of Theorem 7.4. \square

The following should be interpreted as a generalization of Corollary 7.6.

Corollary 8.4. *The minimal number of e_0 in a string of e_i taking b to $u_b^{\ell\Lambda_0}$ is $D(b) - D(u_b^{\ell\Lambda_0})$.*

Proof. This follows as in the proof of Corollary 7.6, using the fact that, for any $b \in B$, $b \otimes u_{\ell\Lambda_0}$ is connected to $u_b^{\ell\Lambda_0} \otimes u_{\ell\Lambda_0}$ in the affine highest weight crystal. \square

Remark 8.5. The statements in this section do not give a relationship with Demazure crystals. However, Naoi [46, Proposition after Theorem B, Proposition 7.6] and [47] has recently proven that $B \otimes u_\Lambda$ is isomorphic to a disjoint union of Demazure crystals inside the various irreducible components of $B \otimes B(\Lambda)$, where Λ is an arbitrary dominant weight of level $\ell \geq 1$ and B is a tensor product of perfect KR crystals each of which has level at most ℓ . It would be interesting to determine whether or not this continues to hold for B nonperfect.

9. APPLICATIONS

In this section we discuss how the relation between the affine grading in the Demazure crystal and the energy function can be used to derive a formula for the Demazure character using the energy function, as well as showing how they are related to nonsymmetric Macdonald polynomials and Whittaker functions.

9.1. Demazure characters. Since the character of a Demazure module $V_w(\lambda)$ can be expressed in terms of the Demazure crystal by (2.9), we have the following immediate corollary of Theorem 6.1.

Corollary 9.1. *Let $B = B^{r_N, \ell c_{r_N}} \otimes \dots \otimes B^{r_1, \ell c_{r_1}}$ be a $U'_q(\mathfrak{g})$ -composite level- ℓ KR crystal, $\lambda = -(c_{r_1} \omega_{r_1^*} + \dots + c_{r_N} \omega_{r_N^*})$, and $t_\lambda = v\tau$ as in Theorem 6.1. Then*

$$(9.1) \quad \text{ch} V_v(\ell \Lambda_{\tau(0)}) = e^{\ell \Lambda_0} \sum_{b \in B} e^{\text{wt}(b) - \delta E^{\text{int}}(b)} = e^{\ell \Lambda_0} \sum_{b \in B} e^{\text{wt}^{\text{aff}}(b)},$$

where $\text{wt}^{\text{aff}}(b) = \text{wt}(b) - \delta E^{\text{int}}(b)$ and $\text{wt}(b)$ is the $U'_q(\mathfrak{g})$ -weight of b .

Proof. Recall that by Theorem 6.1 we have the isomorphism

$$j(B_v(\ell \Lambda_{\tau(0)})) = B \otimes u_{\ell \Lambda_0}.$$

Combining this with (2.9), we obtain the result. \square

As in e.g. [16, 17], the one-dimensional configuration sums are defined as

$$(9.2) \quad X(\mu; B) = \sum_{\substack{b \in B, \text{wt}(b) = \mu \\ e_i(b) = 0 \text{ for } i \in I \setminus \{0\}}} q^{-D(b)},$$

where $B = B^{r_N, s_N} \otimes \dots \otimes B^{r_1, s_1}$. By Theorem 7.5 we can rewrite Corollary 9.1 as follows (compare also with [54, Theorem 1.2]).

Corollary 9.2. *With the same assumptions and notation as in Corollary 9.1 and setting $q = e^\delta$ we have*

$$(9.3) \quad \text{ch} V_v(\ell \Lambda_{\tau(0)}) = e^{\ell \Lambda_0} q^{D(u_B)} \sum_{\mu} X(\mu; B) \text{ch}(V(\mu)),$$

where $V(\mu)$ is the module of highest weight μ for the underlying finite type algebra.

Example 9.3. Let us illustrate various quantities used in this section. Consider $B = (B^{1,1})^{\otimes 3}$ of type $A_2^{(1)}$. This has $u_B = 3 \otimes 2 \otimes 1$. Take $b = 2 \otimes 3 \otimes 1$. Then one can calculate $D(b) = -2$ and $D(u_B) = -3$, so that $E^{\text{int}}(b) = D(b) - D(u_B) = 1$. Also

$$\text{wt}(b) = (\Lambda_2 - \Lambda_1) + (\Lambda_0 - \Lambda_2) + (\Lambda_1 - \Lambda_0) = 0,$$

so that $\text{wt}^{\text{aff}}(b) = -\delta$.

9.2. Nonsymmetric Macdonald polynomials. Fix \mathfrak{g} of affine type. Let $\tilde{P} \subset P$ be the sublattice of level 0 weights. Recall that \tilde{P} is naturally contained in $\overline{P} + \mathbb{Z}\delta$, where δ is the null root and we identify $\overline{P} \otimes_{\mathbb{Z}} \mathbb{R}$ with a subspace of $P \otimes_{\mathbb{Z}} \mathbb{R}$ by identifying the finite type simple roots with their corresponding affine simple roots (and this containment is equality except in type $A_{2n}^{(2)}$). Let t be the collection of

indeterminates t_α for each root α such that $t_\alpha = t_{\alpha'}$ if α and α' have the same length. Consider the following elements of the group algebra $\mathbb{Q}(q, t)\overline{P}$:

$$\Delta := \prod_{\alpha \in R_+^{\text{aff}}} \frac{1 - e^\alpha}{1 - t_\alpha e^\alpha} \Big|_{e^\delta = q}, \quad \text{and} \quad \Delta_1 := \Delta / ([e^0] \Delta),$$

where $[e^0]$ means the coefficient of e^0 and R_+^{aff} is the set of positive affine real roots. Cherednik's inner product [4] on $\mathbb{Q}(q, t)\overline{P}$ is $\langle f, g \rangle_{q, t} = [e^0](f \overline{g} \Delta_1)$, where $\overline{\cdot}$ is the involution $\overline{q} = q^{-1}$, $\overline{t} = t^{-1}$, $\overline{e^\lambda} = e^{-\lambda}$.

The nonsymmetric Macdonald polynomials $E_\lambda(q, t) \in \mathbb{Q}(q, t)\overline{P}$ for $\lambda \in \overline{P}$ were introduced by Opdam [48] in the differential setting and Cherednik [4] in general (although here we follow conventions of Haglund, Haiman, Loehr [14, 15]). They are uniquely characterized by two conditions: (Triangularity): $E_\lambda \in x^\lambda + \mathbb{Q}(q, t)\{x^\mu \mid \mu < \lambda\}$ and (Orthogonality): $\langle E_\lambda, E_\mu \rangle_{q, t} = 0$ for $\lambda \neq \mu$. Here $<$ is the Bruhat ordering on \overline{P} identified with the set of minimal coset representatives in $\widetilde{W}/\overline{W}$, where \widetilde{W} is the extended affine Weyl group and \overline{W} is the classical Weyl group.

Extending Sanderson's work [53] for type A , Ion [21] showed that for all simply laced untwisted affine root systems, the specialization of the nonsymmetric Macdonald polynomial $E_\lambda(q, t)$ at $t = 0$ coincides with a specialization of a Demazure character of a level one affine integrable module (see [53, Theorem 6], [21]): Write $t_\lambda \in \widetilde{W}$ as $t_\lambda = w\tau$, where $w \in W, \tau \in \Sigma$. Then

$$(9.4) \quad E_\lambda(q, 0) = q^c \text{ch}(V_w(\Lambda_{\tau(0)})) \Big|_{e^\delta = q, e^{\Lambda_0} = 1},$$

where c is a specific exponent described in [21, 53] (and described in Corollary 9.5 below in types $A_n^{(1)}$ and $D_n^{(1)}$). For λ a single row, this relation also follows from combining [37, 20].

Remark 9.4. We have used the conventions of Haglund, Haiman, Loehr [14, 15], which differ from those in [21] by the change of variables $q \rightarrow q^{-1}$.

If λ is anti-dominant and $\mathfrak{g} = A_n^{(1)}$ or $D_n^{(1)}$, we can apply Theorem 6.1 and Corollary 7.6 to give a connection with KR crystals, along with their energy.

Corollary 9.5. *Fix $\mathfrak{g} = A_n^{(1)}$ or $D_n^{(1)}$. Fix an anti-dominant weight λ , and write $\lambda = -(\omega_{r_1^*} + \cdots + \omega_{r_N^*})$. Let $B = B^{r_N, 1} \otimes \cdots \otimes B^{r_1, 1}$. Then*

$$P_\lambda(q, 0) = E_\lambda(q, 0) = \sum_{b \in B} q^{-D(b)} e^{\text{wt}(b)}.$$

Proof. The equality $P_\lambda(q, 0) = E_\lambda(q, 0)$ is [21, Theorem 4.2]. By Theorem 6.1 and Corollary 8.3 (noting that $c_r = 1$ for types $A_n^{(1)}$ and $D_n^{(1)}$ as in Figure 4), equation (9.4) implies

$$P_\lambda(q, 0) = q^c \sum_{b \in B} q^{-D(b) + D(u_B)} e^{\text{wt}(b)}.$$

It remains to show that $c = -D(u_B)$. Note that the generator $v = v_{r_N, 1} \otimes \cdots \otimes v_{r_1, 1} \in B$ has weight $\text{wt}(v) = -\lambda^* = w_0(\lambda)$. Since $P_\lambda(q, 0)$ is symmetric, the coefficient of $e^{w_0(\lambda)}$ is the same as the coefficient of e^λ , which by definition of the Macdonald polynomials, is 1. Hence we obtain the condition $c + D(u_B) - D(v) = 0$ which implies by our normalization $D(v) = 0$ that $c = -D(u_B)$. \square

The restriction to types $A_n^{(1)}$ and $D_n^{(1)}$ in Corollary 9.5 stems from the use of Ion's result [21]. An extension of Corollary 9.5 to type $C_n^{(1)}$ follows from [40, 41]. It would be interesting to determine if this result continues to hold in other types.

Remark 9.6. In order to match our notation for nonsymmetric Macdonald polynomials, we index symmetric Macdonald polynomials by anti-dominant weights. Many people (see for example [24, 44]) index Macdonald polynomials by dominant weights. The correct conversion between these conventions is that, for a dominant integral weight λ , P_λ in the above references is denoted by $P_{w_0(\lambda)}$ here.

Corollary 9.5 implies that the coefficients in the expansion of the symmetric Macdonald polynomial $P_\lambda(q, 0)$ at $t = 0$ in terms of the irreducible characters $\text{ch}(V(\mu))$ coincide with $X(\mu; B^{r_N, c_{r_N}} \otimes \dots \otimes B^{r_1, c_{r_1}})$ of (9.2). In formulas

$$P_\lambda(q, 0) = \sum_{\mu} X(\mu; B) \text{ch}(V(\mu)),$$

where $B = B^{r_N, c_{r_N}} \otimes \dots \otimes B^{r_1, c_{r_1}}$ with the r_i determined from λ as in Theorem 6.1.

Remark 9.7. In the case $\mathfrak{g} = A_n^{(1)}$, one can define nonsymmetric Macdonald polynomials $E_\lambda(q, t)$ for any \mathfrak{gl}_n weight λ (although if λ and μ correspond to the same \mathfrak{sl}_n weight, these only differ by a scalar). Letting \overline{P}' denote the lattice of \mathfrak{gl}_n weights, there is a natural injection

$$\begin{aligned} \mathbb{Q}(q, t)\overline{P}' &\rightarrow \mathbb{Q}(q, t)[x_1^{\pm 1}, \dots, x_n^{\pm 1}] \\ e^\lambda &\mapsto x_1^{\lambda_1} x_2^{\lambda_2} \dots x_n^{\lambda_n}, \end{aligned}$$

so we can identify $E_\lambda(q, t)$ with an actual polynomial, and we do so in the examples below by writing $E_\lambda(x; q, t)$.

Example 9.8. The Macdonald polynomial of type $A_2^{(1)}$ indexed by the anti-dominant weight $(0, 0, 2)$ is given by

$$P_{(0,0,2)}(x; q, 0) = x_1^2 + (q+1)x_1x_2 + x_2^2 + (q+1)x_1x_3 + (q+1)x_2x_3 + x_3^2.$$

By Corollary 9.5, this is given by the character of $B^{2^*, 1} \otimes B^{2^*, 1}$, where the power of q counts E^{int} . Here $2^* = 1$, so we consider the KR crystal $B^{1,1} \otimes B^{1,1}$. Under the map from Theorem 6.1, this is isomorphic as a classical crystal to $B_{s_2 s_1 s_0 s_2}(\Lambda_2)$, and there are enough 0 arrows in this Demazure crystal to calculate E^{int} . The KR crystal is then given as follows, where to make the picture cleaner we have only included arrows which survive in the Demazure crystal. Note that the ground state path is $2 \otimes 1$.

$$(9.5) \quad \begin{array}{cccccccc} 2 \otimes 1 & \xrightarrow{2} & 3 \otimes 1 & \xrightarrow{0} & 1 \otimes 1 & \xrightarrow{1} & 1 \otimes 2 & \xrightarrow{1} & 2 \otimes 2 & \xrightarrow{2} & 2 \otimes 3 & \xrightarrow{2} & 3 \otimes 3 \\ & & & \searrow^1 & & & & & \searrow^2 & & 1 \otimes 3 & \nearrow^1 & \\ & & & & 3 \otimes 2 & & & & & & & & \end{array}$$

The black arrows are all arrows that appear in $f_2^{n_1} f_1^{n_2} f_0^{n_3} f_2^{n_4} (2 \otimes 1)$ for some exponents $n_i \geq 0$, which give all vertices of the Demazure crystal, but not all arrows. The red arrows are the additional arrows in the Demazure crystal. We calculate that $D(2 \otimes 1) = D(3 \otimes 1) = D(3 \otimes 2) = -1$, and the rest of the elements of $B^{1,1} \otimes B^{1,1}$ have $D = 0$. This confirms Corollary 9.5 in this case.

The connection with KR crystals can also be used to compute all nonsymmetric Macdonald polynomials specialized at $t = 0$. Fix λ which need not be anti-dominant, and define w, τ as in (9.4). There is a unique minimal $w' \in W_0$ such that

$w'(\lambda)$ is an anti-dominant weight. Then $w'w((w')^{-1})^\tau\tau$ is a translation by an anti-dominant weight, where the superscript τ means conjugation by τ . Also, $B_w(\Lambda_{\tau(0)})$ embeds in $B_{w'w((w')^{-1})^\tau}(\Lambda_{\tau(0)}) = B_{w'w}(\Lambda_{\tau(0)})$, where the final equality follows because $w' \in W_0$. Let $j : B_{w'w}(\Lambda_{\tau(0)}) \rightarrow B$ be the isomorphism from Theorem 6.1, where B is the appropriate composite KR crystal, and let $B' = j(B_{w'w}(\Lambda_{\tau(0)}))$. Then in types $A_n^{(1)}$ and $D_n^{(1)}$ we have

$$(9.6) \quad E_\lambda(q, 0) = \sum_{b \in B'} q^{-D(b)} e^{\text{wt}(b)},$$

where D is calculated in the ambient composite KR crystal B . The subset B' of B can be described as $\{f_{i_k}^{n_k} \cdots f_{i_1}^{n_1} u_B \mid n_1, \dots, n_k \geq 0\}$, where $s_{i_k} \cdots s_{i_1}$ is any reduced expression for w .

Example 9.9. Again consider type $A_2^{(1)}$, and take the nonsymmetric Macdonald polynomial

$$E_{(0,2,0)}(x; q, 0) = x_1^2 + (q+1)x_1x_2 + x_2^2 + qx_1x_3 + qx_2x_3.$$

Then w, τ from (9.4) are $w = s_1s_0s_2s_1$ and $\tau = 0 \rightarrow 2 \rightarrow 1 \rightarrow 0$. The shortest $w' \in W_0$ so that $w'(0, 2, 0)$ is anti-dominant is $w' = s_2$. Thus the above argument along with Example 9.8 shows that $B_w(\Lambda_{\tau(0)}) = B_{s_1s_0s_2s_1}(\Lambda_2) = B_{s_1s_0s_2}(\Lambda_2)$ embeds into $B = B^{1,1} \otimes B^{1,1}$ (as a classical crystal), and a simple verification shows that the image B' is everything but $1 \otimes 3, 2 \otimes 3$ and $3 \otimes 3$. This indeed verifies that $E_{(0,2,0)}(x; q, 0)$ is given by (9.6).

Example 9.10. Consider the Macdonald polynomial of type $A_2^{(1)}$

$$P_{(0,1,2)}(x; q, 0) = x_1^2x_2 + x_1x_2^2 + x_1^2x_3 + (q+2)x_1x_2x_3 + x_2^2x_3 + x_1x_3^2 + x_2x_3^2.$$

Our results say that this is given by the character of the affine crystal B , where $B = B^{2,1} \otimes B^{1,1}$. The ground state path is $\begin{smallmatrix} 3 \\ 2 \end{smallmatrix} \otimes 1$. As a classical crystal, B consists

of two components, with highest weight elements $\begin{smallmatrix} 3 \\ 2 \end{smallmatrix} \otimes 1$ and $\begin{smallmatrix} 2 \\ 1 \end{smallmatrix} \otimes 1$ of weight 0 and $\omega_1 + \omega_2$ respectively. One can check that $D(\begin{smallmatrix} 3 \\ 2 \end{smallmatrix} \otimes 1) = -1$, and $D(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix} \otimes 1) = 0$.

This indeed confirms

$$P_{(0,1,2)}(x, q, 0) = q + \text{ch}(V(\omega_1 + \omega_2)).$$

9.3. q -deformed Whittaker functions. Gerasimov, Lebedev, Oblazin [13, Theorem 3.2] showed that q -deformed \mathfrak{gl}_n -Whittaker functions are Macdonald polynomials specialized at $t = 0$. As above this also gives a link to Demazure characters, and hence by the results in Section 7 to KR crystals graded by their energy functions. It would be interesting to generalize this to other types (in particular type $D_n^{(1)}$, where both Ion's results [21] and our results from Section 7 hold). The q -deformed \mathfrak{gl}_n -Whittaker functions are simultaneous eigenfunctions of a q -deformed Toda chain, which might serve as a starting point for this generalization. See also the recent paper by Braverman and Finkelberg [1]. Brubaker, Bump and Licata [2] constructed a natural basis of the Iwahori fixed vectors in the Whittaker model using Demazure-Lusztig operators. This gives another avenue to the link between Demazure characters and Whittaker functions.

REFERENCES

- [1] A. Braverman, M. Finkelberg, *Weyl modules and q -Whittaker functions*, preprint [arXiv:1203.1583](#).
- [2] B. Brubaker, D. Bump, A. Licata, *Whittaker functions and Demazure operators*, [arXiv:1111.4230](#).
- [3] V. Chari, *On the fermionic formula and the Kirillov-Reshetikhin conjecture*, Internat. Math. Res. Notices **12** (2001), 629–654.
- [4] I. Cherednik, *Nonsymmetric Macdonald polynomials*, Internat. Math. Res. Notices **10** (1995), 483–515.
- [5] V. Chari, A. Pressley, *Quantum affine algebras and their representations*, CMS Conf. Proc., 16, Amer. Math. Soc., Providence, RI, 1995, 59–78.
- [6] V. Chari, A. Pressley, *Twisted quantum affine algebras*, Comm. Math. Phys. **196** (1998), no. 2, 461–476. [arXiv:q-alg/9611002](#)
- [7] M. Demazure, *Désingularisation des variétés de Schubert généralisées*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 53–88.
- [8] G. Fourier, P. Littelmann, *Tensor product structure of affine Demazure modules and limit constructions*, Nagoya Math. J. **182** (2006), 171–198. [arXiv:math/0412432](#)
- [9] G. Fourier, P. Littelmann, *Weyl modules, Demazure modules, KR-modules, crystals, fusion products and limit constructions*, Adv. Math. **211** (2007), no. 2, 566–593. [arXiv:math/0509276](#)
- [10] G. Fourier, M. Okado, A. Schilling *Kirillov–Reshetikhin crystals of nonexceptional type*, Advances in Math. **222** (2009) 1080–1116 [arXiv:0810.5067](#)
- [11] G. Fourier, M. Okado, A. Schilling, *Perfectness of Kirillov–Reshetikhin crystals for nonexceptional type*, Contemp. Math. **506** (2010) 127–143. [arXiv:0811.1604](#)
- [12] G. Fourier, A. Schilling, M. Shimozono, *Demazure structure inside Kirillov–Reshetikhin crystals*, J. Algebra **309** (2007) 386–404. [arXiv:math/0605451](#)
- [13] A. Gerasimov, D. Lebedev, S. Oblezin, *On q -deformed $gl(l+1)$ -Whittaker function III*, Lett. Math. Phys. **97** (2011), no. 1, –24. [arXiv:0805.3754](#)
- [14] M. Haiman, *Cherednik algebras, Macdonald polynomials and combinatorics*, International Congress of Mathematicians. Vol. III, 843–872, Eur. Math. Soc., Zürich, 2006.
- [15] J. Haglund, M. Haiman, N. Loehr, *A combinatorial formula for nonsymmetric Macdonald polynomials*, Amer. J. Math. **130** (2008), no. 2, 359–383. [arXiv:math/0601693](#)
- [16] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, Z. Tsuboi, *Paths, crystals and fermionic formulae*, MathPhys Odyssey 2001, 205–272, Prog. Math. Phys. **23**, Birkhäuser Boston, Boston, MA, 2002. [arXiv:math/0102113](#)
- [17] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, Y. Yamada, *Remarks on fermionic formula* Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998), 243–291, Contemp. Math., **248**, Amer. Math. Soc., Providence, RI, 1999. [arXiv:math/9812022v3](#)
- [18] J. Hong, S.-J. Kang, *Introduction to quantum groups and crystal bases*, Graduate Studies in Mathematics, **42**, American Mathematical Society, Providence, RI, 2002. xviii+307 pp.
- [19] D. Hernandez, *Kirillov-Reshetikhin conjecture: the general case*, Int. Math. Res. Not. IMRN 2010, no. 1, 149–193.
- [20] K. Hikami, *Representation of the Yangian invariant motif and the Macdonald polynomial*, J. Phys. A **30** (1997), no. 7, 2447–2456.
- [21] B. Ion, *Nonsymmetric Macdonald polynomials and Demazure characters*, Duke Math. J. **116**(2) (2003), 299–318. [arXiv:math/0105061](#)
- [22] B. Jones, A. Schilling *Affine structures and a tableau model for E_6 crystals*, J. Algebra **324** (2010) 2512–2542. [arXiv:0909.2442](#)
- [23] V. G. Kac, *Infinite-dimensional Lie algebras*, Third edition, Cambridge University Press, Cambridge, 1990.
- [24] F. Knop, *Integrality of two variable Kostka functions*, J. Reine Angew. Math. **482** (1997), 177189. [arXiv:q-alg/9603027](#)
- [25] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, A. Nakayashiki, *Affine crystals and vertex models*, Int. J. Mod. Phys. A **7** (suppl. 1A) (1992), 449–484.
- [26] S.-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, A. Nakayashiki, *Perfect crystals of quantum affine Lie algebras*, Duke Math. J. **68** (1992) 499–607.

- [27] M. Kashiwara, *The crystal base and Littelmann's refined Demazure character formula*, Duke Math. J. **71** (1993), no. 3, 839–858.
- [28] M. Kashiwara, *On crystal bases of the q -analogue of universal enveloping algebras*, Duke Math. J. **73** (1994), 383–413.
- [29] M. Kashiwara, *Similarity of crystal bases*, Lie algebras and their representations (Seoul, 1995), 177–186, Contemp. Math., 194, Amer. Math. Soc., Providence, RI, 1996.
- [30] M. Kashiwara, *On level-zero representations of quantized affine algebras*, Duke Math. J. **112** (2002), no. 1, 117–175. [arXiv:math/0010293](#)
- [31] M. Kashiwara *Level zero fundamental representations over quantized affine algebras and Demazure modules*, Publ. Res. Inst. Math. Sci. **41** (2005), no. 1, 223–250. [arXiv:math/0309142](#)
- [32] M. Kashiwara, K.C. Misra, M. Okado, D. Yamada *Perfect crystals for $U_q(D_4^{(3)})$* , J. Algebra **317** (2007), no. 1, 392–423. [arXiv:math/0610873](#)
- [33] M. Kashiwara, T. Nakashima, *Crystal graphs for representations of the q -analogue of classical Lie algebras*, J. Algebra **165** (1994), no. 2, 295–345.
- [34] A.N. Kirillov, N.Yu. Reshetikhin *Representations of Yangians and multiplicities of the inclusions of the irreducible components of the tensor product of representations of simple Lie algebras*. Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **160** (1987), Anal. Teor. Chisel i Teor. Funktsii. **8**, 211–221, 301; translation in J. Soviet Math. **52** (1990), no. 3, 3156–3164
- [35] S. Kumar, *Demazure character formula in arbitrary Kac–Moody setting*, Invent. Math. **89** (1987), no. 2, 395–423.
- [36] A. Kuniba, K. C. Misra, M. Okado, J. Uchiyama, *Demazure modules and perfect crystals*, Comm. Math. Phys. **192** (1998), no. 3, 555–567. [arXiv:q-alg/9607011](#)
- [37] A. Kuniba, K. C. Misra, M. Okado, T. Takagi, J. Uchiyama, *Characters of Demazure modules and solvable lattice models*, Nuclear Phys. B **510** (1998), no. 3, 555–576. [arXiv:q-alg/9707004](#)
- [38] A. Kuniba, K. C. Misra, M. Okado, T. Takagi, J. Uchiyama, *Crystals for Demazure modules of classical affine Lie algebras*, J. Algebra **208** (1998), no. 1, 185–215. [arXiv:q-alg/9707014](#)
- [39] C. Lecouvey, M. Okado, M. Shimozono, *Affine crystals, one dimensional sums and parabolic Lusztig q -analogues*, Math. Zeitschrift, to appear, DOI: 10.1007/s00209-011-0892-9. [arXiv:1002.3715v1](#).
- [40] C. Lenart, *From Macdonald polynomials to a charge statistic beyond type A*, J. Combin. Theory Ser. A, **119** (2012), 683–712. [arXiv:1106.3296](#).
- [41] C. Lenart, A. Schilling, *Crystal energy functions via the charge in types A and C*, Math. Zeitschrift, to appear. [arXiv:1107.4169](#).
- [42] P. Littelmann, *Paths and root operators in representation theory*, Ann. of Math. (2) **142** (1995), no. 3, 499–525.
- [43] P. Littelmann, *Crystal graphs and Young tableaux*, J. Algebra **175** (1995), no. 1, 65–87.
- [44] D. Marshall, *Symmetric and nonsymmetric Macdonald polynomials. On combinatorics and statistical mechanics*, Ann. Comb. **3** (1999), no. 2-4, 385–415. [arXiv:math/9812080](#)
- [45] O. Mathieu, *Formules de Demazure–Weyl, et généralisation du théorème de Borel–Weil–Bott*, C. R. Acad. Sci. Paris Sér. I Math. **303** (1986), no. 9, 391–394.
- [46] K. Naoi, *Weyl modules, Demazure modules and finite crystals for non-simply laced type*, preprint [arXiv:1012.5480](#)
- [47] K. Naoi, *Demazure crystals and tensor products of perfect Kirillov–Reshetikhin crystals with various levels*, preprint [arXiv:1108.3139](#)
- [48] E. Opdam, *Harmonic analysis for certain representations of graded Hecke algebras*, Acta Math. **175** (1995), no. 1, 75–121.
- [49] M. Okado *Existence of crystal bases for Kirillov–Reshetikhin modules of type D*, Publ. Res. Inst. Math. Sci. **43** (2007), no. 4, 977–1004. [arXiv:math.QA/0610874](#)
- [50] M. Okado, A. Schilling, *Existence of Kirillov–Reshetikhin crystals for nonexceptional types* Represent. Theory **12** (2008) 186–207. Erratum: Represent. Theory **12** (2008), 499–500. [arXiv:0706.2224](#)
- [51] M. Okado, A. Schilling, M. Shimozono, *Virtual crystals and fermionic formulas of type $D_{n+1}^{(2)}$, $A_{2n}^{(2)}$, and $C_n^{(1)}$* , Representation Theory **7** (2003) 101–163. [arXiv:math/0105017](#)
- [52] M. Okado, A. Schilling, M. Shimozono, *Virtual crystals and Kleber's algorithm*, Commun. Math. Phys. **238** (2003) 187–209. [arXiv:math/0209082](#)

- [53] Y. Sanderson, *On the connection between Macdonald polynomials and Demazure characters*, J. Algebraic Combin. **11** (2000) 269–275.
- [54] M. Shimozono, *Affine type A crystal structure on tensor products of rectangles, Demazure characters, and nilpotent varieties*, J. Algebraic Combin. **15** (2002), no. 2, 151–187.
- [55] A. Schilling, *Combinatorial structure of Kirillov–Reshetikhin crystals of type $D_n^{(1)}$, $B_n^{(1)}$, $A_{2n-1}^{(2)}$* , J. Algebra **319** (2008) 2938–2962. [arXiv:0704.2046](https://arxiv.org/abs/0704.2046)
- [56] Y. Yamane *Perfect crystals of $U_q(G_2^{(1)})$* . J. Algebra 210 (1998), no. 2, 440–486. [arXiv:q-alg/9712012](https://arxiv.org/abs/q-alg/9712012)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, ONE SHIELDS AVENUE, DAVIS, CA 95616-8633, U.S.A.

E-mail address: anne@math.ucdavis.edu

URL: <http://www.math.ucdavis.edu/~anne>

DEPARTMENT OF MATHEMATICS, M.I.T., CAMBRIDGE, MA 02139-4307, U.S.A.

E-mail address: peter.tingley@gmail.com

URL: <http://www-math.mit.edu/~ptingley/>